Psychiatric Symptoms Related to the COVID-19 Pandemic

Christopher Rohde M.D.[1,2]*, Oskar Hougaard Jefsen B.Sc.[1,2]*, Bettina Nørremark M.Sc.[1],

Andreas Aalkjær Danielsen M.D. Ph.D.[1,2], Søren Dinesen Østergaard, M.D. Ph.D.[1,2]

[1] Department of Affective Disorders, Aarhus University Hospital – Psychiatry, Aarhus Denmark

[2] Department of Clinical Medicine, Aarhus University, Aarhus, Denmark

*Equal contribution

Corresponding author:

Professor Søren D. Østergaard, MD PhD
Department of Affective Disorders
Aarhus University Hospital - Psychiatry
Palle Juul-Jensens Boulevard 175
8200 Aarhus N
Phone: + 45 61282753
Email: soeoes@rm.dk

Background

The COVID-19 pandemic and the consequential societal changes are likely to have major health consequences way beyond those caused by the virus infection.[1] One of the medical fields that may experience significant consequences of the pandemic is that of psychiatry.[2-4] Importantly, individuals who already live with mental disorders may be particularly vulnerable to the psychological stress associated with the pandemic.[4] However, data on how the COVID-19 pandemic affects this group of people is largely absent from the literature.

Here, we report data from an investigation conducted at the psychiatric services of the Central Denmark Region (CDR), which has a catchment area of approximately 1.3 million people covered by five psychiatric hospitals providing inpatient and outpatient treatment of all types of mental disorders. Based on clinical observation and evidence of the psychological effects of both the current[5] and past[2] pandemics, we hypothesized that we would find examples of "pandemic-related" symptoms (deterioration) of anxiety, obsessive-compulsive disorder, depression and psychotic disorders (e.g. schizophrenia).

Methods

We extracted all clinical notes from the period from February 1[st] to March 23[rd] 2020 (first verified case of COVID-19 in Denmark was reported on February 26[th] 2020)[6] from the electronic medical record system used by the psychiatric services of the CDR for individuals (patients) aged 18 or above. Subsequently, we conducted an electronic search for clinical notes containing at least one of the following words: "corona", "COVID", "virus", "epidemic", "pandemic", and "contaminate/contamination" (including compound words). Then, all clinical notes containing these words were screened manually to determine whether they described symptoms that seemed plausibly related to the COVID-19 pandemic. Finally, the clinical notes describing pandemic-related psychiatric symptoms were labeled according to the most dominant psychopathology. For a more detailed description of the methods, see the Supplementary Material.

Results

In the period from February 1[st] to March 23[rd] there were 412,804 clinical notes reporting on a total of 14,561 adult patients in the psychiatric services of the CDR. In this dataset, the electronic search resulted in 11,072 notes from 5796 patients that contained at least one of the six COVID-19

pandemic-related words mentioned above. The subsequent manual screening showed that while the majority of these notes described logistics (e.g. "The appointment was rescheduled to a phone consultation due to the risk of COVID-19"), 1357 of the notes from 918 patients (621 females with a mean age of 36.3 years (SD=14.3) and 297 males with a mean age of 40.9 years (SD=13.8)) described pandemic-related symptoms. The main diagnoses of these patients and the type of pandemic-related psychopathology described in the notes are reported in Figure 1, which also shows the cumulative incidence of i) patients with pandemic-related psychopathology, ii) clinical notes describing pandemic-related psychopathology, and iii) confirmed cases of COVID-19 in Denmark[5] in the period from February 1st to March 23rd. The figure shows a clear association between the number of COVID-19 cases in Denmark (and the societal restrictions) and the degree of pandemic-related psychopathology.

Discussion

To our knowledge, this is the first investigation of COVID-19 pandemic-related psychopathology from a large psychiatric treatment setting. The results clearly suggest that this phenomenon exists among individuals with mental illness. However, as the data from this investigation stems from standard clinical practice where the patients were not systematically assessed for pandemic-related psychopathology, we cannot speak to its overall prevalence. Nevertheless, we believe that our findings underline the necessity of taking urgent action to mitigate the negative effects of the COVID-19 pandemic on patients with mental disorders. A low-cost intervention would be to simply discuss the ongoing pandemic and its societal consequences with patients seeking care in psychiatric settings - as this may lead to relief of pandemic-related symptoms via general comforting, and clarification of misunderstandings/false beliefs regarding the pandemic.

Funding

This project is supported by an unconditional grant from the Novo Nordisk Foundation (Grant number: NNF20SA0062874).

Conflicts of interest

None.

References

1. World Health Organization webpage on the novel coronavirus (https://www.who.int/emergencies/diseases/novel-coronavirus-2019). Accessed April 14th 2020.
2. Brooks SK, Webster RK, Smith LE, et al. The psychological impact of quarantine and how to reduce it: rapid review of the evidence. *Lancet.* 2020;395(10227):912-920.
3. Druss BG. Addressing the COVID-19 Pandemic in Populations With Serious Mental Illness. *JAMA Psychiatry.* 2020.
4. Reger MA, Stanley IH, Joiner TE. Suicide Mortality and Coronavirus Disease 2019-A Perfect Storm? *JAMA Psychiatry.* 2020.
5. Wang C, Pan R, Wan X, et al. Immediate Psychological Responses and Associated Factors during the Initial Stage of the 2019 Coronavirus Disease (COVID-19) Epidemic among the General Population in China. *International journal of environmental research and public health.* 2020;17(5).
6. Danish State Serum Institute webpage containing COVID-19 data from Denmark (https://www.ssi.dk/aktuelt/sygdomsudbrud/coronavirus). Accessed April 14th 2020.

Figure 1. The cumulative incidence of COVID-19 pandemic-related psychopathology among patients with mental disorders in the Central Denmark Region.

ICD-10	Diagnosis	Patients
DF20-29	Schizophrenia and related disorders	198
DF30-31	Mania and bipolar disorder	68
DF32-33	Unipolar depression	130
DF40-41	Anxiety disorders	37
DF42	Obsessive-compulsive disorder	36
DF43	Reactive and adjustment disorders	117
DF43.10	Post-traumatic stress disorder	85
DF50	Eating disorders	31
DF60-69	Personality and behavioral disorders	97
DF70-79	Mental retardation	8
DF84	Autism spectrum disorders	13
DF90	Hyperkinetic disorders	34
DZ03.2	Suspected mental disorder	30
	Other diagnosis	32
	No diagnosis	87

Psychopathology	Clinical notes
Substance abuse	16
Delusions	149
Hallucinations	24
Mania / hypomania / mixed states	30
Depression	146
Suicidality	102
Anxiety	539
PTSD-related symptoms	26
Obsessive-compulsive symptoms	85
Feeding behavior symptoms	30
Autism-related symptoms	6
ADHD-related symptoms	10
Aggression	10
Unspecific stress	174
Other symptoms	10

The nationwide lockdown announced on March 11th 2020 involved closure of kindergartens, schools, teaching institutions, restaurants, bars and many other small businesses. On March 23rd 2020, the lockdown was extended to April 14th 2020.

Supplementary material