Updated July 17, 2020

Title

Spatial variability of nitrogen dioxide and formaldehyde and residential exposure of children in the industrial area of Viadana, Northern Italy

Authors

Alessandro Marcon[1], Silvia Panunzi[1], Massimo Stafoggia[2], Chiara Badaloni[2], Kees de Hoogh[3,4], Linda Guarda[5], Francesca Locatelli[1], Caterina Silocchi[6], Paolo Ricci[5], Pierpaolo Marchetti[1]

Affiliations

1. Unit of Epidemiology and Medical Statistics, Department of Diagnostics and Public Health, University of Verona, Verona, Italy
2. Department of Epidemiology, Lazio Region Health Service ASL Roma 1, Rome, Italy
3. Swiss Tropical and Public Health Institute, Basel, Switzerland
4. University of Basel, Basel, Switzerland
5. UOC Osservatorio Epidemiologico, Agenzia di Tutela della Salute della Val Padana, Mantova, Italy
6. UOS Salute e Ambiente, Agenzia di Tutela della Salute della Val Padana, Mantova, Italy

Corresponding author

Alessandro Marcon, PhD, Unit of Epidemiology and Medical Statistics, Department of Diagnostics and Public Health, University of Verona, c/o Istituti Biologici II, strada Le Grazie 8, 37134 Verona, Italy; phone: +39-045-8027668; e-mail: alessandro.marcon@univr.it; ORCID: 0000-0002-2778-658X

Contributorship statement

AM and PR are co-principal investigators of the Viadana III study. AM conceived the present analysis and drafted the first version of the manuscript. SP performed the geographical and statistical analysis. PM supervised the geographical and statistical analyses. CS and PR obtained data from the environmental protection agency (ARPA Lombardia). LG reconstructed the residential history of participants. CB and MS provided exposure estimates from EPISAT. KdH provided exposure estimates from ELAPSE. All the authors contributed in the discussion of the analysis plan, the interpretation of results, and commented on the first draft, they critically reviewed and approved the final version of the manuscript.

Funding

The Viadana III study was funded by the Agenzia Tutela della Salute della Val Padana, Mantova (Decree n. 278, 17/07/2018). A Research Scholarship was co-funded by the Department of Diagnostics and Public Health, University of Verona. Geocoding of participants' addresses was supported by EGON Solutions, via Enrico Fermi 13/C, 37135 Verona.

Disclosure of potential conflicts of interest

The authors declare no conflicts of interest related to this study.

Keywords

Aldehydes; BTEX; environmental monitoring; exposure assessment; geographic information system; nitrogen oxides; passive sampling; particulate matter; wood industry.

ABSTRACT

Chipboard production is a source of ambient air pollution. We assessed the spatial variability of outdoor pollutants and residential exposure of children living in proximity to the largest industry for chipboard production in Italy, aiming at evaluating the reliability of exposure estimates obtained from passive sampling for an ongoing epidemiological study.

We obtained data on NO_2 and formaldehyde, collected by the environmental protection agency of Lombardia region, at 25 passive sampling sites in the municipality of Viadana during 10 weeks (2017-18), and compared NO_2 measurements with average weekly concentrations from continuous monitors. We compared interpolated surfaces of NO_2 and formaldehyde for 2017-18 with previous maps for 2010. We assessed the relationship between residential proximity to the industry and pollutant exposures, assigned both using our maps and available countrywide/continental models based on routine monitoring data on NO_2, PM_{10}, and $PM_{2.5}$.

The correlation between weekly NO_2 concentrations from a continuous monitor and a co-located passive sampler was high (Pearson's r = 0.89), although passive sampling underestimated NO_2 during the cold season. For both 2010 and 2017-18, we observed higher NO_2 and formaldehyde concentrations in the southern part of Viadana, with hot-spots in proximity to the industry. PM_{10} and $PM_{2.5}$ exposures were higher for children at <1km to the industry and NO_2 exposure was higher at 1-1.7 km, compared to the children living at >3.5 km. The density of roads and population was also higher in proximity to the industry.

Findings from a variety of exposure models suggest that children living in proximity to the chipboard industry in Viadana are more exposed to air pollution and that exposure gradients are relatively stable over time.

Abbreviations

BC, black carbon

BTEX, benzene, toluene, ethylbenzene, o-xylene, mp-xylene

ELAPSE, Effects of Low-Level Air Pollution: A study in Europe

EPISAT, Dati satellitari ed uso del territorio per la stima delle esposizioni a livello nazionale

LoQ, limit of quantification

LUR, land use regression

NO_2, nitrogen dioxide

PM, particulate matter

PM_{10}, particulate matter with an aerodynamic diameter of 10 µm or less

$PM_{2.5}$, particulate matter with an aerodynamic diameter of 2.5 µm or less

1. INTRODUCTION

Industrial wood manufacturing and chipboard production are a source of ambient air pollutants, including dust from mechanical woodworking, formaldehyde from the resins used to bond wood particles, and combustion by-products (Dahlgren et al. 2003; Marcon et al. 2014).

The largest industrial park for the production of chipboard in Italy is located in the health district of Viadana (labelled "district" for short). The district is located in the Po Plain, in northern Italy, one of the most polluted regions in Europe (Larsen et al. 2012; Stafoggia et al. 2017). It extends over an area of 363 km^2 and comprises Viadana and 9 other smaller municipalities, counting 47,701 inhabitants overall in 2018 (http://demo.istat.it). The main emission sources of air pollution are two big industries in the south of the district, which include chemical plants for the synthesis of urea-formaldehyde resins, chipboard production and storage facilities, and small incinerators (Marcon et al. 2014). Both industries are under Directive 2010/75/UE for Integrated Pollution Prevention and Control.

We previously showed that living close to the industries is associated with several adverse outcomes in the paediatric population, including respiratory and irritation symptoms (de Marco et al. 2010; Girardi et al. 2012; Rava et al. 2012) and hospital admissions for respiratory diseases (Marchetti et al. 2014; Rava et al. 2011). Using data from *ad hoc* monitoring conducted by passive sampling in 2010, we found that residential proximity to the factories is linked to a higher outdoor exposure to nitrogen dioxide (NO_2) and formaldehyde, and that exposure is associated with increased genotoxicity in mouth mucosa cells among children (Marcon et al. 2014).

NO_2 has been related to a range of adverse effects, including mortality, exacerbations of obstructive airway diseases, and childhood asthma (Curtis et al. 2006). Formaldehyde is

known to cause irritation and neuro-vegetative symptoms, and has mutagenic and carcinogenic properties (IARC 2006, update 2018). In urban environments, NO_2 and formaldehyde mostly derive from combustions linked with road traffic, industrial activities, and domestic heating; formaldehyde is also generated through photo-oxidation of anthropogenic hydrocarbons (Kheirbek et al. 2012). In the district, wood waste incineration and power generation at the industrial premises are additional sources of nitrogen oxides, whereas formaldehyde is emitted during chipboard production and storage (Marcon et al. 2014).

The working principle of passive sampling is uptake of gaseous pollutants from the air by an adsorbing material at a constant uptake rate controlled by diffusion, followed by desorption and titration (Buczynska et al. 2009; Yu et al. 2008). As opposed to continuous monitors, passive samplers are cheap and easy to use since they do not require forced air movement through the device and thus a pump operated using electricity (Santana et al. 2017). However, diffusion rates can vary largely according to environmental conditions thus affecting their performance (Mason et al. 2011).

In the "Viadana III" study (http://biometria.univr.it/viadanastudy/) we are conducting new cohort studies of the population aged 0–20 years at baseline (2013), aimed to assess the relationship between exposure to air pollutants and further health outcomes during follow-up (2013–2017). The main aim of the present analysis is to understand whether the previously mentioned NO_2 and formaldehyde exposure estimates for 2010 can be reliably used in the new follow-up study. To fulfil this aim:

1) we appraised the validity of passive sampling of NO_2 against continuous monitoring (reference data were not available for formaldehyde);

2) we assessed whether living next to the industrial plants is still related to a higher air pollution exposure, by comparing NO_2 and formaldehyde surfaces obtained using

recent (2017-18) monitoring data for the municipality of Viadana, where the largest

chipboard industry is located, with our previous maps for 2010 (Marcon et al. 2014);

3) we evaluated the consistency between exposure estimates assigned through our maps

and other available models that employed different input data, modelling techniques,

and covered wider geographical areas. With the objective of focusing on residential

areas alone, exposures were attributed to home addresses of Viadana III participants.

2. METHODS

2.1. Passive sampling monitoring campaigns

We obtained data from monitoring campaigns conducted using Radiello® passive samplers (Fondazione Salvatore Maugeri, Padova, Italy) both in 2010 and in 2017–2018.

The monitoring campaigns in 2010 were planned for epidemiological purposes and conducted in collaboration between the environmental protection agency of Lombardia region (Agenzia Regionale per la Protezione dell'Ambiente [ARPA] Lombardia, Dipartimento di Cremona e Mantova, https://www.arpalombardia.it/) and our team (Marcon et al. 2014). Briefly, measurements of formaldehyde and NO_2 were conducted during two weeks in the cold season and two weeks in the warm season at 62 sites in the district. Site selection overrepresented densely populated areas close to the two industries. Thirty-three of these sites were located in the eponymous municipality (labelled "Viadana" for short).

The monitoring campaigns in 2017-18 where conducted by ARPA Lombardia. Twenty-five sites were identified over a 10 km^2 area around the centre of Viadana; none of these sites were the same as in 2010. NO_2 and formaldehyde, but also BTEX (benzene, toluene, ethylbenzene, o-xylene, mp-xylene), acetaldehyde, and benzaldehyde were monitored during five weeks in the cold season (from 21/11/2017 to 27/12/2017) and five weeks in the warm season (from 8/5/2018 to 12/6/2018). When measurements were below the limit of quantification (LoQ), we assigned half the LoQ value, except when this occurred during all/most of the weeks (**Table S1**).

2.2. Data from continuous air quality stations

For 2017-18, we obtained hourly concentrations of NO_2 from two air quality stations in Viadana (see maps in **Figure S1**) during the same periods of passive sampling. The first was a routine urban background station located 500 m away from the industry (via

Gianmarco Cavalli, Viadana). The second was a temporary station located in a meadow
close to a graveyard (via Cesare Airoldi, Viadana). One passive sampler was co-located
with this station. The two stations were equipped with similar detectors in agreement with
current legislation (D. Lgs. 155/2010), and measured NO_2 by continuous chemo-
luminescence detectors. There were no traffic-type monitoring stations operating in the
district area.

2.3. Geocoding of residential addresses

We used data from the Viadana III study to derive a list of residential addresses in the
study area. Viadana III includes prospective studies of two paediatric cohorts followed up
between 2013 (baseline) and 2017. The first cohort is made of all the 3,854 children and
adolescents born in 1992–2003 who were attending the schools in the district in 2006,
when their parents were surveyed (de Marco et al. 2010). Of these, 3,711 (96%) subjects
were traced through the healthcare provider records; 3,688 were still living in the district
area in 2013. The second cohort consisted of all the 4,221 children born in the district in
2004–2012 who lived in the district in 2013. Overall, the age range of the two cohorts was
0–20 years at baseline.

We geocoded residential addresses at baseline following the procedure described in
Appendix S1. After excluding 206 and 178 addresses that failed to be geocoded from the
first and second cohorts respectively, the number of available geocodes was 3,482 (94%)
for the first cohort, and 4,043 (96%) for the second. We then combined coordinates from
the two cohorts and removed duplicates (i.e. children living at the same address),
obtaining a list of 4,390 unique locations in the district; 1,814 of these were in Viadana.

2.4. Population and street density indicators

With the aim of characterising non-industrial pollutant sources in the area, we obtained the
number of inhabitants, buildings, and households per census tract by national statistics

census data (ISTAT, 14° Censimento generale della popolazione e delle Abitazioni

(http://dawinci.istat.it/), and derived population density indicators within 500 m buffers

around the passive sampling locations by area-weighting. We also obtained the road

network through Open Street Map (https://www.openstreetmap.org/), and derived total

length of all types of roads in the 500 m buffers as an indicator of street density.

2.5. Exposure assignment

We obtained estimates of exposure to outdoor air pollutants at the 1,814 geocoded

addresses by applying models available from three projects for baseline (or closest year),

and new exposure models for 2017-18.

For baseline, we used the ordinary kriging models of NO_2 and formaldehyde

concentrations that we had previously devised using passive sampling data for 2010

(Marcon et al. 2014). For the same year, we also used the land-use regression (LUR)

models for NO_2, $PM_{2.5}$, and black carbon (BC) concentrations developed for Western

Europe within the ELAPSE study (Effects of Low-Level Air Pollution: A study in Europe)

(de Hoogh et al. 2018). Dependent variables for these models were routine air quality data

for NO_2 and $PM_{2.5}$ and *ad hoc* monitoring data for BC, whereas predictor variables

included satellite data, dispersion model estimates, land cover and traffic indicators.

Moreover, we obtained PM_{10} and $PM_{2.5}$ concentrations for 2012 and 2013, respectively,

using the spatiotemporal models developed for Italy within the EPISAT project (Dati

satellitari ed uso del territorio per la stima delle esposizioni a livello nazionale) (Badaloni et

al. 2018; Stafoggia et al. 2017). EPISAT models used routine air quality data as dependent

variables, and incorporated both spatial (e.g. population density and emission data), and

spatiotemporal predictors (e.g. satellite data and meteorology) as predictor variables. Co-

authors MS and CB derived optimised exposure estimates for the study area aimed to

capture PM variation due to very local sources, i.e. stage 4 modelling in Stafoggia et al. (2017).

For 2017-18, we devised new ordinary kriging models for NO_2 and formaldehyde using passive sampling data. The best-fitting models were chosen by minimizing the root mean squared error (RMSE) by leave-one-out cross-validation (LOOCV) (Pebesma and Wesseling 1998). The spatial variogram providing the best model fit was a Gaussian class model for both pollutants. The variogram parameters were: partial sill = 13.4, range = 3.5 km, nugget = 2.5, and direction in plane = 135° (east) for NO_2; and partial sill = 0.09, range = 2.0 km, nugget = 0.07 (no anisotropy) for formaldehyde. The estimated LOOCV-RMSEs were 1.95 and 0.28, respectively.

2.6. Statistical analysis

For the ten monitoring weeks in 2017-18, we calculated daily NO_2 concentrations by averaging hourly data from each of the two continuous monitoring stations, setting to missing the days with less than 25% of hourly measurements available, and then imputing missing data by k-nearest neighbour algorithm using measurements from the other station as we did in Marchetti et al. (2017). We calculated weekly average concentrations using the imputed time-series. We then compared weekly concentrations from the temporary station and the co-located passive sampler, and we obtained a "recalibration equation" by linear regression. The resulting equation was used to predict NO_2 concentrations at the 25 passive sampling sites.

We reported statistics on air pollution measured concentrations, as well as estimated exposures at the residential addresses, as a function of distance to the industry (categorised according to quantiles, see table legends). Overall comparisons of exposures across groups of children were tested by the Kruskall-Wallis rank test.

The analyses were carried out using Stata Statistical Software, Release 16.1 (College Station, TX: StataCorp LLC) and R version 3.5.1 (http://www.R-project.org/).

2.7. Ethical approval

The Viadana III study has been approved by the Comitato Etico Val Padana (Prot. n. 4813, 12/02/2019).

3. RESULTS AND DISCUSSION

3.1. Comparison between passive sampling and continuous monitoring

All the NO_2 measurements obtained using Radiello® tubes during the cold season, and most of those obtained in the warm season, were below the average concentrations derived by the continuous stations for the corresponding weeks (**Figure 1**). A comparison of weekly concentrations between the temporary station and the co-located sampler (solid and dashed thick lines, respectively) suggests that passive sampling underestimated NO_2 concentrations especially in winter. In a validation study of different diffusive sampler types, Gerboles et al. (2006) found that Radiello® tubes provided lower NO_2 concentrations compared to other samplers and underestimated reference concentrations both in laboratory conditions and in a rural setting. We observed a considerable variability across monitoring weeks, ranging from an overestimation on week 6 to a heavy underestimation on week 1, which supports the common approach of building exposure models from "annual" concentrations (i.e. averaged across a number of monitoring periods) rather than modelling "seasonal" concentrations (Beelen et al. 2013; de Hoogh et al. 2018). The correlation between the temporary station and the co-located sampler was high, as indicated by a Pearson's r correlation coefficient of 0.89 (**Figure S2**). All in all, these findings suggest that NO_2 exposure estimates based on passing sampling data can be reasonably used in the Viadana III epidemiological study, where the main aim is detecting exposure-response relationships. Nonetheless, recalibration may be useful to increase the accuracy of estimates of health effects related to a range of NO_2 exposures.

For formaldehyde, since we did not have reference data and validation studies of passive sampling are few, comparison of concentrations with other studies should be done cautiously. Using Radiello® tubes, Evans and Stuart (2011) observed quite similar winter

concentrations of formaldehyde between active monitors and co-located passive samplers (2.1 vs 2.2 µg/m^3) in Hillsborough County, Florida. However, Mason et al. (2011) documented an excellent performance in controlled experiments (2% difference from a reference method) but also substantial (39%) underestimation of formaldehyde in a real-world setting.

3.2. NO$_2$ concentrations

The mean concentrations of NO$_2$ during the cold and warm seasons in 2017-18 were 19.8 and 9.9 µg/m^3, respectively (**Table 1**). Spatial variability was higher for NO$_2$ during the warm season (coefficient of variation = 0.31), compared to measurements during the cold season as well as to the other pollutants (**Table 2**). The median annual measured concentration was 16.0 (Q1–Q3: 12.4–16.6) µg/m^3. After recalibration, the median concentration was 26.9 (Q1–Q3: 18.5–28.2) µg/m^3; at all the sites, recalibrated annual concentrations were within the EU air quality standard of 40 µg/m^3 (**Figure 2**). For comparison, average NO$_2$ concentrations measured by routine stations in 2017-18 were around 30 µg/m^3 for the urban area of Mantova, and 20 µg/m^3 for the larger province encompassing also suburban/rural areas (ARPA Lombardia Rapporto qualità aria 2018).

A clear relationship with distance was found when looking at observed NO$_2$ concentrations (**Table 2**). The median recalibrated concentration was 11 µg/m^3 higher at <1 km to the industry compared to the area >1.8 km (**Figure 2**).

3.3. Formaldehyde concentrations

As opposed to the campaigns in 2010 (Marcon et al. 2014), we found higher formaldehyde concentrations in the cold season than in the warm season (2.2 vs 1.1 µg/m^3, respectively, **Table 1**), which was unexpected because lower concentrations are usually found in winter (Kheirbek et al. 2012; Villanueva et al. 2014). The median annual concentration of formaldehyde was 1.6 µg/m^3 (Q1–Q3: 1.5–2.0 µg/m^3), which is similar to other studies that

used Radiello® tubes (Evans and Stuart 2011; Kheirbek et al. 2012). There are no ambient air quality standards for aldehydes, and studies on the health effects of outdoor formaldehyde are still few (Dahlgren et al. 2003; Marcon et al. 2014; Morello-Frosch et al. 2000). For long-term exposure, Effects Screening Levels (calculated so that lower ambient levels are unlikely to be of concern for health and the environment) are set to 3.3 µg/m^3 from the Texas Air Monitoring Information System (USA – Texas Commission on Environmental Quality, https://www.tceq.texas.gov/ accessed 11 March 2020) (Santana et al. 2017). However, formaldehyde concentrations in our study were above 0.8 µg/m^3, the benchmark set by the US Environmental Protection Agency for a 1 in 100,000 lifetime cancer risk for inhaled exposures (https://cfpub.epa.gov/ accessed 12 March 2020).

Observed concentrations did not vary appreciably according to proximity to the industry (**Table 2**), and spatial contrasts were small, as we observed during the previous campaigns (Marcon et al. 2014). Using Radiello® tubes, Kheirbek et al. (2012) reported a coefficient of variation of 22% for formaldehyde in New York City at springtime, whereas even lower coefficients of 12-13% were reported in other studies (Evans and Stuart 2011; Marcon et al. 2014).

3.4. Concentrations of other pollutants

In the cold season, mean concentrations of benzene, toluene, and acetaldehyde were 1.5, 2.8, and 0.7 µg/m^3 respectively (**Table S3**), whereas they were mostly below the LoQ during the warm season (**Table S1**). Benzene and toluene concentrations were comparable or lower to concentrations observed at background or semirural sites in other studies using Radiello® tubes (Buczynska et al. 2009; Gaeta et al. 2016; Kerchich and Kerbachi 2012). Ethylbenzene, o-xylene, mp-xylene, and benzaldehyde concentrations were below the LoQ during both seasons.

3.5. Interpolated surfaces of NO$_2$ and formaldehyde

For both NO$_2$ (**Figure 3**) and formaldehyde (**Figure 4**), we observed higher concentrations in the south of Viadana, with hot-spots of concentrations above the 90th percentile in proximity to the industry. However, the north-eastern area of the municipality had higher concentrations in 2010 compared to 2017-18 for both pollutants, which could have been driven by the different location of samplers between the two campaigns. Very few samplers were placed in that area, possibly affecting estimation accuracy and precision. The same holds true for the area at the East of the industry, where no samplers were located in 2017-18.

Areas closer to the industry were also characterised by a higher density of roads, population and buildings (**Table 3**). For example, the median total road length in 500-m buffers around the sampling sites decreased from 8.3 km to 4.6 km as the distance to the factory increased (from <1.2 to >1.8 km). It is therefore likely that road traffic and domestic heating contribute the higher levels of air pollution observed in proximity to the industry.

Overall, estimated concentrations were lower in 2017-18 compared to 2010 (see legends of **Figures 3-4**). This could be linked to the different weeks during the solar year selected for the two campaigns, which captured different seasonal and meteorological conditions. An overall change in industrial emissions or local traffic is an unlikely explanation for this finding, because that would have affected spatial gradients rather than overall concentrations.

3.6. Comparison between air pollution exposure models

Correlations between estimates of residential exposure attributed to children's addresses are reported in **Table 4**. Pearson's r coefficients between pollutants varied widely, from 0.22 (NO$_2$ Viadana II vs PM$_{2.5}$ ELAPSE) to 0.86 (NO$_2$ vs formaldehyde, Viadana III). Most of the highest correlations were observed between pollutants estimated from the same

project, rather than when comparing the same pollutant from different models. To give an example for the models employing routine monitoring data, r coefficient for the correlation between NO_2 and $PM_{2.5}$ from ELAPSE was 0.79, whereas it was 0.42 for $PM_{2.5}$ obtained from ELAPSE and EPISAT.

The three models for NO_2 detected a maximum concentration at a distance of 1–1.7 km to the industry, followed by a decline as distance increased (**Table 5**), possibly reflecting the pattern of dispersion of nitrogen oxides from the 70-m high chimney. The latter finding is less likely to reflect the local distribution of urban emission sources, since the densities of roads, buildings and population were higher in the buffer closest to the industry. Absolute differences in median NO_2 exposures between the children at 1–1.7 km and >3.5 km ranged from 2.4 µg/m^3 (Viadana II) to 5.9 µg/m^3 (Viadana III). For the other pollutants, estimated exposures decreased with distance: differences in medians between children living at <1 km and ≥3.5 km to the industry were 2.2–2.4 for $PM_{2.5}$ and 5.0 µg/m^3 for PM_{10}. Differences between medians for formaldehyde and BC were small.

In summary, despite the differences in methodology (kriging vs LUR), input data (routine vs passive sampling data), extension (municipality vs national/European), and period (2010-13 vs 2017-18), all the exposure models consistently suggest that children living closer to the chipboard industry (<1.7 km) were more exposed than the children living farther away (>3.5). However, it is important to highlight that, given the proximity between the chipboard industry and the most urbanised areas in Viadana, it is not straightforward to disentangle between industrial emissions and other anthropic sources of contaminants.

3.7. Limitations

A direct comparison between passive sampling data from the campaigns conducted in 2010 and 2017-18 is hindered by the different number and location of samplers in the monitoring area. Another drawback is that we had no direct PM measurements in the

study area. PM are one of the most important emissions from the industry in Viadana, and they convey most of formaldehyde mass that is released during production. They are the most critical component of air pollution in the Po Plain due to their regional distribution and long-term atmospheric persistence (Larsen et al. 2012). LUR models based on routine monitors from country-wide or larger areas perform well in the Po Plain (de Hoogh et al. 2018; Stafoggia et al. 2017) but they may not be ideal to capture variability in industrial pollution, due to the relatively small number of industrial monitoring sites, and the inability of LUR predictors (e.g. land use) to accurately reflect industrial density and specific industrial sources.

4. CONCLUSIONS

Using data from recent passive sampling campaigns, we found that residential exposure to NO_2 and formaldehyde was higher in proximity (<1.7 km) to the chipboard industry in Viadana. Hot-spots with higher pollutant concentrations were found both in 2017-18 and in 2010 in the south, in close proximity to the industry, which is also the most densely urbanised zone. This suggests that the increased pollutant exposure for residents in proximity to the industry is expected to be stable during the Viadana III follow-up period (2013-2017). Exposure estimates derived from available large-scale LUR models of NO_2, $PM_{2.5}$ and PM_{10} from routine stations showed associations with distance comparable to our interpolated surfaces. This highlights that people living close to the industry are exposed to a mixture of air pollutants, and it supports the reliability of exposure estimates derived for the Viadana III study. Finally, we point that Radiello® tubes may underestimate NO_2 concentrations, which is important to keep in mind when interpreting association estimates from epidemiological research.

TABLES

Table 1. Distribution of weekly measurements of air pollutants from 25 passive samplers in Viadana, and daily temperature and rain precipitations from the nearest meteorological station for the same weeks (2017-18).

Week n.	Starting date (DD/MM/YY)	NO$_2$, µg/m^3 (mean, SD)	Formaldehyde, µg/m^3 (mean, SD)	Temperature, °C (mean, min–max)	Rain, mm (mean, min–max)
1	21/11/17	19.5 (4.3)	1.6 (1.0)	6.8 (2.9–9.5)	2.6 (0.0–11.4)
2	28/11/17	24.1 (3.1)	3.2 (0.6)	2.8 (0.9–5.0)	0.0 (0.0–0.0)
3	05/12/18	22.3 (5.5)	2.0 (1.5)	1.7 (-0.8–4.3)	1.7 (0.0–9.4)
4	12/12/18	12.2 (5.6)	2.6 (1.2)	2.2 (-0.9–3.4)	0.1 (0.0–0.4)
5	18/12/18	20.8 (4.5)	1.8 (0.9)	1.3 (-0.6–5.2)	0.0 (0.0–0.0)
Overall (cold season)		19.8 (3.5)	2.2 (0.5)	2.9 (-0.9–9.5)	0.8 (0.0–11.4)
6	08/05/18	13.2 (5.9)	0.6 (0.2)	18.7 (16.1–20.4)	2.2 (0.0–7.4)
7	15/05/18	8.5 (2.7)	0.7 (0.4)	17.5 (12.8–19.7)	0.7 (0.0–5.2)
8	22/05/18	10.1 (4.3)	0.6 (0.4)	21.8 (17.3–23.9)	7.8 (0.0–50.6)
9	29/05/18	9.5 (3.9)	0.8 (0.5)	23.2 (22.4–23.8)	0.4 (0.0–2.0)
10	05/06/18	8.3 (2.9)	2.6 (0.8)	23.6 (20.5–25.1)	5.0 (0.0–32.2)
Overall (warm season)		9.9 (3.1)	1.1 (0.3)	21.0 (12.8–25.1)	3.2 (0.0–50.6)

6 **Table 2**. Distribution of measured concentrations of air pollutants at the 25 passive sampling locations in Viadana (2017-18), by
7 groups based on tertiles of distance to the industry.

	Distance to the industrial plant				Coefficient of variation (n=25)
	<1.2 km [a] (n=8)	1.2–1.7 km [a] (n=9)	1.8–15.5 km [a] (n=8)	Overall [a] (n=25)	
NO_2 (cold season), µg/m^3	21.5 (19.8–23.6)	19.4 (18.8–21.2)	16.4 (14.7–20.4)	19.9 (17.5–22.0)	0.18
NO_2 (warm season), µg/m^3	11.6 (9.5–13.4)	9.0 (6.6–13.0)	8.8 (6.5–11.0)	9.9 (6.7–13.0)	0.31
NO_2 (annual), µg/m^3	16.4 (15.8–16.6)	14.4 (13.0–17.1)	11.9 (10.9–16.1)	16.0 (12.4–16.6)	0.19
Formaldehyde (cold season), µg/m^3	2.3 (2.1–2.4)	2.0 (1.71–2.7)	2.2 (2.0–2.3)	2.2 (2.0–2.6)	0.22
Formaldehyde (warm season), µg/m^3	1.1 (1.1–1.4)	1.1 (0.9–1.2)	1.0 (0.9–1.0)	1.0 (0.9–1.2)	0.25
Formaldehyde (annual), µg/m^3	1.7 (1.5–2.0)	1.5 (1.4–2.0)	1.6 (1.4–1.7)	1.6 (1.5–2.0)	0.18
Acetaldehyde (cold season), µg/m^3	0.7 (0.6–0.7)	0.7 (0.6–0.9)	0.7 (0.6–0.8)	0.7 (0.6–0.8)	0.19
Benzene (cold season), µg/m^3	1.6 (1.5–1.6)	1.6 (1.3–1.6)	1.6 (1.5–1.6)	1.6 (1.5–1.6)	0.10
Toluene (cold season), µg/m^3	2.9 (2.8–3.0)	2.9 (2.5–3.1)	2.8 (2.6–3.2)	2.8 (2.6–3.1)	0.11

9 [a] Medians with interquartile ranges reported.

Table 3. Distribution of population and road density indicators in 500 m buffers around the 25 passive sampling sites in Viadana, by groups based on tertiles of distance to the industry. [a]

	Distance to the industrial plant			
	<1.2 km (n=8)	1.2–1.7 km (n=9)	1.8–15.5 km (n=9)	Overall (n=25)
Number of inhabitants	1506 (1048–2701)	713 (536–2702)	527 (337–939)	993 (536–2383)
Number of buildings	332 (236–508)	255 (151–501)	198 (107–255)	254 (185–426)
Number of households	631 (419–1189)	294 (229–1202)	224 (126–381)	393 (229–1053)
Total road length (km)	8.3 (6.9–10.7)	5.6 (3.2–9.9)	4.6 (3.1–6.9)	6.9 (3.8–9.7)

[a] Medians with interquartile ranges reported.

Table 4. Pearson's r coefficients for the correlation between exposures estimated at children's residential addresses (n=1,814). [a]

		Viadana II (2010)		ELAPSE (2010)			EPISAT		Viadana III (2017-18)	
		NO_2	Form-aldehyde	NO_2	$PM_{2.5}$	BC	$PM_{2.5}$ (2013)	PM_{10} (2012)	NO_2	Form-aldehyde
Viadana II (2010)	NO_2	1								
	Formaldehyde	0.42	1							
ELAPSE (2010)	NO_2	0.43	0.51	1						
	$PM_{2.5}$	0.22	0.49	0.79	1					
	BC	0.26	0.54	0.74	0.80	1				
EPISAT	$PM_{2.5}$ (2013)	0.40	0.39	0.44	0.42	0.41	1			
	PM_{10} (2012)	0.35	0.48	0.43	0.42	0.46	0.60	1		
Viadana III (2017-18)	NO_2	0.49	0.68	0.68	0.69	0.74	0.46	0.50	1	
	Formaldehyde	0.57	0.68	0.61	0.57	0.65	0.38	0.50	0.86	1

[a] all p<0.001

Table 5. Distribution of exposures estimated at children's residential addresses, by groups based on quartiles of distance to the industry. [a]

	Distance to the industrial plant				
	<1 km (n=454)	1–1.7 km (n=455)	1.8–3.4 km (n=452)	3.5–15.7 km (n=453)	Overall (n=1,814)
Viadana II study (2010)					
NO_2, µg/m^3	17.9 (16.8–18.4)	19.3 (17.8–20.0)	18.1 (17.4–18.6)	16.9 (14.9–17.4)	17.8 (16.8–18.7)
Formaldehyde, µg/m^3	2.7 (2.6–2.9)	2.7 (2.6–2.7)	2.5 (2.5–2.6)	2.3 (2.2–2.5)	2.6 (2.5–2.7)
ELAPSE study (2010)					
NO_2, µg/m^3	28.7 (27.4–29.9)	29.0 (27.9–29.8)	28.4 (27.0–29.5)	24.6 (23.1–26.7)	28.1 (26.2–29.5)
$PM_{2.5}$, µg/m^3	26.9 (26.4–27.0)	26.8 (26.6–26.9)	26.6 (26.2–26.9)	24.5 (22.9–26.4)	26.6 (26.0–26.9)
BC, 10^{-5} m^{-1}	2.0 (2.0–2.0)	2.0 (2.0–2.0)	2.0 (1.9–2.0)	1.9 (1.8–1.9)	2.0 (1.9–2.0)
EPISAT study (2012-13)					
$PM_{2.5}$ (2013), µg/m^3	25.8 (25.0–26.6)	25.5 (25.2–26.0)	25.9 (25.1–26.8)	23.6 (22.5–24.8)	25.4 (24.6–26.3)
PM_{10} (2012), µg/m^3	38.3 (36.7–40.5)	37.2 (36.1–39.0)	36.6 (34.7–38.8)	33.3 (31.8–34.6)	36.5 (34.0–38.6)
Viadana III study (2017-18)					
NO_2, µg/m^3	16.5 (15.4–16.8)	17.1 (16.7–17.2)	16.0 (13.9–16.8)	11.2 (10.8–12.1)	16.2 (13.6–16.9)
Formaldehyde, µg/m^3	1.7 (1.6–1.8)	1.8 (1.7–1.8)	1.7 (1.6–1.8)	1.6 (1.5 1.6)	1.7 (1.6–1.8)

[a] Medians with interquartile range reported. p-values for overall comparisons of concentrations across groups were all <0.001 (Kruskall-Wallis rank test)

FIGURES

Figure 1. Weekly concentrations of NO$_2$ measured by continuous monitors and passive
samplers during the cold (panel A) and warm seasons (panel B). Municipality of Viadana,
2017-18. [a]

[a] the solid thin line represents weekly average concentrations by the routine station; the solid and dashed
thick lines represent the weekly average concentrations by the temporary station and the co-located passive
sampler, respectively; the dotted line is the median concentration by the 25 passive samplers (coloured
lines)

Figure 2. Box & whiskers plot of recalibrated NO_2 concentrations at passive sampling
locations in Viadana (2018-19), by groups based on tertiles of distance to the industry.

Figure 3. Spatial distribution of annual NO$_2$ concentrations measured by passive samplers in 2010 (panel A) and 2017-18 (panel B). [a]

[a] Obtained using ordinary kriging. Black contour lines represent the 90[th] percentile. The white triangle represents the industry in Viadana. Blue triangles represent smaller wood factories. Cyan dots represent passive sampling sites.

Figure 4. Spatial distribution of annual formaldehyde concentrations measured by passive samplers in 2010 (panel A) and 2017-18 (panel B). [a]

[a] Obtained using ordinary kriging. Black contour lines represent the 90[th] percentile. The white triangle represents the industry in Viadana. Blue triangles represent smaller wood factories. Cyan dots represent passive sampling sites.

REFERENCES

Badaloni C, Cattani G, De' Donato F, Gaeta A, Leone G, Michelozzi P, et al. 2018. Big data in environmental epidemiology. Satellite and land use data for the estimation of environmental exposures at national level. Epidemiol Prev 42:46-59.

Beelen R, Hoek G, Vienneau D, Eeftens M, Dimakopoulou K, Pedeli X, et al. 2013. Development of NO2 and NOX land use regression models for estimating air pollution exposure in 36 study areas in Europe - the ESCAPE project. Atmospheric Environment 72:10-23.

Buczynska AJ, Krata A, Stranger M, Godoi AFL, Kontozova-Deutsch V, Bencs L, et al. 2009. Atmospheric btex-concentrations in an area with intensive street traffic. Atmos Environ 43:311-318.

Curtis L, Rea W, Smith-Willis P, Fenyves E, Pan Y. 2006. Adverse health effects of outdoor air pollutants. Environ Int 32:815-830.

Dahlgren J, Warshaw R, Thornton J, Anderson-Mahoney CP, Takhar H. 2003. Health effects on nearby residents of a wood treatment plant. Environ Res 92:92-98.

de Hoogh K, Chen J, Gulliver J, Hoffmann B, Hertel O, Ketzel M, et al. 2018. Spatial pM2.5, NO2, O3 and BC models for western Europe - evaluation of spatiotemporal stability. Environ Int 120:81-92.

de Marco R, Marcon A, Rava M, Cazzoletti L, Pironi V, Silocchi C, et al. 2010. Proximity to chipboard industries increases the risk of respiratory and irritation symptoms in children: The Viadana study. Sci Total Environ 408:511-517.

Evans AM, Stuart AL. 2011. A passive sampling study of small-scale variations in ambient acetaldehyde and formaldehyde concentrations. Air, Soil and Water Research 4:ASWR. S7582.

Gaeta A, Cattani G, Di Menno di Bucchianico A, De Santis A, Cesaroni G, Badaloni C, et al. 2016. Development of nitrogen dioxide and volatile organic compounds land use regression models to estimate air pollution exposure near an italian airport. Atmos Environ 131:254-262.

Gerboles M, Buzica D, Amantini L, Lagler F. 2006. Laboratory and field comparison of measurements obtained using the available diffusive samplers for ozone and nitrogen dioxide in ambient air. J Environ Monit 8:112-119.

Girardi P, Marcon A, Rava M, Pironi V, Ricci P, de Marco R. 2012. Spatial analysis of binary health indicators with local smoothing techniques: The Viadana study. Sci Total Environ 414:380-386.

IARC. 2006, update 2018. Formaldehyde, 2-butoxyethanol and 1-tert-butoxypropan-2-ol. IARC Monogr Eval Carcinog Risks Hum 88:1-478.

Kerchich Y, Kerbachi R. 2012. Measurement of BTEX (benzene, toluene,
ethybenzene, and xylene) levels at urban and semirural areas of Algiers city using passive
air samplers. Journal of the Air & Waste Management Association 62:1370-1379.

Kheirbek I, Johnson S, Ross Z, Pezeshki G, Ito K, Eisl H, et al. 2012. Spatial variability
in levels of benzene, formaldehyde, and total benzene, toluene, ethylbenzene and xylenes
in New York City: A land-use regression study. Environmental Health 11:51.

Larsen BR, Gilardoni S, Stenström K, Niedzialek J, Jimenez J, Belis CA. 2012.
Sources for pm air pollution in the Po plain, Italy: II. Probabilistic uncertainty
characterization and sensitivity analysis of secondary and primary sources. Atmos Environ
50:203-213.

Marchetti P, Marcon A, Pesce G, Girardi P, Guarda L, Pironi V, et al. 2014. Children
living near chipboard and wood industries are at an increased risk of hospitalization for
respiratory diseases: A prospective study. Int J Hyg Environ Health 217:95-101.

Marchetti P, Pesce G, Villani S, Antonicelli L, Ariano R, Attena F, et al. 2017. Pollen
concentrations and prevalence of asthma and allergic rhinitis in Italy: Evidence from the
geird study. Sci Total Environ 584:1093-1099.

Marcon A, Fracasso ME, Marchetti P, Doria D, Girardi P, Guarda L, et al. 2014.
Outdoor formaldehyde and NO2 exposures and markers of genotoxicity in children living
near chipboard industries. Environ Health Perspect 122:639-645.

Mason JB, Fujita EM, Campbell DE, Zielinska B. 2011. Evaluation of passive samplers
for assessment of community exposure to toxic air contaminants and related pollutants.
Environmental Science & Technology 45:2243-2249.

Morello-Frosch RA, Woodruff TJ, Axelrad DA, Caldwell JC. 2000. Air toxics and health
risks in california: The public health implications of outdoor concentrations. Risk Anal
20:273-291.

Pebesma EJ, Wesseling CG. 1998. Gstat: A program for geostatistical modelling,
prediction and simulation. Computers & Geosciences 24:17-31.

Rava M, Marcon A, Girardi P, Pironi V, Silocchi C, Ricci P, et al. 2011. Proximity to
wood factories and hospitalizations for respiratory diseases in children. Sci Total Environ
410:80-86.

Rava M, Crainicianu C, Marcon A, Cazzoletti L, Pironi V, Silocchi C, et al. 2012.
Proximity to wood industries and respiratory symptoms in children: A sensitivity analysis.
Environ Int 38:37-44.

Santana FO, Campos VP, Cruz LP, Luz SR. 2017. Formaldehyde and acetaldehyde in
the atmosphere of Salvador-Ba, Brazil, using passive sampling. Microchemical Journal
134:78-86.

Stafoggia M, Schwartz J, Badaloni C, Bellander T, Alessandrini E, Cattani G, et al.
2017. Estimation of daily PM10 concentrations in italy (2006-2012) using finely resolved
satellite data, land use variables and meteorology. Environ Int 99:234-244.

Villanueva F, Tapia A, Notario A, Albaladejo J, Martínez E. 2014. Ambient levels and
temporal trends of VOCs, including carbonyl compounds, and ozone at Cabañeros
national park border, Spain. Atmos Environ 85:256-265.

Yu CH, Morandi MT, Weisel CP. 2008. Passive dosimeters for nitrogen dioxide in
personal/indoor air sampling: A review. J Expo Sci Env Epid 18:441-451.

136