Analysis of the impact of lockdown on the evolution of Covid-19 epidemics in Spain

Alexandre Hyafil[1,2*], David Moriña[2,3]

[1]Centre de Recerca Matemàtica (CRM)

[2]Barcelona Graduate School of Mathematics (BGSMath), Departament de Matemàtiques, Universitat Autònoma de Barcelona (UAB), 08193 Cerdanyola del Vallès, Barcelona (Spain)

[3]Department of Econometrics, Statistics and Applied Economics, Riskcenter-IREA, Universitat de Barcelona (UB), 08034 Barcelona (Spain)

* Corresponding author:

Alexandre Hyafil

Centre de Recerca Matemàtica, Edifici C, Campus de Bellaterra, Universitat Autònoma de Barcelona, 08193 Cerdanyola del Vallès, Barcelona (Spain)

ahyafil@crm.cat

Abstract

Objective: The late 2019 Covid-19 disease outbreak has put the health systems of many countries to the limit of their capacity. The most affected European countries are, so far, Italy and Spain. In both countries (and others), the authorities decreed a lockdown, with local specificities. The objective of this work is to evaluate the impact of the measures undertaken in Spain to deal with the pandemic. ***Method:*** We estimated the number of cases and the impact of lockdown on the reproducibility number based on the hospitalization reports. ***Results:*** The estimated number of cases shows a sharp increase until the lockdown, followed by a slowing down and then a decrease after full quarantine was implemented. Differences in the basic reproduction ratio are also very significant, dropping from de 5.89 (95% CI: 5.46-7.09) before the lockdown to 0.48 (95% CI: 0.15-1.17) afterwards. ***Conclusions:*** Handling a pandemic like Covid-19 is very complex and requires quick decision making. The large differences found in the speed of propagation of the disease show us that being able to implement interventions at the earliest stage is crucial to minimise the impact of a potential infectious threat. Our work also stresses the importance of reliable up to date epidemiological data in order to accurately assess the impact of Public Health policies on viral outbreak.

Keywords: Covid-19, intervention analysis, public health, infections, virus

Análisis del impacto del confinamiento en la evolución de la epidemia de Covid-19 en España

Resumen

Objetivo: El brote de la enfermedad Covid-19 a finales de 2019 ha puesto los sistemas de salud de muchos países al límite de su capacidad. Los países europeos más afectados son, hasta ahora, Italia y España. En ambos países (y en otros), las autoridades decretaron un confinamiento, con especificidades locales. El objetivo de este trabajo es evaluar el impacto de las medidas adoptadas en España para hacer frente a la pandemia. ***Método:*** Estimamos el número de casos y el impacto del confinamiento en el número básico de reproducción según los informes de hospitalización. ***Resultados:*** El número estimado de casos muestra un fuerte aumento hasta el bloqueo, seguido de una desaceleración y luego una disminución después de la implementación del confinamiento total. Las diferencias en el número básico de reproducción también son muy significativas, cayendo de 5.89 (95% IC: 5.46-7.09) antes del bloqueo a 0.48 (95% IC: 0.15-1.17) después. ***Conclusiones:*** Gestionar una pandemia como la Covid-19 es muy complejo y requiere una rápida toma de decisiones. Las grandes diferencias encontradas en la velocidad de propagación de la enfermedad nos muestran que poder implementar intervenciones en la etapa más temprana de la misma es crucial para minimizar el impacto de una potencial amenaza. Nuestro trabajo también muestra la importancia de contar con datos epidemiológicos actualizados y confiables para evaluar con precisión el impacto de las políticas de Salud Pública en el brote.

Palabras clave: Covid-19, análisis de intervención, salud pública, infecciones, virus

1. Introduction

By late 2019 an outbreak of Covid-19 disease -caused by SARS-Cov-2 virus- started in the region of Hubei (China), more specifically in the city of Wuhan. Since then, the disease has spread all over the world, being declared pandemic by the World Health Organization (WHO) on 2020 March 11th. The rapid propagation of the virus around the world has stressed the health systems of many countries to their limit. In Europe, the most affected countries in terms of number of detected cases, hospitalizations and deaths are, to the date, Italy and Spain. This unprecedented situation for the public health systems of these countries forced decision makers to act very quickly in order to minimize the impact of the disease and to avoid collapse. In Spain, the emergency state (*estado de alarma*) was declared on 2020 March 14th and was hardened with mandatory home confinement except for vital sectors workers (including health professionals, food supply, etc.) on March 30th. Under this situation, it is urgent to assess the efficiency of social distancing and other non-pharmaceutical interventions undertaken to control the pandemics of Covid-19 (Niud and Xu 2020; Anderson et al. 2020). One of the main challenges in evaluating the impact of these actions is that data are only partially available, as many of the cases are asymptomatic or with mild symptoms (Huang et al. 2020), and shortage of testing kits prevent testing all patients with possible Covid-19 symptoms. Therefore, the number of cases might be severely underestimated. This issue is common in epidemiology and several methods have been recently proposed to address it (see (Fernández-Fontelo et al. 2016, 2019)) under specific circumstances. Another issue is that, when no massive viral testing of the population is performed, new cases are only detected once symptoms appear, at a variable delay after contamination. This complicates the assessment of the impact of interventions to reduce/slow down the spread of a virus.

Here, we aimed at evaluating the impact of the Covid-19 related non-pharmaceutical interventions undertaken in Spain over the basic reproduction ratio R_0, considering three periods of time in 2020: No intervention (until March 13th), emergency state (March 16-30th and April 13th-15th) and mandatory confinement (March 31st to April 12th). We inferred retrospectively the number of contamination in each Spanish region (*Comunidad*

Autónoma, or CCAA) from the patterns of hospitalizations, deaths and detected cases. The mandatory confinement situation was softened on 2020/04/13 (2020/04/14 on some regions) by allowing non-essential sectors to return to work, so a projection of the situation in the next few weeks was also included. Additionally, the results are compared to a scenario where no measures are undertaken.

2. Methods

We modelled the dynamics of infection with a discrete-time SIR model with time-varying R_0 defined for each region r and day d. We also assumed the stock of susceptible population $S(r, d)$ is almost constant through time at $S(r, d) \approx N_d$, (i.e. a small percentage of the population is contaminated) as is probably valid up to now. The dynamics of infected people $I(r, d)$ in region r at day d thus varies as:

$$I(r, d + 1) = I(r, d)e^{(R_0(r,d)-1)\gamma}$$

R_0 is a stochastic variable whose expected value is defined by the ongoing social distancing measures (no measure, state emergency or mandatory confinement). Formally, the prior over R_0 is defined as a Gaussian Processes independently defined between regions. The prior mean is defined by the distancing measures, while the prior covariance isSquared Exponential covariance K with variance $\sigma^2 = 0.1^2$ and length $l = 1$ day (Bishop 2006). Such prior allows to capture differences in the spread between different regions as well as temporary change of R_0 within a region (Gaussian Processes enforce that these fluctuations are smooth in time). The variance term σ^2 was set so it is unlikely that the infected population goes down in a single day by a proportion larger than the recovery rate γ. Based on the mean 20 days of contagious before recovery (duration of viral shedding (Zhou et al. 2020)), we set $\gamma = 0.05 \text{day}^{-1}$.

We assumed that the value of infected people at the beginning of the period studied (20th of February 2020) was drawn from a log-normal distribution with mean $x_i(r)$ and variance 1, where x_i is a parameter specific to each CCAA. This variability allowed to captured initial variations in the spread of the epidemics at the beginning of the period of study.

The number of new cases per day is $N(c,d) = I(r,d) - I(r,d-1) + \gamma I(r,d-1) = I(r,d) - (1-\gamma)I(r,d-1)$. This true number of cases however cannot be observed directly. Here we use a latent-state approach: we estimated the evolution of the number of true cases in each CCAA based on the recorded accumulated number of detected cases, hospitalizations and deaths provided by Instituto de Salud Carlos III (https://portalcne.isciii.es/covid19/). We used estimates for the proportion of cases p_C that are detected by the Spanish Health system, the proportion of cases p_H that are hospitalized, and the lethality rate p_D, as the well as the distribution of latency between infection and this three types of events $l_i(d)$ (figure 1). We used the following estimates: 15% of contaminated persons are hospitalized; 1% die from disease (Lauer et al. 2020; Zhou et al. 2020); 30% of cases get detected (this latter number was defined arbitrarily, since the number of detected cases is roughly twice the number of hospitalizations). Note that these percentages affect the estimated number of true cases by a scaling factor, but do not affect the estimation of reproduction parameters. We used estimates of the distribution of duration of infection-to-detection, infection-to-hospitalization based on published literature (Lauer et al. 2020; Zhou et al. 2020). Lauer and colleagues describe that the incubation period can be captured by log-normal distribution with median 5.1 days. To capture the extra time from the apparition of symptoms to case detection/hospitalization, we increased the μ parameter (to yield an increase in median time of 50% for case detection and 100% for hospitalization, that is $\mu = 2.03$ and $\mu = 2.314$, respectively) while keeping the same dispersion parameter σ. For infection-to-death, we used the same distribution as in (Flaxman et al. 2020), i.e. the sum of two gamma distributions for infection-to-detection and detection-to-death with overall mean 23.9 days. Thus the expected numbers of events $y_i(r,d)$ in a certain CCAA is given by convolving the pattern of new cases with the distribution of infection-to-event.

$$E[y_i(r,d)] = p_i \sum_{t<d} N(c,t) l_i(d-t)$$

$$E[y_i(r,d)] = \sum_{t<d} w_{d-t} I(c,t)$$

with $w_d = p_i\big(l_i(d) - (1-\gamma)l_i(d-1)\big)$. We assumed that the actual number of events recorded at that day and time was drawn from a negative binomial distribution with mean $E[y_i(r,d)]$ as defined above and parameter $r = 2$ (Flaxman et al. 2020). The reports provide accumulated time series, so in principle new events correspond to the difference between two successive days. However new events are sometimes reported later than on the day of occurrence (especially during weekends). We estimated that 30% of events were reported on the following day, 10 % two days later. Assigning such proportion of events to one or two days before they are reported allowed to smooth the event time series. Finally, for the case of 2 CCAA (Madrid and Castilla y La Mancha, until April 12[th] for the latter), the reported number of hospitalizations was not cumulative but corresponded to the current number of hospitalized patients related to Covid-19. For Castilla y La Mancha, we found that we could recover the reported cumulative number of hospitalizations on April 12[th] by assuming that the duration of hospitalization is distributed uniformly between 5 and 15 days. We used this rule to estimate the cumulative number of hospitalizations for this CCAA, and applied it similarly for Madrid.

There is a debate about the reliability of these data. It is believed that hospitalizations in the most reliable of these indicators,[1] as many cases go undetected (some patients are asymptomatic or suffer mild symptoms; saturation of health systems have led to testing only more severely affected patients in some CCAA), and some deaths are not integrated in the official count because of the lack of viral charge testing. There is indeed a large variability between the fraction of hospitalizations per reported case between CCAA: it is 31% in Galicia but 82% in Comunidad de Madrid. This difference is more likely to be due to differences in detecting and reporting cases rather than in the true proportion of infected people requiring hospitalization. It is also more difficult to assess reliably the number of cases from death reports as the latency from infection to death is long and can be very variable across individuals.

---

[1]   https://elpais.com/sociedad/2020-04-03/una-epidemia-en-la-que-todas-las-cifras-son-una-incognita.html

Parameters of the model, including the value of R_0 in the different conditions, were estimated from the data using an Expectation-Maximization algorithm (see Appendix 5 for details).

Figure 1: Probability of the three types of events used to infer number of cases: detected cases, hospitalizations and deaths. Left panel: overall probability for each event. Right panel: normalized distribution for the latency from contamination to event.

3. Results

We present the number of Covid-19 cases in all 19 CCCAA inferred from the pattern of hospitalizations in Figure 2. The estimated cases display a sharp increase until the lockdown followed by a plateau, and then a decrease. We estimated the R_0 before state of emergency, during state emergency and during enforced lockdown. R_0 was found to drop from 5.89 (95% CI: 5.46-7.09) to 1.86 (95% CI: 1.10-2.63) after state of emergency, and down to 0.48 (95% CI: 0.15-1.17) after full lockdown (Figure 3). We estimate a total number of 0.871 million Covid-19 cases in Spain by April 15th 2010, including 0.294 active cases, 0.559 recovered and 0.018 deceased (see Figure 4 for breakdown by CCAA).

Figure 2: Number of estimated Covid-19 cases in all Spanish Comunidades Autónomas, inferred from the number of hospitalizations. The posterior mean of inferred cases is plotted in blue. Cumulative number of detected cases, hospitalizations and deaths are plotted in green, red and black curves, respectively. Grey curves represent the inferred number of cases in other CCAA. The onset of state of emergency and mandatory confinement are indicated by a vertical bar.

Figure 3: Estimated R_0 during the three periods of time (prior to state of emergency; state of emergency; during mandatory confinement). Error bars denote 95% confidence intervals, obtained from fits of bootstrap estimates to a gamma distribution.

Figure 4: Cumulative number of cases in each CCAA, and breakdown into active cases, recovered cases and deceased patients.

Estimates were similar when we used detected cases rather than hospitalizations. Using detected case reports, we estimated the R_0 to be 6.91 (95% CI: 6.75-7.39) before the state of urgency, 2.22 (95% CI: 1.92-2.74) during the state of emergency and 0.85 (95% CI: 0.5-1.05) during the full lockdown (Figure 5). The estimate for the cumulative number of cases were 0.823 million overall in Spain (0.351 million active).

Because deaths occur after a long and variable interval after contamination, we could not reliably estimate the value of R_0 separately for the state emergency and full lockdown measures, so we simply estimated R_0 before and after state of emergency. We estimated the R_0 to be 6.48 before the lockdown (95% CI: 5.5-7.51), and 0.49 afterwards (95% CI: 0.16 - 1.57) (Figure 6). The estimate for the cumulative number of cases were 2.82 millions overall in Spain (0.72 millions active).

Finally, we used the reproducibility parameters estimated from hospitalizations to evaluate the death toll in different scenarios (Appendix B). First, we estimated retrospectively how many deaths have been avoided by the state of emergency (and temporary mandatory confinement). We estimate that, without state of emergency, the new coronavirus would have cause 0.180 million deaths by April 15th (Figure 7, left panels), i.e. 10 times the number of deaths actually reported. In that scenario, Spain would have been approaching herd immunity by mid-April.

We also performed two prospective analyses for the period April 16th -May 15th, assuming either that the state of emergency is maintained as currently in place, or the stricter mandatory confinement with closure of all non-essential activities is implemented again (Figure 7, right panels). The analysis predicts a total of 23.900 deaths (95% CI: 23.560-24.239) in the former case, and 21.423 (95% CI: 21.298-21.548) in the latter. It should be noted however that only the latter predicts a gradual extinction of cases, while our analyses predict that the state of emergency would lead to a new surge of cases.

Figure 5: R_0 before and after lockdown estimated from death reports. Error bars denote 95% confidence intervals, obtained from fits of bootstrap estimates to a gamma distribution.

Figure 6: R_0 before and after lockdown estimated from detected cases reports. Error bars denote 95% confidence intervals, obtained from fits of bootstrap estimates to a gamma distribution.

Figure 7: Projections of deaths in each region for different scenarios. On left plots, number of predicted deaths if no state of emergency had been issued (black lines) vs. actual number of deaths (orange line) for period April 16th-May 15th 2020. On right plots, predictions for next 30 days if state of emergency is maintained (black lines) vs. going back to mandatory confinement (red lines). Shaded area represent standard deviation (i.e. 68% confidence interval).

4. Discussion and limitations

We found similar estimates of reproducibility number and the proportion of the Spanish population contaminated by the new coronavirus, whether they were estimated from hospitalization numbers or detected cases. Both estimates from case reports and hospitalizations suggest that only mandatory quarantine achieved R_0 lower than 1, while R_0 during state of emergency before non-essential services were shut down was estimated to be well beyond 1. This predicts that the opening of the non-essential services by April 13 may lead to a new surge of cases. Based on this empirical study, mandatory confinement is the only measure that effectively reduces the number of contaminations.

Estimated based on death reports differed considerably from those based on either hospitalizations or detected cases. This suggests that some of the assumptions and data our modeling is based on may not be accurate (although commonly used in previous studies), and shows that this can induce very large biases in the estimation of the propagation of the new coronavirus in Spain. Since hospitalizations reports are believed to be more reliable than either reported cases or deaths reports, we further comment on the results obtained with these reports.

Our approach is very similar to a study by an Imperial College team published last week inferring the impact of non-pharmaceutical measures (including lockdown) on propagation of the new coronavirus in 11 European countries (Flaxman et al. 2020). Both studies rely on fitting a model of infection dynamics to observed data (here hospitalizations). This contrasts with other approaches based on fitting a curve to the observed time series, (e.g. for patterns of fatalities in United States (Team and Murray 2020) or patterns of cases in China (Zhao et al. 2020)), or to model simulations studies that capture how the pattern of contacts in different scenarios (with or without social distancing measures) affect virus propagation (Lin et al. 2020; Hellewell et al. 2020; Ferguson et al. 2020; Lai et al. 2020). Other studies have also estimated the number of cases from the reported deaths, assuming a fixed duration from infection to death.[2]

2   https://biocomsc.upc.edu/en/covid-19/

Figure 8: Increase in number of cases (in percent) for each CCAA in the 10 days following lockdown versus number of cases per 100.000 habitants at the time of lockdown

Our modeling approach include stochasticity in the reproducibility number in each area. This notably allows to capture distinct trajectories of infection in different areas. We noted a negative correlation across CCAA between the proportion of infected people at the time of lockdown and the subsequent increase in infection (Figure 8, Pearson coefficient: $r = -0.46$, $p = 0.048$). This could be due to a series of factors. First, the communities with largest proportion of cases could start developing herd immunity, hence limiting the propagation of the infection (which seems unlikely given that in the most affected CCAA, only a few percent have been infected according to our estimations). It could also be the result of local policies taken before the national lockdown in the most affected CCAA, or a better compliance of lockdown and social distancing in most affected areas. Another factor could be the migration from the most affected (especially Madrid) to less affected region before lockdown was implemented,[3] or some distortions in the reporting (under-reporting) of cases in saturated health systems.

---

[3]   see https://elpais.com/espana/madrid/2020-04-08/los-madrilenos-que-huyeron-a-la-espana-rural-durante-el-confinamiento.html

Our study has several important limitations. First, as noted previously, it is not clear how reliable is the data the modeling is based on: both epidemiological reports, and infectious estimate probability and latency of symptoms, case detection, hospitalization and death). It should be stressed that the unreliability of epidemiologic data (with changes of criterion of inclusion along time and between CCAA) induces important biases which impedes an accurate estimation of the impact of political measures on the propagation of the new coronavirus. A faster and more reliable tracking of the epidemics could be performed if cases were reported in a systematic way dated by the onset of symptoms rather than detection, as the incubation period has been well characterized.

Second, the model captures the number of new cases each day as a proportion from the pool of infected people in the same area. It does not take into account how the age distribution in each area affects that each infection leads to hospitalizations or death, nor how the probability of infecting depends on the days from infection as in (Flaxman et al. 2020). Nor do it take into account mobility between regions, whose impact is believed to be more important at the initiation of the epidemics. It is also worth noting that in locked down environment where most contacts are compartmentalized in households, it is possible that at beginning the virus continues to spread rapidly within households, but less so between households. As immunity was not taken care here, R_0 may decrease significantly more without further policies, after this first wave of within-household contamination is over. Finally, we only modeled impact of lockdown, not of other measures which were taken too close apart (banning public events, closing schools, etc.), and simultaneously in most regions, so it is not possible here to disentangle their effects precisely.

In conclusion, the greatest interest should be focused on the trends in R_0 found in this work, while concrete numerical predictions should be taken with extreme caution, as there is a lack of reliable epidemiological data we can base the models on.

Appendix A: Parameter estimation procedure

Parameters of the model θ include the prior mean of R_0 for the three different lockdown conditions $(\beta_0, \beta_1, \beta_2)$, as well the expected value of the log-infected population at initial point $x_i(r)$. Parameters were fitted from the data by Maximum Likelihood estimation, using an Expectation-Maximization procedure (Bishop 2006). The procedure also allowed to recover the posterior distribution of true cases $p(I|y; \theta)$ for each day and CCAA. We convert the infected population to the log-scale, defining $x(r, d) = \log I(r, d)$:

$$x(r, d+1) = x(r, d) - (R_0(r, d) - 1)\gamma$$

This can be turned into:

$$x(r, d) = x_i(r) + \gamma\left(\sum_{t<d} R_0(r, d) - d\right)$$

In vectorial terms, we have $x = T[x(0, r), R_0(r, :)]$, where T is an upper triangular matrix of 1 that implements the summing operation. Since both $x_i(r)$ and $R_0(r, :)$ have multivariate normal prior distribution, the prior over x is normal itself with mean $\mu_r = T[x_i(r), \Phi\beta]$ and covariance $K_x = T\widetilde{K}T'$ where Φ is an D-by-3 indicator matrix indicating the lockdown state for each day, and $\widetilde{K}$ is block diagonal with submatrices σ_0^2 and K. In the Expectation step, we estimate the posterior distribution over log-infected population using a Laplace approximation $p(x|y, \theta) \approx N(m, V)$. We first identified the maximum-a-posteriori variable m through gradient search, and then computed V as the inverse of the negative of the hessian joint-log-probability evaluated at m.

Parameters were updated in the M-step by maximizing the objective function analytically. Confidence intervals for parameters were estimated using parametric bootstrapping using 20 bootstraps. All analyses were implemented in Matlab with custom codes, which will be uploaded on a public repository upon publication of the manuscript.

Appendix B: Projection analyses

We performed three projection analyses: two projective analyses for period April 16th –
May 15th and one retrospective analysis for the period March 14th-April 15th. For the
projective analysis, we simulated the model with a R0 whose prior mean was set to the
estimated value either for the state of emergency or for mandatory confinement. We found
that we could better predict the death patterns up to April 15th by sampling infection-to-
death duration from a log-normal distribution with median 15 days and dispersion
parameter 0.5 rather than the one previously, so we used this distribution to predict the
patterns of deaths from the number of simulated cases. We ran 100 different simulations
with the same parameters and averaged the results of death over simulations.

We applied a similar approach to predict retrospectively the number of deaths if the state
of emergency would not have been declared, by applying a mean R0 for the whole period
until April 15th equal to our estimate for our pre-state of emergency period. However, we
simulated a full SIR model, since in this case we could no longer approximate the
Susceptible population as being nearly constant in time.

Acknowledgements

The authors thank J. Barbosa and P. Puig for useful comments on the manuscript.

References

[1] Anderson, Roy M., Hans Heesterbeek, Don Klinkenberg, and T. Déirdre Hollingsworth.
2020. "How will country-based mitigation measures influence the course of the COVID-19
epidemic?" Lancet Publishing Group. https://doi.org/10.1016/S0140-6736(20)30567-5.

[2] Bishop, Christopher M. 2006. *Pattern Recognition and Machine Learning*.

[3] Ferguson, Neil M, Daniel Laydon, Gemma Nedjati-Gilani, Natsuko Imai, Kylie Ainslie,
Marc Baguelin, Sangeeta Bhatia, et al. 2020. "Impact of non-pharmaceutical interventions
(NPIs) to reduce COVID-19 mortality and healthcare demand." *Imperial.ac.uk*, 3–20.
https://doi.org/10.25561/77482.

[4] Fernández-Fontelo, A, A Cabaña, H Joe, and others. 2019. "Untangling Serially Dependent Underreported Count Data for Gender-Based Violence." *Statistics in Medicine*. https://doi.org/https://doi.org/10.1002/sim.8306.

[5] Fernández-Fontelo, A, A Cabaña, P Puig, and D Moriña. 2016. "Under-Reported Data Analysis with Inar-Hidden Markov Chains." *Statistics in Medicine*. https://doi.org/https://doi.org/10.1002/sim.7026.

[6] Flaxman, S, S Mishra, A Gandy, and others. 2020. "Estimating the Number of Infections and the Impact of Non-Pharmaceutical Interventions on Covid-19 in 11 European Countries." *Imperial College Preprint*. https://doi.org/https://doi.org/10.25561/77731.

[7] Hellewell, Joel, Sam Abbott, Amy Gimma, Nikos I. Bosse, Christopher I. Jarvis, Timothy W. Russell, James D. Munday, et al. 2020. "Feasibility of controlling COVID-19 outbreaks by isolation of cases and contacts." *The Lancet Global Health* 8 (4): e488–e496. https://doi.org/10.1016/S2214-109X(20)30074-7.

[8] Huang, L, X Zhang, X Zhang, and others. 2020. "Rapid Asymptomatic Transmission of Covid-19 During the Incubation Period Demonstrating Strong Infectivity in a Cluster of Youngsters Aged 16-23 Years Outside Wuhan and Characteristics of Young Patients with Covid-19: A Prospective Contact-Tracing Study." *Journal of Infection*. https://doi.org/https://doi.org/10.1016/j.jinf.2020.03.006.

[9] Lai, Shengjie, Nick W Ruktanonchai, Liangcai Zhou, Olivia Prosper, Wei Luo, Jessica R Floyd, Amy Wesolowski, et al. 2020. "Effect of non-pharmaceutical interventions for containing the COVID-19 outbreak in China." *medRxiv*, March, 2020.03.03.20029843. https://doi.org/10.1101/2020.03.03.20029843.

[10] Lauer, Stephen A, Kyra H Grantz, Qifang Bi, Forrest K Jones, Qulu Zheng, Hannah R Meredith, Andrew S Azman, Nicholas G Reich, and Justin Lessler. 2020. "The Incubation Period of Coronavirus Disease 2019 (COVID-19) From Publicly Reported Confirmed Cases: Estimation and Application." *Annals of Internal Medicine*, March. https://doi.org/10.7326/M20-0504.

[11] Lin, Qianying, Shi Zhao, Daozhou Gao, Yijun Lou, Shu Yang, Salihu S. Musa, Maggie H. Wang, et al. 2020. "A conceptual model for the coronavirus disease 2019 (COVID-19) outbreak in Wuhan, China with individual reaction and governmental action." *International Journal of Infectious Diseases* 93 (April): 211–16. https://doi.org/10.1016/j.ijid.2020.02.058.

[12] Niud, Yan, and Fujie Xu. 2020. "Deciphering the power of isolation in controlling COVID-19 outbreaks." Elsevier Ltd. https://doi.org/10.1016/S2214-109X(20)30085-1.

[13] Team, IHME COVID-19 health service utilization forecasting, and Christopher JL Murray. 2020. "Forecasting COVID-19 impact on hospital bed-days, ICU-days, ventilator-days and deaths by US state in the next 4 months." *medRxiv*, March, 2020.03.27.20043752. https://doi.org/10.1101/2020.03.27.20043752.

[14] Zhao, Shi, Qianyin Lin, Jinjun Ran, Salihu S. Musa, Guangpu Yang, Weiming Wang, Yijun Lou, et al. 2020. "Preliminary Estimation of the Basic Reproduction Number of Novel Coronavirus (2019-nCoV) in China, from 2019 to 2020: A Data-Driven Analysis in the Early Phase of the Outbreak." *International Journal of Infectious Diseases* 92: 214–17. https://doi.org/https://doi.org/10.1016/j.ijid.2020.01.050.

[15] Zhou, Fei, Ting Yu, Ronghui Du, Guohui Fan, Ying Liu, Zhibo Liu, Jie Xiang, et al. 2020. "Clinical course and risk factors for mortality of adult inpatients with COVID-19 in Wuhan, China: a retrospective cohort study." *The Lancet* 395 (10229): 1054–62. https://doi.org/10.1016/S0140-6736(20)30566-3.