# Predicting mortality due to SARS-CoV-2: A mechanistic score relating obesity and diabetes to COVID-19 outcomes in Mexico

Omar Yaxmehen Bello-Chavolla[1,2*], Jessica Paola Bahena-López[3], Neftali Eduardo Antonio-Villa[1,3], Arsenio Vargas-Vázquez[1,3], Armando González-Díaz[4], Alejandro Márquez-Salinas[2,3], Carlos A. Fermín-Martínez[1,3], J. Jesús Naveja[5], Carlos A. Aguilar-Salinas[1,6,7]

[1]Unidad de Investigación de Enfermedades Metabólicas, Instituto Nacional de Ciencias Médicas y Nutrición Salvador Zubirán. [2]Division of Research, Instituto Nacional de Geriatría. [3]MD/PhD (PECEM), Faculty of Medicine, National Autonomous University of Mexico. [4]Universidad Nacional Autónoma de México. [5]Instituto de Química, Universidad Nacional Autónoma de México. [6]Department of Endocrinolgy and Metabolism. Instituto Nacional de Ciencias Médicas y Nutrición Salvador Zubirán. [7]Tecnologico de Monterrey, Escuela de Medicina y Ciencias de la Salud.

**Short title:** COVID-19 mortality, diabetes and obesity in Mexico

**Corresponding authors**

Omar Yaxmehen Bello-Chavolla. Division of Research. Instituto Nacional Geriatría. Anillo Perif. 2767, San Jerónimo Lídice, La Magdalena Contreras, 10200, Mexico City, Mexico. Phone: +52 (55) 5548486885. E-mail: oyaxbell@yahoo.com.mx

Carlos A. Aguilar-Salinas. Unidad de Investigación de Enfermedades Metabólicas. Instituto Nacional de Ciencias Médicas y Nutrición Salvador Zubirán. Vasco de Quiroga 15. CP 14080; Tlalpan, Distrito Federal, México. Phone: +52(55)54870900, 5703. E-mail: caguilarsalinas@yahoo.com

**CONFLICT OF INTERESTS:** Nothing to disclose.

**FUNDING:** This research received no specific funding

**ABSTRACT (250 WORDS)**

**BACKGROUND:** The SARS-CoV-2 outbreak poses challenge to healthcare systems due to

high complication rates in patients with cardiometabolic diseases. Here, we identify risk

factors and propose a clinical score to predict COVID-19 lethality, including specific factors

for diabetes and obesity and its role in improving risk prediction.

**METHODS:** We obtained data of confirmed and negative COVID-19 cases and their

demographic and health characteristics from the General Directorate of Epidemiology of

Mexican Ministry of Health. We investigated specific risk factors associated to COVID-19

positivity and mortality and explored the impact of diabetes and obesity on modifying COVID-

19 related lethality. Finally, we built a clinical score to predict COVID-19 lethality.

**RESULTS:** Among 71,103 subjects at April 27[th], 2020, we observed 15,529 subjects with

SARS-CoV-2 and 1,434 deaths. Risk factors for lethality in COVID-19 includes early-onset

diabetes obesity, COPD, advanced age, immunosuppression, and CKD; we observed that

obesity mediates 45.5% of the effect of diabetes on COVID-19 lethality. Early-onset diabetes

conferred an increased risk of hospitalization and obesity conferred an increased risk for ICU

admission and intubation. Our predictive score for COVID-19 lethality included age ≥65 years,

diabetes, early-onset diabetes, obesity, age <40 years, CKD, hypertension, pregnancy and

immunosuppression and significantly discriminates lethal from non-lethal COVID-19 cases (c-

statistic=0.830).

**CONCLUSIONS:** Here, we propose a mechanistic approach to evaluate risk for

complications and lethality attributable to COVID-19 considering the effect of obesity and

diabetes in Mexico. Our score offers a clinical tool for quick determination of high-risk

susceptibility patients in a first contact scenario.

**Keywords:** COVID-19; SARS-CoV-2; Diabetes; Obesity, Lethality; Mexico

**INTRODUCTION**

The first cases of SARS-CoV-2 infection in Mexico were reported at the end of February [1]; since then, the number of COVID-19 cases have been steadily increasing, with most fatal cases being associated with the presence of comorbidity and, particularly, cardiometabolic comorbidities. A high prevalence of cardiometabolic diseases worldwide represents a challenge during the COVID-19 epidemic. An elevated number of patients with SARS-CoV-2 infection have a preexisting disease such as obesity, hypertension, cardiovascular disease, diabetes, chronic respiratory disease or cancer [2,3]. Diabetes mellitus and obesity represent a large share of the cardiometabolic morbidity burden of the region [4]; moreover, most cases of diabetes remain either undiagnosed or lack adequate glycemic control, posing them at risk of increased COVID-19 severity. Despite several reports evaluating the burden of comorbidities including obesity, diabetes and hypertension on the clinical course of COVID-19, the joint role of obesity and diabetes in modifying COVID-19 outcomes has not been fully explored [5].

Several studies have demonstrated a higher susceptibility to acute respiratory infectious diseases in people with diabetes [6]. Moreover, diabetes and obesity have been described as independent risk factors for severe pulmonary infection [7,8]. Obesity influences the clinical outcomes during an acute severe respiratory distress syndrome (ASRDS); it has been proposed as a protective factor for mortality following lung injury (due to reverse causality) or as a cause of mortality and adverse clinical outcomes for severe influenza cases (due to mechanical and immunologic factor). In the cases of the COVID-19 outbreak, obesity has been consistently associated with adverse outcomes [9,10]. Furthermore, a large proportion of obesity cases in Mexico lives in geographical areas of increased social vulnerability, which poses a structural inequality that might also increase mortality for COVID-19 associated to both diabetes and obesity [11]. Chronic inflammation in obesity might worsen the acute inflammatory response triggered by SARS-Cov2 infection, which might be associated to a

cytokine release syndrome [5,12]. Here, we investigate the role of both diabetes and obesity

in determining propensity for SARS-CoV-2 infection and its associated clinical outcomes

including disease severity and COVID-19 lethality; using these associations, we further

construct a clinically useful predictive model for COVID-19 mortality using national

epidemiological surveillance data from Mexico.

**METHODS**

*Data sources*

We extracted this dataset from the General Directorate of Epidemiology of the Mexican

Ministry of Heath, which is an open-source dataset comprising daily updated data of

suspected COVID-19 cases which have been confirmed with a positive test for SARS-CoV-2

certified by the National Institute for Diagnosis and Epidemiological Referral [13].

*Definitions of suspected and confirmed COVID-19 cases*

The Ministry of Health defines a suspected COVID-19 case as an individual of any age whom

in the last 7 days has presented cough, fever or headache (at least two), accompanied by

either dyspnea, arthralgias, myalgias, sore throat, rhinorrhea, conjunctivitis or chest pain.

Amongst these suspected cases, the Ministry of Health establishes two protocols for case

confirmation: 1) SARS-CoV-2 testing is done widespread for suspected COVID-19 cases with

severe acute respiratory infection with signs of breathing difficulty or deaths in suspected

COVID-19 cases, 2) for all other suspected cases, a sentinel surveillance model is being

utilized, whereby 475 health facilities comprise a nationally representative sample which

samples ~10% of mild outpatient cases and all suspected severe acute respiratory infection

[14]. Demographic and health data are collected and uploaded to the epidemiologic

surveillance database by personnel from the corresponding individual facility.

*Variables definitions*

Available information for all confirmed, negative and suspected COVID-19 cases includes

age, sex, nationality, state and municipality where the case was detected, immigration status

as well as identification of individuals who speak indigenous languages from Mexico. Health

information includes status of diabetes, obesity, chronic obstructive pulmonary disease

(COPD), immunosuppression, pregnancy, arterial hypertension, cardiovascular disease,

chronic kidney disease (CKD) and asthma. Date of symptom onset, hospital admission and

death are available for all cases as well as treatment status (outpatient or hospitalized),

information regarding diagnosis of pneumonia, ICU admission and whether the patient

required invasive mechanical ventilation.

*Statistical analysis*

*Comorbidities associated to SARS-CoV-2 positivity*

We investigated the association of demographic and health data associated with SARS-CoV-

2 positivity using logistic regression analyses, excluding individuals who were only suspected

but unconfirmed cases of COVID-19. Next, we stratified these analyses for individuals with

only diabetes or only obesity to identify specific risk factors within these populations,

especially focusing on individuals who were <40 years and likely acquired the disease early.

*COVID-19 mortality risk*

In order to investigate risk factors predictive of COVID-19 related 30-day lethality, we fitted

Cox Proportional risk regression models estimating time from symptom onset up to death or

censoring, whichever occurred first in cases with confirmed positivity for SARS-CoV-2. To

identify diabetes and obesity-specific risk factors, we carried out stratified analyses. Given the

availability of SARS-CoV2 negative cases within the dataset, we fitted Cox models for 30-day

mortality which included SARS-CoV-2 positivity as an interaction term with different

comorbidities, hypothesizing that some factors increase mortality risk specifically for COVID-

19. Finally, we fitted a logistic regression model only for mortality cases to evaluate

associations with lethality rates in COVID-19 related and non-related deaths.

*Influence of obesity and diabetes in COVID-19 related outcomes*

Finally, we estimated factors associated to admission to hospital facilities, intensive care units

(ICUs) and requirements for mechanical ventilation in all confirmed COVID-19 cases using

logistic regression. To identify specific factors for all COVID-19 and non-COVID-19 patients

related to outcomes, we included interaction effects with comorbidities factors; we also

performed Kaplan-Meier analyses to identify the role of comorbidities in modifying lethality

risk in individuals with diabetes and obesity and compared across categories using Breslow-

Cox tests. Finally, we performed causal-mediation analyses with causally-ordered mediators

using a previously validated approach to investigate whether obesity mediates the decreases

COVID-19 survival attributable to diabetes, particularly in early-onset cases <40 years [15].

*Mechanistic mortality risk score for COVID-19*

Finally, we constructed a clinically useful model to predict 30-day mortality in COVID-19

cases that might be useful to apply in primary care facilities, including variables and

interactions which were identified in mortality analyses. Points were assigned by

standardizing all $\beta$ coefficients with the minimum absolute $\beta$ coefficient obtained from Cox

regression. Points were stratified according to categories of Low risk ($\leq$0), Mild risk (1-3),

Moderate risk (4-6), High risk (7-9) and very high risk ($\geq$10). Risk across categories was

verifies using Kaplan-Meier analyses. C-statistics and $D_{xy}$ values were corrected for over-

optimism using k-fold cross-validation (k=10) using the *rms* R package. A p-value <0·05 was

considered as statistical significance threshold. All analyses were performed using R software

version 3.6.2.

**RESULTS**

*COVID-19 cases in Mexico*

At the time of writing this report (April 27[th], 2020), a total of 71,103 subjects had been treated

initially as suspected COVID-19 cases. Amongst them, 15,529 had been confirmed with as

positive and 46,079 tested negatives for SARS-CoV-2 infection; additionally, 8,477 cases

were still being studied as suspected cases pending testing reports. Amongst confirmed

cases, 1,434 deaths were reported (9.2%) whilst 881 deaths were SARS-CoV-2 negative cases (1.9%) and 137 deaths of suspected but unconfirmed cases (1.6%) had been reported. Compared to SARS-CoV-2 negative cases, confirmed cases were older, predominantly male (1.35:1 ratio), had higher rates of hospitalization and showed a higher prevalence of diabetes, hypertension and obesity, but less CKD. SARS-CoV-2 cases were also more likely to be treated as inpatients and had higher rates of ICU admission and requirements for invasive ventilation compared to negative cases (**Table 1**).

*Factors associated with COVID-19 positivity*

We investigated cases related to COVID-19 positivity within Mexico. We found that risk of SARS-CoV-2 positivity was higher with diabetes, hypertension, COPD, immunosuppression, CKD, obesity, age >65 and male sex (**Supplementary Figure 1**). When assessing age, we observed reduced odds of SARS-CoV-2 positivity in patients younger than age 40 and, in contrast, when exploring its interaction with diabetes, we observed an increased probability of SARS-CoV-2 infection. In stratified models, we observed that for patients with diabetes, SARS-CoV-2 positivity was associated with obesity, male sex and an interaction effect was observed between obesity and having less than 40 years; for patients with obesity, diabetes and male sex were also significant.

*Predictors for COVID-19 related 30-day mortality*

We identified that COVID-19 cases were associated with a near four-fold increase in mortality due to acute respiratory infection (HR 3.864, 95%CI 3.501-4.264) compared to non-COVID-19 cases. Of interest, the only comorbidity which conferred increased lethality risk exclusively for COVID-19 compared to non-COVID-19 was obesity (HR 1.567, 95%CI 1.273-1.928, **Figure 1A**). Factors associated to increased lethality in COVID-19 cases were age >65 years, diabetes mellitus, obesity, CKD, COPD, immunosuppression and hypertension (**Figure 1B**). We searched for an interaction between diabetes mellitus and age <40 years adjusted

for sex and obesity; a higher mortality risk was found for early onset diabetes cases (HR

2.764, 95%CI 1.718-4.447).

*COVID-19 in patients with diabetes mellitus*

Confirmed COVID-19 cases with diabetes had a mean age of 56·9 (±12·9) years and were

predominantly male. This population had particularly higher mortality rate (10.0% vs. 1.6%),

hospitalization and confirmed pneumonia compared with those without diabetes. When

stratifying mortality for those with early onset diabetes (<40 years) had higher mortality rates

compared to individuals <40 years without diabetes (17.1% vs. 4.6%); similarly, those aged

>65 years had lower rates compared to those <65 years with diabetes (23.8% vs. 29.7%). As

expected, obesity, hypertension, COPD, CKD, CVD, immunosuppression and asthma were

also more prevalent in this population. Interestingly, we observed no significant differences in

ICU admission and requirements for mechanical ventilation between subjects with and

without diabetes (**Supplementary table 1**).  In patients with diabetes mellitus, COVID-19

related mortality was higher in those with concomitant immunosuppression, COPD, CKD and

those aged >65 years (**Figure 1C)**. When assessing the role of comorbidities, COVID-19

patients with diabetes, coexistent obesity, those with early onset diabetes (<40 years) or an

increase in the number of comorbidities had increased risk of COVID-19 lethality (Breslow

test p<0.001, **Figure 2**).  When comparing cases with and without COVID-19 amongst

patients with diabetes mellitus, only obesity displayed a significant interaction with COVID-19

(HR 1.449 95%CI 1.071-1.960) with diabetes after adjustment for age, sex and comorbidities

(**Supplementary Figure 2**).

*COVID-19 in patients with obesity*

Similar to patients with diabetes, confirmed COVID-19 cases with obesity had particularly

higher rates of mortality (13.6% vs. 8.17%), hospitalization and confirmed pneumonia.

Furthermore, patients with obesity also had higher rates of ICU admission (13.2% vs. 10.5%)

and were more likely to be intubated (13.1% vs. 10.3%). As expected, diabetes,

hypertension, COPD, smoking, CVD and asthma were also more prevalent in patients with

obesity (**Supplementary table 2**). As previously mentioned, obesity was identified as a risk

factor which displayed differential risk for COVID-19 infection, specific risk factors for lethality

in obese patients with COVID-19 infection immunosuppression and age >65 (**Figure 1D**);

overall, COVID-19 increased risk of mortality in obesity nearly six-fold (HR 5.954, 95%CI

4.939-7.178). The addition of obesity to any number of comorbidities significantly increased

the risk for COVID-19 lethality (Breslow test p<0.001, **Figure 2-C**). Using causally ordered

mediation analysis we investigated whether the effect of diabetes (E) on COVID-19 related

lethality was partially mediated by obesity (M); the direct effect of diabetes on COVID-19

lethality was significant ($\Delta_{E1 \to Y}$=1.72, 95%CI 1.54-1.93) as was the indirect effect of obesity

($\Delta_{M1 \to Y}$ =1.44, 95%CI: 1.32-1.58), representing 45.5% of the total effect of diabetes.

*COVID-19 outcomes and comorbidities*

Given the increased risk of diabetes and obesity in modifying COVID-19 related lethality, a

secondary objective of our study was to investigate its associations with inpatient outcomes,

including hospitalization rate, ICU admission and requirement for mechanical ventilation. In

general, early-onset diabetes patients and those with obesity had higher risk of

hospitalization, whilst patients with obesity also had increased risk for ICU admission and

required intubation. Patients with diabetes mellitus overall had higher risk of hospitalization

with no significant additional risk of ICU admission and intubation (**Figure 3**).

*Mechanistic score for mortality in COVID-19*

Using the identified predictors for mortality and the observed interaction for early-onset

diabetes, we designed a predictive score for COVID-19 mortality using Cox regression using

a random split of 80% of the dataset stratified my mortality (n=12,424, deaths=1,137). We

identified as significant predictors age>65 years, diabetes mellitus, obesity, CKD, COVID-19

related pneumonia, COPD, pregnancy and immunosuppression (**Table 2**); age<40 was a

protective factor which was modified by its interaction with T2D ($R^2$=0.160, C-statistic 0.828,

$D_{xy}$=0.656); assigning the point system did not significantly reduce the model's performance

($R^2$=0.159, C-statistic=0.829, $D_{xy}$=0.657). Finally, category stratification reduced only

moderately performance statistics ($R^2$=0.157, C-statistic=0.816, $D_{xy}$=0.631), and were not

significantly modified after cross-validation correction ($R^2$=0.182, $D_{xy}$=0.657). The score was

then validated using the remaining 20% of the population (n=3,105, deaths=297); we

observed that the score retained its predictive and discriminative ability ($R^2$=0.141, C-

statistic=0.801, $D_{xy}$=0.602) as did the categories ($R^2$=0.144, C-statistic=0.792, $D_{xy}$=0.585).

Distribution of the score significantly discriminates between lethal and non-lethal COVID-19

cases (**Figure 4**).

**DISCUSSION**

Our results demonstrate that diabetes, particularly early onset diabetes mellitus, obesity and

comorbidity burden modify risk profiles in patients with COVID-19 infection in Mexico and

significantly improve mortality prediction related to COVID-19 lethality. These findings

position the notion that early-onset type 2 diabetes might carry a higher risk of mortality in

younger patients, which is similar to older patients with comorbidities and only higher in older

patients with diabetes. Furthermore, our results suggest that obesity is a COVID-19 specific

risk factor for mortality, risk of ICU admission, tracheal intubation and hospitalization and

even increases risk in patients with diabetes and COVID-19 infection. Overall, this positions

the co-existence of obesity and diabetes, particularly early-onset diabetes, as a considerable

risk factor for COVID-19 mortality in Mexicans, whom have reported an alarmingly high

burden of both conditions in recent health surveys.

The relationship between increased risk of mortality attributable to acute severe respiratory

infections in patients with diabetes mellitus has been extensively reported, particularly for the

acute respiratory syndrome caused by SARS-CoV-1 [16–18]. Evidence relating SARS-CoV-2

infections in China demonstrated increased rates of diabetes mellitus in hospitalized patients

and in those with increased disease severity as assessed by ICU admission and requirement

for invasive ventilation. Additionally, hospitalized patients have shown increased rates of both

obesity and diabetes for COVID-19 compared to non-hospitalized cases in the US, China and

Italy [5,19,20]. Increased susceptibility for COVID-19 in patients with diabetes may be

explained for several potential mechanisms including an increased lung ACE2 expression

and elevated circulating levels of furin, a protease involved in viral entry to cells, and a

decreased clearance of SARS-CoV-2 viral particles in subjects with diabetes and/or

hypertension associated with ACE2 expression [21–24]. Impairments in immunity observed in

patients with diabetes are characterized by initial delay in activation of Th1 cell-mediated

immunity and late hyper-inflammatory response and are consistent with the increased risk

associated with additional immunosuppression observed with our data [25]. Additional factors

which have been proposed to modify COVID-19 mortality risk and worsen glycemic control in

diabetes include corticosteroid therapy, inadequate glucose monitoring, the effect of social

distancing on diabetes care and the use of antihypertensive medication; however these

factors remain to be confirmed by clinical evidence [26]. Given the large proportion of

undiagnosed diabetes cases in Mexican and poor glycemic control reported by recent

estimates, the burden of COVID-19 might be higher than expected in Mexico and poses a

challenge for the Mexican healthcare system  to give particular attention to this sector as the

epidemic moves forward [27–29].

Diabetes mellitus is one of the main causes of morbidity and it accounts for a large proportion

of mortality risk in Mexican population [4]. Of relevance, Mexicans have increased risk of

diabetes and diabetes-related obesity attributable to genetic variants associated to its

Amerindian ancestry, and an earlier age of onset independent of body-mass index [30,31].

Data on the incidence of early-onset type 2 diabetes in Mexican population position obesity

and insulin resistance as significant risk factors, which are also highly prevalent in younger

patients and increase metabolic risk [32,33]. These associations partly explain the increased

risk of COVID-19 lethality in younger patients within our cohort despite the younger average

age of Mexican population and poses early-onset diabetes mellitus as a significant risk factor

for COVID-19 mortality and increased severity of infection in younger patients [5,34].

In our work, we demonstrate that compared to non-COVID-19 infections, obesity significantly

modifies the risk of mortality attributable to COVID-19 infection. Obesity and, in particular,

abdominal obesity, is one of Mexico's main public health problems; in recent years the socio-

economic burden of obesity as well as its impact on mortality have increased drastically, with

the Ministry of Health declaring a state of epidemiological emergency [35]. Evidence from

different regions has supported the notion that obesity increases mortality risk and severity of

COVID-19 infections, which holds particularly true for younger patients [2,36]. Obesity is

characterized by a low-grade state of inflammation and may impair immune response which

in turn also may affect the lung parenchyma, increasing the risk for inflammatory lung

diseases [37,38]. In particular, abdominal obesity reduces the compliance of lung, chest wall

and entire respiratory system, resulting in impaired ventilation of the base of the lungs and

reduced oxygen saturation of blood [39]. Recently, Simonnet et al. explored the high

prevalence of obesity in patients with COVID-19 reporting that obesity is a risk factor for

SARS-CoV-2 infection severity independent of age, diabetes and hypertension. Notably,

ACE2 expression in adipose tissue is higher than in the lung and its expression profile is not

different in obese and non-obese subjects; however, obese subjects have more adipocytes;

thus, they have a greater number of ACE2-expressing cells and thus higher likelihood of

SARS-CoV-2 entry [2,40]. Our data shows that obesity is a specific risk factor for COVID-19

related outcomes and that it partly mediates the risk associated with diabetes mellitus.  Public

health efforts by the Mexican government in epidemiological surveillance have largely

focused in identifying patient's at highest risk of complications; these findings could inform

public health decisions and increase awareness on the role of obesity in modifying risk of

COVID-18 outcomes.

Our study had some strengths and limitations. First, we analyzed a large dataset which

included information on both confirmed positive and negative SARS-CoV-2 cases, which

provides a unique opportunity to investigate COVID-19 specific risk factors and develop a

predictive model for COVID-19 mortality. Additionally, with the database being nationally-

representative it allows for reasonable estimates on the impact of both diabetes and obesity

despite the possibility of important regional differences in cardiometabolic risk which might

influence risk estimates. A potential limitation of this study is the use of data collected from a

sentinel surveillance system model, which is skewed towards investigating high risk cases or

only those with specific risk factors which on one hand increases power to detect the effect of

comorbidities and on the other hand might not be representative of milder cases of the

disease; this is demonstrated in the risk of COVID-19 positivity, which is higher for high risk

cases. The updating daily estimates of COVID-19 cases are unlikely to change the direction

of the identified associations though it might modify numeric estimates. Implementation of our

proposed model might be useful to allocate prompt responses to high risk cases and improve

stratification of disease severity.

In conclusion, we show that both diabetes and obesity increase the risk of SARS-CoV-2

infection in Mexico. In particular, diabetes increases the risk of COVID-19 related mortality

and, specifically, increases mortality risk in early-onset cases. Obesity is a COVID-19 specific

risk factor for mortality and for increased disease severity; obesity also is a partial mediator

on the effect of diabetes in decreasing survival associated with COVID-19 infection. This

mechanistic interpretation on the risk of comorbidities allowed the development of a model

with good performance to predict mortality in COVID-19 cases. Given the burden of obesity

and diabetes in Mexico, COVID-19 lethality might be higher in younger cases. Special

attention should be given to susceptible individuals and screening should be conducted for all

symptomatic cases with either obesity and/or diabetes to decrease the burden associated

with COVID-19 in Mexico.

**Conflict of interest:** Nothing to disclose.

**Role of the funding source:** This research received no funding.

**ACKNOWLEDGMENTS**

JPBL, AVV, NEAV, AMS and CAFM are enrolled at the PECEM program of the Faculty of

Medicine at UNAM. JPBL and AVV are supported by CONACyT. The authors would like to

acknowledge the invaluable work of all of Mexico's healthcare community in managing the

COVID-19 epidemic. Its participation in the COVID-19 surveillance program has made this

work a reality, we are thankful for your effort.

**DATA AVAILABILITY**

All data sources and R code are available for reproducibility of results at

https://github.com/oyaxbell/covid_diabetesmx.

**AUTHOR CONTRIBUTIONS**

Research idea and study design OYBC, JPBL, CAAS; data acquisition: OYBC, AGD; data

analysis/interpretation: OYBC, JPBL, NEAV, AVV, AGD, JJN, CAAS; statistical analysis:

OYBC, NEAV; manuscript drafting: OYBC, NEAV, AVV, JPBL, AMS, CAFM, JJN; supervision

or mentorship: OYBC, CAAS. Each author contributed important intellectual content during

manuscript drafting or revision and accepts accountability for the overall work by ensuring

that questions pertaining to the accuracy or integrity of any portion of the work are

appropriately investigated and resolved.

**FUNDING:** No funding was received.

**CONFLICT OF INTEREST/FINANCIAL DISCLOSURE:** Nothing to disclose.

**REFERENCES**

1.    Información Internacional y Nacional sobre nuevo Coronavirus (COVID-2019) |

Secretaría de Salud | Gobierno | gob.mx [Internet]. [cited 19 Apr 2020]. Available:

https://www.gob.mx/salud/documentos/informacion-internacional-y-nacional-sobre-

nuevo-coronavirus-2019-ncov

2.  Simonnet A, Chetboun M, Poissy J, Raverdy V, Noulette J, Duhamel A, et al. High

prevalence of obesity in severe acute respiratory syndrome coronavirus-2 (SARS-CoV-

2) requiring invasive mechanical ventilation. Obesity (Silver Spring). 2020;

doi:10.1002/oby.22831

3.  Zhou F, Yu T, Du R, Fan G, Liu Y, Liu Z, et al. Clinical course and risk factors for

mortality of adult inpatients with COVID-19 in Wuhan, China: a retrospective cohort

study. Lancet. 2020;395: 1054–1062. doi:10.1016/S0140-6736(20)30566-3

4.  Bello-Chavolla OY, Rojas-Martinez R, Aguilar-Salinas CA, Hernández-Avila M.

Epidemiology of diabetes mellitus in Mexico. Nutr Rev. 2017;75: 4–12.

doi:10.1093/nutrit/nuw030

5.  Grasselli G, Zangrillo A, Zanella A, Antonelli M, Cabrini L, Castelli A, et al. Baseline

Characteristics and Outcomes of 1591 Patients Infected With SARS-CoV-2 Admitted to

ICUs of the Lombardy Region, Italy. JAMA. 2020; doi:10.1001/jama.2020.5394

6.  Muller LMAJ, Gorter KJ, Hak E, Goudzwaard WL, Schellevis FG, Hoepelman AIM, et al.

Increased risk of common infections in patients with type 1 and type 2 diabetes mellitus.

Clin Infect Dis. 2005;41: 281–288. doi:10.1086/431587

7.  Van Kerkhove MD, Vandemaele KAH, Shinde V, Jaramillo-Gutierrez G, Koukounari A,

Donnelly CA, et al. Risk factors for severe outcomes following 2009 influenza A (H1N1)

infection: a global pooled analysis. PLoS Med. 2011;8: e1001053.

doi:10.1371/journal.pmed.1001053

8.  Luzi L, Radaelli MG. Influenza and obesity: its odd relationship and the lessons for

COVID-19 pandemic. Acta Diabetol. 2020; doi:10.1007/s00592-020-01522-8

9.  Jose RJ, Manuel A. Does COVID-19 Disprove the Obesity Paradox in ARDS? Obesity

(Silver Spring). 2020; doi:10.1002/oby.22835

10. Dietz W, Santos-Burgoa C. Obesity and its Implications for COVID-19 Mortality. Obesity

(Silver Spring). 2020; doi:10.1002/oby.22818

384 11. Shamah-Levy T, Campos-Nonato I, Cuevas-Nasu L, Hernández-Barrera L, Morales-

385   Ruán MDC, Rivera-Dommarco J, et al. [Overweight and obesity in Mexican vulnerable

386   population. Results of Ensanut 100k]. Salud Publica Mex. 2019;61: 852–865.

387   doi:10.21149/10585

388 12. Moore BJB, June CH. Cytokine release syndrome in severe COVID-19. Science. 2020;

389   eabb8925. doi:10.1126/science.abb8925

390 13. Datos Abiertos - Dirección General de Epidemiología | Secretaría de Salud | Gobierno |

391   gob.mx [Internet]. [cited 18 Apr 2020]. Available:

392   https://www.gob.mx/salud/documentos/datos-abiertos-152127?idiom=es

393 14. Secretaría de Salud Subsecretaría de Prevención y Promoción de la Salud Dirección

394   General de Epidemiología. LINEAMIENTOESTANDARIZADO PARA LA VIGILANCIA

395   EPIDEMIOLÓGICA Y POR LABORATORIO DE LAENFERMEDAD RESPIRATORIA

396   VIRAL. Apr 1, 2020.

397 15. Cho S-H, Huang Y-T. Mediation analysis with causally ordered mediators using Cox

398   proportional hazards model. Stat Med. 2019;38: 1566–1581. doi:10.1002/sim.8058

399 16. Yang JK, Feng Y, Yuan MY, Yuan SY, Fu HJ, Wu BY, et al. Plasma glucose levels and

400   diabetes are independent predictors for mortality and morbidity in patients with SARS.

401   Diabet Med. 2006;23: 623–628. doi:10.1111/j.1464-5491.2006.01861.x

402 17. Bello-Chavolla OY, Bahena-Lopez JP, Garciadiego-Fosass P, Volkow P, Garcia-Horton

403   A, Velazquez-Acosta C, et al. Bloodstream infection caused by S. aureus in patients

404   with cancer: a 10-year longitudinal single-center study. Support Care Cancer. 2018;26:

405   4057–4065. doi:10.1007/s00520-018-4275-1

406 18. Akbar DH. Bacterial pneumonia: comparison between diabetics and non-diabetics. Acta

407   Diabetol. 2001;38: 77–82. doi:10.1007/s005920170017

408 19. CDC COVID-19 Response Team. Preliminary Estimates of the Prevalence of Selected

409   Underlying Health Conditions Among Patients with Coronavirus Disease 2019 - United

States, February 12-March 28, 2020. MMWR Morb Mortal Wkly Rep. 2020;69: 382–

386. doi:10.15585/mmwr.mm6913e2

20.    Liang W-H, Guan W-J, Li C-C, Li Y-M, Liang H-R, Zhao Y, et al. Clinical characteristics

and outcomes of hospitalised patients with COVID-19 treated in Hubei (epicenter) and

outside Hubei (non-epicenter): A Nationwide Analysis of China. Eur Respir J. 2020;

doi:10.1183/13993003.00562-2020

21.    Chen X, Hu W, Ling J, Mo P, Zhang Y, Jiang Q, et al. Hypertension and Diabetes Delay

the Viral Clearance in COVID-19 Patients. medRxiv. 2020;

doi:10.1101/2020.03.22.20040774

22.    Rao S, Lau A, So H-C. Exploring diseases/traits and blood proteins causally related to

expression of ACE2, the putative receptor of 2019-nCov: A Mendelian Randomization

analysis. medRxiv. 2020; doi:10.1101/2020.03.04.20031237

23.    Fernandez C, Rysä J, Almgren P, Nilsson J, Engström G, Orho-Melander M, et al.

Plasma levels of the proprotein convertase furin and incidence of diabetes and

mortality. J Intern Med. 2018;284: 377–387. doi:10.1111/joim.12783

24.    Wrapp D, Wang N, Corbett KS, Goldsmith JA, Hsieh C-L, Abiona O, et al. Cryo-EM

structure of the 2019-nCoV spike in the prefusion conformation. Science. 2020;367:

1260–1263. doi:10.1126/science.abb2507

25.    Hodgson K, Morris J, Bridson T, Govan B, Rush C, Ketheesan N. Immunological

mechanisms contributing to the double burden of diabetes and intracellular bacterial

infections. Immunology. 2015;144: 171–185. doi:10.1111/imm.12394

26.    Klonoff DC, Umpierrez GE. COVID-19 in patients with diabetes: risk factors that

increase morbidity. Metab Clin Exp. 2020; 154224. doi:10.1016/j.metabol.2020.154224

27.    Singh AK, Gupta R, Ghosh A, Misra A. Diabetes in COVID-19: Prevalence,

pathophysiology, prognosis and practical considerations. Diabetes Metab Syndr.

2020;14: 303–310. doi:10.1016/j.dsx.2020.04.004

28.  Campos-Nonato I, Ramírez-Villalobos M, Flores-Coria A, Valdez A, Monterrubio-Flores

E. Prevalence of previously diagnosed diabetes and glycemic control strategies in

Mexican adults: ENSANUT-2016. PLoS One. 2020;15: e0230752.

doi:10.1371/journal.pone.0230752

29.  Puig-Domingo M, Marazuela M, Giustina A. COVID-19 and endocrine diseases. A

statement from the European Society of Endocrinology. Endocrine. 2020;

doi:10.1007/s12020-020-02294-5

30.  SIGMA Type 2 Diabetes Consortium, Williams AL, Jacobs SBR, Moreno-Macías H,

Huerta-Chagoya A, Churchhouse C, et al. Sequence variants in SLC16A11 are a

common risk factor for type 2 diabetes in Mexico. Nature. 2014;506: 97–101.

doi:10.1038/nature12828

31.  Almeda-Valdes P, Gómez Velasco DV, Arellano Campos O, Bello-Chavolla OY, Del

Rocío Sevilla-González M, Viveros Ruiz T, et al. The SLC16A11 risk haplotype is

associated with decreased insulin action, higher transaminases and large-size

adipocytes. Eur J Endocrinol. 2019;180: 99–107. doi:10.1530/EJE-18-0677

32.  Arellano-Campos O, Gómez-Velasco DV, Bello-Chavolla OY, Cruz-Bautista I,

Melgarejo-Hernandez MA, Muñoz-Hernandez L, et al. Development and validation of a

predictive model for incident type 2 diabetes in middle-aged Mexican adults: the

metabolic syndrome cohort. BMC Endocr Disord. 2019;19: 41. doi:10.1186/s12902-

019-0361-8

33.  Aguilar-Salinas CA, Rojas R, Gómez-Pérez FJ, García E, Valles V, Ríos-Torres JM, et

al. Prevalence and characteristics of early-onset type 2 diabetes in Mexico. Am J Med.

2002;113: 569–574. doi:10.1016/s0002-9343(02)01314-1

34.  Jiménez-Corona A, Rojas R, Gómez-Pérez FJ, Aguilar-Salinas CA. Early-onset type 2

diabetes in a Mexican survey: results from the National Health and Nutrition Survey

2006. Salud Publica Mex. 2010;52 Suppl 1: S27–35. doi:10.1590/s0036-

36342010000700006

35.  Nieto C, Tolentino-Mayo L, Monterrubio-Flores E, Medina C, Patiño SR-G, Aguirre-

Hernández R, et al. Nutrition Label Use Is Related to Chronic Conditions among

Mexicans: Data from the Mexican National Health and Nutrition Survey 2016. J Acad

Nutr Diet. 2019; doi:10.1016/j.jand.2019.07.016

36.  Lighter J, Phillips M, Hochman S, Sterling S, Johnson D, Francois F, et al. Obesity in

patients younger than 60 years is a risk factor for Covid-19 hospital admission. Clin

Infect Dis. 2020; doi:10.1093/cid/ciaa415

37.  Huttunen R, Syrjänen J. Obesity and the risk and outcome of infection. Int J Obes.

2013;37: 333–340. doi:10.1038/ijo.2012.62

38.  Zhang X, Zheng J, Zhang L, Liu Y, Chen GP, Zhang HP, et al. Systemic inflammation

mediates the detrimental effects of obesity on asthma control. Allergy Asthma Proc.

2017; doi:10.2500/aap.2017.38.4096

39.  Dixon AE, Peters U. The effect of obesity on lung function. Expert Rev Respir Med.

2018;12: 755–767. doi:10.1080/17476348.2018.1506331

40.  Jia X, Yin C, Lu S, Chen Y, Liu Q, Bai J, et al. Two Things about COVID-19 Might Need

Attention. 2020; doi:10.20944/preprints202002.0315.v1

**Table 1.** Descriptive statistics comparing negative, positive and suspected cases for SARS-CoV-2 in Mexico at 04/27/2020.

*Abbreviations*: ICU= intense care unit; COPD= chronic obstuctive pulmonary disease; CKD= chronic kidney disease; CVD=

cardiovascular disease**.**

| Parameter | Positive for SARS-CoV-2 n=15529 | Negative for SARS-CoV-2 n=46960 | Suspected for SARS-CoV-2 n=8614 |
|---|---|---|---|
| Age (mean ± sd) | 46.55±15.51 | 39.56±17.9 | 43.22±17.27 |
| Male sex (%) | 8977(57.8) | 21955(46.8) | 4664(54.1) |
| Mortality (%) | 1434(9.2) | 881(1.9) | 137(1.6) |
| Hospitalization (%) | 6042(38.9) | 9974(21.2) | 2607(30.3) |
| Pneumonia (%) | 4588(29.5) | 6341(13.5) | 1841(21.4) |
| ICU admission (%) | 676(4.4) | 774(1.6) | 211(2.4) |
| Invasive ventilation (%) | 669(4.3) | 598(1.3) | 197(2.3) |
| Diabetes (%) | 2831(18.2) | 5163(11) | 1323(15.4) |
| COPD (%) | 389(2.5) | 1330(2.8) | 214(2.5) |
| Asthma (%) | 542(3.5) | 2571(5.5) | 337(3.9) |
| Immunosuppression (%) | 296(1.9) | 1402(3) | 183(2.1) |
| Hypertension (%) | 3370(21.7) | 7353(15.7) | 1611(18.7) |

| Parameter | Positive for SARS-CoV-2 n=15529 | Negative for SARS-CoV-2 n=46960 | Suspected for SARS-CoV-2 n=8614 |
|---|---|---|---|
| Obesity (%) | 3215(20.7) | 6570(14) | 1532(17.8) |
| CKD (%) | 359(2.3) | 1130(2.4) | 262(3) |
| CVD (%) | 453(2.9) | 1610(3.4) | 267(3.1) |
| Smoking (%) | 1374(8.8) | 4835(10.3) | 811(9.4) |

485 **Table 2.** Cox proportional risk models for lethality using the mechanistic COVID-19 lethality score in confirmed cases of COVID-19

486 using individual components, single score point and risk stratification categories. Abbreviations:  COPD= chronic obstructive

487 pulmonary disease; CKD= chronic kidney disease; CVD= cardiovascular disease.

| Model | Parameter | B | Score | SE | Wald | HR | 95% CI | P Value |
|---|---|---|---|---|---|---|---|---|
| | Age ≥65 years | 0.753 | 3 | 0.065 | 11.568 | 2.12 | 1.87-2.41 | <0.001 |
| | Pregnancy | 0.963 | 4 | 0.451 | 2.133 | 2.62 | 1.08-6.35 | 0.033 |
| | Diabetes | 0.251 | 1 | 0.066 | 3.802 | 1.29 | 1.13-1.46 | <0.001 |
| Individual variables *C-statistic= 0.828* | Diabetes*Age <40 years | 1.092 | 4 | 0.265 | 4.119 | 2.98 | 1.77-5.01 | <0.001 |
| | Age <40 years | -1.397 | -6 | 0.136 | -10.283 | 0.25 | 0.19-0.32 | <0.001 |
| | Obesity | 0.374 | 1 | 0.066 | 5.680 | 1.45 | 1.28-1.65 | <0.001 |
| | Pneumonia | 1.697 | 7 | 0.072 | 23.444 | 5.46 | 4.74-6.29 | <0.001 |

| | | Beta | Points | SE | Wald | OR | 95% CI | p-value |
|---|---|---|---|---|---|---|---|---|
| | CKD | 0.568 | 2 | 0.120 | 4.179 | 1.76 | 1.39-2.23 | <0.001 |
| | COPD | 0.277 | 1 | 0.117 | 2.361 | 1.32 | 1.05-1.66 | 0.018 |
| | Immunosuppression | 0.500 | 2 | 0.137 | 3.365 | 1.65 | 1.26-2.16 | <0.001 |
| Score point training *C-statistic= 0.829* | 1-Point increment | 0.246 | | 0.007 | 36.15 | 1.28 | 1.26-1.30 | <0.001 |
| | Low-Risk (0 pts) | | | | Reference | | | |
| Risk Categories training *C-statistic= 0.816* | Mild-Risk (1-3 pts) | 1.873 | | 0.136 | 13.73 | 6.51 | 4.98-8.50 | <0.001 |
| | Moderate-Risk (4-6 pts) | 2.727 | | 0.131 | 20.84 | 15.28 | 11.83-19.75 | <0.001 |
| | High-Risk (7-9 pts) | 3.287 | | 0.125 | 26.38 | 26.76 | 20.96-34.16 | <0.001 |
| | Very High-Risk (≥10 pts) | 3.824 | | 0.131 | 29.07 | 45.78 | 35.38-59.25 | <0.001 |

 **FIGURE LEGENDS**

**Figure 1.** Cox proportional risk regression analysis to evaluate lethality of SARS-CoV-2 in Mexico, compared to SARS-CoV2 negative

cases for all suspected cases with SARS-CoV2 status available (A) and stratified by diabetes mellitus (B) and obesity (C).

*Abbreviations*: ICU= intense care unit; COPD= chronic obstructive pulmonary disease; CKD= chronic kidney disease; CVD=

cardiovascular disease, HR= Hazard ratio.

**Figure 2.** Kaplan-Meier survival curves to evaluate lethality of SARS-CoV-2 positivity in patients with diabetes and comorbidities (A),

499    diabetes and obesity (B), obesity and comorbidities (C) and diabetes with age 40 years (D). Abbreviations: DM= Diabetes

500    mellitus; OB= Obesity; Comorb= Comorbidities.

**Figure 3.** Logistic regression analyses to evaluate COVID-19 related outcomes in all patients with SARS-CoV2 positivity for

admission to ICU (A), mechanical ventilation (B) and hospital admission risk (C).

**Figure 4.** Symptom onset among lethal and non-lethal cases in new-confirmed COVID 19 cases, stratified by diabetes and obesity status (A), density histogram of scores of the mechanistic COVID-19 score (B). Points and score intervals considered for clinical score scale (C) and Kaplan-Meir Survival analysis curves to evaluate lethality using risk categories in the training (D) and validation cohorts (E). Abbreviations: OB= Obesity; DM= Diabetes mellitus; CKD= chronic kidney disease; COPD= Chronic obstructive pulmonary disease.