Indexing Cerebrovascular Health Using Transcranial Doppler Ultrasound

R. Ghorbani Afkhami[1], R. Wong[2], S. Ramadan[3], F. R. Walker[2], S. J. Johnson[1]

[1] School of Electrical Engineering & Computing, University of Newcastle, Australia

[2] School of Biomedical Sciences and Pharmacy, University of Newcastle, Australia

[3] School of Health Sciences, University of Newcastle, Australia

Author for correspondence

Mr. Rashid Ghorbani Afkhami, School of Electrical Engineering & Computing, University of Newcastle, Australia

Email: rashid.ghorbaniafkhami@uon.edu.au

Abstract

Transcranial Doppler (TCD) blood flow velocity has been extensively used in biomedical research as it provides a cost-effective and relatively simple approach to assess changes in cerebral blood flow dynamics and track cerebrovascular health status. In this paper, we introduce a new TCD based timing index, TI_{TCD}, as an indicator of vascular stiffening and vascular health. We use two different datasets, D1 and D2 with 38 and 55 subjects, respectively to investigate the correlations of the new index and the existing indices, namely the pulsatility index (PI_{TCD}) and the augmentation index (AI_{TCD}), with age, cardiorespiratory fitness (CRF) and magnetic resonance imaging (MRI) blood flow pulsatility index (PI_{MRI}). Comparing the correlation results, AI_{TCD} and TI_{TCD} both showed strong correlations with age ($r_{D1} = 0.71$, $r_{D2} = 0.58$ and $r_{D1} = 0.69$, $r_{D2} = 0.50$ respectively). While, PI_{TCD} had a significant correlation with age only in one of the datasets ($r_{D2} = 0.41$). The new TI_{TCD} index also showed better correlations with CRF ($r_{D1} = -0.77$) and PI_{MRI} ($r_{D1} = 0.50$) compared to AI_{TCD} ($r_{D1} = -0.65$ and $r_{D1} = 0.33$, respectively) and PI_{TCD} (no significant correlations with CRF or PI_{MRI}). The proposed index has a much clearer relationship with vascular aging factors such as stiffness and cerebrovascular resistance (CVR) and therefore had the strongest correlations with CRF.

1 Introduction

Changes in vascular stiffness is an important indicator for vascular health independent of blood pressure [1] and the American Heart Association recommends the clinical measurement of vascular stiffness to predict future cardiovascular disease risk [1]. Vessel compliance describes the process of localized distension of the vascular wall in response to the pulsatile delivery of blood driven by the contraction of the heart. A stiff vessel is one which has lost its compliance. In a healthy vascular system, the compliant vessels act to smooth out the pressure waveform coming from the heart and reduce the peak pressure experienced by more distal vessels. Conversely, a loss of compliance (i.e. increased stiffening), results in increased pressure due to less efficient smoothing of the pressure waveform. This increase in pressure can compound other causes of elevated pressure such as vascular resistance caused by vessel narrowing.

As vessels lose their compliance, the speed of the pressure wave, called its pulse wave velocity (PWV), increases. PWV is measured by the time taken for the pressure wave, caused by the ejection of blood from the heart, to move down a given length of vessel. At a local level, when the pressure wave reaches a vessel bifurcation, a part of the wave is reflected towards the heart. Due to its fast speed the reflected wave can arrive back sufficiently close in time to the original wave front to augment the peak pressure exerted on vessels proximal to the recombination point. As a result of this process, small arteries, arterioles and capillaries face higher pressure and flow pulsations in each cardiac cycle. Over time these higher pressures can result in tearing of their endothelial and smooth muscle cells, producing a condition known as called small vessel disease (SVD) [2, 3]. SVD is especially important in the brain leading to cognitive decline, dementia, age-related disabilities and stroke [4].

Due to the well documented association between changes in compliance and serious brain pathology, there has been considerable interest research into techniques that measure changes in cerebrovascular compliance. Currently, the most common methodology to assess changes in the cerebrovascular compliance is Transcranial Doppler (TCD) ultrasound. TCD is a non-

invasive technique that has been preferred because it is considered to be safe, cost-effective and relatively fast to perform. TCD is based on the "Doppler effect" according to which the ultrasound waves reflected back from the moving blood cells within the vessels have a different frequency from the emitted waves and this difference in frequency is directly proportional to the speed of those blood cells [5]. TCD is recognized to have excellent temporal resolution, making it ideal to capture information on blood flow dynamics.

Importantly, TCD can not measure compliance directly. However, several indirect indices have been developed including the pulsatility index (PI) and the augmentation index (AI) which are considered to provide reasonable approximations. The pulsatility index is the most commonly reported indirect measure of vascular health in the literature. PI_{TCD}, first defined by Gosling and King in 1974, computes the peak to peak height of the flow velocity waveform divided by the mean flow velocity [6]. Consistent with the idea that PI_{TCD} indexes vascular health, it has been shown to correlate robustly with aging [20–22] and aortic PWV [21]. Studies have reported correlations between PI_{TCD} and white matter disease [10,19], diabetes mellitus [23] and dementia [9].

There have been different interpretations of which particular vessel properties influence PI_{TCD} measurements. Some have proposed that PI_{TCD} measures cerebrovascular resistance (CRV) at the reflection site [7–11], while other studies have reported inconsistent results [12–15]. Following the discovery of vascular compliance, it has been hypothesized that as the vessels lose their compliance and become stiffer the peak-to-peak flow velocity increase also elevates calculated PI_{TCD} values [19]. More recently PI_{TCD} has been modeled as combination of resistance, compliance, PWV, and heart rate [20]. Consequently, PI_{TCD} is now often reported as an index of cerebrovascular health in general.

The second commonly used index of compliance is the augmentation index. AI was initially defined for the pressure waveform [24] and has been correlated with vascular aging and aortic PWV [25]. AI captures the reflected wave augmentation of the peak pressure by calculating the ratio of the reflected wave hight to the systolic wave height with higher values

said to result from earlier arrival of the reflected wave due to stiffer arteries. The AI index has been applied to TCD flow velocity waveforms in several studies [26, 27] and positive and strong correlation between middle cerebral arterial (MCA) AI_{TCD} and age has been reported, $r = 0.88$ ($n = 25$) [28]. However, several studies have questioned the reliability of AI as an indicator of vascular compliance [29–31]. Studies have shown that AI also correlates with numerous biological factors not directly related to vessel properties including sex, heart rate, food intake, hydration status, height, weight and body composition [30].

An alternative TCD index to PI and AI that may provide a more direct measure of vessel health is to use a time-based index which directly measures the timing of the reflected wave. Evidence in an analysis of pressure waves suggests that a timing index can more accurately reflect vascular properties as it has a clearer relationship with vascular resistance, vascular compliance and PWV [32] and may be less susceptible to being influenced by biological factors that are independent of the vessel properties. In this paper we define a TCD timing index, TI_{TCD}, as the inverse of the time between the systolic peak and the reflected waveform peak in the TCD signal. After introducing the new timing index, we investigate the correlation of the three TCD indices, PI, AI and TI, with other accepted measures of vascular aging such as magnetic resonance imaging (MRI) pulsatility index, age and cardiorespiratory fitness (CRF).

2 Methods

Data from two different sets of experiments have been used in this study. Experiment one is a new study including Doppler, MRI and CRF whereas the second experiment re-analysis Doppler recordings in existing data.

2.1 Experiment 1

Data from 38 adult volunteers (23 female and 14 male, age range = 24-66 years, mean age = 41.7 years) was recorded, which we will refer to as dataset D1. Participants were recruited from the local Novocastrian community, and signed an informed consent prior to any assessment. The study protocol is approved by the University of Newcastle Human Research Ethics Committee and registered in the Australian New Zealand Clinical Trials Registry (ACTRN12619000144112). Height, weight, age, gender and resting heart rate were recorded and the participants were asked to complete a physical activity questionnaire. Data acquisition was carried out over two imaging sessions on two consecutive days for each participant. Participants were asked to refrain from caffeine consumption before their scans. However, it was not an exclusion criteria in the study.

TCD ultrasound (DopplerBox X; Compumedics DWL, Singen, Germany) was used to record cerebral blood flow velocity from the right and left middle cerebral arteries. The participants wore a headpiece with bilateral Doppler probes which stayed in place for the 300 seconds of resting-state recording. In this session, the resting-state heart rate was measured using OMRON HEM-7320 blood pressure monitor device. After five minutes of sitting during the headpiece setup, heart rate was measured three times every one minute while the TCD recordings were taking place. The three measurements were averaged later and used as the resting-state heart rate.

The participants were also scanned on a 3T MRI scanner (Magnetom Prisma, Siemens Healthineers, Erlangen, Germany), equipped with 64-channel receive only head coil, while a standard built in dual channel body coil was used for RF transmission. Blood flow was quantified using phase contrast flow quantification sequence (TR = 26.5ms, TE = 6.9ms, slice thickness = 5mm, matrix 256×256). A single excitation with the velocity encoding value 120 cm/s was used to quantify blood flow in middle cerebral arteries [34, p. 161-175]. The TCD and MRI scan sessions were held on two consecutive days.

Two subjects did not have usable Doppler recordings and were excluded from further

Figure 1: Two different samples from D1. (a) Averaged MCA, subject #22, age 25 years (b) Averaged MCA, subject #17, age 66 years

analysis. A further three did not have have MRI data and were excluded from the Doppler versus MRI comparison.

2.2 Experiment 2

This experiment was carried out using previously recorded TCD experiments examining cognitive performance with partial data being published previously in [35, 36]. The aim of using this dataset, which we will refer to as D2, is to uncover the relationship of TCD indices with age in a population with wider age range including older participants as consistently lower correlations have been reported for PI_{TCD} in narrower and younger population age ranges [21–23, 37–39]. D2 has 55 subjects (34 female and 21 male, age range = 21-80 years, mean age = 45.8 years), 3 subjects were excluded from the initial data as peak detection was not possible due to noise levels.

2.3 Calculated Indices

PI$_{\text{TCD}}$: Indices were calculated for both experiments. PI$_{\text{TCD}}$ is a device output with values averaged per subject. The device calculates PI$_{\text{TCD}}$ as defined by [6] i.e.,

$$\text{PI}_{\text{TCD}} = \frac{V_{\max} - V_{\min}}{\overline{V}}, \tag{1}$$

where $V_{\max}$, $V_{\min}$ and $\overline{V}$ are maximum, minimum and mean flow velocities, respectively. An alternative definition, PI$^{*}_{\text{TCD}}$, uses $V_{\text{sys}} - V_{\text{dia}}$ in the numerator where V_{sys} and V_{dia} are peak systolic and end-diastolic velocities, respectively. The two definitions report the same value in most cases, the difference arises when V_{sys} and $V_{\max}$ have different values , see Fig. 1b for an example. This is the case in older populations and in stiffer arteries where the forward and the reflected waves meet early and V_{sys} hides on an inflection point whereas $V_{\max}$ becomes the reflected wave peak, V_{refl}, see Fig. 1b. Nonetheless, we compared PI$_{\text{TCD}}$ values given by the device with manually calculated PI$^{*}_{\text{TCD}}$ values for our experimental data and there was no notable difference. Therefore, only PI$_{\text{TCD}}$ values directly calculated by the device are reported in this paper.

AI$_{\text{TCD}}$: In order to calculate the augmentation index, first end-diastolic points were extracted automatically using Matlab by finding the minimum points of each waveform and then the signal was averaged giving us a single heartbeat per vessel of measurement. Next, the systolic peak, V_{sys}, and reflected peak, V_{refl}, were located manually on the waveform and AI$_{\text{TCD}}$ was calculated as

$$\text{AI}_{\text{TCD}} = \frac{V_{\text{refl}} - V_{\text{dia}}}{V_{\text{sys}} - V_{\text{dia}}}. \tag{2}$$

AI$_{\text{TCD}}$ values were averaged across both MCAs, providing a single number per subject.

PI$_{\text{MRI}}$: The MRI pulsatility index, is defined similar to the TCD pulsatility index as [40, 41]:

$$\text{PI}_{\text{MRI}} = \frac{F_{\max} - F_{\min}}{\overline{F}}, \tag{3}$$

where F is the blood flow (in ml/sec) and the bar sign indicates the mean value. MRI scanner software (Siemens Syngo) was used to quantify flow values by placing a region of interest around the middle cerebral artery. The flow values for every repetition time, TR, produced a waveform for the length of the heart beat for which the maximum, minimum and the mean values were used to calculate PI_{MRI} following the method in [40]. From $n = 33$ subjects of D1, PI_{MRI} was calculated either for both MCAs, $n = 20$, or a single MCA, $n = 13$, based on data quality. All the indices from right and left MCA were averaged for each subject.

Cardiorespiratory Fitness (CRF): CRF was estimated based on a non-exercise method developed by [42], using information such as age, gender, body mass index, resting state heart rate and physical activity score. See the *supplementary materials* for physical activity questionnaire and CRF calculation.

Proposed Index: The proposed index relies on the information from both forward and reflected waves, however, unlike AI which is amplitude based, the new index only uses the time information. Here, we define the TCD reflection time, T_{refl} as the time difference between the occurrence of peak systole and the reflected shoulder of peak systole, in cases of augmented waveforms, and the time difference between two distinct positive peaks, in cases of separated forward and reflected waves, see Fig. 1. Then, we define TCD reflection index, TI_{TCD}, as

$$TI_{TCD} = \frac{1}{T_{refl}}. \tag{4}$$

So that TI_{TCD} will be an index of vascular stiffness. TI_{TCD} was calculated for both MCAs using custom Matlab software and then the values were averaged for each subject.

2.4 Statistical Analysis

Matlab software was used to calculate p-values for testing the hypothesis of no correlation against the alternative, non-zero correlation case. Bonnferroni corrections with initial $\alpha = 0.05$ were made for multiple comparisons, resulting in the adjusted level of significance set

at 0.0056 for experiment one and 0.0167 for experiment two.

3 Results

The correlation matrix for experiment one and experiment two are shown in Table 1. Significant correlations have been marked after adjustment for multiple comparisons.

In experiment one, PI_{TCD} did not have any significant correlation with age, CRF or PI_{MRI} and the observed weak correlation with age was negative. AI_{TCD} and TI_{TCD} both correlated significantly with age. Although the correlation coefficients of AI_{TCD} and TI_{TCD} are similar with age, TI_{TCD} has higher correlation coefficient with CRF. Among the three Doppler indices only TI_{TCD} has significant correlation with PI_{MRI}.

In experiment two, all three Doppler indices, PI_{TCD}, AI_{TCD} and TI_{TCD} correlated significantly with age. The correlation coefficient was higher for AI_{TCD} and TI_{TCD} compared to PI_{TCD}.

4 Discussion

PI_{TCD} is the most common and easy-to-calculate Doppler index. To understand the controversy in the interpretation of PI_{TCD}, it is useful to consider a mathematical model developed in [20] extending the single-element model in [43] to a 3-element Windkessel which describes

	Age	PI_{MRI}	CRF
PI_{TCD}	$-0.13(0.45)$	$0.06(0.76)$	$0.10(0.55)$
AI_{TCD}	$0.71^*(0.11e{-}5)$	$0.33(0.057)$	$-0.65^*(0.15e{-}4)$
TI_{TCD}	$0.69^*(0.38e{-}5)$	$0.50^*(0.003)$	$-0.77^*(0.32e{-}7)$

	Age
PI_{TCD}	$0.41^*(0.0019)$
AI_{TCD}	$0.58^*(0.12e{-}4)$
TI_{TCD}	$0.50^*(0.15e{-}3)$

Table 1: Correlation matrix for experiment one (left) and experiment two (right) reported as $r(p\text{-value})$. Correlations found to be significant after Bonferroni correction for multiple comparisons are marked by $*$.

PI_{TCD} as,

$$PI_{TCD} = \frac{p_{HR}}{CPP} \times \sqrt{1 + (2\pi f_{HR} \times CVR \times C)^2}, \tag{5}$$

where PI_{TCD} is defined as the fundamental harmonic of the flow velocity divided by the mean flow velocity. p_{HR} is the fundamental harmonic of the arterial pulse pressure, f_{HR} is the heart rate, C is the cerebrovascular compliance and CPP is the cerebral perfusion pressure, the driving source of blood inside the cerebrum [20,44]. The main point to consider in (5) is that although CVR and C affect PI_{TCD} positively, they each change in different directions with age i.e., CVR increases whereas C decreases. For younger adults these effects may cancel out each other hence the lower correlation between PI_{TCD} and both age ($r_{D1} = -0.13$) and CRF ($r_{D1} = 0.10$). However, for older adults the increased stiffness, or decreased C, may have a dominant effect resulting in a strong decrease in PI_{TCD} even though CVR is increasing. This can explain the low correlations of PI_{TCD} with age for younger adults reported in the literature [38,39]. Our results confirm this for D1, which has a relatively young population, where the correlation between PI_{TCD} and age is negative ($r_{D1} = -0.13$, insignificant). For D2 ($r_{D2} = 0.41$), PI_{TCD} experiences a bigger increase after the age of 65, as shown in the *supplementary materials.*

PI_{MRI} is a different predictor of vascular aging than PI_{TCD} as it considers blood volume changes rather than blood velocity changes. The compliance is the change in blood volume relative to a change in pressure. MRI which measures blood flow, i.e., volume per second, more accurately tracks volume changes than does TCD. If vascular cross-sectional area was known, or if it was constant, it would be possible to derive blood flow from blood velocity. However, changes in vascular cross-section area can not assume to be constant as it is actually these changes which derive compliance. In fact, a subtle assumption in deriving (5) is that the vascular cross-sectional area is constant, which is particularly problematic in younger population with compliant arteries. Thus, the uncorrelated PI_{TCD} and PI_{MRI} results ($r_{D1} = 0.06$, insignificant) can be explained by the younger population of D1 and also the lack of cross-sectional area information in PI_{TCD}.

AI is an index strongly related to vascular aging factors [33]. Assuming a transmission line with a three-element Windkessel at the load, AI in the pressure waveform can be formulated as [33]

$$\text{AI} \approx \frac{2}{3}\left(\cos\left(2\pi f_{\text{HR}} T_{\text{refl}}\right) + |\Gamma(0)| + \frac{1}{2}\right), \tag{6}$$

where $\Gamma(0)$ is called the reflection coefficient and equals the amplitude of the reflected wave divided by the amplitude of the forward wave, measured at the load site [33]. As seen in (6), AI includes T_{refl} and the reflection coefficient, both are reported to have significant changes with age [32, 45]. For AI calculated using the Doppler waveform these factors are assumed to contributed in the same way as we have showed a high correlation of AI_{TCD} with age ($r_{\text{D1}} = 0.71$ and $r_{\text{D2}} = 0.58$), CRF ($r_{\text{D1}} = -0.65$) and PI_{MRI} ($r_{\text{D1}} = 0.33$). However, heart rate is another deciding factor in AI_{TCD} (see (6)) for which the dependency has also been reported in practice [46]. This is a factor which might change AI independent of vascular compliance or resistance [29].

The timing based index, TI has a strong relationship with vascular compliance. In the context of the pressure waveform, evidence of change in the time of reflected waves with age have been presented practically [32] and theoretically [33]. As initially suggested in [47] and then mathematically modeled in [33], the time difference between the forward and the backward traveling pressure waves i.e., T_{refl}, can be given as

$$T_{\text{refl}} \approx \underbrace{\frac{2Z_0 \times \text{CVR} \times C}{\text{CVR} + 2Z_0}}_{\Delta t_{\text{load}}} + \underbrace{\frac{2d_0}{\text{PWV}}}_{\Delta t_{\text{line}}}, \tag{7}$$

where Δt_{line} is the time delay between the two waves caused by the length of the arteries and Δt_{load} is the delay at the reflection site due to the compliance of the vessels beyond that point and Z_0 is the characteristic impedance. The line delay is inversely proportional to the pulse wave velocity i.e., $\Delta t_{\text{line}} \propto 1/\text{PWV}$. In addition, the load delay is proportional to the compliance of the vascular bed i.e., $\Delta t_{\text{load}} \propto C$, and will also decrease as the compliance

Figure 2: Simulated flow velocity signal; blue, green and red dashed lines represent the incident forward traveling wave, the re-reflected wave and the reflected wave, respectively. Solid line is the sum of the three waveforms i.e., the recorded waveform. t_{sys}, t_{refl}, t_{i}, t_{r} and t_{rr} correspond respectively to observed systolic, observed reflected, incident, reflected and re-reflected peak times.

of the vessels distal to the measurement site reduces with aging. Therefore, similarly we would expect TI_{TCD} to be highly dependent on vascular compliance. The high correlations of TI_{TCD} with age ($r_{\text{D1}} = 0.69$ and $r_{\text{D2}} = 0.50$), CRF ($r_{\text{D1}} = -0.77$) and with PI_{MRI} ($r_{\text{D1}} = 0.50$) support this assumption. TI_{TCD} correlates better with CRF than age and by looking specifically at older subjects with high CRF (see the *supplementary materials*) we hypothesize that the timing index is a predictor of vascular health independent of age, while AI more closely predicts age than CRF. Moreover, the relationship of T_{refl} with age is not linear. T_{refl} and consequently TI_{TCD} do not show a considerable change after the age of 65 years which is consistent with results reported for timing metrics taken from the pressure waveform [48, 49].

In general, the application of pressure waveform indices to the Doppler waveform must be done with care due to the differences between the two waveforms. While we often think of a single reflected wave, in reality there are many reflected and re-reflected waveforms adding to form an observed signal. In order to investigate the influence of these factors a simulated flow waveform is shown in Fig. 2 (a simpler form of the ones reported in

the literature [50]) which illustrates the underlying reflected and re-reflected waveforms. Reflected flow velocity, i.e., blood which travels towards the heart has a negative amplitude and thus is subtracted from the forward waves. The re-reflected flow, however, reflected twice during the cardiac cycle and travels in the same direction as the original forward wave. This waveform has a positive amplitude when its velocity is read by the Doppler device. Note that both reflected and re-reflected waveforms have positive amplitudes in pressure readings. Therefore, Doppler waves will have a different shape to pressure waves and the use of the Doppler indices imported from pressure studies must be done with care. However, based on Fig. 2, we can see that an observed reflected peak is more influenced by the re-reflected wave. In other words, the reflected wave plays part in forming a negative dip between the observed forward and reflected peaks and V_{refl} is mainly determined by the re-reflected wave. Further, in practice, comparison of simultaneously recorded pressure and flow velocity waveforms has shown that the characteristic time points used in the definitions of AI and TI are similar in both waveforms [51]. This suggests that in both pressure and flow velocity waveforms the reflection time, the pulsatility index and augmentation index account for not just the reflected wave but a combination of reflected and re-reflected waves similarly in both technologies. It also suggests that the stiffness / compliance measured through the existing, and the proposed indices, may be equally effected by the arteries proximal to the heart (from the point the measurement) rather than just the more distal cerebral arteries. Given the similarity of timing of the second peak for pressure and flow velocity waveforms we have continued to use this second peak for our derivation of both the AI and TI indices in our calculations. Nonetheless, to our knowledge, there has been no TCD timing index to investigate the effects of the negative reflected wave in the flow velocity waveform and this could be considered in future work.

5 Conclusion

The most commonly used TCD index, PI_{TCD}, should be interpreted with caution due to the complex relationships with CVR and vascular stiffness / compliance, each having opposite effects on this index resulting in a weaker correlation between PI_{TCD} and age and no correlation with cardiorespiratory fitness. On the other hand, AI_{TCD} and the new proposed index, TI_{TCD}, both showed strong correlations with age linking the two indices to vascular aging. In addition, TI_{TCD} had the strongest correlation with cardiorespiratory fitness and MRI among the other Doppler indices suggestion that this index may be a suitable replacement for PI_{TCD} and AI_{TCD} when attempting to index cardiovascular health.

References

[1] R. R. Townsend, I. B. Wilkinson, E. L. Schiffrin, A. P. Avolio, J. A. Chirinos, J. R. Cockcroft, K. S. Heffernan, E. G. Lakatta, C. M. McEniery, G. F. Mitchell, S. S. Najjar, W. W. Nichols, E. M. Urbina, and T. Weber, "Recommendations for improving and standardizing vascular research on arterial stiffness," *Hypertension*, vol. 66, no. 3, pp. 698–722, 2015.

[2] M. M. Poels, K. Zaccai, G. C. Verwoert, and et. al., "Arterial stiffness and cerebral small vessel disease," *Stroke*, vol. 43, no. 10, pp. 2637–2642, 2012.

[3] M. F. O'Rourke and M. E. Safar, "Relationship between aortic stiffening and microvascular disease in brain and kidney," *Hypertension*, vol. 46, no. 1, pp. 200–204, 2005.

[4] T. M. De Silva and F. M. Faraci, "Contributions of aging to cerebral small vessel disease," *Annual Review of Physiology*, vol. 82, no. 1, pp. 275–295, 2020.

[5] S. Purkayastha and F. Sorond, "Transcranial doppler ultrasound: Technique and application," *Seminars in Neurology*, vol. 32, no. 4, pp. 411–420, 2012.

[6] R. Gosling and D. King, "Arterial assessment by doppler-shift ultrasound," *Proceedings of The Royal Society of Medicine-London*, vol. 67, no. 6, pp. 447–449, 1974.

[7] J. Legarth and E. Thorup, "Characteristics of doppler blood-velocity waveforms in a cardiovascular in vitro model i the model and the influence of pulse rate," *Scandinavian Journal of Clinical and Laboratory Investigation*, vol. 49, no. 5, p. 451–457, Jan 1989.

[8] L. Legarth and E. Thorup, "Characteristics of doppler blood-velocity waveforms in a cardiovascular in vitro model ii the influence of peripheral resistance, perfusion pressure and blood flow," *Scandinavian Journal of Clinical and Laboratory Investigation*, vol. 49, no. 5, Sep 1989.

[9] M. Altmann, B. Thommessen, O. M. Ronning, J. S. Benth, A. S. Reichenbach, and B. Fure, "Middle cerebral artery pulsatility index is associated with cognitive impairment in lacunar stroke," *Journal of Neuroimaging*, vol. 26, no. 4, pp. 431–435, 2016.

[10] C. S. Kidwell, S. El-Saden, Z. Livshits, N. A. Martin, T. C. Glenn, and J. L. Saver, "Transcranial doppler pulsatility indices as a measure of diffuse small-vessel disease," *Journal of Neuroimaging*, vol. 11, no. 3, pp. 229–235, 7 2001.

[11] M. E. Wagshul, P. K. Eide, and J. R. Madsen, "The pulsating brain: A review of experimental and clinical studies of intracranial pulsatility," *Fluids and Barriers of the CNS*, vol. 8, no. 1, p. 5, Jan 2011.

[12] M. Czosnyka, H. K. Richards, H. E. Whitehouse, and J. D. Pickard, "Relationship between transcranial doppler-determined pulsatility index and cerebrovascular resistance: an experimental study," *Journal of Neurosurgery*, vol. 84, no. 1, pp. 79–84, 1996.

[13] L. A. Lipsitz, S. Mukai, J. Hamner, M. Gagnon, and V. Babikian, "Dynamic regulation of middle cerebral artery blood flow velocity in aging and hypertension," *Stroke*, vol. 31, no. 8, pp. 1897–1903, 2000.

[14] H.-Y. Hsu, C.-M. Chern, J.-S. Kuo, T. B.-J. Kuo, Y.-T. Chen, and H.-H. Hu, "Correlations among critical closing pressure, pulsatility index and cerebrovascular resistance," *Ultrasound in Medicine & Biology*, vol. 30, no. 10, pp. 1329 – 1335, 2004.

[15] L. A. Calviello, F. A. Zeiler, J. Donnelly, A. Uryga, N. [de Riva], P. Smielewski, and M. Czosnyka, "Estimation of pulsatile cerebral arterial blood volume based on transcranial doppler signals," *Medical Engineering & Physics*, vol. 74, pp. 23 – 32, 2019.

[16] N. Westerhof, J.-W. Lankhaar, and B. E. Westerhof, "The arterial windkessel," *Medical and Biological Engineering and Computing*, vol. 47, no. 2, pp. 131–141, 2008.

[17] G. F. Mitchell, L. A. Moyé, E. Braunwald, J.-L. Rouleau, V. Bernstein, E. M. Geltman, G. C. Flaker, M. A. Pfeffer, and for the SAVE Investigators, "Sphygmomanometrically determined pulse pressure is a powerful independent predictor of recurrent events after myocardial infarction in patients with impaired left ventricular function," *Circulation*, vol. 96, no. 12, pp. 4254–4260, 1997.

[18] A. Benetos, A. Rudnichi, M. Safar, and L. Guize, "Pulse pressure and cardiovascular mortality in normotensive and hypertensive subjects," *Hypertension*, vol. 32, no. 3, pp. 560–564, 1998.

[19] A. J. Webb, M. Simoni, S. Mazzucco, W. Kuker, U. Schulz, and P. M. Rothwell, "Increased cerebral arterial pulsatility in patients with leukoaraiosis," *Stroke*, vol. 43, no. 10, pp. 2631–2636, 2012.

[20] N. de Riva, K. P. Budohoski, P. Smielewski, M. Kasprowicz, C. Zweifel, L. A. Steiner, M. Reinhard, N. Fábregas, J. D. Pickard, and M. Czosnyka, "Transcranial doppler pulsatility index: What it is and what it isn't," *Neurocritical Care*, vol. 17, no. 1, pp. 58–66, Aug 2012.

[21] T.-Y. Xu, J. A. Staessen, F.-F. Wei, J. Xu, F.-H. Li, W.-X. Fan, P.-J. Gao, J.-G. Wang, and Y. Li, "Blood Flow Pattern in the Middle Cerebral Artery in Relation to Indices of Arterial Stiffness in the Systemic Circulation," *American Journal of Hypertension*, vol. 25, no. 3, pp. 319–324, 03 2012.

[22] J. Y. Kim, C. D. Bushnell, J. H. Park, S. M. Han, J. H. Im, S. W. Han, J. S. Baik, and J. H. Park, "Central aortic pressure and pulsatility index in acute ischemic stroke," *Journal of Neuroimaging*, vol. 25, no. 3, pp. 438–442, 2015.

[23] Y. Ohta, K. Fujii, H. Arima, K. Matsumura, T. Tsuchihashi, M. Tokumoto, K. Tsuruya, H. Kanai, M. Iwase, H. Hirakata *et al.*, "Increased renal resistive index in atherosclerosis and diabetic nephropathy assessed by doppler sonography," *Journal of hypertension*, vol. 23, no. 10, pp. 1905–1911, 2005.

[24] J. P. Murgo, N. Westerhof, J. P. Giolma, and S. A. Altobelli, "Aortic input impedance in normal man: relationship to pressure wave forms." *Circulation*, vol. 62, no. 1, pp. 105–116, 1980.

[25] K. Kohara, Y. Tabara, A. Oshiumi, Y. Miyawaki, T. Kobayashi, and T. Miki, "Radial augmentation index: A useful and easily obtainable parameter for vascular aging," *American Journal of Hypertension*, vol. 18, no. S1, pp. 11S–14S, 01 2005.

[26] K. Hirata, T. Yaginuma, M. F. O'Rourke, and M. Kawakami, "Age-related changes in carotid artery flow and pressure pulses," *Stroke*, vol. 37, no. 10, pp. 2552–2556, 2006.

[27] G. F. Mitchell, M. A. van Buchem, S. Sigurdsson, J. D. Gotal, M. K. Jonsdottir, O. Kjartansson, M. Garcia, T. Aspelund, T. B. Harris, V. Gudnason, and L. J. Launer, "Arterial stiffness, pressure and flow pulsatility and brain structure and function: the Age, Gene/Environment Susceptibility – Reykjavik Study," *Brain*, vol. 134, no. 11, pp. 3398–3407, 11 2011.

[28] K. Alyakhan, C. Debert, W. A. Whitelaw, J. M. Rawling, R. Frayne, and M. J. Poulin, "Differences between middle cerebral artery blood velocity waveforms of young and postmenopausal women," *The Journal of The North American Menopause Society*, vol. 13, no. 2, pp. 303–313, 2006.

[29] M. Vyas, J. L. Izzo, Y. Lacourcière, J. M. O. Arnold, M. E. Dunlap, J. L. Amato, M. A. Pfeffer, and G. F. Mitchell, "Augmentation Index and Central Aortic Stiffness in Middle-Aged to Elderly Individuals*: ," *American Journal of Hypertension*, vol. 20, no. 6, pp. 642–647, 06 2007.

[30] G. F. Mitchell, H. Parise, E. J. Benjamin, M. G. Larson, M. J. Keyes, J. A. Vita, R. S. Vasan, and D. Levy, "Changes in arterial stiffness and wave reflection with advancing age in healthy men and women," *Hypertension*, vol. 43, no. 6, pp. 1239–1245, 2004.

[31] K. S. Heffeernan, W. K. Lefferts, and J. A. Augustine, "Hemodynamic correlates of late systolic flow velocity augmentation in the carotid artery," *International Journal of Hypertension*, vol. 2013, no. 6, 2013.

[32] A. J. Baksi, T. A. Treibel, J. E. Davies, N. Hadjiloizou, R. A. Foale, K. H. Parker, D. P. Francis, J. Mayet, and A. D. Hughes, "A meta-analysis of the mechanism of blood pressure change with aging," *Journal of the American College of Cardiology*, vol. 54, no. 22, pp. 2087–2092, 2009.

[33] R. Ghorbani Afkhami and S. J. Johnson, "Wave reflection: More than a "round trip"," *medRxiv*, 2020.

[34] J. Ramalho and M. Castillo, *Vascular Imaging of the Central Nervous System: Physical Principles, Clinical Applications, and Emerging Techniques.* Wiley-Blackwell, 2014.

[35] R. Wong, M. Al-Omary, D. Baker, N. Spratt, A. Boyle, N. Baker, P. Howe, and N. Collins, "Cognitive dysfunction is associated with abnormal responses in cerebral blood flow in patients with single ventricular physiology: Novel insights from transcranial doppler ultrasound," *Congenital Heart Disease*, vol. 14, no. 4, pp. 638–644, 2019.

[36] R. S. Nealon, P. R. C. Howe, L. Jansen, M. Garg, and R. H. X. Wong, "Impaired cerebrovascular responsiveness and cognitive performance in adults with type 2 diabetes," *Journal of Diabetes and its Complications*, vol. 31, no. 2, pp. 462 – 467, 2017.

[37] A. Kwater, J. Gąsowski, B. Gryglewska, B. Wizner, and T. Grodzicki, "Is blood flow in the middle cerebral artery determined by systemic arterial stiffness?" *Blood Pressure*, vol. 18, no. 3, pp. 130–134, 2009.

[38] M. P. Pase, N. A. Grima, C. Stough, A. Scholey, and A. Pipingas, "Association of pulsatile and mean cerebral blood flow velocity with age and neuropsychological performance," *Physiology & Behavior*, vol. 130, pp. 23 – 27, 2014.

[39] W. K. Lefferts, J. P. DeBlois, B. B. Gump, and K. S. Heffernan, "Carotid artery stiffness and cerebral pulsatility in children," *Artery Research*, vol. 22, pp. 64 – 67, 2018.

[40] G. Bateman, "Pulse-wave encephalopathy: a comparative study of the hydrodynamics of leukoaraiosis and normal-pressure hydrocephalus," *Neuroradiology*, vol. 44, no. 9, pp. 740–748, Sep 2002.

[41] M. Holmgren, A. Wåhlin, T. Dunås, J. Malm, and A. Eklund, "Assessment of cerebral blood flow pulsatility and cerebral arterial compliance with 4d flow mri," *Journal of Magnetic Resonance Imaging*, vol. n/a, no. n/a, November 2019.

[42] R. Jurca, A. S. Jackson, M. J. LaMonte, J. R. Morrow, S. N. Blair, N. J. Wareham, W. L. Haskell, W. van Mechelen, T. S. Church, J. M. Jakicic, and R. Laukkanen, "Assessing cardiorespiratory fitness without performing exercise testing," *American Journal of Preventive Medicine*, vol. 29, no. 3, pp. 185–193, 2005.

[43] R. Aaslid, T. Lundar, K. Lindegaard, and H. Nornes, "Estimation of cerebral perfusion pressure from arterial blood pressure and transcranial doppler recordings," *Intracranial Pressure VI*, pp. 226–229, 1986.

[44] G. V. Varsos, M. Kasprowicz, P. Smielewski, and M. Czosnyka, "Model-based indices describing cerebrovascular dynamics," *Neurocritical Care*, vol. 20, no. 1, pp. 142–157, Feb 2014.

[45] P. Segers, E. R. Rietzschel, M. L. D. Buyzere, S. J. Vermeersch, D. D. Bacquer, L. M. V. Bortel, G. D. Backer, T. C. Gillebert, and P. R. Verdonck, "Noninvasive (input) impedance, pulse wave velocity, and wave reflection in healthy middle-aged men and women," *Hypertension*, vol. 49, no. 6, pp. 1248–1255, 2007.

[46] C. D. Gatzka, J. D. Cameron, A. M. Dart, K. L. Berry, B. A. Kingwell, E. M. Dewar, C. M. Reid, and G. L. Jennings, "Correction of carotid augmentation index for heart rate in elderly essential hypertensives," *American Journal of Hypertension*, vol. 14, no. 6, pp. 573–577, 06 2001.

[47] B. E. Westerhof and N. Westerhof, "Uniform tube models with single reflection site do not explain aortic wave travel and pressure wave shape," *Physiological Measurement*, vol. 39, no. 12, p. 124006, dec 2018.

[48] J. Sugawara, K. Hayashi, and H. Tanaka, "Distal shift of arterial pressure wave reflection sites with aging," *Hypertension*, vol. 56, no. 5, pp. 920–925, 2010.

[49] T. S. Phan, J. K. Li, P. Segers, M. Reddy-Koppula, S. R. Akers, S. T. Kuna, T. Gislason, A. I. Pack, and J. A. Chirinos, "Aging is associated with an earlier arrival of reflected waves without a distal shift in reflection sites," *Journal of American Heart Association*, vol. 5, no. 9, 2016.

[50] M. Masuda, T. Emoto, A. Suzuki, M. Akutagawa, T. Kitawaki, K. Kitaoka, H. Tanaka, S. Obara, K. Yoshizaki, S. Konaka, and Y. Kinouchi, "Evaluation of blood flow velocity waveform in common carotid artery using multi-branched arterial segment model of human arteries," *Biomedical Signal Processing and Control*, vol. 8, no. 6, pp. 509 – 519, 2013.

[51] K. Hirata, T. Yaginuma, M. F. O'Rourke, and M. Kawakami, "Age-related changes in carotid artery flow and pressure pulses," *Stroke*, vol. 37, no. 10, pp. 2552–2556, 2006.