

Title: A national landscaping survey of critical care services in hospitals accredited for training in a lower-middle income country: Pakistan

Authors

Madiha Hashmia^a, Arshad Taqib^b, Muhammad Iqbal Memon^c, Syed Muneeb Alic^c, Saleh Khaskheli^d, Muhammad Sheharyar^e, Muhammad Hayat^f, Mohiuddin Shiekh^g, Chamira Kodippily^h, Dilanthi Gamage^h, Arjen M Dondorpⁱ, Rashan Haniffah^{h,i,j}, Abi Beane^{h,i,k*}.

Affiliations

^aZiauddin University, 4/B, Saharah-e-Ghalib, Block 6, Clifton Karachi, 7500 Sindh, Pakistan.

^bKaul Associates, 14 AbuBaker Block, New Garden Town, Lahore, Punjab, Pakistan.

^cPakistan Institute of Medical Sciences, Ibn-e-Sina Road, G-8/3, Islamabad, Islamabad Capital Territory, Pakistan.

^dPeople's University of Medical & Health Sciences for Women, Nawabshah, Shaheed Benazirabad, 67480 Sindh, Pakistan.

^eLady Reading Hospital, Soekarno Road, Peshawar, Khyber Pakhtunkhwa 25000, Pakistan.

^fNorth West General Hospital, Sector A-3, Phase 5, Hayatabad, Peshawar, Khyber Pakhtunkhwa 25100, Pakistan.

^gSouth East Asian Research in Critical care and Health, Remedial Centre Hospital, D-9, Block-I, North Nazimabad, Karachi 74700, Pakistan.

^hNetwork for Improving Critical Care Systems and Training, 2nd floor, YMBA Building, Colombo 08, Sri Lanka.

ⁱDepartment of Malaria and Critical Illness, Mahidol Oxford Tropical Medicine Research Unit, 3/F, 60th Anniversary Chalermprakit Building, 420/6 Rajvithi Road, Bangkok 10400, Thailand.

^jBloomsbury Institute for Intensive Care Medicine, Division of Medicine, University College London, Gower Street, London WC1E 6BT, United Kingdom.

^kAmsterdam Institute for Global Health and Development, University of Amsterdam, Paasheuvelweg 25, 1105 BP Amsterdam, Netherlands.

***Corresponding author.** Address. Department of Malaria and Critical Illness, Mahidol Oxford Tropical Medicine Research Unit, 3/F, 60th Anniversary Chalermprakit Building, 420/6 Rajvithi Road, Bangkok 10400, Thailand. Email. abi@nicsslk.com

Abstract

Purpose: To describe the extent and variation of critical care services in Pakistan

Materials and methods: A cross-sectional survey was conducted in all CCUs recognised for postgraduate training to determine administration, infrastructure, equipment, staffing, and training.

Results: There were 220 CCUs registered for training, providing 2166 CCU beds and 1473 ventilators. Regional distribution of CCU beds per 100,000 population ranged from 1.0 in Sindh to none in Gilgit Baltistan (median 0.7). A senior clinician trained in critical care was available in 19 (12.1%) of units, giving a ratio of one trained intensivist for every 82 CCU beds and 0.009 trained intensivists per 100,000 population. One to one nurse to bed ratio during the day was available in 84 (53.5%) of units, dropping to 75 (47.8%) at night. Availability of 1:1 nursing also varied between provinces, ranging from 56.5% in Punjab compared to 0% in Azad Jamu Kashmir. All CCUs had basic infrastructure (electricity, running water, piped oxygen) and basic equipment (electronic monitoring and infusion pumps).

Conclusion: Pakistan, a lower middle-income country has an established network of critical care facilities with access to basic equipment, but inequalities in its distribution. Investment in critical care training for doctors and nurses is needed.

Keywords: Critical care resources; Critical care services; Critical care staffing; Lower middle-income country.

Background

Demand for critical care services continue to grow internationally. Resources remain limited, most notably in low and lower-middle income countries (LMICs). In South Asia, overall improved public health and primary healthcare services in the region, the growing burden of noncommunicable disease, and with it a demand for surgical and trauma care has resulted in a shift in health systems priorities [1,2]. There is thus an increasing demand for critical care services, and the associated manpower, infrastructure and equipment requirements in LMICs.

Understanding the landscape of existing infrastructure, equipment and staffing both between and within countries provides valuable information for those seeking to strengthen critical care services and inform disaster and pandemic planning. Furthermore, mapping critical care services to the clinical characteristics of the patient it serves is a fundamental step in evaluating quality of existing service provision and to identify priorities for research and quality improvement.

Sri Lanka was the first country in South Asia to undertake a comprehensive national survey of critical care services [3]. Since then, regional efforts to map critical care services in Asia have contributed valuable information regarding critical care unit (CCU) bed availability in the region [9]. However, information regarding skills, training and organisational processes (essential to developing strategies for improving the quality of care) remains absent. The Pakistan Registry of Intensive Care (PRICE) [3], a cloud-based surveillance platform, currently supports a network of 43 CCUs in Pakistan recording over 2000 monthly critical care admissions. PRICE provides near real-time reporting on the epidemiology, severity of illness, treatment, microbiology and outcomes of CCU patients, alongside information regarding work force, unit occupancy, unit acuity, and resource utilisation. This information is used to drive local service evaluation and quality improvement interventions. PRICE is a founding member of the recently established Wellcome-MORU-CRIT Care Asia (CCA).

This paper details a national survey of critical care services in Pakistan including organisational structures, equipment, infrastructure and training capacity.

Setting

Pakistan consists of four provinces (Balochistan, Khyber Pakhtunkhwa, Punjab, and Sindh), two autonomous territories (Azad Jammu Kashmir, Gilgit-Baltistan) and one federal territory (Islamabad Capital Territory) [4]. Islamabad was included in the province of Punjab for the purposes of this study.

Methods

A CCU was defined as a clinical area (excluding operating theatres) which had the ability to provide organ support for in-patients, including mechanical ventilation. All hospitals recognised by the Pakistan Medical and Dental Council (PMDC) for internship training and the College of Physicians and Surgeons Pakistan (CPSP) for postgraduate residency training in anaesthesia, internal medicine, general surgery, cardiac surgery, pulmonology, nephrology, cardiology and critical care medicine were contacted by telephone by MH. All such hospitals were invited to participate in the survey if they reported the presence of at least one CCU. Eligible hospitals were asked for the number of CCUs, number of ventilators and asked to nominate a senior CCU doctor or sister in charge to respond to the survey questions. If a nominated contact was unavailable, at least one follow-up call was made for each CCU. The surveys were administered by telephone or online between February 2017 and December 2018. All responses were included in the

analysis. The survey instrument including characteristics and organizational structure, infrastructure and human resources was based on the tool pioneered in South Asia by our group [5]. Population per region was obtained from the government census up to January 2018 from publicly available sources [4]. CCUs were defined as open, closed or semi-closed [6].

Results

One hundred and fifty-one hospitals were identified, of which 30 did not have a CCU and were therefore excluded. All 121 eligible hospitals participated in reporting bed and ventilatory capacity. Two hundred and twenty CCUs were identified in these hospitals providing 2166 critical beds and 1473 ventilators. Of these 220 CCUs, 157 (71.4%), completed the full survey of organisational structure, infrastructure, equipment and human resources. The designated contact was not reachable in 60 CCUs and a further 3 were not available for interview (Figure 1). Table 1 summarizes the main characteristics of the hospitals surveyed. The density and distribution of CCU beds within teaching institutions (total and per 100 000 population) by administrative regions is described in Table 2 and Figure 2. Figure 3 reports population density as a reference.

Access and organisational structure.

Average beds within teaching institutions per 100,000 population was 0.7 (total number of CCU beds in participating institutions divided by the total population of Pakistan), ranging from 0 in Gilgit Baltistan to 1.0 in Sindh. The median number of critical care beds per unit was 9 (range 0-30). A total of 58 (47.9%) CCUs were managed directly by the government and 42 (34.7%) of units were managed by the private sector, with the remainder being administered by not-for-profit organisations (17.4%). Seventeen (10.8%) of the CCUs surveyed reported a ‘closed’ model of care whereby intensivists were the consultant in charge of care. Unrestricted visiting for families was practiced in 40 (25.5%) of units (Table 3).

Infrastructure.

Ventilator to bed ratio of 1:1 was observed in 82 (52.2%) of the CCUs (table 4), with Punjab province having the greatest number 49 (59.8%) and Azad Jamu Kashmir the lowest 0. All CCUs had a telephone line, however, only 52.2% had access to the internet. Table 5 summarises the availability of equipment to monitor critically ill patients. Almost all CCUs (95.5%) had access to 1:1 non-invasive multiparameter monitoring. Invasive arterial monitoring and capnography was available in 69 (44.0%) and 9 (5.7%) CCUs respectively. In addition, 51 (32.5%) and 39 (24.8%) units had access to point-of-care haemoglobin and lactate measurement respectively. Isolation rooms essential for management of infectious diseases, including severe acute respiratory infections, were available in 45 (28.7%) CCUs.

Human resources, team structure and training opportunities.

Table 6 summarises the human resources, team structure and training opportunities. A senior clinician trained in critical care was available in only 19 (12.1%) of units. This is a ratio of one trained intensivist for every 82 CCU beds and 0.009 trained intensivists per 100,000 population. The majority of units (86, 54.8%) was overseen by a consultant anaesthetist. In the remainder, 38.8% were overseen by a consultant physician and 6.4% by a consultant surgeon. A non-consultant doctor was assigned to CCU round-the-clock with no other work commitments in 140 (89.1%) of CCUs. Of the 121 institutions surveyed, 101 (83.5%) were recognised by the College of Physicians and Surgeons for speciality training (residency training). Critical care medicine (CCM) fellowship training was offered by 6 (4.9%) of institutions.

The majority of critical care units were managed by registered nurses with general training 149 (94.9%), with the remaining 8 (5.1%) being managed by technicians trained in anaesthesia or critical care. One to one nurse to bed ratio during the day for ventilated patients was available in 84 (53.5%) of units, and in 19 (12.1%) of units for self-ventilated patients. At night this availability dropped to 75 (47.8%) and 14 (8.9%) respectively. Availability of 1:1 nursing also varied between provinces, ranging from 56.5% having a 1:1 availability in the Punjab compared to 0% in Azad Jamu Kashmir. Microbiologists and haematologists were accessible in 96 (61.1%) and 103 (65.6%) of units respectively. Health care assistants or trained technicians were part of the care provision team in 94 (59.9%) of CCUs. Radiology technicians were available in 150 (95.5%) units and a further 128 (81.5%) CCUs had access to physiotherapy services.

Discussion

This national survey from Pakistan reports very limited critical care bed availability but where available CCUs are well resourced with basic equipment for invasive ventilation and monitoring. It further highlights the lack of critical care trained staff and the need for urgent investment in critical care services to address this gap in training capacity if care is to be improved.

Critical care capacity

The number of critical care beds in LMICs are known to be lower when compared to high-income countries [7,8], this disparity is pronounced in Pakistan in comparison to neighbouring countries: At 0.71 per 100 000 population, it is lower than Sri Lanka (2.3 critical care beds per 100,000), Nepal (2.8) and India (2.3) [9]. The survey further identified a wide disparity in access to critical care beds between the provinces (Figure 3). Punjab, whilst being the most densely populated province of the country, has lower availability of critical care beds than neighbouring Sindh. Similar disparity exists between major cities in each province (Table 2). As urbanisation and migration to cities for employment continues in Pakistan, and as the burden of non-communicable disease rises - including road traffic accidents and multimorbidities, it is a national priority to address the disparity in access to critical care services [2].

CCUs in both public and private sector institutions (including not-for-profit) had the basic **infrastructure** (electricity and a backup generator, piped oxygen, medical air and suction, infusion and syringe pumps), and basic monitoring (non-invasive multiparameter monitor, mercury thermometer, and manual CVP measurement). Overall ventilator to bed ratio was 1:1.3, meaning 3 out of every 4 CCU beds have the facility to mechanically ventilate. Availability of these resources is reassuring, and suggests that the provision of the mainstays of critical care organ support- ventilation therapy, basic cardiovascular monitoring and support, and delivery of fluids is possible. The safe and effective delivery of these therapies, however, relies not only on the availability of equipment, but on specialist trained staff with the skills to instigate, titrate and troubleshoot treatment.

In contrast to the specific resources of critical care described above, sinks for hand washing were absent in 29.3% of CCUs and access to isolation rooms or cubicles to control cross infection with negative/positive air exchange mechanism was available in just 28.7% of CCUs, the majority of which were private sector tertiary care hospitals in the major cities. Addressing the absence of facilities for infection control is perhaps a key priority for those seeking to improve critical care services in the country, given the increasingly important role critical care plays in the preparation and management of seasonal epidemics (including

severe acute respiratory infections-SARI) and in the rising burden of drug resistant infections. Furthermore, access to point-of-care (POC) measurements including lactate haemoglobin, and availability of invasive haemodynamic monitoring, which are increasingly seen as essential resources for the management of critical illness, is lacking [10,11]. Only 39 (24.8%) of units had access to POC lactate and just 14 (8.9%) could invasively monitor haemodynamics. As Pakistan seeks to improve diagnosis and management of critically ill patients with SARI, sepsis and following trauma, better access to POC services and invasive monitoring, along with specially trained staff to interpret and respond to this information, is essential.

Capacity for training in critical care

The ratio of **trained intensivists** for each CCU bed in this survey is one to 82, much lower than estimates from South Asia, Latin and North America [12–14]. Critical care has been a recognised speciality in Pakistan with a structured training programme since 2004, however, at the time of this survey, only six institutions out of 121 teaching institutions (excluding military sites) were recognized by the College of Physicians and Surgeons of Pakistan for Critical Care Medicine training [15]. To date, just twenty-seven fellows have obtained the fellowship from the College in Critical Care Medicine [16]. Many currently practising intensivists in CCUs still have to travel outside of Pakistan for their higher training fellowships. Low intensivist to patient ratios (< 1:14) in academic medical CCUs have been cited as a barrier to delivery of quality of care and having a detrimental effect on staff well-being, specifically to the quality of pastoral and professional mentorship available for rotating trainees, who may consider specialising in CCM [17]. Lack of specialist training opportunities may be perpetuating the low numbers of designated critical care doctors on-call in CCUs and the low percentage of CCUs which are led by a trained intensivist. Lack of training opportunities extends beyond doctors, with only 40.1% of nurses in charge of CCUs having received any formal training in intensive, critical or cardiac care. Access to microbiologists, specialists who are increasingly considered fundamental to the interdisciplinary management of critically ill patients, was limited (61.1%).

Whilst there is growing evidence to support that intensivist-led patient management is associated with better patient outcomes and greater compliance with broadly accepted indicators of critical care quality [18], a closed model was uncommon in the CCUs surveyed (10.8%). A further 25.5% of CCUs reported a semi-closed structure, but without a trained intensivist as their clinical lead. Investment in the reorganisation of critical care services to improve operational efficiency and patient outcomes (length of stay, duration of mechanical ventilation) in CCUs has resulted in a shift toward closed organisational structures whereby admission to and management of patients within the CCU is coordinated by designated critical care clinicians [18]. Given the paucity of critical care services in Pakistan, such a model of management may promote effective resource utilisation. However, such models require national level investment in specialist training, and institution level investment in hiring and retaining such a specialist workforce. Other settings have demonstrated how investment in critical care trained clinician staff to lead units and investment in training for nurses working in critical care has positively impacted on safety within CCUs and outcomes for critical care patients. Without this investment, efforts to strengthen specialist capacity and improve quality of critical care services through research and implementation will be hindered.

Limitations.

This survey only approached institutions recognised for specialist teaching. Consequently, the number of CCU beds per 100,000 population is underestimated. A recent multicountry snapshot of critical care bed availability [9], to which Pakistan contributed, reported a national average in Pakistan of 1.3 beds per 100,000 population. However, such estimates included units which may have no recognised affiliation with critical care training and no support from critical care societies. Whether considering the numbers reported here, or the estimates from those with broader inclusion, CCU bed availability and trained, skilled staff is still lower than neighbouring countries. Pakistan has no central register or standard definition for CCUs and as such units may be operating without the support of trained intensivists.

Conclusion.

This survey provides a detailed landscape of critical care resources and training in Pakistan. Pakistan has an established network of critical care facilities with access to basic equipment but inequalities in access within and between provinces is prominent. Investment in critical care training for doctors and nurses is a key priority for the country. Investment in training for health care staff will likely enable efforts to improve safety within CCUs, accelerate opportunities for research and quality improvement.

Ethics approval and consent to participate

A waiver from ethical review was obtained from the National Bioethics Committee of Pakistan.

Financial Disclosure statement

This survey was supported by the Network for Improving Critical care Systems and Training (NICST), and a Wellcome funded innovations in Technology Partnership Award.

Conflict of Interest statement

The authors declare that they have no competing interests.

Acknowledgments

The authors would like to thank all members of the critical care community in Pakistan who contributed to this work with the shared goal of improving critical care services in the country. We would also like to thank Dr T Tolppa for his support with manuscript preparation, and Mr T Rashan for support with figures.

References

- [1] Murthy S, Leligdowicz A, Adhikari NKJ. Intensive care unit capacity in low-income countries: A systematic review. *PLoS One* 2015;10:1–12. <https://doi.org/10.1371/journal.pone.0116949>.
- [2] Hirshon JM, Risko N, Calvellido EJB, de Ramirez SS, Narayan M, Theodosios C, et al. Health systems and services: the role of acute care. *Bulletin of the World Health Organization* 2013;91:386–8. <https://doi.org/10.2471/blt.12.112664>.
- [3] Hashmi M, Beane A, Taqi A, Memon MI, Athapattu P, Khan Z, et al. Pakistan Registry of Intensive Care (PRICE): Expanding a lower middle-income, clinician-designed critical care registry in South Asia. *Pediatr Crit Care Med* 2019;20:190–5. <https://doi.org/10.1177/1751143718814126>.
- [4] Central Intelligence Agency. Pakistan. *The World Factbook* 2020. <https://www.cia.gov/library/publications/the-world-factbook/geos/pk.html> (accessed April 2, 2020).
- [5] Haniffa R, De Silva AP, Iddagoda S, Batawalage H, De Silva STGR, Mahipala PG, et al. A cross-sectional survey of critical care services in Sri Lanka: A lower middle-income country. *J Crit Care* 2014;29:764–8. <https://doi.org/10.1016/j.jcrc.2014.04.021>.
- [6] Juneja D, Nasa P, Singh O. Physician staffing pattern in intensive care units: Have we cracked the code? *Pediatr Crit Care Med* 2012;1:10–4. <https://doi.org/10.5492/wjccm.v1.i1.10>.
- [7] Rhodes A, Ferdinande P, Flaatten H, Guidet B, Metnitz PG, Moreno RP. The variability of critical care bed numbers in Europe. *Intensive Care Med* 2012;38:1647–53. <https://doi.org/10.1007/s00134-012-2627-8>.
- [8] Halpern NA, Goldman DA, Tan KS, Pastores SM. Trends in Critical Care Beds and Use Among Population Groups and Medicare and Medicaid Beneficiaries in the United States: 2000-2010. *Crit Care Med* 2016;44:1490–9. <https://doi.org/10.1097/CCM.0000000000001722>.
- [9] Phua J, Faruq MO, Kulkarni AP, Redjeki IS, Detleuxay K, Mendsaikhan N, et al. Critical Care Bed Capacity in Asian Countries and Regions. *Crit Care Med* 2020:1. <https://doi.org/10.1097/ccm.0000000000004222>.
- [10] Cecconi M, Evans L, Levy M, Rhodes A. Sepsis and septic shock. *Lancet* 2018;392:75–87. [https://doi.org/10.1016/S0140-6736\(18\)30696-2](https://doi.org/10.1016/S0140-6736(18)30696-2).
- [11] Spahn DR, Bouillon B, Cerny V, Duranteau J, Filipescu D, Hunt BJ, et al. The European guideline on management of major bleeding and coagulopathy following trauma: fifth edition. *Crit Care* 2019;23:98. <https://doi.org/10.1186/s13054-019-2347-3>.
- [12] Halpern NA, Tan KS, DeWitt M, Pastores SM. Intensivists in U.S. Acute Care Hospitals. *Crit Care Med* 2019;47:517–25. <https://doi.org/10.1097/CCM.0000000000003615>.
- [13] Arabi YM, Phua J, Koh Y, Du B, Faruq MO, Nishimura M, et al. Structure, Organization, and Delivery of Critical Care in Asian ICUs. *Crit Care Med* 2016;44:e940–8. <https://doi.org/10.1097/CCM.0000000000001854>.
- [14] Estenssoro E, Alegría L, Murias G, Friedman G, Castro R, Nin Vaeza N, et al. Organizational issues, structure, and processes of care in 257 ICUs in Latin America: A study from the Latin America intensive care network. *Crit Care Med* 2017;45:1325–36. <https://doi.org/10.1097/CCM.0000000000002413>.
- [15] College of Physicians and Surgeons Pakistan. Accredited Institutions. College of Physicians and Surgeons Pakistan n.d. <https://www.cpsp.edu.pk/accredited-institutes-fcps.php> (accessed April 1, 2020).
- [16] College of Physicians and Surgeons Pakistan. List of Good Standing Fellows. College of Physicians and Surgeons Pakistan 2020. <https://www.cpsp.edu.pk/online-directories.php> (accessed April 1, 2020).
- [17] Ward NS, Afessa B, Kleinpell R, Tisherman S, Ries M, Howell M, et al. Intensivist/patient ratios in closed ICUs: a statement from the Society of Critical Care Medicine Taskforce on ICU Staffing. *Crit Care Med* 2013;41:638–45. <https://doi.org/10.1097/CCM.0b013e3182741478>.
- [18] Nates JL, Nunnally M, Kleinpell R, Blosser S, Goldner J, Birriel B, et al. ICU Admission, Discharge, and Triage Guidelines: A Framework to Enhance Clinical Operations, Development of

Institutional Policies, and Further Research. *Crit Care Med* 2016;44:1553–602.
<https://doi.org/10.1097/CCM.0000000000001856>.

Table 1: Institutional location and profile.

Demographic	Institutions, no.(%) N=121	Functioning no.(%) N=1566	beds,
Provinces and autonomous territories (AT)			
Punjab	71 (58.7)	916 (58.5)	
Sindh	35 (28.9)	484 (30.9)	
Khyber Pakhtunkhwa	12 (9.9)	145 (9.3)	
Balochistan	2 (1.7)	15 (1.0)	
Gilgit Baltistan (AT)	0	0	
Azad Jammu and Kashmir (AT)	1 (0.8)	6 (0.4)	
Institution category			
Government	58 (47.9)	710 (45.3)	
Private	42 (34.7)	594 (37.9)	
Not-for-profit hospital	21 (17.4)	262 (16.7)	
Affiliated hospital recognised for training by:			
PMDC for Internship training	121 (100)	1566 (100)	
PMDC & CPSP for Internship & Residency training (Anaesthesia, Medicine, Surgery, Cardiac surgery, Cardiology, Nephrology)	101 (83.5)	1395 (89.1)	
CPSP for CCM Fellowship training	6 (5.0)	181 (11.6)	

Table 2: Institutions and bed availability per 100 000 population.

Administrative Regions	Population as per 2017 census survey	Functioning beds per 100,000 population (total)
Pakistan (all 4 provinces and 2 independent territories)	221,613,314	0.71
Punjab	120,012,442	0.76
Sindh	47,886,051	1.01
Khyber Pakhtunkhwa	35,525,047	0.41
Balochistan	12,344,408	0.12
Gilgit Baltistan	1,800,000	0
Azad Jammu and Kashmir	4,045,366	0.14

Table 3: Critical care unit profile

Demographic	CCUs, N=157	CCU beds, N=1566	Ventilators, N=1175
Type of CCU			
Medical	34 (21.7)	390 (24.9)	257 (21.9)
Surgical	51 (32.5)	443 (28.3)	375 (31.9)
General/Mixed	56 (35.7)	556 (35.5)	415 (35.3)
Specialized	16 (10.2)	177 (11.3)	131 (11.1)
Model of care			
Open	100 (63.7)	994 (63.5)	706 (60.1)
Close	17 (10.8)	204 (13.0)	169 (14.4)
Semi-closed	40 (25.5)	368 (23.5)	300 (25.5)

Table 4: Available infrastructure for each CCU

Facilities	CCUs, no. (%)
Piped oxygen	154 (98.1)
Wall suction units	132 (84.1)
Piped medical air	147 (93.6)
1:1 bed:ventilator	82 (52.2)
Paediatric mode in all ventilators	118 (75.2)
Syringe pumps	132 (84.1)
Infusion pumps	125 (79.6)
Difficult Airway Trolley (DAT)	115 (73.2)
Backup automatic electricity generator	
Yes	157 (100)
Hand washing facilities in the CCU	
Yes	111 (70.7)
CCU isolation rooms	
Yes	45 (28.7)
Access to arterial blood gas (ABG) analysis	
Yes	142 (90.4)
Location of ABG machine (n=142)	
Hospital lab	70 (49.3)
Within CCU	52 (36.6)
Another CCU or Operating theatre	20 (14.1)
CCUs reported a POC ABG service.*	72
Access to external internet	
Yes	82 (52.2)
Telephone	
Yes	157 (100)
*ABG machines located in the CCU or OR are considered point of care (POC) services.	

Table 5: Available monitoring facilities

Facilities	CCUs, no. (%)
Non-invasive monitoring (including manual central pressure monitoring)	157 (100)
Invasive arterial blood pressure	110 (70.1)
Capnogram	63 (40.1)
Cardiac output (haemodynamics)	14 (8.9)
Point-of-care haemoglobin measurement	51 (32.5%)
Point-of-care lactate measurement	39 (24.8%)

Table 6: CCU Medical Staffing

Availability of CCU Medical Staff	CCUs, no. (%)
CCU Consultant primary specialty	
Anaesthesia	86 (54.8)
Pulmonology	21 (13.4)
Medicine	30 (19.1)
Surgeon	10 (6.4)
Nephrologist	7 (4.5)
Cardiologist	3 (1.9)
CCU Consultant trained in CCM	
Yes	19 (12.1)
Availability of specialists for Consultant	
Anaesthesiologist	154 (98.1)
General Physician	152 (96.8)
General Surgeon	152 (96.8)
Obstetrician and gynaecologist	141 (89.8)
Cardiologist	145 (92.4)
Nephrologist	128 (81.5)
Pulmonologist	141 (89.8)
Gastroenterologist	113 (72.0)
Neurologist	119 (75.8)
Microbiologist	96 (61.1)
Haematologist	103 (65.6)
Orthopaedic	126 (80.3)
Urologist	128 (81.5)
Respiratory	125 (79.6)
Pathologist	117 (74.5)
Paediatrician	134 (85.4)
Cardiothoracic	79 (50.3)
Non-consultant doctors assigned only to CCU with no other work commitment round the clock	
Yes	140 (89.1)
Availability of paramedical staff/support staff	
Primary Training of Incharge CCU as;	
Registered Nurse	149 (94.9)
Anaesthesia/CCU Technician	8 (5.1)
Specialized training of incharge CCU in CCM/Coronary care	63 (40.1)
1:1 nursing of ventilated patients during day	84 (53.5)
1:1 nursing of ventilated patients during night	75 (47.8)

1:1 nursing of self-ventilated patients during day	19 (12.1)
1:1 nursing of self-ventilated patients during night	14 (8.9)
Healthcare Assistants (HCA) and Technicians	94 (59.9)
Physiotherapist	128 (81.5)
Radiology technician for portable x-ray	150 (95.5)





