SARS-CoV-2 RNA titers in wastewater anticipated

COVID-19 occurrence in a low prevalence area

Walter Randazzo[1,2], Pilar Truchado[3], Enric Cuevas-Ferrando[2], Pedro Simón[4], Ana

Allende[3], Gloria Sánchez[2*]

[1]Department of Microbiology and Ecology, University of Valencia, Av. Dr. Moliner,

50, Burjassot, 46100 Valencia, Spain;

[2]Department of Preservation and Food Safety Technologies, Institute of Agrochemistry

and Food Technology, IATA-CSIC, Av. Agustín Escardino 7, Paterna, 46980, Valencia,

Spain;

[3]Research Group on Quality, Safety and Bioactivity of Plant Foods, Department of

Food Science and Technology, CEBAS-CSIC, Campus Universitario de Espinardo, 25,

30100, Murcia, Spain;

[5]ESAMUR, Avenida Juan Carlos, s/n - Edificio Torre Jemeca, Murcia, Spain.

Running title: First detection of SARS-CoV-2 in untreated wastewater in Spain.

*Correspondence to: Gloria Sánchez, gloriasanchez@iata.csic.es.

Abstract

Severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) has caused more than

200,000 reported COVID-19 cases in Spain resulting in more than 20,800 deaths as of

April 21, 2020. Faecal shedding of SARS-CoV-2 RNA from COVID-19 patients has

extensively been reported. Therefore, we investigated the occurrence of SARS-CoV-2

RNA in six wastewater treatments plants (WWTPs) serving the major municipalities

within the Region of Murcia (Spain), the area with the lowest COVID-19 prevalence

within Iberian Peninsula. Firstly, an aluminum hydroxide adsorption-precipitation

concentration method was tested using a porcine coronavirus (Porcine Epidemic

Diarrhea Virus, PEDV) and mengovirus (MgV). The procedure resulted in average

recoveries of $10.90 \pm 3.54\%$ and $10.85 \pm 2.11\%$ in influent water and $3.29 \pm 1.58\%$ and

$6.19 \pm 1.00\%$ in effluent water samples for PEDV and MgV, respectively. Then, the

method was used to monitor the occurrence of SARS-CoV-2 from March 12 to April

14, 2020 in influent, secondary and tertiary effluent water samples. By using the real-

time RT-PCR (RT-qPCR) Diagnostic Panel validated by US CDC that targets three

regions of the virus nucleocapsid (N) gene, we estimated quantification of SARS-CoV-

2 RNA titers in untreated wastewater waters of 5.38 ± 0.21 log genomic copies/L on

average. Two secondary water samples resulted positive (2 out of 18) and all tertiary

water samples tested as negative (0 out 12). This environmental surveillance data were

compared to declared COVID-19 cases at municipality level, revealing that SARS-

CoV-2 was circulating among the population even before the first cases were reported

by local or national authorities in many of the cities where wastewaters have been

sampled. The detection of SARS-CoV-2 in wastewater in early stages of the spread of

COVID-19 highlights the relevance of this strategy as an early indicator of the infection

within a specific population. At this point, this environmental surveillance could be

implemented by municipalities right away as a tool, designed to help authorities to

coordinate the exit strategy to gradually lift its coronavirus lockdown.

Keywords: environmental surveillance, influent water, reclaimed water, concentration

protocol, RNA virus, coronavirus.

1. Introduction

Coronaviruses (CoVs) are a family of viruses pathogenic for humans and animals associated to respiratory and gastro-intestinal infections. CoVs used to be considered as minor pathogens for humans as they were responsible of common cold or mild respiratory infections in immunocompetent people. Nonetheless, the emergence of novel and highly pathogenic zoonotic diseases caused by CoVs such as Severe Acute Respiratory Syndrome (SARS), Middle East Respiratory Syndrome (MERS) and most recently SARS-CoV-2 arises questions to be addressed to guide public health response. CoVs are mainly transmitted through respiratory droplets. However, as for SARS and MERS, SARS-CoV-2 RNA has been detected in stool samples from patients suffering COVID-19 and from asymptomatic carriers (He et al., 2020; Pan et al., 2020; Woelfel et al., 2020; Young et al., 2020; Zhang et al., 2020). The duration of viral shedding has been observed to vary among patients with means of 14-21 days (Y. Wu et al., 2020; Xu et al., 2020), as well as the magnitude of shedding varies from 10^2 up to 10^8 RNA copies per gram (Lescure et al., 2020; Pan et al., 2020; Woelfel et al., 2020).

Infectious viruses deriving from fecal specimen have been cultured in Vero E6 cells and observed by electron microscopy (Wang et al., 2020). In addition, gastric, duodenal, and rectal epithelial cells are infected by SARS-CoV-2 and the release of the infectious virions to the gastrointestinal tract supports the possible fecal-oral transmission route (Xiao et al., 2020). Even though the possibility of faecal-oral transmission has been hypothesized, the role of secretions in the spreading of the disease is not clarified yet (Wang et al., 2020; Y. Wu et al., 2020; Xu et al., 2020; Yeo et al., 2020).

Wastewater monitoring has been a successful strategy pursued to track chemical and biological markers of human activity including illicit drugs consumption, pharmaceuticals use/abuse, water pollution, and occurrence of antimicrobial resistance

genes (Choi et al., 2018; de Oliveira et al., 2020; Lorenzo and Picó, 2019; Mercan et al.,

2019). Viral diseases have been also surveilled by the detection of genetic material into

wastewater as for enteric viruses (Hellmer et al., 2014; Prevost et al., 2015; Santiso-

Bellón et al., 2020), re-emerging zoonotic hepatitis E virus (Cuevas-Ferrando et al.,

2020; Miura et al., 2016), and poliovirus during the global eradication programme

(Asghar et al., 2014).

Currently, various studies detected SARS-CoV-2 RNA in wastewater worldwide

(Ahmed et al., 2020; Lodder and de Roda Husman, 2020; Medema et al., 2020; F. Wu

et al., 2020; Wurtzer et al., 2020), and wastewater testing has been suggested as a non-

invasive early-warning tool for monitoring the status and trend of COVID-19 infection

and as an instrument for tuning public health response (Daughton, 2020; Mallapaty,

2020; Naddeo and Liu, 2020). Under current circumstance, this environmental

surveillance could be implemented in wastewater treatment plants as a tool, designed to

help authorities to coordinate the exit strategy to gradually lift its coronavirus lockdown.

Here, we report the first detection of SARS-CoV-2 RNA in untreated wastewater

samples in Spain collected from six different wastewater treatment plants (WWTPs) in

Murcia, the lowest prevalence area in Iberian Peninsula. Additionally, the efficacy of

the secondary and tertiary treatments implemented in the WWPTs against SARS-CoV-2

has been confirmed. The outcomes of the environmental surveillance reflect the

epidemiological data in a low COVID-19 diagnosed cases setting, thus supporting the

need of developing and implementing advanced models for wastewater-based

epidemiology (WBE).

2. Material and methods

2.1. Sampling sites and samples collection

Influent and effluent water samples were collected from six WWTPs located in the main cities of the Region of Murcia, Spain (Figure 1). Technical data on WWTPs are provided in Table 1.

A total of 42 influent, and 18 secondary and 12 tertiary treated effluent water samples were collected from 12 March to 14 April 2020 and investigated for the occurrence of SARS-CoV-2 RNA. Collected samples were transferred on ice to the laboratory and concentrated on the same day of sampling or the day after.

2.2. Wastewater and effluent water concentration

The porcine epidemic diarrhea virus (PEDV) strain CV777 and the mengovirus (MgV) vMC0 (CECT 100000) were preliminary used to evaluate the aluminum hydroxide adsorption-precipitation method previously described for concentrating enteric viruses from wastewater and effluent water (AAVV, 2011; Randazzo et al., 2019). In brief, 200 mL of biobanked influent (n=2) and effluent water samples (n=2) were artificially inoculated with PEDV and MgV. Then pH was adjusted to 6.0 and $Al(OH)_3$ precipitate formed by adding 1 part 0.9N $AlCl_3$ (Acros organics, Geel, Belgium) solution to 100 parts of sample. The pH was readjusted to 6.0 and sample mixed using an orbital shaker at 150 rpm for 15 min at room temperature. Then, viruses were concentrated by centrifugation at $1,700 \times g$ for 20 min. The pellet was resuspended in 10 mL of 3% beef extract pH 7.4, and samples were shaken for 10 min at 150 rpm. Concentrate was recovered by centrifugation at $1,900 \times g$ for 30 min and pellet resuspended in 1 mL of PBS.

All wastewater and effluent water samples included in this study were processed as described and MgV (5 log PCRU) was spiked as process control.

2.3. Viral extraction, detection and quantification

Viral RNA was extracted from concentrates using the NucleoSpin RNA virus kit (Macherey-Nagel GmbH & Co., Düren, Germany) according to the manufacturer's instructions with some modifications. Briefly, 150 μL of the concentrated sample was mixed with 25 μL of Plant RNA Isolation Aid (Thermo Fisher Scientific, Vilnius, Lithuania) and 600 μL of lysis buffer from the NucleoSpin virus kit and subjected to pulse-vortexing for 1 min. Then, the homogenate was centrifuged for 5 min at $10,000 \times$ g to remove the debris. The supernatant was subsequently processed according to the manufacturer's instructions.

Viral RNA was detected by TaqMan real-time RT-PCR (RT-qPCR) on LightCycler 480 instrument (Roche Diagnostics, Germany). MgV RNA was detected by using UltraSense One-Step kit (Invitrogen, SA, US) and the RT-qPCR assay as in ISO 15216-1:2017 (Costafreda et al., 2006; ISO 15216-1, 2017). Undiluted and ten-fold diluted MgV RNA was tested to check for RT-qPCR inhibitors.

PEDV RNA was detected by using PrimeScript™ One Step RT-PCR Kit (Takara Bio, USA) and the TaqMan RT-qPCR assay described by (Zhou et al., 2017).

SARS-CoV-2 RNA was detected by using PrimeScript™ One Step RT-PCR Kit and the RT-qPCR diagnostic panel assays validated by the US Centers for Disease Control and Prevention (CDC, 2020). The first version of the kit with three sets of oligonucleotide primers and probes was used to target three different SARS-CoV-2 regions of the nucleocapsid (N) gene. The sets of primer and probes (2019-nCoV RUO Kit) as well as the positive control (2019-nCoV_N_Positive Control) were provided by IDT (Integrated

DNA Technologies, Leuven, Belgium). Each RNA was analyzed in duplicate and every

RT-qPCR assay included negative (nuclease-free water) and positive controls.

Biobanked samples (n=4) collected in October 2019, before the first COVID-19 case

was documented, were used as relevant negative control to exclude false positive

reactions.

SARS-CoV-2 RNA was quantified as genome copies (gc) by plotting the quantification

cycles (Ct) to an external standard curve built with 10-fold serial dilution of a quantified

plasmid control (IDT). MgV and PEDV RNA were quantified by plotting the Cts to an

external standard curve generated by serial end-point dilution method.

MgV recovery rates were calculated and used as quality assurance parameters according

to ISO 15216-1:2017 (ISO 15216-1, 2017).

3. Results and Discussion

3.1. Performance of the concentration method

The aluminum hydroxide adsorption-precipitation method was tested by spiking

influent and effluent samples with MgV and PEDV. On average, MgV was recovered at

ranges of 10.85 ± 2.11% in influent and 6.19 ± 1.00% in effluent water. PEDV was

recovered at ranges of 10.90 ± 3.54% in influent and 3.29 ± 1.58% in effluent water.

These results are in line with the MgV recoveries reported for enteric viruses

concentration in water samples by the same aluminum-based method (Cuevas-Ferrando

et al., 2020; Randazzo et al., 2019) and higher than the 1% as the quality assurance

parameter indicated for bottled water into ISO 15216-1:2017 (ISO 15216-1, 2017).

Similarly, MgV was successfully used as recovery control for hepatitis E virus

concentration from influent and effluent water samples (5-13%) by applying a

polyethylene glycol (PEG) precipitation method (Miura et al., 2016). A similar PEG-

based protocol was recently used to recover SARS-CoV-2 from wastewater, although recovery control was not included in the study (F. Wu et al., 2020).

Moreover, filtration through 10 kDa Centricon® Plus-70 centrifugal device successfully recovered SARS-CoV-2 in wastewater with recovery efficiencies of F-specific RNA phages of 73% (Medema et al., 2020). However, concentration by electropositive membrane should be further evaluated given a SARS-CoV recovery from wastewater of 1% (Wang et al., 2005).

Rigorous limits of detection should be established by spiking SARS-CoV-2 cell-culture adapted strain or positive COVID-19 fecal samples in influent and effluent wastewater samples to be concentrated following the aluminum hydroxide adsorption-precipitation method. Nonetheless, the need of a BSL3 laboratory facility to handle SARS-CoV-2 represents the main limitation of this experiment.

3.2. SARS-CoV-2 titers in wastewater and effluent water

A total of 42 influent, and 18 secondary and 12 tertiary treated effluent water samples were collected from 12 March to 14 April 2020 and investigated for the occurrence of SARS-CoV-2 RNA. Samples were considered positive for Ct below 40 (as in Medema et al., 2020 and F. Wu et al., 2020) and titrated by using the quantified plasmid control for each of the RT-qPCR targets.

The 83.3% (35 positive samples out of 42) influent samples and the 11.1% (2 out of 18) secondary treated water samples were tested positive for at least one SARS-CoV-2 RT-qPCR target. None of the tertiary effluent samples (0 out of 12) tested positive for any of the SARS-CoV-2 RT-qPCR target (Figure 2). A relevant number of influent water samples (12%) showed Ct ranging between 37 and 40, even though lower Ct of 34-37

were observed (29%). In all samples, MgV recoveries were above 1% (10.05 ±

14.10%).

On average, SARS-CoV-2 RNA titers of 5.15 ± 0.25, 5.53 ± 0.24, and 5.49 ± 0.27 log

gc/L were quantified in wastewater by using N1, N2 and N3 primer/probe mixes,

respectively. Titers of 4 and 5 to more than 6 log gc/L have been reported in

Massachusetts and France, respectively (F. Wu et al., 2020; Wurtzer et al., 2020).

A secondary effluent sample resulted positive for N2 and quantified as 5.40 log gc/L.

An additional secondary effluent sample was positive for the three molecular targets and

below the quantification limit.

Detection of SARS-CoV-2 RNA in influent water has been reported worldwide (Ahmed

et al., 2020; Lodder and de Roda Husman, 2020; Medema et al., 2020; F. Wu et al.,

2020), and only one study tested treated wastewater that resulted positive (Paris)

(Wurtzer et al., 2020). We observed discrepancies among RT-qPCR N1, N2 and N3

assays for several water samples in agreement to a previous report (Medema et al.,

2020). This could be due to the different analytical sensitivity among the assays as well

as the detection of possible false positive samples by RT-qPCR N3 in low concentrated

samples (Jung et al., 2020; Vogels et al., 2020). The latter possibility has been solved by

excluding the N3 primers/probe set from the US CDC 2019-nCoV RT-qPCR

diagnostic panel in its last revision (March, 30) (CDC, n.d.). In addition, a partial

inhibitory effect of the matrix is not to be completely excluded despite the controls

included in the assays. A more sensitive estimation of SARS-CoV-2 loads in

wastewater should be studied by digital RT-qPCR (dRT-qPCR). dRT-qPCR could be

used to quantify samples with low viral loads, even though it may not be the best

practical and economically sustainable option for environmental surveillance.

Even though the SARS-CoV-2 RNA detection in wastewater is functional for WBE

purposes, the risk for human health associated to the water cycle is still under debate as

infectivity of viral particles remain to be confirmed as well as its potential fecal-oral

transmission.

In spite of the high concentration of viral RNA in specimen and the evidence of

gastrointestinal infection (Xiao et al., 2020), infectious viruses from stools have been

isolated in one study (Wang et al., 2020) while another attempt resulted without success

(Woelfel et al., 2020).

The potential transmission of SARS-CoV-2 via wastewater has not been proven (CDC,

n.d.; WHO, 2020) and it seems unlikely given the poor viral stability in environmental

conditions and its elevated sensitivity to disinfectants (Chin et al., 2020; Haas, 2020;

Lodder and de Roda Husman, 2020).

3.3. Environmental surveillance

Epidemiological data on COVID-19 in the Murcia Region have been retrieved from the

publically available repository of the "Servicio de epidemiologia" of the "Consejería de

Salud de la Región de Murcia" (available at http://www.murciasalud.es/principal.php)

(Table 2) and plotted to the SARS-CoV-2 RNA mean loads as detected by three RT-

qPCR assays (Figure 3).

In general, RT-qPCR amplification signals have been detected in wastewaters when

cases were diagnosed within the municipality. Positive wastewater samples have been

detected with at least two out of three RT-qPCR assays in low prevalence municipalities

as in Murcia (96 cases, 21.18 cases per 100,000 inhabitants), Cartagena (36 cases,

16.76) and Molina de Segura (12 cases, 16.69). Of note, positive wastewater samples

were detected 12-16 days before COVID-19 cases were declared in Lorca, Cieza and

Totana municipalities.

A similar study conducted in Paris (France) demonstrated the detection of viral genome

before the exponential phase of the epidemic (Wurtzer et al., 2020). However, our

results indicate that SARS-CoV-2 can be detected weeks before the first confirmed

case. The early detection of SARS-CoV-2 RNA in wastewater could have alerted about

the imminent danger, giving a valuate time to the managers to coordinate and

implement actions to slow the spread of the disease. Therefore, our outcomes support

that WBE could be used as an early warning tool to monitor the status and the trend of

COVID-19 infection within a community.

On the other hand, we believe that this environmental surveillance could be used as an

instrument to drive the right decisions to reduce the risk of lifting restrictions too early.

For instance, a key question is how to reduce the risk of a "second wave" and/or

recurring local outbreaks. Massive population tests are the first choice, but in their

absence, wastewater monitorization of SARS-CoV-2 RNA can give a reliable picture of

the current situation. Our wastewater data do not quantitatively resemble the prevalence

of COVID-19 confirmed cases. To this end, a quantitative model that includes and

corrects all the variables affecting these wastewater surveillance data would be useful

for a better interpretation. For instance, not all COVID-19 positive patients excrete

SARS-CoV-2 RNA in faeces, and when it occurs, the titers and the duration of shedding

vary among individuals and across time (He et al., 2020; Pan et al., 2020; Woelfel et al.,

2020; Xu et al., 2020). On the other hand, the real number of positive cases within the

Murcia Region remains unknown because of the large number of mild or asymptomatic

carriers that have not been included in epidemiological statistics.

These aspects together with environmental variables (e.g., rainfall events, temperature)

increase the uncertainties linked to the correlation between SARS-CoV-2 RNA

detection in wastewater samples and the prevalence of COVID-19 that could be

explored by using complex models.

4. Conclusion

Overall, wastewater surveillance and WBE may represent a complementary approach to

estimate the presence and even the prevalence of COVID-19 in communities. This

represents an effective tool that needs to be further explored in order to direct public

health response, especially in cases of limited capacity for clinical testing.

Funding

The study was funded by the projects 20180705 of ESAMUR, 202070E101 of CSIC

and "VIRIDIANA" AGL2017-82909 (AEI/FEDER, UE) of MICIU. WR is supported

by APOSTD/2018/150 postdoctoral fellowship of Generalitat Valenciana. EC-F is

recipient of a predoctoral contract from the MICINN, Call 2018. PT is holder of the

RYC2018- 025510-I Ramón y Cajal contract from the MICIU.

Acknowledgments

The authors acknowledge the "Entidad Regional de Saneamiento y Depuración de

Aguas Residuales (ESAMUR)" for authorizing the sampling and Prof. Ana Carvajal

(Faculty of Veterinary Medicine, University of Leon, Spain) for kindly providing

PEDV.

References

AAVV, 2011. Standard Methods For the Examination of Water and Wastewater (9510),
in: Detection of Enteric Viruses. American Public Health Association.

Ahmed, W., Angel, N., Edson, J., Bibby, K., Bivins, A., O'Brien, J.W., Choi, P.M.,
Kitajima, M., Simpson, S.L., Li, J., Tscharke, B., Verhagen, R., Smith, W.J.M.,
Zaugg, J., Dierens, L., Hugenholtz, P., Thomas, K. V, Mueller, J.F., 2020. First
confirmed detection of SARS-CoV-2 in untreated wastewater in Australia: A proof
of concept for the wastewater surveillance of COVID-19 in the community. Sci.
Total Environ. 138764.
https://doi.org/https://doi.org/10.1016/j.scitotenv.2020.138764

Asghar, H., Diop, O.M., Weldegebriel, G., Malik, F., Shetty, S., El Bassioni, L.,
Akande, A.O., Al Maamoun, E., Zaidi, S., Adeniji, A.J., Burns, C.C., Deshpande,
J., Oberste, M.S., Lowther, S.A., 2014. Environmental surveillance for
polioviruses in the Global Polio Eradication Initiative. J. Infect. Dis. 210 Suppl,
S294-303. https://doi.org/10.1093/infdis/jiu384

CDC, 2020. CDC 2019-Novel Coronavirus (2019-nCoV) Real-Time RT-PCR
Diagnostic Panel [WWW Document]. URL
https://www.fda.gov/media/134922/download

CDC, n.d. CDC 2019-Novel Coronavirus (2019-nCoV) Real-Time RT-PCR Diagnostic
Panel [WWW Document]. URL https://www.fda.gov/media/134922/download

CDC, n.d. Water Transmission and COVID-19: Questions and Answers [WWW
Document]. URL https://www.cdc.gov/coronavirus/2019-ncov/php/water.html
(accessed 4.21.20b).

Chin, A.W.H., Chu, J.T.S., Perera, M.R.A., Hui, K.P.Y., Yen, H.-L., Chan, M.C.W.,
Peiris, M., Poon, L.L.M., 2020. Stability of SARS-CoV-2 in different
environmental conditions. The Lancet Microbe.
https://doi.org/https://doi.org/10.1016/S2666-5247(20)30003-3

Choi, P.M., Tscharke, B.J., Donner, E., O'Brien, J.W., Grant, S.C., Kaserzon, S.L.,
Mackie, R., O'Malley, E., Crosbie, N.D., Thomas, K. V, Mueller, J.F., 2018.
Wastewater-based epidemiology biomarkers: Past, present and future. TrAC -
Trends Anal. Chem. https://doi.org/10.1016/j.trac.2018.06.004

Costafreda, M.I., Bosch, A., Pintó, R.M., 2006. Development, evaluation, and
standardization of a real-time TaqMan reverse transcription-PCR assay for
quantification of hepatitis A virus in clinical and shellfish samples. Appl. Environ.
Microbiol. 72, 3846–3855. https://doi.org/10.1128/AEM.02660-05

Cuevas-Ferrando, E., Randazzo, W., Pérez-Cataluña, A., Sánchez, G., 2020. HEV
Occurrence in Waste and Drinking Water Treatment Plants   . Front. Microbiol. .

Daughton, C., 2020. The international imperative to rapidly and inexpensively monitor
community-wide Covid-19 infection status and trends. Sci. Total Environ. 138149.
https://doi.org/https://doi.org/10.1016/j.scitotenv.2020.138149

de Oliveira, M., Frihling, B.E.F., Velasques, J., Filho, F.J.C.M., Cavalheri, P.S.,
Migliolo, L., 2020. Pharmaceuticals residues and xenobiotics contaminants:
Occurrence, analytical techniques and sustainable alternatives for wastewater

treatment. Sci. Total Environ. https://doi.org/10.1016/j.scitotenv.2019.135568

Haas, C., 2020. Coronavirus and Risk Analysis. Risk Anal. 40, 660–661.
https://doi.org/10.1111/risa.13481

He, X., Lau, E.H.Y., Wu, P., Deng, X., Wang, J., Hao, X., Lau, Y.C., Wong, J.Y.,
Guan, Y., Tan, X., Mo, X., Chen, Y., Liao, B., Chen, W., Hu, F., Zhang, Q.,
Zhong, M., Wu, Y., Zhao, L., Zhang, F., Cowling, B.J., Li, F., Leung, G.M., 2020.
Temporal dynamics in viral shedding and transmissibility of COVID-19. Nat. Med.
https://doi.org/10.1038/s41591-020-0869-5

Hellmer, M., Paxeus, N., Magnius, L., Enache, L., Arnholm, B., Johansson, A.,
Bergstrom, T., Norder, H., 2014. Detection of pathogenic viruses in sewage
provided early warnings of hepatitis A virus and norovirus outbreaks. Appl.
Environ. Microbiol. 80, 6771–6781. https://doi.org/10.1128/AEM.01981-14

ISO 15216-1, 2017. Microbiology of the Food Chain – Horizontal Method for
Determination of Hepatitis A Virus and Norovirus Using Real-Time RT-PCR –
Part 1: Method for Quantification. ISO 15216-1:2017.

Jung, Y.J., Park, G.-S., Moon, J.H., Ku, K., Beak, S.-H., Kim, Seil, Park, E.C., Park, D.,
Lee, J.-H., Byeon, C.W., Lee, J.J., Maeng, J.-S., Kim, S.J., Kim, Seung Il, Kim, B.-
T., Lee, M.J., Kim, H.G., 2020. Comparative analysis of primer-probe sets for the
laboratory confirmation of SARS-CoV-2. bioRxiv 2020.02.25.964775.
https://doi.org/10.1101/2020.02.25.964775

Lescure, F.-X., Bouadma, L., Nguyen, D., Parisey, M., Wicky, P.-H., Behillil, S.,
Gaymard, A., Bouscambert-Duchamp, M., Donati, F., Le Hingrat, Q., Enouf, V.,
Houhou-Fidouh, N., Valette, M., Mailles, A., Lucet, J.-C., Mentre, F., Duval, X.,
Descamps, D., Malvy, D., Timsit, J.-F., Lina, B., van-der-Werf, S., Yazdanpanah,
Y., 2020. Clinical and virological data of the first cases of COVID-19 in Europe: a
case series. Lancet Infect. Dis. https://doi.org/10.1016/S1473-3099(20)30200-0

Lodder, W., de Roda Husman, A.M., 2020. SARS-CoV-2 in wastewater: potential
health risk, but also data source. Lancet Gastroenterol. Hepatol.
https://doi.org/10.1016/S2468-1253(20)30087-X

Lorenzo, M., Picó, Y., 2019. Wastewater-based epidemiology: current status and future
prospects. Curr. Opin. Environ. Sci. Heal.
https://doi.org/10.1016/j.coesh.2019.05.007

Mallapaty, S., 2020. How sewage could reveal true scale of coronavirus outbreak.
Nature. https://doi.org/10.1038/d41586-020-00973-x

Medema, G., Heijnen, L., Elsinga, G., Italiaander, R., Brouwer, A., 2020. Presence of
SARS-Coronavirus-2 in sewage. medRxiv 2020.03.29.20045880.
https://doi.org/10.1101/2020.03.29.20045880

Mercan, S., Kuloglu, M., Asicioglu, F., 2019. Monitoring of illicit drug consumption
via wastewater: development, challenges, and future aspects. Curr. Opin. Environ.
Sci. Heal. https://doi.org/10.1016/j.coesh.2019.05.002

Miura, T., Lhomme, S., Le Saux, J.-C., Le Mehaute, P., Guillois, Y., Couturier, E.,
Izopet, J., Abranavel, F., Le Guyader, F.S., 2016. Detection of Hepatitis E Virus in
Sewage After an Outbreak on a French Island. Food Environ. Virol. 8, 194–9.

376 https://doi.org/10.1007/s12560-016-9241-9

Naddeo, V., Liu, H., 2020. Editorial Perspectives: 2019 novel coronavirus (SARS-CoV-2): what is its fate in urban water cycle and how can the water research community respond? Environ. Sci. Water Res. Technol. https://doi.org/10.1039/D0EW90015J

Pan, Y., Zhang, D., Yang, P., Poon, L.L.M., Wang, Q., 2020. Viral load of SARS-CoV-2 in clinical samples. Lancet Infect. Dis. 20, 411–412. https://doi.org/10.1016/S1473-3099(20)30113-4

Prevost, B., Lucas, F.S., Goncalves, A., Richard, F., Moulin, L., Wurtzer, S., 2015. Large scale survey of enteric viruses in river and waste water underlines the health status of the local population. Environ. Int. 79, 42–50. https://doi.org/10.1016/j.envint.2015.03.004

Randazzo, W., Piqueras, J., Evtoski, Z., Sastre, G., Sancho, R., Gonzalez, C., Sanchez, G., 2019. Interlaboratory Comparative Study to Detect Potentially Infectious Human Enteric Viruses in Influent and Effluent Waters. Food Environ. Virol. https://doi.org/10.1007/s12560-019-09392-2

Santiso-Bellón, C., Randazzo, W., Pérez-Cataluña, A., Vila-Vicent, S., Gozalbo-Rovira, R., Muñoz, C., Buesa, J., Sanchez, G., Díaz, J.R., 2020. Epidemiological surveillance of norovirus and rotavirus in sewage (2016–2017) in Valencia (Spain). Microorganisms. https://doi.org/10.3390/microorganisms8030458

Vogels, C.B.F., Brito, A.F., Wyllie, A.L., Fauver, J.R., Ott, I.M., Kalinich, C.C., Petrone, M.E., Casanovas-Massana, A., Muenker, M.C., Moore, A.J., Klein, J., Lu, P., Lu-Culligan, A., Jiang, X., Kim, D.J., Kudo, E., Mao, T., Moriyama, M., Oh, J.E., Park, A., Silva, J., Song, E., Takehashi, T., Taura, M., Tokuyama, M., Venkataraman, A., Weizman, O.-E., Wong, P., Yang, Y., Cheemarla, N.R., White, E., Lapidus, S., Earnest, R., Geng, B., Vijayakumar, P., Odio, C., Fournier, J., Bermejo, S., Farhadian, S., Dela Cruz, C., Iwasaki, A., Ko, A.I., Landry, M.-L., Foxman, E.F., Grubaugh, N.D., 2020. Analytical sensitivity and efficiency comparisons of SARS-COV-2 qRT-PCR assays. medRxiv 2020.03.30.20048108. https://doi.org/10.1101/2020.03.30.20048108

Wang, W., Xu, Y., Gao, R., Lu, R., Han, K., Wu, G., Tan, W., 2020. Detection of SARS-CoV-2 in Different Types of Clinical Specimens. JAMA. https://doi.org/10.1001/jama.2020.3786

Wang, X.-W., Li, J.-S., Jin, M., Zhen, B., Kong, Q.-X., Song, N., Xiao, W.-J., Yin, J., Wei, W., Wang, G.-J., Si, B.-Y., Guo, B.-Z., Liu, C., Ou, G.-R., Wang, M.-N., Fang, T.-Y., Chao, F.-H., Li, J.-W., 2005. Study on the resistance of severe acute respiratory syndrome-associated coronavirus. J. Virol. Methods 126, 171–177. https://doi.org/10.1016/j.jviromet.2005.02.005

WHO, W.H.O., 2020. Water, sanitation, hygiene and waste management for COVID-19 [WWW Document]. URL https://www.who.int/publications-detail/water-sanitation-hygiene-and-waste-management-for-covid-19 (accessed 4.21.20).

Woelfel, R., Corman, V.M., Guggemos, W., Seilmaier, M., Zange, S., Mueller, M.A., Niemeyer, D., Vollmar, P., Rothe, C., Hoelscher, M., Bleicker, T., Bruenink, S., Schneider, J., Ehmann, R., Zwirglmaier, K., Drosten, C., Wendtner, C., 2020. Clinical presentation and virological assessment of hospitalized cases of

coronavirus disease 2019 in a travel-associated transmission cluster. medRxiv
2020.03.05.20030502. https://doi.org/10.1101/2020.03.05.20030502

Wu, F., Xiao, A., Zhang, J., Gu, X., Lee, W.L., Kauffman, K., Hanage, W., Matus, M.,
Ghaeli, N., Endo, N., Duvallet, C., Moniz, K., Erickson, T., Chai, P., Thompson,
J., Alm, E., 2020. SARS-CoV-2 titers in wastewater are higher than expected from
clinically confirmed cases. medRxiv 2020.04.05.20051540.
https://doi.org/10.1101/2020.04.05.20051540

Wu, Y., Guo, C., Tang, L., Hong, Z., Zhou, J., Dong, X., Yin, H., Xiao, Q., Tang, Y.,
Qu, X., Kuang, L., Fang, X., Mishra, N., Lu, J., Shan, H., Jiang, G., Huang, X.,
2020. Prolonged presence of SARS-CoV-2 viral RNA in faecal samples. Lancet
Gastroenterol. Hepatol. 5, 434–435. https://doi.org/10.1016/S2468-
1253(20)30083-2

Wurtzer, S., Marechal, V., Mouchel, J.-M., Moulin, L., 2020. Time course quantitative
detection of SARS-CoV-2 in Parisian wastewaters correlates with COVID-19
confirmed cases. medRxiv 2020.04.12.20062679.
https://doi.org/10.1101/2020.04.12.20062679

Xiao, F., Tang, M., Zheng, X., Li, C., He, J., Hong, Z., Huang, S., Zhang, Z., Lin, X.,
Fang, Z., Lai, R., Chen, S., Liu, J., Huang, J., Xia, J., Li, Z., Jiang, G., Liu, Y., Li,
X., Shan, H., 2020. Evidence for gastrointestinal infection of SARS-CoV-2.
medRxiv 2020.02.17.20023721. https://doi.org/10.1101/2020.02.17.20023721

Xu, Y., Li, X., Zhu, B., Liang, H., Fang, C., Gong, Y., Guo, Q., Sun, X., Zhao, D.,
Shen, J., Zhang, H., Liu, H., Xia, H., Tang, J., Zhang, K., Gong, S., 2020.
Characteristics of pediatric SARS-CoV-2 infection and potential evidence for
persistent fecal viral shedding. Nat. Med. 26, 502–505.
https://doi.org/10.1038/s41591-020-0817-4

Yeo, C., Kaushal, S., Yeo, D., 2020. Enteric involvement of coronaviruses: is
faecal–oral transmission of SARS-CoV-2 possible? Lancet Gastroenterol.
Hepatol. 5, 335–337. https://doi.org/10.1016/S2468-1253(20)30048-0

Young, B.E., Ong, S.W.X., Kalimuddin, S., Low, J.G., Tan, S.Y., Loh, J., Ng, O.-T.,
Marimuthu, K., Ang, L.W., Mak, T.M., Lau, S.K., Anderson, D.E., Chan, K.S.,
Tan, T.Y., Ng, T.Y., Cui, L., Said, Z., Kurupatham, L., Chen, M.I.-C., Chan, M.,
Vasoo, S., Wang, L.-F., Tan, B.H., Lin, R.T.P., Lee, V.J.M., Leo, Y.-S., Lye, D.C.,
Team, for the S. 2019 N.C.O.R., 2020. Epidemiologic Features and Clinical
Course of Patients Infected With SARS-CoV-2 in Singapore. JAMA.
https://doi.org/10.1001/jama.2020.3204

Zhang, W., Du, R.-H., Li, B., Zheng, X.-S., Yang, X.-L., Hu, B., Wang, Y.-Y., Xiao,
G.-F., Yan, B., Shi, Z.-L., Zhou, P., 2020. Molecular and serological investigation
of 2019-nCoV infected patients: implication of multiple shedding routes. Emerg.
Microbes Infect. 9, 386–389. https://doi.org/10.1080/22221751.2020.1729071

Zhou, X., Zhang, T., Song, D., Huang, T., Peng, Q., Chen, Y., Li, A., Zhang, F., Wu,
Q., Ye, Y., Tang, Y., 2017. Comparison and evaluation of conventional RT-PCR,
SYBR green I and TaqMan real-time RT-PCR assays for the detection of porcine
epidemic diarrhea virus. Mol. Cell. Probes 33, 36–41.
https://doi.org/https://doi.org/10.1016/j.mcp.2017.02.002

465   **Figure 1.** Maps of the sampling location. Symbols represents WWTPs and are sized

466   according to the number of equivalent inhabitants.

468   **Table 1.** Operating characteristics of WWTPs.

	Population		Capacity (m³/a)		Reclamation processes	Reuse
	Connected	Equivalent	Designed	Current		
Murcia	370,893	530,499	36,500,000	36,952,999	Activated sludge (A2O process), Disinfection, NaClO	Public domain
Cartagena	175,870	163,969	12,775,000	8,625,103	Activated sludge, Disinfection	Irrigation
Molina de Segura	67,455	150,545	9,125,000	5,699,930	Activated sludge, Decantation, Coagulation, Flocculation, Sand filtration, Disinfection, UV, NaClO	Irrigation
Lorca	73,057	101,161	7,300,000	3,366,919	Activated sludge, Coagulation, Flocculation, Sand filtration, Disinfection, UV, NaClO	Irrigation
Cieza	33,744	69,502	3,650,000	2,338,673	Activated sludge (Extended aeration), Disinfection, Coagulation, Flocculation, Sand filtration, Disinfection, UV	Irrigation
Totana	29,113	28,289	2,190,000	1,440,463	Activated sludge (Extended aeration), Disinfection, UV	Irrigation

Sampling site	Water sample	Molecular Target	12 March	16 March	18 March	26 March	02 April	07 April	14 April
Murcia	Influent	N1	-		-				
		N2	-		-		-		-
		N3	-	-			-		
	Secondary treated	N1, N2, N3					-	-	-
	Tertiary treated	N1, N2, N3					-	-	-
Cartagena	Influent	N1	-	-	-			-	
		N2	-	-	-			-	
		N3	-			-			
	Secondary treated	N1, N2, N3					-	-	-
Molina de Segura	Influent	N1	-	-	-	-			
		N2			-	-			
		N3		-					
	Secondary treated	N1, N2, N3					-	-	-
	Tertiary treated	N1, N2, N3					-	-	-
Lorca	Influent	N1	-	-		-			
		N2		-	-			-	
		N3		-		-			
	Secondary treated	N1				-		-	-
		N2						-	-
		N3						-	-
	Tertiary treated	N1, N2, N3					-	-	-
Cieza	Influent	N1	-						
		N2	-						
		N3	-				-		-
	Secondary treated	N1, N3					-	-	-
		N2					-		-
	Tertiary treated	N1, N2, N3					-	-	-
Totana	Influent	N1	-		-			-	-
		N2	-		-	-		-	-
		N3	-		-			-	-
	Secondary treated	N1, N2, N3					-	-	-

Ct scale: | 34 | 35 | 36 | 37 | 38 | 39 | 40 |

477   **Table 2.** Epidemiological data[1] summary of COVID-19 cases in the area of study.

	20/03/2020		25/03/2020		30/03/2020		08/04/2020		15/04/2020	
	Cases	Prevalence[2]	Cases	Prevalence	Cases	Prevalence	Cases	Prevalence	Cases	Prevalence
Murcia	96	21.18	210	46.33	332	73.2	551	121.6	622	137.2
Cartagena	36	16.76	64	29.79	111	51.7	163	75.9	190	88.5
Molina de Segura	12	16.69	26	36.17	40	55.6	60	83.5	70	97.4
Lorca	-	-	8	8.47	18	19.1	29	30.7	31	32.8
Cieza	-	-	12	34.30	22	62.9	45	128.6	49	140.0
Totana	-	-	-	-	7	21.9	13	40.6	14	43.7

479   [1,] Data retrieved from the public repository of the "Servicio de epidemiologia" of the "Consejería de Salud de la Región de Murcia" (available at

480   http://www.murciasalud.es/principal.php).

481   [2,] Prevalence, percentage of diagnosed cases per 100.000 inhabitants.

482   **Figure 3.** Epidemiological surveillance of COVID-19 by wastewater SARS-CoV-2 RT-

483   qPCR in six municipalities.