Extensive weight loss can reduce immune age by altering IgG N-glycosylation

Valentina L Greto, PhD[1]*, Ana Cvetko, MSc[2]*, Tamara Štambuk, PhD[2,3]*, Niall J Dempster, MD[4], Domagoj Kifer, MSc[2], Helena Deriš, MSc[3], Ana Cindrić, MSc[3], Frano Vučković, PhD[3], Mario Falchi, PhD[5], Richard S Gillies, MD[6], Jeremy W Tomlinson, PhD[4], Olga Gornik, PhD[2,3], Bruno Sgromo, MD[6], Tim D Spector, PhD[5], Cristina Menni, PhD[5]*, Alessandra Geremia, PhD[1]*, Carolina V Arancibia-Cárcamo, PhD[1]*, Gordan Lauc, PhD[2,3]*

[1]Translational Gastroenterology Unit and NIHR Oxford Biomedical Research Centre, Nuffield Department of Medicine, University of Oxford, United Kingdom

[2]Faculty of Pharmacy and Biochemistry, University of Zagreb, Zagreb, Croatia

[3]Genos Glycoscience Research Laboratory, Zagreb, Croatia

[4]Oxford Centre for Diabetes and NIHR Oxford Biomedical Research Centre, Endocrinology and Metabolism, Radcliffe Department of Medicine, University of Oxford, United Kingdom

[5]The Department of Twin Research, King's College London, St Thomas' Hospital, London, United Kingdom

[6]Department of Upper GI Surgery, Oxford University Hospitals, United Kingdom

*The authors equally contributed to this work

Running title: Weight loss improves aging and IgG N-glycosylation

Corresponding author:
Gordan Lauc
glauc@pharma.hr
Faculty of Pharmacy and Biochemistry, University of Zagreb
A. Kovačića 1, 10 000 Zagreb, Croatia

AUTHORS` DECLARATION OF INTEREST

Tamara Štambuk Declaration of interest: Dr. Štambuk reports that she is an employee of

Genos Glycoscience Research Laboratory which offers commercial service of glycomic

analysis and has several patents in the field.

Helena Deriš Declaration of interest: Helena Deriš reports that she is an employee of

Genos Glycoscience Research Laboratory which offers commercial service of glycomic

analysis and has several patents in the field.

Ana Cindrić Declaration of interest: Ana Cindrić reports that she is an employee of Genos

Glycoscience Research Laboratory which offers commercial service of glycomic analysis

and has several patents in the field.

Frano Vučković Declaration of interest: Dr. Vučković reports that he is an employee of

Genos Glycoscience Research Laboratory which offers commercial service of glycomic

analysis and has several patents in the field.

Olga Gornik Declaration of interest: Dr. Gornik reports that she is an employee of Genos

Glycoscience Research Laboratory which offers commercial service of glycomic analysis

and has several patents in the field.

Alessandra Geremia Declaration of interest: Dr. Geremia reports grants from Wellcome

Trust, grants from NIHR research capability fund, during the conduct of the study; other

from UCB Pharma, outside the submitted work.

Gordan Lauc Declaration of interest: Dr. Lauc reports that he is founder and owner of

Genos LTD Zagreb; In addition, Dr. Lauc has a multiple patents in the field of glycoscience

pending or issued.

Valentina L Greto, Ana Cvetko, Niall J Dempster, Mario Falchi, Cristina Menni, Jeremy W

Tomlinson, Domagoj Kifer, Bruno Sgromo, Richard S Gillies, Tim Spector, Cristina Menni,

Carolina V Arancibia-Cárcamo Declarations of interest: none

ABSTRACT

Background

Obesity is a major global health problem, and is associated with increased cardiometabolic

morbidity and mortality. Protein glycosylation is a frequent postranslational modification,

highly responsive to numerous pathophysiological conditions and ageing. The prospect of

biological age reduction, by reverting glycosylation changes through metabolic intervention,

opens many possibilities. We have investigated whether weight loss interventions affect

inflammation- and ageing-associated IgG glycosylation changes, in a longitudinal cohort of

bariatric surgery patients. To support potential findings, BMI-related glycosylation changes

were monitored in a longitudinal twins cohort.

Methods

IgG N-glycans were chromatographically profiled in 37 obese patients, subjected to low-

calorie diet, followed by bariatric surgery, across multiple timepoints. Similarly, plasma-

derived IgG N-glycan traits were longitudinally monitored in 1,680 participants from the

TwinsUK cohort.

Results

Low-calorie diet induced a marked decrease in the levels of IgG N-glycans with bisecting

GlcNAc, whose higher levels are usually associated with ageing and inflammatory conditions.

Bariatric surgery resulted in extensive alterations of the IgG glycome that accompanied

progressive weight loss during one-year follow-up. We observed a significant increase in

digalactosylated and sialylated glycans, and a substantial decrease in agalactosylated and core

fucosylated IgG glycans. In general, this IgG glycan profile is associated with a younger

biological age and reflects an enhanced anti-inflammatory IgG potential. Loss of BMI over a

20 year period in the TwinsUK cohort validated a weight loss-associated agalactosylation

decrease and an increase in digalactosylation.

Conclusions

Altogether, these findings highlight that weight loss substantially affects IgG N-glycosylation,

resulting in reduced biological and immune age.

GRAPHICAL ABSTRACT

76
77 **HIGHLIGHTS**
78
79 • Obesity is associated to inflammation-related agalactosylated and bisected IgG
80   glycoforms

81
82 • IgG galactosylation and sialylation increase after bariatric surgery-induced weight loss

83
84 • Progressive decrease of BMI is associated to increased IgG galactosylation, implying a
85   reduction of biological age

86 **KEYWORDS**

87 N-glycosylation; bariatric surgery; weight loss; immunoglobulin G; biological age

INTRODUCTION

The global prevalence of obesity has risen dramatically in the past decades, and it is now considered a pandemic [1]. According to the World Health Organization, over 650 million individuals are obese, accounting for 13% of the world's adult population. Obesity confers a risk for metabolic syndrome, contributing to type 2 diabetes and cardiovascular disease (CVD) development [2]. Metabolic syndrome is linked to a chronic systemic low-grade inflammation, which contributes to the aging of the immune system denoted as inflammaging [3,4]. Obesity-related inflammaging results in impaired innate and adaptive immune function, and is characterized by high serum levels of IL-6, TNF-α and CRP [5]. Altered protein N-glycosylation is one of the hallmarks of inflammaging [4,6]. The human circulating N-glycome represents the entire set of glycans that are covalently attached to plasma proteins through a nitrogen on an asparagine residue. N-glycans are essential for life and are involved in many physiological processes [7], including signal transduction, protein trafficking and folding, receptor regulation and cell adhesion. Glycosylation has a fundamental role in the innate and adaptive immune responses, accentuated by the fact that all five classes of immunoglobulins (Ig) bear N-glycans. In this regard, IgG is probably the most investigated glycoprotein, whose effector functions are controlled by its Fc-bound glycans [8].

Inter-individual differences in the pace of biological ageing is an intriguing concept which may contribute to why some people stay healthy until their late chronological age, while others age faster and have shorter life expectancy. Progressive age-related changes of IgG glycosylation have been extensively studied [7,9,10] and the GlycanAge model has been proposed to express the difference between chronological and IgG glycome ageing [11]. The age of the IgG glycome might be estimated through the levels of agalactosylated species, which increase with ageing and are associated with enhanced immune activation [12]. The

opposite applies for digalactosylated IgG glycoforms, which are usually related to a younger age. Besides age-related changes, specific IgG glycosylation patterns have been already associated with CVD risk score and subclinical atherosclerosis in two large independent UK cohorts [13]. Moreover, a prospective follow-up of the EPIC-Potsdam cohort confirmed that changes in plasma N-glycome composition are predictive of future CVD events, with comparable predictive power to the American Heart Association (AHA) score in men and even better predictive power in women [14]. The link between a proinflammatory IgG N-glycome and hypertension has also been extensively studied [15–17], and similar IgG glycosylation patterns were associated with increased body mass index (BMI) and measures of central adiposity [18,19].

Studies in mouse models further corroborate the importance of differential IgG glycoforms in CVD pathogenesis. It has been shown that hyposialylated IgG (corresponding to an old-like IgG glycome) can induce obesity-related hypertension and insulin resistance in B-cell-deficient mice, through activation of the endothelial FcγRIIB [20,21]. These findings indicate that the IgG N-glycome could represent more than a biomarker of inflammation and aging, since distinctive IgG glycoforms act as effector molecules in certain pathologies. Furthermore, supplementation with *N*-acetylmannosamine (ManNAc), a precursor of sialic acid, protects obese mice from hypertension and insulin resistance induction by reverting an IgG N-glycome associated with old age into an IgG N-glycome associated with young age [21,22]. However, studies exploring the possibilities of converting an old-like into a young-like IgG glycome by metabolic intervention in humans are limited. Of note, only one small study indicated that high-intensity interval training can rejuvenate the IgG N-glycome [23].

Bariatric surgery is very effective for the treatment of severe obesity [24]. The resulting weight loss impacts energy balance and metabolism, contributing to the increased insulin

response, improved glycaemic control and reduction of total body fat, leading to decreased

CVD risk and mortality [25].

In this study we aimed to determine whether weight loss affects glycan markers related to

inflammation and ageing, in a longitudinally-monitored cohort of obese individuals

undergoing low-calorie diet and then bariatric surgical interventions. We also investigated

BMI-related glycosylation changes in the longitudinal TwinsUK cohort, the largest cohort of

adult twins with the most detailed clinical database in the world.

METHODS

Study populations

Bariatric cohort. The cohort included 37 participants, recruited at Oxford University Hospitals

to the Gastrointestinal Illnesses study (Ref: 16/YH/0247). All patients were characterised by

metabolic status and medical history. Bariatric patients were considered eligible in accordance

to National Institute for Health and Care Excellence (NICE) and local guidelines.

Patients with a history of alcoholism and/or ongoing anticoagulant treatment were excluded.

Patients were also excluded in case of pregnancy, active substance abuse or uncontrolled

psychiatric condition including eating disorders. Participants were sampled at baseline and

subjected to 3-week low calorie carbohydrate-restricted diet (900 cal, maximum 100 g of

carbohydrates per day), followed by bariatric surgery. The sequential follow-up timepoints

included the day of the surgery (baseline), at 20% of weight loss after 6.54 ± 3.4 months (mean

± IQR) and 12.47 ± 6.55 months post-op. Characteristics of the bariatric cohort are shown in

Table 1.

TwinsUK cohort. We have analysed a total of 6,032 plasma samples from 2,146 participants of

the TwinsUK study, collected at multiple timepoints over a 20 year-period [26]. These included

1,865 individuals sampled at 3 timepoints, 156 individuals sampled at 2 timepoints and 125

individuals sampled only once. Following the plasma N-glycome analysis, glycan data

underwent quality control (see *Statistical analysis* section), which decreased the dataset to

5,889 samples (measurements). Out of these 5,889 measurements, we have proceeded with

statistical analysis on a subset of 3,742 samples (measurements) which had information on

BMI available. Description of the TwinsUK cohort is provided in Table 2.

Ethical statement

Ethical approval for the study was obtained by the National Research Ethics Committees of

the UK National Health Service (NHS) under the reference number 16/YH/0247. All individuals

participating in this study gave written informed consent. The TwinsUK study was approved

by NRES Committee London–Westminster, and all twins provided informed written consent.

N-glycome analysis

Isolation of IgG from human plasma

IgG was isolated from plasma samples by affinity chromatography as described previously

[27]. In brief, IgG was isolated in a high-throughput manner, using 96-well protein G

monolithic plates (BIA Separations, Slovenia), starting from 100 µl of plasma. Plasma was

diluted 7× with phosphate buffered saline (PBS; Merck, Germany) and applied to the protein

G plate. IgG was eluted with 1 ml of 0.1 M formic acid (Merck, Germany) and immediately

neutralized with 1 M ammonium bicarbonate (Acros Organics, USA).

N-glycan release from IgG and total plasma proteins

Isolated IgG samples were dried in a vacuum centrifuge. After drying, IgG was denatured with

the addition of 30 µl of 1.33% SDS (w/v) (Invitrogen, USA) and by incubation at 65 °C for 10

min. Plasma samples (10 µl) were denatured with the addition of 20 µl of 2% (w/v) SDS

(Invitrogen, USA) and by incubation at 65 °C for 10 min. From this point on, the procedure was

identical for both IgG and plasma samples. After denaturation, 10 µl of 4% (v/v) Igepal-CA630

(Sigma Aldrich, USA) was added to the samples, and the mixture was shaken 15 min on a plate

shaker (GFL, Germany). N-glycans were released with the addition of 1.2 U of PNGase F

(Promega, USA) and overnight incubation at 37 °C.

Fluorescent labelling and HILIC SPE clean-up of released N glycans

The released N-glycans were labelled with 2-aminobenzamide (2-AB). The labelling mixture

consisted of 2-AB (19.2 mg/ml; Sigma Aldrich, USA) and 2-picoline borane (44.8 mg/ml; Sigma

Aldrich, USA) in dimethyl sulfoxide (Sigma Aldrich, USA) and glacial acetic acid (Merck,

Germany) mixture (70:30 v/v). To each sample 25 µl of labelling mixture was added, followed

by 2 h incubation at 65 °C. Free label and reducing agent were removed from the samples

using hydrophilic interaction liquid chromatography solid-phase extraction (HILIC-SPE). After

incubation samples were brought to 96% of acetonitrile (ACN) by adding 700 µl of ACN (J.T.

Baker, USA) and applied to each well of a 0.2 µm GHP filter plate (Pall Corporation, USA).

Solvent was removed by application of vacuum using a vacuum manifold (Millipore

Corporation, USA). All wells were prewashed with 70% ethanol (Sigma-Aldrich, St. Louis, MO,

USA) and water, followed by equilibration with 96% ACN. Loaded samples were subsequently

washed 5× with 96% ACN. N-glycans were eluted with water and stored at − 20 °C until usage.

Hydrophilic interaction liquid chromatography of N-glycans

Fluorescently labelled N-glycans were separated by hydrophilic interaction liquid

chromatography (HILIC) on Acquity UPLC H-Class instrument (Waters, USA) consisting of a

quaternary solvent manager, sample manager and a fluorescence detector, set with excitation

and emission wavelengths of 250 and 428 nm, respectively. The instrument was under the

control of Empower 3 software, build 3471 (Waters, Milford, USA). Labelled N-glycans were

separated on a Waters BEH Glycan chromatography column, with 100 mM ammonium

formate, pH 4.4, as solvent A and ACN as solvent B. In the case of IgG N-glycans, separation

method used linear gradient of 75–62% acetonitrile at flow rate of 0.4 ml/min in a 27-min

analytical run. For plasma protein N-glycans separation method used linear gradient of 70–

53% acetonitrile at flow rate of 0.561 ml/min in a 25-min analytical run. The system was

calibrated using an external standard of hydrolysed and 2-AB labelled glucose oligomers from

which the retention times for the individual glycans were converted to glucose units (GU).

Data processing was performed using an automatic processing method with a traditional

integration algorithm after which each chromatogram was manually corrected to maintain

the same intervals of integration for all the samples. The chromatograms were all separated

in the same manner into 24 peaks (GP1– GP24) for IgG N-glycans and 39 peaks (GP1–GP39)

for plasma protein N-glycans and are depicted in Supplementary Figure 1 and Supplementary

Figure 2, respectively. Detailed description of glycan structures corresponding to each glycan

peak is presented in Supplementary Table 1. Glycan peaks were analysed based on their

elution positions and measured in glucose units, then compared to the reference values in the

"GlycoStore" database (available at: https://glycostore.org/) for structure assignment. The

amount of glycans in each peak was expressed as a percentage of the total integrated area.

For IgG glycans, in addition to 24 directly measured glycan traits, 8 derived traits were

calculated (Supplementary Table 2). In the case of TwinsUK cohort, IgG N-glycan traits were

calculated from plasma protein glycan profiles, based on known elution positions of

predominat IgG glycan structures (Supplementary Table 3). In general, derived glycan traits

average particular glycosylation features, such as galactosylation, fucosylation, bisecting

GlcNAc, and sialylation.

Statistical analysis

Bariatric cohort. In order to remove experimental variation from the measurements, normalization and batch correction were performed on the UPLC glycan data. To make measurements across samples comparable, normalization by total area was performed. Prior to batch correction, normalized glycan measurements were log-transformed due to right-skewness of their distributions and the multiplicative nature of batch effects. Batch correction was performed on log-transformed measurements using the ComBat method (R package sva [28], where the technical source of variation (which sample was analysed on which plate) was modelled as batch covariate. To correct measurements for experimental noise, estimated batch effects were subtracted from log-transformed measurements.

Longitudinal analysis of patient samples through their observation period was performed by implementing a linear mixed effects model, where time was modelled as fixed effect, while the individual ID was modelled as random effect. Prior to the analyses, glycan variables were all transformed to standard Normal distribution by inverse transformation of ranks to Normality (R package "GenABEL", function rntransform). Using rank transformed variables makes estimated effects of different glycans comparable, as these will have the same standardized variance. False discovery rate (FDR) was controlled by the Benjamini-Hochberg procedure at the specified level of 0.05. Data was analysed and visualized using R programming language (version 3.5.2)[29].

TwinsUK cohort. Normalization of peak intensities to the total chromatogram area was performed for each measured sample separately. Calculated proportions were then batch corrected using ComBat method (R package *sva*)[28]. After the batch correction the first 11 peaks, which predominantly originate from IgG [30], were used to calculate 6 derived glycan traits – agalactosylation (G0), monogalactosylation (G1), digalactosylation (G2), bisecting

GlcNAc (B), core fucosylation (CF) and high mannose structures (HM). Mixed models were

fitted to estimate the effect of BMI change on IgG N-glycome (R package *lme4*)[31]. Directly

measured or derived glycan trait was used as a dependent variable in the mixed model. To

differentiate between BMI change and the absolute BMI value, the variable was separated to

BMI_{baseline} and BMI_{difference} (calculated according to the following equation: $BMI_{difference} =$

$BMI_{follow\ up\ age} - BMI_{baseline\ age}$), and both were used in the model as a fixed effect. Since

IgG N-glycome is affected by aging, age was included both as a fixed effect and a random

slope. Finally, to meet the independency criteria, family ID and individual ID (nested within

family) were included in the model as a random intercept. Due to multiple model fitting (for

11 directly measured and 6 derived glycan traits) false discovery rate was controlled using

Benjamini-Hochberg method. All statistical analyses were performed using R programming

language (version 3.6.3)[29].

RESULTS

Impact of pre-surgical low-calorie diet on IgG glycosylation

We chromatographically profiled the IgG N-glycome in a cohort of bariatric surgery-candidate

patients before and after the pre-operative diet. By employing a linear mixed model, we

observed significant change in only one out of eight examined IgG derived glycan traits.

Namely, the levels of bisecting GlcNAc (B) were substantially decreased after the low-calorie

diet intervention (Table 3), indicating a decreased proinflammatory potential of the circulating

IgG. The other IgG glycosylation features did not exhibit significant alterations, possibly due

to rather short follow-up period (3 weeks) and limited number of participants (n=8) (Table

3).Graphical representation of the longitudinal alterations in IgG N-glycome after low-calorie

diet are depicted in Supplementary Figure 3.

IgG N-glycosylation markedly changes after weight loss surgery

Using the same chromatographic approach, we analysed samples from patients who underwent bariatric surgery. The plasma samples were collected on the day of surgery (month 0), approximately 6 months post-surgery and 12 months post-surgery. IgG N-glycans were profiled in each of these timepoints, and the obtained values were used for derived glycan traits calculations. Statistical analysis revealed extensive changes in IgG N-glycome following the bariatric procedure. Namely, four out of eight tested derived traits showed marked changes: core fucosylated (CF) and agalactosylated (G0) glycans decreased, while digalactosylated (G2) and monosialylated (S1) glycans increased after the surgery (Table 4). The IgG glycans whose abundances were increased after bariatric surgery are major components of a young IgG glycome, as they are typically associated with a younger age. The opposite applies to agalactosylated structures, which are usual denominators of an old-like IgG glycome profile. We also examined the correlation of patients' clinical data with IgG N-glycome features using multivariate analysis, but found no statistically significant associations (Supplementary Table 4). Finally, the type of bariatric surgery (either sleeve gastrectomy or Roux-en-Y gastric bypass) did not affect IgG glycome composition. Graphical representations of the longitudinal alterations in IgG glycosylation features are depicted in Figure 1.

Figure 1 Bariatric surgery-related alterations in IgG N-glycome over time (months). Red line – significant decrease; green line – significant increase; blue line – non-significant change.

Weight loss induces a shift towards young-like IgG N-glycome

Using the same chromatographic approach, we profiled the plasma protein N-glycome from 1,680 TwinsUK study participants sampled at several timepoints over a 20-year-period. This served as a replication of the findings from the bariatric cohort, whose participants exhibited the reversal from old- to young-likeIgG N-glycome due to weight loss. Due to the fact that for TwinsUK cohort plasma glycome was profiled, we calculated derived traits and performed statistical analysis based only on the first 11 plasma glycan peaks, corresponding to the glycans predominantly originating from IgG [30]. We examined IgG glycome alterations associated with changes in BMI using a mixed model on a subset of 3,742 samples. Out of six examined IgG glycosylation features (derived traits), three displayed significant BMI-decrease-related

changes – agalactosylation (G0), digalactosylation (G2) and incidence of high mannose

structures (HM) (Table 5). Namely, the abundance of digalactosylated (G2) glycans increased

with the BMI decrease, while the abundance of agalactosylated (G0) and high mannose

glycans (HM) decreased with the weight loss, estimated by the BMI drop. These findings are

in line with the results observed in the bariatric surgery cohort. Graphical representation of

the longitudinal BMI-dependent alterations of IgG glycosylation are depicted at Figure 2.

Figure 2 BMI-associated alterations in IgG N-glycosylation across multiple timepoints. Changes in IgG

derived traits are presented with lineplots of hypothetical ageing of TwinsUK participants (all women).

Black dot represents a starting point of a 30-year-old woman, black triangle of a 40-year-old woman

and black square of a 50-year-old woman. All of these women have a baseline BMI of 25 kg m^{-2}. Blue

lines represent age-related IgG glycosylation changes attributed to stabile BMI. Green lines represent

age-related IgG glycosylation changes attributed to increasing BMI (0.5 kg m^{-2} per year, through a

period of 10 years). Red lines represent age-related IgG glycosylation changes attributed to decreasing

BMI (0.5 kg m^{-2} per year, through a period of 10 years). Highlighted areas represent 95% confidence

intervals. The effect of age on IgG glycosylation is represented with the curve slope, while the effect

of BMI change is represented with the distance of green/red line from the blue line.

DISCUSSION

In this study, we have observed extensive changes in the IgG N-glycome associated with

weight loss following a low-calorie diet, bariatric surgery, or a decrease of BMI throughout

time. To the best of our knowledge, this is the first study to investigate IgG N-glycome

alterations in patients who underwent a low-calorie diet followed by bariatric surgery.

Prior to bariatric surgery, patients were subjected to a 3-week low-calorie diet which induced

a single significant change to the IgG N-glycome. Specifically, the levels of bisecting GlcNAc

were found to be reduced after the dieting period. In general, higher levels of bisecting GlcNAc

are associated with enhanced affinity for FcγRs and, consequently, with enhanced antibody-

dependent cellular cytotoxicity (ADCC) and other effector functions of the immune cells [9].

Hence, the reduction of bisecting GlcNAc levels on IgG decreases IgG inflammatory potential.

Furthermore, several studies have reported a sex-independent increase in the levels of

bisecting GlcNAc with age [32,33], suggesting that this diet-related decrease of its levels also

contributes to the reduction of the biological age. Lastly, increased abundance of bisecting

GlcNAc on IgG has been previously associated with type 2 diabetes [34] and with higher

cardiovascular risk [13], implying that dieting improves individual's cardiometabolic health

through altered IgG glycosylation as well. As for the other examined IgG glycosylation traits,

no significant changes were found to be associated with a low-calorie diet, probably due to

the relatively short follow-up period (three weeks), which matches IgG serum half-life.

Moreover, only 8 participants were subjected to the low-calorie diet intervention, hence,

further analysis on a larger sample size might reveal additional relevant changes.

We have also analysed IgG N-glycome from individuals who underwent bariatric surgery, in a

longitudinal manner. We observed several significant changes in IgG N-glycome, such as a

marked decrease in agalactosylated (G0) IgG. Elevated levels of G0 IgG glycoforms are

typically associated with aging, pro-inflammatory IgG glycan profile and various inflammatory

diseases [9]. On the other hand, the levels of digalactosylated (G2) glycans increased after

bariatric surgery and at sequential timepoints, in accordance with a reduced inflammatory

potential of the circulating IgG. The increased levels of IgG galactosylation were previously

associated with a younger biological age and are considered, in a way, as a measure of an

individual's well-being [9,12]. Converesely, IgG galactosylation levels substantially decrease

with ageing and during inflammation [9,11]. Our results demonstrate that weight loss,

resulting from bariatric surgery, can initiate the reversal from an old-like to a young-like IgG

N-glycome, potentially reversing the clock for the immune/biological age. Furthermore,

bariatric surgery also resulted in significant IgG glycome alteration inducing a decrease in core

fucosylation. The vast majority of circulating IgG molecules bears core fucose (approximately

95%), which profoundly decreases IgG binding affinity to FcγRIIIA receptor and sequential

ADCC [35]. ADCC is largely mediated by natural killer cells that can lyse target cells and fight

viral infections. This would suggest that an extensive weight loss ameliorates immune

responses against pathogens, by altering IgG glycosylation and modulating its effector

functions. Lastly, bariatric surgery-related weight loss led to an increase in IgG sialylation,

which is the main modulator of the IgG anti-inflammatory actions [36]. In addition to its anti-

inflammatory actions, the level of IgG sialylation has been implicated in the pathogenesis of

obesity-induced insulin resistance and hypertension, as already mentioned [20,21]. Inhibitory

IgG receptor FcγRIIB was found to be expressed in the microvascular endothelium. Moreover,

it was shown that hyposialylated IgG acts as its operating ligand, leading to the induction of

obesity-related insulin resistance and hypertension. On the other hand, the sialylated

glycoform is not activating the signalling pathways responsible for the insulin resistance and

hypertension development, but is rather preserving insulin sensitivity and normal vasomotor

tone, even in obese mice. Interestingly, the same group made another significant discovery –

promotion of IgG sialylation breaks the link between obesity and hypertension/insulin

resistance [21,22]. Namely, supplementation with the sialic acid precursor restored IgG

sialylation, highlighting a potential approach to improve both metabolic and cardiovascular

health in humans, with a single intervention. Our data suggest that a similar effect might be

achieved by weight loss interventions, through restoring of IgG sialylation levels

In order to confirm the effects of weight loss on the biological immune age, we investigated

how a decreasing BMI affects the IgG N-glycome during a 20-year-period. We observed the

prominent inverse changes of agalactosylated (G0) and digalactosylated (G2) IgG glycans.

Namely, agalactosylated IgG glycans significantly decreased, while digalactosylated ones

substantially increased as the BMI decreased. These observations corroborated our findings

from the bariatric patients, confirming that the body weight reduction reverses IgG glycome

from old-like to young-like, implying at the same time a likely reduction in the biological and

immune age. Nonetheless, TwinsUK participants have not experienced such an extensive

weight loss, which potentially influenced the replication of other significant glycan changes

from the bariatric cohort; second – herein, the weight loss was approximated by BMI

decrease, which is usually a legitimate assumption, however, it does not have to apply to all

cases; and third –we profiled plasma glycome, while the IgG glycan traits were approximated

and the information on IgG sialylation was confounded by other plasma glycoproteins. Ideally,

these issues could be circumvented in the future studies whose experimental design would

allow the simultaneous, multi-centre follow-up of larger groups of patients.

Intense physical exercise can also shift IgG N-glycome towards a young-like profile by

increasing IgG galactosylation [23]. Although another study reported that prolonged intensive

exercise can have the opposite effect and promote pro-inflammatory changes of IgG N-

glycome [37], its findings are not surprising since it recruited healthy, normal-weight female

participants, subjected to the intense energy deprivation and exercise levels, to induce

substantial fat loss in a rather short time period. The authors also highlighted that starting

weight and the way in which weight loss is achieved could be crucial for the final effect on the

immune system [37]. Therefore, it seems that exercise overall has a positive impact on the

immune system and biological clock, but its intensity and duration should be personalized in

order to achieve the optimal results.

To summarize, our results indicate that both dieting and bariatric surgery have an impact on

inflammation and biological aging by altering IgG N-glycan patterns. All of the observed

weight-loss-associated alterations in IgG glycosylation are suggesting a decreased

inflammatory potential of the circulating IgG and a reduction of biological age. Hence,

improving metabolic and endocrine health through weight loss consequently also contributes

to the preservation of a healthy immune system.

ACKNOWLEDGMENTS

This research was funded by the National Institute for Health Research (NIHR) Oxford

Biomedical Research Centre (BRC). The views expressed are those of the author(s) and not

necessarily those of the NHS, the NIHR or the Department of Health. The authors thank Rachel

Franklin, Michelle Haylock, James Chivenga, Roxanne Williams and the BRC Oxford GI Biobank

for sample collection. The authors thank all the patients who took part in this study.

AUTHORS` CONTRIBUTION

Cristina Menni, Alessandra Geremia, Carolina V Arancibia-Cárcamo and Gordan Lauc:

Conceptualization, Supervision, Project administration, Funding acquisition. Valentina L

Greto, Ana Cvetko, Tamara Štambuk, Niall J Dempster, Helena Deriš, Ana Cindrić, Mario Falchi,

Cristina Menni: Investigation. Domagoj Kifer, Frano Vučković: Methodology, Software, Formal

analysis. Valentina L Greto, Ana Cvetko, Tamara Štambuk, Niall J Dempster, Domagoj Kifer,

Frano Vučković , Mario Falchi, Jeremy W Tomlinson, Olga Gornik, Tim Spector, Cristina Menni,

Alessandra Geremia, Carolina V Arancibia-Cárcamo, Gordan Lauc: Data curation and

visualization: Valentina L Greto, Ana Cvetko, Tamara Štambuk, Niall J Dempster, Domagoj

Kifer, Frano Vučković, Gordan Lauc: Writing-original draft. Valentina L Greto, Ana Cvetko,

Tamara Štambuk, Niall J Dempster, Domagoj Kifer, Helena Deriš, Ana Cindrić, Frano Vučković,

Mario Falchi, Richard S Gillies, Jeremy W Tomlinson, Olga Gornik, Bruno Sgromo, MD6, Tim D

Spector, Cristina Menni, Alessandra Geremia, Carolina V Arancibia-Cárcamo, Gordan Lauc:

Review & Editing.

REFERENCES

[1]   Obesity and overweight https://www.who.int/news-room/fact-sheets/detail/obesity-and-overweight (accessed April 16, 2020).

[2]   Han TS, Lean ME. A clinical perspective of obesity, metabolic syndrome and cardiovascular disease. JRSM Cardiovasc Dis 2016;5:2048004016633371. https://doi.org/10.1177/2048004016633371.

[3]   Alpert A, Pickman Y, Leipold M, Rosenberg-Hasson Y, Ji X, Gaujoux R, et al. A clinically meaningful metric of immune age derived from high-dimensional longitudinal monitoring. Nat Med 2019;25:487–95. https://doi.org/10.1038/s41591-019-0381-y.

[4]   Franceschi C, Garagnani P, Parini P, Giuliani C, Santoro A. Inflammaging: a new immune–metabolic viewpoint for age-related diseases. Nature Reviews Endocrinology 2018;14:576–90. https://doi.org/10.1038/s41574-018-0059-4.

[5]   Touch S, Clément K, André S. T Cell Populations and Functions Are Altered in Human Obesity and Type 2 Diabetes. Curr Diab Rep 2017;17:81. https://doi.org/10.1007/s11892-017-0900-5.

[6]   Lauc G, Sinclair D. Biomarkers of biological age as predictors of COVID-19 disease severity. Aging 2020. https://doi.org/10.18632/aging.103052.

[7]   Lauc G, Pezer M, Rudan I, Campbell H. Mechanisms of disease: The human N-glycome. Biochimica et Biophysica Acta - General Subjects 2016;1860:1574–82. https://doi.org/10.1016/j.bbagen.2015.10.016.

[8]   Gornik O, Pavić T, Lauc G. Alternative glycosylation modulates function of IgG and other proteins - implications on evolution and disease. Biochim Biophys Acta 2012;1820:1318–26. https://doi.org/10.1016/j.bbagen.2011.12.004.

[9]   Gudelj I, Lauc G, Pezer M. Immunoglobulin G glycosylation in aging and diseases. Cellular Immunology 2018. https://doi.org/10.1016/j.cellimm.2018.07.009.

[10] Dall'Olio F. Glycobiology of aging. Subcellular Biochemistry, 2018. https://doi.org/10.1007/978-981-13-2835-0_17.

[11] Krištić J, Vučković F, Menni C, Klarić L, Keser T, Beceheli I, et al. Glycans Are a Novel Biomarker of Chronological and Biological Ages. The Journals of Gerontology: Series A 2014;69:779–89. https://doi.org/10.1093/gerona/glt190.

[12] Štambuk J, Nakić N, Vučković F, Pučić-Baković M, Razdorov G, Trbojević-Akmačić I, et al. Global variability of the human IgG glycome. Biochemistry; 2019. https://doi.org/10.1101/535237.

[13] Menni C, Gudelj I, MacDonald-Dunlop E, Mangino M, Zierer J, Bešić E, et al. Glycosylation Profile of Immunoglobulin G Is Cross-Sectionally Associated with Cardiovascular Disease Risk Score and Subclinical Atherosclerosis in Two Independent Cohorts. Circulation Research 2018;122:1555–64. https://doi.org/10.1161/CIRCRESAHA.117.312174.

[14] Wittenbecher C, Štambuk T, Kuxhaus O, Rudman N, Vučković F, Štambuk J, et al. Plasma N-Glycans as Emerging Biomarkers of Cardiometabolic Risk: A Prospective Investigation in the EPIC-Potsdam Cohort Study. Diabetes Care 2020;43:661–8. https://doi.org/10.2337/dc19-1507.

[15] Gao Q, Dolikun M, Stambuk J, Wang H, Zhao F, Yiliham N, et al. Immunoglobulin G N-Glycans as Potential Postgenomic Biomarkers for Hypertension in the Kazakh Population. Omics 2017;21:380–9. https://doi.org/10.1089/omi.2017.0044.

[16] Liu J, Dolikun M, Štambuk J, Trbojević-Akmačić I, Zhang J, Wang H, et al. The association between subclass-specific IgG Fc N-glycosylation profiles and hypertension in the

490   Uygur, Kazak, Kirgiz, and Tajik populations. J Hum Hypertens 2018;32:555–63.
491   https://doi.org/10.1038/s41371-018-0071-0.

[17]   Wang Y, Klarić L, Yu X, Thaqi K, Dong J, Novokmet M, et al. The Association Between
493   Glycosylation of Immunoglobulin G and Hypertension: A Multiple Ethnic Cross-Sectional
494   Study. Medicine 2016;95:e3379. https://doi.org/10.1097/MD.0000000000003379.

[18]   Nikolac Perkovic M, Pucic Bakovic M, Kristic J, Novokmet M, Huffman JE, Vitart V, et al.
496   The association between galactosylation of immunoglobulin G and body mass index.
497   Progress in Neuro-Psychopharmacology and Biological Psychiatry 2014;48:20–5.
498   https://doi.org/10.1016/j.pnpbp.2013.08.014.

[19]   Russell AC, Kepka A, Trbojević-Akmačić I, Ugrina I, Song M, Hui J, et al. Increased central
500   adiposity is associated with pro-inflammatory immunoglobulin G N-glycans.
501   Immunobiology 2019;224:110–5. https://doi.org/10.1016/j.imbio.2018.10.002.

[20]   Sundgren NC, Vongpatanasin W, Boggan BMD, Tanigaki K, Yuhanna IS, Chambliss KL, et
503   al. IgG receptor FcγRIIB plays a key role in obesity-induced hypertension. Hypertension
504   2015;65:456–62. https://doi.org/10.1161/HYPERTENSIONAHA.114.04670.

[21]   Tanigaki K, Sacharidou A, Peng J, Chambliss KL, Yuhanna IS, Ghosh D, et al.
506   Hyposialylated IgG activates endothelial IgG receptor FcγRIIB to promote obesity-
507   induced insulin resistance. J Clin Invest 2018;128:309–22.
508   https://doi.org/10.1172/JCI89333.

[22]   Peng J, Vongpatanasin W, Sacharidou A, Kifer D, Yuhanna IS, Banerjee S, et al.
510   Supplementation with the Sialic Acid Precursor N-acetyl-D-Mannosamine Breaks the
511   Link Between Obesity and Hypertension. Circulation 2019.
512   https://doi.org/10.1161/circulationaha.119.043490.

[23]   Tijardović M, Marijančević D, Bok D, Kifer D, Lauc G, Gornik O, et al. Intense Physical
514   Exercise Induces an Anti-inflammatory Change in IgG N-Glycosylation Profile. Frontiers
515   in Physiology 2019;10:1–10. https://doi.org/10.3389/fphys.2019.01522.

[24]   Nguyen NT, Kim E, Vu S, Phelan M. Ten-year Outcomes of a Prospective Randomized
517   Trial of Laparoscopic Gastric Bypass Versus Laparoscopic Gastric Banding. Ann Surg
518   2018;268:106–13. https://doi.org/10.1097/SLA.0000000000002348.

[25]   O'Brien P. Bariatric surgery and type 2 diabetes: a step closer to defining an optimal
520   approach. The Lancet Diabetes & Endocrinology 2019;7:889–91.
521   https://doi.org/10.1016/S2213-8587(19)30352-3.

[26]   Verdi S, Abbasian G, Bowyer RCE, Lachance G, Yarand D, Christofidou P, et al. TwinsUK:
523   The UK Adult Twin Registry Update. Twin Res Hum Genet 2019;22:523–9.
524   https://doi.org/10.1017/thg.2019.65.

[27]   Pavić T, Dilber D, Kifer D, Selak N, Keser T, Ljubičić Đ, et al. N-glycosylation patterns of
526   plasma proteins and immunoglobulin G in chronic obstructive pulmonary disease. J
527   Transl Med 2018;16. https://doi.org/10.1186/s12967-018-1695-0.

[28]   Leek JT, Johnson WE, Parker HS, Jaffe AE, Storey JD. The sva package for removing
529   batch effects and other unwanted variation in high-throughput experiments.
530   Bioinformatics 2012;28:882–3. https://doi.org/10.1093/bioinformatics/bts034.

[29]   R Core Team (2020). R: A language and environment for statistical computing. R
532   Foundation for Statistical Computing, Vienna, Austria. URL   https://www.R-
533   project.org/. n.d.

[30]   Clerc F, Reiding KR, Jansen BC, Kammeijer GSM, Bondt A, Wuhrer M. Human plasma
protein N-glycosylation. Glycoconj J 2016;33:309–43. https://doi.org/10.1007/s10719-
015-9626-2.

[31]   Bates D, Mächler M, Bolker B, Walker S. Fitting Linear Mixed-Effects Models Using
lme4. Journal of Statistical Software 2015;67:1–48.
https://doi.org/10.18637/jss.v067.i01.
[32]   Ruhaak LR, Uh H-W, Beekman M, Koeleman CAM, Hokke CH, Westendorp RGJ, et al.
Decreased Levels of Bisecting GlcNAc Glycoforms of IgG Are Associated with Human
Longevity. PLOS ONE 2010;5:e12566. https://doi.org/10.1371/journal.pone.0012566.
[33]   Pučić M, Knežević A, Vidič J, Adamczyk B, Novokmet M, Polašek O, et al. High
Throughput Isolation and Glycosylation Analysis of IgG–Variability and Heritability of
the IgG Glycome in Three Isolated Human Populations. Mol Cell Proteomics 2011;10.
https://doi.org/10.1074/mcp.M111.010090.
[34]   Lemmers RFH, Vilaj M, Urda D, Agakov F, Šimurina M, Klaric L, et al. IgG glycan patterns
are associated with type 2 diabetes in independent European populations. Biochim
Biophys Acta Gen Subj 2017;1861:2240–9.
https://doi.org/10.1016/j.bbagen.2017.06.020.
[35]   Iida S, Kuni-Kamochi R, Mori K, Misaka H, Inoue M, Okazaki A, et al. Two mechanisms of
the enhanced antibody-dependent cellular cytotoxicity (ADCC) efficacy of non-
fucosylated therapeutic antibodies in human blood. BMC Cancer 2009;9:58.
https://doi.org/10.1186/1471-2407-9-58.
[36]   Kaneko Y, Nimmerjahn F, Ravetch JV. Anti-Inflammatory Activity of Immunoglobulin G
Resulting from Fc Sialylation. Science 2006;313:670–3.
https://doi.org/10.1126/science.1129594.
[37]   Sarin HV, Gudelj I, Honkanen J, Ihalainen JK, Vuorela A, Lee JH, et al. Molecular
Pathways Mediating Immunosuppression in Response to Prolonged Intensive Physical
Training, Low-Energy Availability, and Intensive Weight Loss. Front Immunol
2019;10:907. https://doi.org/10.3389/fimmu.2019.00907.

574 **Table 1 Demographic characteristics of the bariatric cohort**

Characteristics	*Bariatric cohort*			
Total No. of participants (N)	37			
No. of participants Sleeve Gastrectomy (SG), N (%)	25 (68%)			
No. Roux-en-Y Gastric Bypass RYGB N (%)	12 (32%)			
No. of participants each timepoint (N)	Pre-op low-calorie diet	Time of surgery	1st post-op timepoint	2nd post-op timepoint
	8	37	30	24
BMI* each timepoint, mean ± SD, kg/m2	Before diet	End of diet	Time of surgery	1st post-op timepoint	2nd post-op timepoint
	48.53 ± 4.13	46.60± 4.21	46.21 ± 4.75	36.01 ± 5.07	32.82 ± 5.17
Female, sex, N (%)	33 (89%)			
Type 2 diabetes, N (%)	6 (16%)			

576 *BMI reference values: <18.5 (underweight), 18.5–24.9 (normal weight), 25–29.9

577 (overweight), >30 (obese).

578
579

580　**Table 2 Demographic characteristics of the TwinsUK cohort**

Characteristics	TwinsUK cohort (BMI subset)
No. of participants (N)	1,680
No. of glycan measurements (N)	3,742
Baseline age, mean ± SD, years	53.23 ± 10.86
Follow up time, mean ± SD, years	7.90 ± 5.66
Female sex, N (%)	1,680 (100)
Baseline BMI, mean ± SD, kg/m^2	25.45 ± 4.53

581
582

583　**Table 3 Impact of pre-surgical low-calorie diet on IgG glycosylation.** Longitudinal analysis

584　was performed by implementing a linear mixed effects model, with time as a fixed effect and

585　the individual sample measurement as a random effect. False discovery rate was controlled

586　using Benjamini-Hochberg method at the specified level of 0.05.

Derived IgG glycan trait	Time_effect	Time_SE	Time_p-value	Adjusted p-value
bisecting GlcNAc (B)	-0.2801	0.0743	0.0043	0.0341
agalactosylation (G0)	-0.0493	0.1876	0.7930	0.7930
monogalactosylation (G1)	0.1122	0.1869	0.5526	0.7930
digalactosylation (G2)	-0.0539	0.1387	0.6991	0.7930
total sialylation (S)	0.0941	0.2090	0.6544	0.7930
monosialylation (S1)	0.1354	0.1878	0.4779	0.7930
disialylation (S2)	-0.0869	0.2474	0.7266	0.7930
core fucosylation (CF)	0.0810	0.2416	0.7385	0.7930

Red – significant decrease; blue – non-significant change. GlcNAc – *N*-acetylglucosamine; SE –

standard error

Table 4 Bariatric surgery induces significant changes in IgG N-glycome. Longitudinal analysis

was performed by implementing a linear mixed effects model, with time as a fixed effect and

the individual sample measurement as a random effect. False discovery rate was controlled

using Benjamini-Hochberg method at the specified level of 0.05.

Derived IgG glycan trait	*Time_effect*	*Time_SE*	*Time_p-value*	*Adjusted p-value*
agalactosylation (G0)	-0.0339	0.0078	9.23×10^{-05}	7.38×10^{-04}
digalactosylation (G2)	0.0275	0.0072	3.75×10^{-04}	1.50×10^{-03}
monosialylation (S1)	0.0193	0.0080	1.97×10^{-02}	3.94×10^{-02}
core fucosylation (CF)	-0.0155	0.0064	1.74×10^{-02}	3.94×10^{-02}
total sialylation (S)	0.0171	0.0083	4.70×10^{-02}	6.27×10^{-02}
bisecting GlcNAc (B)	0.0173	0.0083	4.01×10^{-02}	6.27×10^{-02}
monogalactosylation (G1)	0.0206	0.0107	8.18×10^{-02}	9.35×10^{-02}
disialylation (S2)	0.0048	0.0080	5.48×10^{-01}	5.48×10^{-01}

594   Red – significant decrease; green – significant increase; blue – non-significant change.

595   GlcNAc – *N*-acetylglucosamine; SE – standard error

604 **Table 5 Longitudinally monitored weight loss results with significant changes of IgG N-**

605 **glycosylation.** Longitudinal analysis was performed by implementing a mixed model, fitted

606 to estimate the effect of BMI change on IgG N-glycome. False discovery rate was controlled

607 using Benjamini-Hochberg method at the specified level of 0.05.

Derived IgG glycan trait	BMI difference effect (glycan abundance (%) change per 1 kgm^{-2} decrease in BMI)	BMI difference SE (glycan abundance (%) change per 1 kgm^{-2} decrease in BMI)	p-value	Adjusted p-value
digalactosylation (G2)	0.2004	0.0403	6.88×10^{-07}	5.85×10^{-06}
high mannose (HM)	-0.0519	0.0119	1.33×10^{-05}	5.66×10^{-05}
agalactosylation (G0)	-0.1048	0.0397	8.43×10^{-03}	1.79×10^{-02}
bisecting GlcNAc (B)	0.0526	0.0262	4.49×10^{-02}	6.94×10^{-02}
monogalactosylation (G1)	-0.0573	0.0343	9.53×10^{-02}	1.25×10^{-01}
core fucosylation (CF)	-0.0059	0.0285	8.36×10^{-01}	8.89×10^{-01}

608 Red – significant decrease; green – significant increase; blue – non-significant change.

609 GlcNAc – *N*-acetylglucosamine; SE – standard error