Extensive weight loss can reduce immune age by altering IgG N-glycosylation

Valentina L Greto[1]*, Ana Cvetko[2]*, Tamara Štambuk[3]*, Niall J Dempster[4], Domagoj Kifer[2], Helena Deriš[3], Ana Cindrić[3], Frano Vučković[3], Mario Falchi[5], Richard S Gillies[6], Jeremy W Tomlinson[4], Olga Gornik[2,3], Bruno Sgromo[6], Tim D Spector[5], Cristina Menni[5]*, Alessandra Geremia[1]*, Carolina V Arancibia-Cárcamo[1]*, Gordan Lauc[2,3]*

[1]Translational Gastroenterology Unit and NIHR Oxford Biomedical Research Centre, Nuffield Department of Medicine, University of Oxford, United Kingdom

[2]Faculty of Pharmacy and Biochemistry, University of Zagreb, Zagreb, Croatia

[3]Genos Glycoscience Research Laboratory, Zagreb, Croatia

[4]Oxford Centre for Diabetes and NIHR Oxford Biomedical Research Centre, Endocrinology and Metabolism, Radcliffe Department of Medicine, University of Oxford, United Kingdom

[5]The Department of Twin Research, King's College London, St Thomas' Hospital, London, United Kingdom

[6]Department of Upper GI Surgery, Oxford University Hospitals, United Kingdom

*The authors equally contributed to this work

Corresponding author:

Gordan Lauc

glauc@pharma.hr

Faculty of Pharmacy and Biochemistry, University of Zagreb

A. Kovačića 1, 10 000 Zagreb, Croatia

ABSTRACT

Background

Obesity represents a global health threat, and is associated not only with exponentially

increased cardiometabolic morbidity and mortality, but with adverse clinical outcomes in

patients infected with SARS-CoV-2 as well. Enzymatic attachment of complex oligosaccharides

to proteins (glycosylation) is highly responsive to numerous (patho)physiological conditions

and ageing, which is perhaps best exemplified on IgG. The prospect of immune age reduction,

by reverting induced glycosylation changes through metabolic intervention, opens many

possibilities. Herein, we have investigated whether weight loss interventions affect

inflammation- and ageing-related glycosylation alterations, in a longitudinal cohort of

bariatric-surgery patients. To support potential findings, BMI-related glycosylation changes

were monitored in a longitudinal TwinsUK cohort.

Methods

IgG and plasma N-glycans were chromatographically profiled in 37 obese patients, subjected

to low-calorie diet and then to bariatric surgery, across multiple timepoints. Similarly, plasma

glycome was analysed in 1,680 TwinsUK participants and longitudinally monitored during a

20-year follow-up.

Findings

Low-calorie diet induced marked increase in low branched and significant decrease in highly

branched, more complex plasma N-glycans – the change opposite to the one typically

observed in inflammatory conditions. Bariatric surgery resulted in extensive, gradual

alterations in IgG glycome, that accompanied progressive weight loss during one year follow-

up. We observed significant increase in digalactosylated and sialylated, and substantial

decrease in agalactosylated and core fucosylated IgG glycans. In general, such IgG glycan

profile is associated with a younger biological age and reflects enhanced anti-inflammatory

IgG potential. The TwinsUK cohort replicated weight loss-associated agalactosylation decrease

and digalactosylation increase, estimated through BMI decrease over a 20-year-period.

Interpretation

Altogether, these findings highlight that weight loss substantially affects both plasma and IgG

N-glycosylation, resulting in improved biological and immune age.

GRAPHICAL ABSTRACT

HIGHLIGHTS

- Obesity is associated to circulating pro-inflammatory high branched N-glycans and IgG agalactosylation

- High branching of N-glycans from total plasma proteins decreases after 3-week low-calorie diet

- IgG galactosylation and sialylation increase after bariatric surgery-induced weight loss

- Decrease of BMI over time is associated to increased IgG galactosylation and a reduction of biological age

KEYWORDS

N-glycosylation; bariatric surgery; weight loss; imunoglobulin G; COVID-19; biological age

INTRODUCTION

The global prevalence of obesity has risen dramatically, to the point that it is now considered a pandemic (1). In the current collision with the emergent coronavirus disease (COVID-19 pandemic), there is a strong association of obesity and older age with COVID-19 complications and severity (2). Moreover, US Centre for Disease Control and Prevention recently reported that obesity represents the highest risk factor for COVID-19 mortality in people younger than 65 years (3). According to the World Health Organization, over 650 million individuals are obese, accounting for 13% of the world's adult population. Obesity confers a risk for metabolic syndrome, contributing to type 2 diabetes and cardiovascular disease (CVD) development (4). Metabolic syndrome is linked to a chronic systemic low-grade inflammation, which contributes to the aging of the immune system denoted as inflammaging (5,6). Obesity-related inflammaging results in impaired innate and adaptive immune function, and is characterized by high serum levels of IL-6, TNF-α and CRP (7). Moreover, this persistent inflammatory state may potentially lead to increased viral shedding and to delayed, blunted antiviral responses to SARS-CoV-2 infection (8,9).

Changes in protein N-glycosylation are one of the hallmarks of inflammaging (6,9). The human circulating N-glycome represents the entire set of glycans that are covalently attached to plasma proteins through a nitrogen on an asparagine residue. N-glycans are essential for life and are involved in many physiological processes (10), including signal transduction, protein trafficking and folding, receptor regulation and cell adhesion. Glycosylation has a fundamental role in the innate and adaptive immune responses, accentuated by the fact that all five classes of immunoglobulins (Ig) bear N-glycans. In this regard, IgG is probably the most investigated glycoprotein, whose effector functions are controlled by its Fc-bound glycans (11).

Inter-individual differences in the pace of biological ageing is an intriguing concept that tries to explain why some people stay healthy until very late chronological age, while others age faster and have shorter life expectancy. The same may apply for the risk of severe COVID-19. Progressive age-related changes of IgG glycosylation have been extensively studied (10,12,13), and the GlycanAge model has been proposed to express the difference between chronological ageing and IgG glycome ageing (14). The "age" of the IgG glycome might be estimated through the levels of agalactosylated species, which are increasing with ageing and are associated with enhanced immune activation (15). The opposite applies for digalactosylated IgG glycoforms, which are usually related to a younger age. Besides age-related changes, specific IgG

glycosylation patterns have been already associated with CVD risk score and subclinical atherosclerosis in two large independent UK cohorts (16). Moreover, a prospective follow-up of the EPIC-Potsdam cohort confirmed that changes in plasma N-glycome composition are predictive of future CVD events, with comparable predictive power to the American Heart Association (AHA) score in men and even better predictive power in women (17). The link between an "old", proinflammatory IgG N-glycome and hypertension has also been profoundly studied (18–20), and similar IgG glycosylation patterns were associated with increased body mass index (BMI) and measures of central adiposity (21,22).

Studies on mouse models further corroborate the importance of differential IgG glycoforms in CVD pathogenesis. Namely, it has been shown that hyposialylated IgG (corresponding to an "old" IgG glycome) can induce obesity-related hypertension and insulin resistance in B-cell-deficient mice, through activation of the endothelial FcγRIIB (23,24). These findings indicate that the IgG N-glycome could represent more than a biomarker of inflammation and aging, since distinctive IgG glycoforms act as effector molecules in certain pathologies. Furthermore, supplementation with *N*-acetylmannosamine (ManNAc), a precursor of sialic acid, protects obese mice from hypertension and insulin resistance induction by reverting an "old" IgG N-glycome into a "young" one (24,25). However, studies exploring the possibilities of converting an "old" IgG glycome into a "young" one by metabolic intervention in humans are limited. Of note, only one small study indicated that high-intensity interval training can "rejuvenate" the IgG N-glycome (26).

Finally, severe obesity can be resolved only through bariatric surgery and subsequent weight loss (27). The resulting weight loss impacts energy balance and metabolism, contributing to the increased insulin response, improved glycaemic control and reduction of total body fat, leading to decreased CVD risk and mortality (28). In this study we aimed to determine whether weight loss affects glycan markers related to inflammation and ageing, in a longitudinally-monitored cohort of obese individuals undergoing low-calorie diet and then bariatric surgical interventions. In order to support potential findings, we also investigated the BMI-related glycosylation changes in a longitudinal TwinsUK cohort. The analysis of samples obtained from the largest longitudinal twins cohort in the UK allowed us to observe the changes of BMI in relation to the IgG N-glycosylation alterations over time.

METHODS

Study populations

Bariatric cohort. The cohort included 37 participants, recruited at Oxford University Hospitals to the Gastrointestinal Illnesses study (Ref: 16/YH/0247). All patients were characterised by metabolic status and medical history. Bariatric patients were considered eligible in accordance to National Institute for Health and Care Excellence (NICE) and local guidelines. The choice of bariatric operation was decided according to the Oxford Bariatric Unit regular practice. Patients with a history of alcoholism and/or ongoing anticoagulant treatment were excluded. Patients were also excluded in case of pregnancy, active substance abuse or uncontrolled psychiatric condition including eating disorders. Participants were sampled at baseline and subjected to 3-week low calorie carbohydrate-restricted diet (900 cal, maximum 100 g of carbohydrates per day), followed by bariatric surgery. The sequential follow-up timepoints included the day of the surgery (baseline), after 6.54 ± 3.4 months (mean ± IQR) and 12.47 ± 6.55 months post-op. Characteristics of the bariatric cohort are shown in Table 1.

Table 1 Demographic characteristics of the bariatric cohort

Characteristics	*Bariatric cohort*			
Total No. of participants (N)	37			
No. of participants Sleeve Gastrectomy (SG), N (%)	25 (68%)			
No. Roux-en-Y Gastric Bypass RYGB N (%)	12 (32%)			
No. of participants each timepoint (N)	Pre-op low-calorie diet	Time of surgery	1st post-op timepoint	2nd post-op timepoint
	8	37	30	24
BMI* each timepoint, mean ± SD, kg/m2	Before diet	End of diet	Time of surgery	1st post-op timepoint	2nd post-op timepoint
	48.53 ± 4.13	46.60± 4.21	46.21 ± 4.75	36.01 ± 5.07	32.82 ± 5.17
Female, sex, N (%)	33 (89%)			
Type 2 diabetes, N (%)	6 (16%)			

*BMI reference values: <18.5 (underweight), 18.5–24.9 (normal weight), 25–29.9 (overweight), >30 (obese).

TwinsUK cohort. We have analysed a total of 6,032 plasma samples from 2,146 participants of the TwinsUK study, collected at multiple timepoints over a 20 year-period (29). These included 1,865 individuals sampled at 3 timepoints, 156 individuals sampled at 2 timepoints and 125 individuals sampled only once. Following the plasma N-glycome analysis, glycan data

142 underwent quality control (see *Statistical analysis* section), which decreased the dataset to

143 5,889 samples (measurements). Out of these 5,889 measurements, we have proceeded with

144 statistical analysis on a subset of 3,742 samples (measurements) which had information on

145 BMI available. Description of the TwinsUK cohort is provided in Table 2.

146 **Table 2 Demographic characteristics of the TwinsUK cohort**

Characteristics	*TwinsUK cohort (BMI subset)*
No. of participants (N)	1,680
No. of glycan measurements (N)	3,742
Baseline age, mean ± SD, years	53.23 ± 10.86
Follow up time, mean ± SD, years	7.90 ± 5.66
Female sex, N (%)	1,680 (100)
Baseline BMI, mean ± SD, kg/m^2	25.45 ± 4.53

Ethical statement

Ethical approval for the study was obtained by the National Research Ethics Committees of

the UK National Health Service (NHS) under the reference number 16/YH/0247. All individuals

participating in this study gave written informed consent. The TwinsUK study was approved

by NRES Committee London–Westminster, and all twins provided informed written consent.

N-glycome analysis

Isolation of IgG from human plasma

IgG was isolated from plasma samples by affinity chromatography as described previously

(30). In brief, IgG was isolated in a high-throughput manner, using 96-well protein G

monolithic plates (BIA Separations, Slovenia), starting from 100 µl of plasma. Plasma was

diluted 7× with phosphate buffered saline (PBS; Merck, Germany) and applied to the protein

G plate. IgG was eluted with 1 ml of 0.1 M formic acid (Merck, Germany) and immediately

neutralized with 1 M ammonium bicarbonate (Acros Organics, USA).

N-glycan release from IgG and total plasma proteins

Isolated IgG samples were dried in a vacuum centrifuge. After drying, IgG was denatured with

the addition of 30 µl of 1.33% SDS (w/v) (Invitrogen, USA) and by incubation at 65 °C for 10

min. Plasma samples (10 µl) were denatured with the addition of 20 µl of 2% (w/v) SDS

(Invitrogen, USA) and by incubation at 65 °C for 10 min. From this point on, the procedure was

identical for both IgG and plasma samples. After denaturation, 10 µl of 4% (v/v) Igepal-CA630

(Sigma Aldrich, USA) was added to the samples, and the mixture was shaken 15 min on a plate

shaker (GFL, Germany). N-glycans were released with the addition of 1.2 U of PNGase F

(Promega, USA) and overnight incubation at 37 °C.

Fluorescent labelling and HILIC SPE clean-up of released N glycans

The released N-glycans were labelled with 2-aminobenzamide (2-AB). The labelling mixture

consisted of 2-AB (19.2 mg/ml; Sigma Aldrich, USA) and 2-picoline borane (44.8 mg/ml; Sigma

Aldrich, USA) in dimethyl sulfoxide (Sigma Aldrich, USA) and glacial acetic acid (Merck,

Germany) mixture (70:30 v/v). To each sample 25 µl of labelling mixture was added, followed

by 2 h incubation at 65 °C. Free label and reducing agent were removed from the samples

using hydrophilic interaction liquid chromatography solid-phase extraction (HILIC-SPE). After

incubation samples were brought to 96% of acetonitrile (ACN) by adding 700 µl of ACN (J.T.

Baker, USA) and applied to each well of a 0.2 µm GHP filter plate (Pall Corporation, USA).

Solvent was removed by application of vacuum using a vacuum manifold (Millipore

Corporation, USA). All wells were prewashed with 70% ethanol (Sigma-Aldrich, St. Louis, MO,

USA) and water, followed by equilibration with 96% ACN. Loaded samples were subsequently

washed 5× with 96% ACN. N-glycans were eluted with water and stored at − 20 °C until usage.

Hydrophilic interaction liquid chromatography of N glycans

Fluorescently labelled N-glycans were separated by hydrophilic interaction liquid

chromatography (HILIC) on Acquity UPLC H-Class instrument (Waters, USA) consisting of a

quaternary solvent manager, sample manager and a fluorescence detector, set with excitation

and emission wavelengths of 250 and 428 nm, respectively. The instrument was under the

control of Empower 3 software, build 3471 (Waters, Milford, USA). Labelled N-glycans were

separated on a Waters BEH Glycan chromatography column, with 100 mM ammonium

formate, pH 4.4, as solvent A and ACN as solvent B. In the case of IgG N-glycans, separation

method used linear gradient of 75–62% acetonitrile at flow rate of 0.4 ml/min in a 27-min

analytical run. For plasma protein N-glycans separation method used linear gradient of 70–

53% acetonitrile at flow rate of 0.561 ml/min in a 25-min analytical run. The system was

calibrated using an external standard of hydrolysed and 2-AB labelled glucose oligomers from

which the retention times for the individual glycans were converted to glucose units (GU).

Data processing was performed using an automatic processing method with a traditional

integration algorithm after which each chromatogram was manually corrected to maintain

the same intervals of integration for all the samples. The chromatograms were all separated in the same manner into 24 peaks (GP1– GP24) for IgG N-glycans and 39 peaks (GP1–GP39) for plasma protein N-glycans and are depicted in Supplementary Figure 1 and Supplementary Figure 2, respectively. Detailed description of glycan structures corresponding to each glycan peak is presented in Supplementary Table 1. Glycan peaks were analysed based on their elution positions and measured in glucose units, then compared to the reference values in the "GlycoStore" database (available at: https://glycostore.org/) for structure assignment. The amount of glycans in each peak was expressed as a percentage of the total integrated area. For IgG glycans, in addition to 24 directly measured glycan traits, 8 derived traits were calculated, while for plasma glycans, in addition to 39 directly measured glycan traits, 16 derived traits were also calculated. These derived traits average particular glycosylation features, such as galactosylation, fucosylation, bisecting GlcNAc, and sialylation. Formulas for IgG and plasma protein N-glycan derived traits for bariatric cohort are presented in Supplementary Table 2 and Supplementary Table 3, respectively. Derived trait calculations for TwinsUK cohort are presented in Supplementary Table 4.

Statistical analysis

Bariatric cohort. In order to remove experimental variation from the measurements, normalization and batch correction were performed on the UPLC glycan data. To make measurements across samples comparable, normalization by total area was performed. Prior to batch correction, normalized glycan measurements were log-transformed due to right-skewness of their distributions and the multiplicative nature of batch effects. Batch correction was performed on log-transformed measurements using the ComBat method (R package sva (31), where the technical source of variation (which sample was analysed on which plate) was modelled as batch covariate. To correct measurements for experimental noise, estimated batch effects were subtracted from log-transformed measurements.

Longitudinal analysis of patient samples through their observation period was performed by implementing a linear mixed effects model, where time was modelled as fixed effect, while the individual ID was modelled as random effect. Prior to the analyses, glycan variables were all transformed to standard Normal distribution by inverse transformation of ranks to Normality (R package "GenABEL", function rntransform). Using rank transformed variables makes estimated effects of different glycans comparable, as these will have the same standardized variance. False discovery rate (FDR) was controlled by the Benjamini-Hochberg

procedure at the specified level of 0.05. Data was analysed and visualized using R programming language (version 3.5.2)(32).

TwinsUK cohort. Normalization of peak intensities to the total chromatogram area was performed for each measured sample separately. Calculated proportions were then batch corrected using ComBat method (R package *sva*)(31). After the batch correction the first 11 peaks, which predominantly originate from IgG (33), were used to calculate 6 derived glycan traits – agalactosylation (G0), monogalactosylation (G1), digalactosylation (G2), bisecting GlcNAc (B), core fucosylation (CF) and high mannose structures (HM). Mixed models were fitted to estimate the effect of BMI change on IgG N-glycome (R package *lme4*)(34). Directly measured or derived glycan trait was used as a dependent variable in the mixed model. To differentiate between BMI change and the absolute BMI value, the variable was separated to BMI$_{baseline}$ and BMI$_{difference}$ (calculated according to the following equation: $BMI_{difference} = BMI_{follow\ up\ age} - BMI_{baseline\ age}$), and both were used in the model as a fixed effect. Since IgG N-glycome is affected by aging, age was included both as a fixed effect and a random slope. Finally, to meet the independency criteria, family ID and individual ID (nested within family) were included in the model as a random intercept. Due to multiple model fitting (for 11 directly measured and 6 derived glycan traits) false discovery rate was controlled using Benjamini-Hochberg method. All statistical analyses were performed using R programming language (version 3.6.3)(32).

RESULTS

Pre-surgical low-calorie diet extensively changes plasma N-glycome

We profiled both plasma and IgG N-glycome in a cohort of bariatric surgery-candidate patients at the beginning and at the end of their pre-operative diet. We calculated corresponding derived glycan traits, which average particular glycosylation features. By employing a linear mixed model, we observed ample changes in plasma N-glycome, as 10 out of 16 calculated derived traits displayed significant alterations after the low-calorie diet. Namely, low branched (LB), agalactosylated (G0), monogalactosylated (G1), asialylated (S0), monosialylated (S1), core fucosylated (CF) and glycans bearing a bisecting GlcNAc (B) showed a substantial increase in their abundances (Table 3). Conversely, high branched (HB), trisialylated (S3) and trigalactosylated (G3) glycan abundances decreased (Table 3). Interestingly, IgG glycome was not affected by the low-calorie diet, as its derived traits did not

display any significant changes after the dieting period. Graphical representation of the

longitudinal alterations in plasma N-glycome after low calorie diet are depicted in Figure 1.

Table 3 Low-calorie diet induces significant changes in plasma N-glycome. Longitudinal
analysis was performed by implementing a linear mixed effects model, with time as a fixed
effect and the individual sample measurement as a random effect. False discovery rate was
controlled using Benjamini-Hochberg method at the specified level of 0.05.

Derived glycan trait	Time_effect	Time_SE	Time_p-value	Adjusted p-value
monogalactosylation (G1)	0.6020	0.0796	4.18×10^{-05}	6.69×10^{-04}
trisialylation (S3)	-1.1512	0.2008	3.04×10^{-04}	2.43×10^{-03}
agalactosylation (G0)	0.4903	0.0948	6.09×10^{-04}	2.44×10^{-03}
core fucosylation (CF)	0.5431	0.1011	4.73×10^{-04}	2.44×10^{-03}
high branching (HB)	-1.1925	0.2468	9.43×10^{-04}	2.52×10^{-03}
asialylation (S0)	0.6336	0.1310	9.40×10^{-04}	2.52×10^{-03}
trigalactosylation (G3)	-1.1925	0.2581	1.24×10^{-03}	2.84×10^{-03}
bisecting GlcNAc (B)	0.5457	0.1350	2.84×10^{-03}	5.69×10^{-03}
low branching (LB)	0.9839	0.3060	9.99×10^{-03}	1.78×10^{-02}
monosialylation (S1)	0.5024	0.1981	2.98×10^{-02}	4.77×10^{-02}
disialylation (S2)	-0.4314	0.1941	4.99×10^{-02}	7.25×10^{-02}
high mannose (HM)	0.7545	0.3619	6.24×10^{-02}	8.33×10^{-02}
antennary fucosylation (AF)	-0.1804	0.1102	1.28×10^{-01}	1.58×10^{-01}
digalactosylation (G2)	0.3107	0.2309	2.01×10^{-01}	2.30×10^{-01}
tetragalactosylation (G4)	-0.3446	0.3477	3.36×10^{-01}	3.58×10^{-01}
tetrasialylation (S4)	-0.3049	0.3547	4.00×10^{-01}	4.00×10^{-01}

268 Red – significant decrease; green – significant increase; blue – non-significant change.
269 GlcNAc – *N*-acetylglucosamine; SE – standard error

Figure 1 Low calorie diet-related alterations in plasma N-glycome after a 3-week follow-up. Red line – significant decrease; green line – significant increase; blue line – non-significant change.

Bariatric surgery markedly alters IgG N-glycosylation

Using the same chromatographic approach, we analysed samples from patients who underwent bariatric surgery. The plasma samples were collected on the day of surgery (month 0), approximately 6 months post-surgery and 12 months post-surgery. Both plasma and IgG N-glycans were profiled in each of these timepoints, and the obtained values were used for derived glycan traits calculations. Of the examined plasma glycosylation features, only agalactosylation (G0) displayed significant decrease after bariatric surgery (Table 4). On the other hand, IgG N-glycome showed more extensive changes, following the bariatric procedure. Namely, four out of eight tested derived traits showed marked changes: core fucosylated (CF) and agalactosylated (G0) glycans decreased, while digalactosylated (G2) and monosialylated (S1) glycans increased after the surgery (Table 4). The IgG glycans whose

abundances were increased after bariatric surgery are major components of a "young" IgG
glycome, as they are typically associated with a younger age. The opposite applies to
agalactosylated structures , which are usual denominators of an "old" IgG glycome profile. We
also examined the correlation of patients' clinical data with plasma and IgG glycome features
using multivariate analysis, but found no statistically significant associations (data not shown).
Finally, the type of bariatric surgery (either sleeve gastrectomy or Roux-en-Y gastric bypass)
did not affect plasma nor IgG glycome composition. Graphical representations of the
longitudinal alterations in IgG and plasma glycosylation features are depicted in Figure 2 and
Supplementary Figure 3, respectively.

Table 4 Bariatric surgery induces significant changes in plasma and IgG N-glycomes.
Longitudinal analysis was performed by implementing a linear mixed effects model, with
time as a fixed effect and the individual sample measurement as a random effect. False
discovery rate was controlled using Benjamini-Hochberg method at the specified level of
0.05.

N-glycome	Derived glycan trait	Time_effect	Time_SE	Time_p-value	Adjusted p-value
plasma	agalactosylation (G0)	-0.0273	0.0071	5.26×10^{-04}	8.42×10^{-03}
IgG	agalactosylation (G0)	-0.0339	0.0078	9.23×10^{-05}	7.38×10^{-04}
IgG	digalactosylation (G2)	0.0275	0.0072	3.75×10^{-04}	1.50×10^{-03}
IgG	monosialylation (S1)	0.0193	0.0080	1.97×10^{-02}	3.94×10^{-02}
IgG	core fucosylation (CF)	-0.0155	0.0064	1.74×10^{-02}	3.94×10^{-02}

299 Red – significant decrease; green – significant increase. SE – standard error.

Figure 2 Bariatric surgery-related alterations in IgG N-glycome over time (months). Red line
– significant decrease; green line – significant increase; blue line – non-significant change.

Weight loss induces shift towards ''young'' IgG N-glycome

Using the same chromatographic approach, we profiled plasma protein N-glycome from 1,680
TwinsUK study participants sampled at several timepoints over a 20-year-period. This served
as a replication of the findings from the bariatric cohort, whose participants exhibited the
reversal from ''old'' to ''young'' IgG N-glycome due to weight loss. As these results showed
prominent changes in IgG composite glycan traits only, herein we focused on IgG glycosylation
changes. Due to the fact that for TwinsUK cohort plasma glycome was profiled, we calculated
derived traits and performed statistical analysis only on the first 11 glycan peaks,
corresponding to the glycans predominantly originating from IgG (33). Six derived traits were
calculated – agalactosylation (G0), monogalactosylation (G1), digalactosylation (G2),
incidence of bisecting GlcNAc (B), core fucosylation (CF) and incidence of high mannose
structures (HM). We examined IgG glycome alterations associated with changes in BMI using
a mixed model on a subset of 3,742 samples. Out of six examined IgG glycosylation features
(derived traits), three displayed significant BMI-related changes – agalactosylation (G0),

digalactosylation (G2) and incidence of high mannose structures (HM) (Table 5). Namely, the abundance of digalactosylated (G2) glycans increased with the BMI decrease, while the abundance of agalactosylated (G0) and high mannose glycans (HM) decreased with the weight loss, estimated by the BMI drop. These findings are in line with the results observed in the bariatric surgery cohort. Graphical representation of the longitudinal BMI-dependent alterations of IgG glycosylation are depicted at Figure 3.

Table 5 Longitudinally monitored weight loss results with significant changes of IgG N-glycosylation. Longitudinal analysis was performed by implementing a mixed model, fitted to estimate the effect of BMI change on IgG N-glycome. False discovery rate was controlled using Benjamini-Hochberg method at the specified level of 0.05.

Derived glycan trait	BMI difference effect (glycan abundance (%) change per 1 kgm^{-2} decrease in BMI)	BMI difference SE (glycan abundance (%) change per 1 kgm^{-2} decrease in BMI)	p-value	Adjusted p-value
digalactosylation (G2)	0.2004	0.0403	6.88×10^{-07}	5.85×10^{-06}
high mannose (HM)	-0.0519	0.0119	1.33×10^{-05}	5.66×10^{-05}
agalactosylation (G0)	-0.1048	0.0397	8.43×10^{-03}	1.79×10^{-02}
bisecting GlcNAc (B)	0.0526	0.0262	4.49×10^{-02}	6.94×10^{-02}
monogalactosylation (G1)	-0.0573	0.0343	9.53×10^{-02}	1.25×10^{-01}
core fucosylation (CF)	-0.0059	0.0285	8.36×10^{-01}	8.89×10^{-01}

Red – significant decrease; green – significant increase; blue – non-significant change.

GlcNAc – *N*-acetylglucosamine; SE – standard error

Figure 3 BMI-associated alterations in IgG N-glycosylation across multiple timepoints.
Changes in IgG derived traits are presented with lineplots of hypothetical ageing of TwinsUK
participants (all women). Black dot represents a starting point of a 30-year-old woman, black
triangle of a 40-year-old woman and black square of a 50-year-old woman. All of these women
have a baseline BMI of 25 kg m^{-2}. Blue lines represent age-related IgG glycosylation changes
attributed to stabile BMI. Green lines represent age-related IgG glycosylation changes
attributed to increasing BMI (0.5 kg m^{-2} per year, through a period of 10 years). Red lines
represent age-related IgG glycosylation changes attributed to decreasing BMI (0.5 kg m^{-2} per
year, through a period of 10 years). Highlighted areas represent 95% confidence intervals. The
effect of age on IgG glycosylation is represented with the curve slope, while the effect of BMI
change is represented with the distance of green/red line from the blue line.

DISCUSSION

In this study, we have observed extensive changes in both plasma and IgG N-glycome
associated with weight loss following low-calorie diet and bariatric surgery, or expressed
through BMI decrease. To the best of our knowledge, this is the first study to investigate

plasma and IgG N-glycome alterations in patients who underwent a low-calorie diet followed by bariatric surgery.

Prior to bariatric surgery, patients were subjected to a low-calorie diet which caused significant and extensive changes to the plasma protein N-glycome. The observed changes can be summarized as significant increase in low-branched, biantennary structures (S0, S1, G0, G1…) and, conversely, substantial decrease in high-branched, triantennary, more complex structures (S3, G3). The glycosylation alterations induced by a low-calorie diet are showing exactly the opposite direction of change than those seen in various inflammatory conditions, such as type 2 diabetes, chronic obstructive pulmonary disease and inflammatory bowel disease (17,30,35,36), suggesting a gradual attenuation of inflammation. Interestingly, no significant changes were found for the IgG N-glycome alone after the diet, probably due to the relatively short follow-up period (3 weeks), which matches IgG serum half-life. Altogether, our glycomic data suggest that the overall inflammatory body status improves after a short time of low-calorie diet, reflecting a quickly noticeable and beneficial metabolic effect.

We have analysed both plasma and IgG N-glycome from individuals who underwent bariatric surgery, in a longitudinal manner. We observed significant changes in both plasma and IgG N-glycome. In particular, plasma N-glycome showed significant alterations only in agalactosylated (G0) glycans, whose abundance decreased during the follow-up period. Since the majority of agalactosylated species in plasma originates from IgG (33), it is no surprise that this glycosylation feature was also found to be significantly decreased in IgG glycome analysis. Elevated levels of G0 IgG glycoforms are typically associated with aging, pro-inflammatory IgG glycan profile and various inflammatory diseases (12). On the other hand, the levels of digalactosylated (G2) glycans increased after bariatric surgery and at sequential timepoints, in accordance with a reduced inflammatory potential of the circulating IgG. Interestingly, the increased levels of IgG galactosylation are associated with a younger biological age and are considered, in a way, as a measure of an individual's well-being (15). Converesely, IgG galactosylation levels substantially decrease with ageing and during inflammation (12,14). Our results demonstrate that weight loss, resulting from bariatric surgery, can initiate the reversal from an ''old'' to a ''young'' IgG N-glycome, potentially reversing the clock for the immune/biological age. Furthermore, bariatric surgery also resulted in significant IgG glycome alteration inducing a decrease in core fucosylation. The vast majority of circulating IgG

molecules bears core fucose (approximately 95%), which profoundly decreases IgG binding affinity to FcγRIIIA receptor and sequential antibody-dependent cellular cytotoxicity (ADCC) (37). ADCC is largely mediated by natural killer cells that can lyse target cells and fight viral infections. This would suggest that an extensive weight loss ameliorates immune responses dedicated to combat viral infections, by altering IgG glycosylation and modulating its effector functions. Lastly, bariatric surgery-related weight loss led to an increase in IgG sialylation, which is the main modulator of the IgG anti-inflammatory actions (38). In addition to its anti-inflammatory actions, the level of IgG sialylation has been implicated in the pathogenesis of obesity-induced insulin resistance and hypertension, as already mentioned (23,24). Namely, inhibitory IgG receptor FcγRIIB was found to be expressed in the microvascular endothelium. Sequentially, it was shown that hyposialylated IgG acts as its operating ligand, leading to the induction of obesity-related insulin resistance and hypertension. On the other hand, the sialylated glycoform is not activating the signalling pathways responsible for the insulin resistance and hypertension development, but is rather preserving insulin sensitivity and normal vasomotor tone, even in obese mice. Interestingly, the same group made another significant discovery – promotion of IgG sialylation breaks the link between obesity and hypertension/insulin resistance (24,25). Namely, supplementation with the sialic acid precursor restored IgG sialylation, highlighting a potential approach to improve both metabolic and cardiovascular health in humans, with a single intervention. Our data suggest that a similar effect might be achieved by weight loss interventions, through restoring of IgG sialylation levels. Altogether, it is remarkable that such profound effects were observed on a rather small number of participants, which further highlights their significance in a given biological setting.

Lastly, in order to confirm the effects of weight loss on the biological/immune age, we investigated how BMI decrease affects IgG N-glycome during a 20-year-period. We observed the prominent inverse changes of agalactosylated (G0) and digalactosylated (G2) IgG glycans. Namely, agalactosylated IgG glycans significantly decreased, while digalactosylated ones substantially increased as the BMI decreased. These observations corroborated our earlier findings, confirming that the body weight reduction reverses IgG glycome from "old" to "young", implying at the same time a likely reduction in the biological/ immune age. Nonetheless, we have to acknowledge the following limitations of TwinsUK as a replication

cohort: first – TwinsUK participants have not experienced such an extensive weight loss, which

potentially influenced the replication of other significant glycan changes from the bariatric

cohort; second – herein, the weight loss was approximated by BMI decrease, which is usually

a legitimate assumption, however, it does not have to apply to all cases; and third – herein we

profiled plasma glycome, while the IgG glycan traits were only approximated and the

information on IgG sialylation was confounded by other plasma glycoproteins. Ideally, these

issues could be circumvented in the future studies whose experimental design would allow

the simultaneous, multi-centre follow-up of larger groups of patients.

Intense physical exercise can also shift IgG N-glycome towards a "younger" profile by

increasing IgG galactosylation (26). Although another study reported that prolonged intensive

exercise can have the opposite effect and promote pro-inflammatory changes of IgG N-

glycome (39), its findings are not surprising since it recruited healthy, normal-weight female

participants, subjected to the intense energy deprivation and exercise levels, to induce

substantial fat loss in a rather short time period. The authors also highlighted that starting

weight and the way in which weight loss is achieved could be crucial for the final effect on the

immune system. Therefore, it seems that exercise overall has a positive impact on the immune

system and biological clock, but its intensity and duration should be personalized in order to

provide the optimal results.

To summarize, our results indicate that weight loss has impact on inflammation and biological

aging by altering plasma and IgG N-glycosylation patterns. Glycosylation changes can be

reliably quantified and used to estimate the "age" of the immune system. This could represent

a valuable tool in the context of COVID-19 patient stratification. Since inter-individual

variation in protein glycosylation is extensive, its functional implications on membrane

expression of various proteins (including ACE2), functional properties of newly synthesized

SARS-CoV-2 virus particles and regulation of the immune responses may also explain severe

cases of COVID-19 in apparently healthy individuals. A previous research showed that

impairment of ACE2 N-glycosylation affects its ability to allow the transduction of SARS Human

coronavirus NL63 (HCoV-NL63) particles, by disrupting the viral endocytosis (40).

Future glycomic studies of patients infected with SARS-CoV-2 would help to elucidate whether

ACE2 N-glycosylation differs in patients with severe COVID-19, compared to asymptomatic

patients or healthy individuals. For the time being, it appears that severity of COVID-19

associates with the majority of obesity-related comorbidity risk factors (41). Therefore, by

improving metabolic and endocrine health through weight loss, which consequently

contributes to the preservation of a healthy immune system, one has a best chance to fight

infections.

ACKNOWLEDGMENTS

This research was funded by the National Institute for Health Research (NIHR) Oxford

Biomedical Research Centre (BRC). The views expressed are those of the author(s) and not

necessarily those of the NHS, the NIHR or the Department of Health. The authors thank Rachel

Franklin, Michelle Haylock, James Chivenga, Roxanne Williams and the BRC Oxford GI Biobank

for sample collection. The authors thank all the patients who took part in this study.

FUNDING

Glycosylation analysis was performed in Genos Glycoscience Research Laboratory and partly

supported by the European Union's Horizon 2020 grants Backup (grant# 777090), European

Structural and Investment Funds IRI grant (#KK.01.2.1.01.0003), Centre of Competence in

Molecular Diagnostics (#KK.01.2.2.03.0006) and Croatian National Centre of Research

Excellence in Personalized Healthcare grant (#KK.01.1.1.01.0010). Bariatric research was

supported by the National Institute for Health Research (NIHR) Oxford Biomedical Research

Centre Gastroenterology and Mucosal Immunity Theme to CVA-C and VLG, Oxford Biomedical

Research Centre Diabetes and Metabolism theme for JWT, MRC programme grant to JWT (ref.

MR/P011462/1), the NIHR Research Capability Funding to VLG and AG, the Wellcome Trust

(101734/Z/13/Z) to AG. NJD is supported by a University of Oxford Novo Nordisk Clinical

Research Fellowship. TwinsUK received funding from the Wellcome Trust European

Community's Seventh Framework Programme (FP7/2007-2013 to TwinsUK); the National

Institute for Health Research (NIHR) Clinical Research Facility at Guy's & St Thomas' NHS

Foundation Trust and the Nottingham NIHR Biomedical Research Centre based at Guy's and

St Thomas' NHS Foundation Trust and King's College London. CM is funded by the MRC AimHy

(MR/M016560/1) project grant and by the Chronic Disease Research Foundation.

AUTHOR`S CONTRIBUTION

Study design: CM, AG, CVA-C, GL. Data collection: VLG, ACv, TŠ, NJD, ACi, HD, MF, CM. Data

analysis: DK, FV. Data interpretation: VLG, ACv, TŠ, NJD, DK, FV, MF, JWT, OG, TS, CM, AG,

CVA-C, GL. Drafting manuscript: VLG, ACv, TŠ, NJD, DK, FV, GL. Revising and approving final

version of manuscript: ALL. CM, AG, CVA-C and GL take responsibility for the integrity of the

data analysis

REFERENCES

1.   Obesity and overweight [Internet]. [cited 2020 Apr 16]. Available from:
https://www.who.int/news-room/fact-sheets/detail/obesity-and-overweight

2.   Petrilli CM, Jones SA, Yang J, Rajagopalan H, O'Donnell LF, Chernyak Y, et al. Factors
associated with hospitalization and critical illness among 4,103 patients with COVID-19
disease in New York City. medRxiv. 2020 Apr 11;2020.04.08.20057794.

3.   Garg S. Hospitalization Rates and Characteristics of Patients Hospitalized with
Laboratory-Confirmed Coronavirus Disease 2019 — COVID-NET, 14 States, March 1–30,
2020. MMWR Morb Mortal Wkly Rep [Internet]. 2020 [cited 2020 Apr 26];69. Available
from: https://www.cdc.gov/mmwr/volumes/69/wr/mm6915e3.htm

4.   Han TS, Lean ME. A clinical perspective of obesity, metabolic syndrome and
cardiovascular disease. JRSM Cardiovasc Dis. 2016 Dec;5:2048004016633371.

5.   Alpert A, Pickman Y, Leipold M, Rosenberg-Hasson Y, Ji X, Gaujoux R, et al. A clinically
meaningful metric of immune age derived from high-dimensional longitudinal
monitoring. Nat Med. 2019 Mar;25(3):487–95.

6.   Franceschi C, Garagnani P, Parini P, Giuliani C, Santoro A. Inflammaging: a new
immune–metabolic viewpoint for age-related diseases. Nature Reviews Endocrinology.
2018 Oct;14(10):576–90.

7.   Touch S, Clément K, André S. T Cell Populations and Functions Are Altered in Human
Obesity and Type 2 Diabetes. Curr Diab Rep. 2017 Sep;17(9):81.

8.   Dhurandhar NV, Bailey D, Thomas D. Interaction of obesity and infections. Obes Rev.
2015 Dec;16(12):1017–29.

9.   Lauc G, Sinclair D. Biomarkers of biological age as predictors of COVID-19 disease
severity. Aging [Internet]. 2020 Apr 8 [cited 2020 Apr 14]; Available from:
http://www.aging-us.com/article/103052/text

10.   Lauc G, Pezer M, Rudan I, Campbell H. Mechanisms of disease: The human N-glycome.
Biochimica et Biophysica Acta - General Subjects. 2016;1860(8):1574–82.

11.   Gornik O, Pavić T, Lauc G. Alternative glycosylation modulates function of IgG and other
proteins - implications on evolution and disease. Biochim Biophys Acta. 2012
Sep;1820(9):1318–26.

12.   Gudelj I, Lauc G, Pezer M. Immunoglobulin G glycosylation in aging and diseases.
Cellular Immunology. 2018;

13.   Dall'Olio F. Glycobiology of aging. In: Subcellular Biochemistry. 2018.

14.   Krištić J, Vučković F, Menni C, Klarić L, Keser T, Beceheli I, et al. Glycans Are a Novel
Biomarker of Chronological and Biological Ages. The Journals of Gerontology: Series A.
2014 Jul;69(7):779–89.

15. Štambuk J, Nakić N, Vučković F, Pučić-Baković M, Razdorov G, Trbojević-Akmačić I, et al.
Global variability of the human IgG glycome [Internet]. Biochemistry; 2019 Feb [cited
2020 Jan 31]. Available from: http://biorxiv.org/lookup/doi/10.1101/535237

16. Menni C, Gudelj I, MacDonald-Dunlop E, Mangino M, Zierer J, Bešić E, et al.
Glycosylation Profile of Immunoglobulin G Is Cross-Sectionally Associated with
Cardiovascular Disease Risk Score and Subclinical Atherosclerosis in Two Independent
Cohorts. Circulation Research. 2018;122(11):1555–64.

17. Wittenbecher C, Štambuk T, Kuxhaus O, Rudman N, Vučković F, Štambuk J, et al. Plasma
N-Glycans as Emerging Biomarkers of Cardiometabolic Risk: A Prospective Investigation
in the EPIC-Potsdam Cohort Study. Diabetes Care. 2020 Jan 7;43(3):661–8.

18. Gao Q, Dolikun M, Stambuk J, Wang H, Zhao F, Yiliham N, et al. Immunoglobulin G N-
Glycans as Potential Postgenomic Biomarkers for Hypertension in the Kazakh
Population. Omics. 2017;21(7):380–9.

19. Liu J, Dolikun M, Štambuk J, Trbojević-Akmačić I, Zhang J, Wang H, et al. The association
between subclass-specific IgG Fc N-glycosylation profiles and hypertension in the
Uygur, Kazak, Kirgiz, and Tajik populations. J Hum Hypertens. 2018 Sep;32(8–9):555–63.

20. Wang Y, Klarić L, Yu X, Thaqi K, Dong J, Novokmet M, et al. The Association Between
Glycosylation of Immunoglobulin G and Hypertension: A Multiple Ethnic Cross-Sectional
Study. Medicine. 2016 Apr;95(17):e3379.

21. Nikolac Perkovic M, Pucic Bakovic M, Kristic J, Novokmet M, Huffman JE, Vitart V, et al.
The association between galactosylation of immunoglobulin G and body mass index.
Progress in Neuro-Psychopharmacology and Biological Psychiatry. 2014 Jan;48:20–5.

22. Russell AC, Kepka A, Trbojević-Akmačić I, Ugrina I, Song M, Hui J, et al. Increased central
adiposity is associated with pro-inflammatory immunoglobulin G N-glycans.
Immunobiology. 2019;224(1):110–5.

23. Sundgren NC, Vongpatanasin W, Boggan BMD, Tanigaki K, Yuhanna IS, Chambliss KL, et
al. IgG receptor FcγRIIB plays a key role in obesity-induced hypertension. Hypertension.
2015 Feb 21;65(2):456–62.

24. Tanigaki K, Sacharidou A, Peng J, Chambliss KL, Yuhanna IS, Ghosh D, et al.
Hyposialylated IgG activates endothelial IgG receptor FcγRIIB to promote obesity-
induced insulin resistance. J Clin Invest. 2018 Jan 2;128(1):309–22.

25. Peng J, Vongpatanasin W, Sacharidou A, Kifer D, Yuhanna IS, Banerjee S, et al.
Supplementation with the Sialic Acid Precursor N-acetyl-D-Mannosamine Breaks the
Link Between Obesity and Hypertension. 2019.

26. Tijardović M, Marijančević D, Bok D, Kifer D, Lauc G, Gornik O, et al. Intense Physical
Exercise Induces an Anti-inflammatory Change in IgG N-Glycosylation Profile. Frontiers
in Physiology. 2019;10(December):1–10.

27.   Nguyen NT, Kim E, Vu S, Phelan M. Ten-year Outcomes of a Prospective Randomized
Trial of Laparoscopic Gastric Bypass Versus Laparoscopic Gastric Banding. Ann Surg.
2018;268(1):106–13.

28.   O'Brien P. Bariatric surgery and type 2 diabetes: a step closer to defining an optimal
approach. The Lancet Diabetes & Endocrinology. 2019 Dec 1;7(12):889–91.

29.   Verdi S, Abbasian G, Bowyer RCE, Lachance G, Yarand D, Christofidou P, et al. TwinsUK:
The UK Adult Twin Registry Update. Twin Res Hum Genet. 2019 Dec;22(6):523–9.

30.   Pavić T, Dilber D, Kifer D, Selak N, Keser T, Ljubičić Đ, et al. N-glycosylation patterns of
plasma proteins and immunoglobulin G in chronic obstructive pulmonary disease. J
Transl Med. 2018 Nov 21;16.

31.   Leek JT, Johnson WE, Parker HS, Jaffe AE, Storey JD. The sva package for removing
batch effects and other unwanted variation in high-throughput experiments.
Bioinformatics. 2012 Mar 15;28(6):882–3.

32.   R Core Team (2020). R: A language and environment for statistical computing. R
Foundation for Statistical Computing, Vienna, Austria. URL   https://www.R-
project.org/.

33.   Clerc F, Reiding KR, Jansen BC, Kammeijer GSM, Bondt A, Wuhrer M. Human plasma
protein N-glycosylation. Glycoconj J. 2016 Jun 1;33(3):309–43.

34.   Bates D, Mächler M, Bolker B, Walker S. Fitting Linear Mixed-Effects Models Using
lme4. Journal of Statistical Software. 2015 Oct 7;67(1):1–48.

35.   Clerc F, Novokmet M, Dotz V, Reiding KR, de Haan N, Kammeijer GSM, et al. Plasma N-
Glycan Signatures Are Associated With Features of Inflammatory Bowel Diseases.
Gastroenterology. 2018;155(3):829–43.

36.   Novokmet M, Lukić E, Vučković F, –Durić Ž, Keser T, Rajšl K, et al. Changes in IgG and
total plasma protein glycomes in acute systemic inflammation. Scientific Reports. 2014
Mar 11;4(1):1–10.

37.   Iida S, Kuni-Kamochi R, Mori K, Misaka H, Inoue M, Okazaki A, et al. Two mechanisms of
the enhanced antibody-dependent cellular cytotoxicity (ADCC) efficacy of non-
fucosylated therapeutic antibodies in human blood. BMC Cancer. 2009 Feb 18;9:58.

38.   Kaneko Y, Nimmerjahn F, Ravetch JV. Anti-Inflammatory Activity of Immunoglobulin G
Resulting from Fc Sialylation. Science. 2006 Aug 4;313(5787):670–3.

39.   Sarin HV, Gudelj I, Honkanen J, Ihalainen JK, Vuorela A, Lee JH, et al. Molecular
Pathways Mediating Immunosuppression in Response to Prolonged Intensive Physical
Training, Low-Energy Availability, and Intensive Weight Loss. Front Immunol.
2019;10:907.

40.   Zhao X, Guo F, Comunale MA, Mehta A, Sehgal M, Jain P, et al. Inhibition of
endoplasmic reticulum-resident glucosidases impairs severe acute respiratory

syndrome coronavirus and human coronavirus NL63 spike protein-mediated entry by
altering the glycan processing of angiotensin I-converting enzyme 2. Antimicrob Agents
Chemother. 2015 Jan;59(1):206–16.

41.   Maffetone PB, Laursen PB. The Perfect Storm: Coronavirus (Covid-19) Pandemic Meets
Overfat Pandemic. Front Public Health [Internet]. 2020 [cited 2020 Apr 24];8. Available
from:
https://www.frontiersin.org/articles/10.3389/fpubh.2020.00135/full?utm_source=F-
NTF&utm_medium=EMLX&utm_campaign=PRD_FEOPS_20170000_ARTICLE