The nexus of travel restriction, air pollution and COVID-19 infection: Investigation from a megacity of the southern China

Wei Li[1,2*], Xiaohong Chen[1], Chunxing Chen[2], Guowei Liao[2]

[1] *School of Civil Engineering, Sun Yat-Sen University, 519082, Zhuhai, China.*

[2] *Shenzhen Academy of Environmental Sciences, 518001, Shenzhen, China.*

* Correspondence: liwei@meeb.sz.gov.cn; Tel.: +86-0755-2598-0351

Abstract: To control and prevent the spread of COVID-19, generalized social distancing measures, such as traffic control and travel restriction acted in China. Previous studies indicated that the traffic conditions had significant influence on the air quality, and which was related to the respiratory diseases. This study aimed to reveal the nexus of travel restriction, air pollution and COVID-19. Shenzhen, one of the top 4 megacities in China was considered as the study area, statistical analysis methods, including linear/nonlinear regression and bivariate correlation was conducted to evaluate the relationship of the traffic and passenger population, travel intensity, NO_2, PM_{10}, $PM_{2.5}$ and the number of COVID-19 confirmed cases. The results suggested that traffic control and travel restriction had a significant correlation with the number of COVID-19 confirmed cases, which shown negative correlation with the traffic intensity of the city, NO_2, PM_{10} and $PM_{2.5}$ show significant positive correlation with the traffic intensity, traffic control and travel restriction would slow down and prevent the spread of the viruses at the outbreak period. Different study scale might results in different results, thus the research focused on the nexus of traffic control and travel restriction, air pollution and COVID-19 should been enhanced in future, and differentiated epidemic control and prevention measures should be considered according to the different situation of cities as well as countries.

Keywords: travel restriction, traffic control, air pollutants, COVID-19 infection, correlation analysis

1. Introduction

Since 31 December 2019 China reported a cluster of cases of pneumonia in Wuhan, Hubei Province (WHO, 2020a; Chen et al., 2020; Xie and Zhu., 2020), a novel coronavirus disease, namely COVID-19 spread rapidly and became a world pandemic (WHO, 2020a,b; Vellingiri et al.. 2020), there have been 2,164,111 confirmed cases of COVID-19 in 213 countries, areas or territories, reported to WHO as of 2:00 am CEST, 18 April 2020 (WHO, 2020b). To prevent and

reduce spread of the infection, strong and immediate measures, such as detect disease early, isolate

and treat cases, trace contacts and social distancing acted in China at national scale (Pan et al.,

2020). Modeling studies indicated that the travel restrict of Wuhan delayed the epidemic

progression by 3 to 5 days in mainland China (Chinazzi et al., 2020; Tian et al., 2020; Pan et al.,

2020), while reducing case importations to other countries by nearly 80% through mid-February

(Chinazzi et al., 2020; Pan et al., 2020).

Previous studies indicated that the respiratory disease was highly related to the air pollution

(Phosri et al., 2019; Zhu et al., 2019), and which could be considered as a putative disease risk

factor of the novel COVID-19 disease (Martelletti and Martelletti, 2020). Study conducted by Zhu

et al., 2020 in China reveled that there was a significant relationship between COVID-19 infection

and air pollutants such as PM_{10}, $PM_{2.5}$ and NO_2. To prevent and control the spread of the novel

coronavirus disease, social distancing was encouraged in China, people were encouraged to stay at

home and limit outdoor activities, the worse epidemic regions took traffic control measures to

restrict the travel. Consequently, social distancing was not only a simple polite behavior that

keeping distance with people, but also multiple measures, such as traffic control and travel

restriction, that controlled and prevented the contact among people and more over cut down the

spread of the viruses (Prin and Bartels, 2020; Yezli and Khan, 2020). Different traffic and travel

management strategies had a direct effect on air pollution (Pasquier and André, 2017; Krecl et al.,

2020), however, to our knowledge, no study has yet comprehensively evaluated the nexus of

traffic control and travel restriction, air pollution and COVID-19 infection at city scale, how travel

restriction influences human living habit, and the impact on air environment as well as COVID-19

infection should be well concerned.

In this paper, a contradistinctive study was conducted among the changing of intracity and

intercity travel restriction, air environment quality and COVID-19 infection in Shenzhen, one of

the top 4 megacity of China, from January to March in 2019 and 2020, and the relationship of

which was evaluated using statistical methods. The authors hope the findings will provide useful

enlightenment for the control and prevention of this novel coronavirus disease.

2. Data acquisition and methods

This study was conducted in Shenzhen, southern China, with longitude from 113°43'E to

114°38'E, latitude from 22°24'N to 22°52'N (Fig.1). Shenzhen is one of the top 4 megacities of

China, with 1997.47 km^2 urban area and 13.03 million permanent resident populations. In 2018

the total GDP of Shenzhen was 2422.2 billion CNY and the economic aggregate ranked the top 5

of the Asian cities.

<Fig.1. Location of the study area and the air quality monitoring stations>

The air quality data were adapted from the Shenzhen national air quality automatic

monitoring stations, hourly data of NO_2 (ug/m^3), PM_{10} (ug/m^3) and $PM_{2.5}$ (ug/m^3), from 11 stations

were collected during 0:00 1 January and 24:00 31 March in 2019 and 2020, and which were

calculated to daily the mean value. The traffic data, including intracity traffic (the monthly

passenger population, 10^4 person/time) and intercity traffic (the monthly public transport

population, 10^4 person/time) were adapted from the Transport Bureau of Shenzhen Municipality

(http://jtys.sz.gov.cn/zwgk/sjfb/index.htm). The travel intensity (TI) of Shenzhen was collected

from Baidu Qianxi (http://qianxi.baidu.com/), an open data platform based on the location based

service (LBS) big data technology, and the daily TI was the logarithmic results of the ratio of the

travel population and the resistant population (cited from the data declaration of the Baidu Qianxi

website). The data of COVID-19 confirmed cases were collated from the Shenzhen Municipal

Health Commission (http://wjw.sz.gov.cn/yqxx/202004/t20200420_19176831.htm).

The descriptive statistical analysis of the intercity and intracity traffic and travel, air

pollutants and COVID-19 confirmed cases was conducted firstly and nonparametric test

(Kolmogorov-Smirnov method) (Zhao et al., 2017) was used to reveal the data distribution

characters. Secondly, linear/nonlinear regression and bivariate correlation analysis (Grange et al.,

2016) was conducted to evaluate the relationship of these variables, and the correlation was

considered significant at the 0.01 level (2-tailed) (Zhu et al., 2020).

3. Results

3.1 data descriptive

Fig.2 illustrated the temporal variability of the number of COVID-19 confirmed cases and

the travel intensity of Shenzhen in 2019 and 2020. With the developments on the novel

coronavirus disease in Shenzhen, the time series of January to March 2020 were divided into 3

parts: the normal period, the social distancing period and the reopened period. The normal period

was from 1 January to 23 January. During this period, the variation tendency of travel intensity in

2020 was similar with that in 2019. The average value of travel intensity in 2020 was 4.34, with

25% increased from 2019, indicating a more active travel desire of the population in 2020. The

social distancing period was from 24 January to 9 February. With the rapidly increasing confirmed

cases, Shenzhen strengthened social distancing and the control of traffic and travel, for instance,

people were encouraged to stay at home, real-name registration with residential address and

personal health condition was conducted by mobile phone APP, and the electric health QR code

was checked as the certificate for entering a housing estate (Fig.3 (a)), some bus routes were

closed, the subway and taxi adjusted the serving time and encouraged the passengers to scan the

code to record the travel information (Fig.3 (b) and (c)). During the social distancing period, a

sharp decrease of travel intensity appeared, the average value of which was 0.87 (2.21 in 2019 at

the same time), decreasing nearly 80% from the normal period. The reopened period was from 10

February to 31 March, the economy and travel was reopened, the travel intensity was slightly

increasing with average value of 2.87 (2.3 times of that in the social distancing period), but which

was only about 59% of the value in the same time period in 2019, indicating people's travel desire

was not recovered yet.

<Fig.2 Temporal variability of the confirmed cases and travel intensity>
113
114 <Fig.3 Social distancing, traffic control and travel restriction measures conducted in Shenzhen
115 (a) real-name registration and information collection
116 (b) registration and information collection on the subway
117 (c) registration and information collection on the texi>
118

The intracity traffic (the monthly passenger population) and intercity traffic (the monthly

public transport population) revealed the similar situation of the traffic and travel conditions of

Shenzhen. In January 2020, the total volume of passenger population (including passengers

transferred by civil aviation, highway, rail way and waterway) was about 17.73 million

person/times, decreasing about 6.04% compared to 2019, while which was 3.75 million

person/times in February 2020, decreasing about 78.82% and 39.48% compared to January in

2020 and February in 2019, respectively. In March 2020, the total volume of passenger population

was 6.79 million person/times, increasing about 80.66% compared to February in 2020, but which

decreased about 45.89% compared to March in 2019. The total volume of public transport

population (including intercity passengers transferred by bus, subway taxi and tramcars) changed

rapidly in the same period. In January 2020 the total volume was about 250 million person/times,

while it was 41.77 million person/times in February 2020, decreasing about 83.29% and 48.23%

compared to January 2020 and February 2019, respectively.

< Fig.4 The volume and variable ratio of monthly passenger population and public transport

population in Shenzhen from January to March in 2020>

With rapidly decreasing of the volume of transport population, the air quality of Shenzhen

improved in 2020. Fig.5 illustrated the concentration variability of NO_2, PM_{10} and $PM_{2.5}$, both of

which decreased compared to 2019. The mean concentrations of NO_2, PM_{10} and $PM_{2.5}$ in 2020

from January to March were 21.82, 37.54 and 22.53 ug/m^3, while which were 28.51, 44.52 and

28.02 ug/m^3, respectively (Fig.5 (a)). More specifically, the concentrations of air pollutants varied

in the similar tendency of the volume of transport population in different time period of 2020. In

normal period, the concentrations of NO_2, PM_{10} and $PM_{2.5}$ were relatively higher with mean value

of 28.77, 52.77 and 30.86 ug/m^3, while in social distancing period the mean value of which were

12.39, 26.83 and 19.33 ug/m^3, with an average decreasing ratio about 50%. In reopened period,

the concentrations of NO_2, PM_{10} and $PM_{2.5}$ slightly increased with mean value of 22.16, 34.75 and

20.06 ug/m^3, and which were both lower than the monitored results of the normal period in 2020

and the same time period in 2019.

< Fig.5 The concentration variability of NO_2, PM_{10} and $PM_{2.5}$ ((a) the comparative results in 2020

and 2019; (b) the concentration of the air pollutants in different time period in 2020)>

152 3.2 correlation and regression analysis of traffic control and travel restriction, air pollution

153 and COVID-19 infection

To reveal the relationship of traffic control and travel restriction, air pollution and

COVID-19 infection, both regression and correlation analysis among the travel intensity,

concentration of NO_2, PM_{10}, $PM_{2.5}$ and the number of confirmed cased were conducted. The one

sample Kolmogorov-Smirnov method test was used to evaluate the variable data distribution. The

results indicated that the distribution of NO_2, PM_{10}, $PM_{2.5}$ and travel intensity was normal with

standard deviation of 8.5, 14.85, 8.58 and 1.52, respectively, while the distribution of the number

of COVID-19 confirmed cases was not normal. Thus, Spearman's correlation test, a wildly used

correlation method was conducted to analyze the relationship of the variables (Wang et al., 2019;

Ma et al., 2020). The results indicated that the number of COVID-19 confirmed cases shown

significant negative correlation with travel intensity (Table 1), with correlation coefficient -0.549

(0.01 level, 2 tailed), indicating reverse variable tendency of travel intensity and the number of

COVID-19 confirmed cases, which mean that traffic control and travel restriction would slow

down and prevent the spread of the viruses at the outbreak period. The concentration of NO_2 also

shown a significant negative correlation with the number of COVID-19 confirmed cases

(correlation coefficient -0.433, 0.01 level, 2 tailed), a possible reason was that the increasing

number of COVID-19 confirmed cases inhibited the travel desire, and decreased volume of traffic

resulting in the reduction of NO_2 emissions. The travel intensity shown significant positive

correlation with the concentrations of NO_2, PM_{10}, and $PM_{2.5}$, and the correlation coefficients were

0.765, 0.569 and 0.354 (0.01 level, 2 tailed), respectively, suggesting traffic-related air pollution in

Shenzhen. The air pollutants also had shown significant positive correlation with each other. The

correlation coefficient between NO_2 and PM_{10}, NO_2 and $PM_{2.5}$ were 0.556 and 0.411 (0.01 level, 2

tailed), while the correlation coefficient between PM_{10} and $PM_{2.5}$ was 0.906 (0.01 level, 2 tailed).

177   <Table 1 Spearman's correlation test results of the variables>

179   The nonlinear regression of travel intensity and COVID-19 confirmed cased was illustrated

in Fig.6, the results fitted with cubic equation and the R^2 equaled to 0.502 (n=73). The travel

intensity and the air pollutants fitted with linear regression, and the slopes were positive while R^2

equaled to 0.34 (travel intensity with PM_{10}), 0.52 (travel intensity with NO_2) and 0.14 (travel

intensity with $PM_{2.5}$), indicating a relatively strong correlation, and these results were in

accordance with the Spearman's test results. The number of COVID-19 confirmed cases and the

air pollutants fitted with nonlinear regression, with R^2 equaled to 0.25 (confirmed cases with NO_2),

0.01 (confirmed cases with PM_{10}) and 0.07 ($PM_{2.5}$), indicating a relatively weak correlation.

188   <Fig.6 Cubic regression result of COVID-19 confirmed cased and travel intensity>
189
190   <Fig.7 Regression result of air pollutants, COVID-19 confirmed cased and travel intensity>
191

192   4. Discussion and perspectives

193   The high agglomeration of air pollutants could facilitate the spread of the virus (Martelletti

and Martelletti, 2020), the study conducted by Cui et al., 2006 indicated that the mortality rate of

SARS shown linear relationship with the API (Air Pollution Index), increasing PM_{10}, $PM_{2.5}$, NO_2, and

O_3 shown associated increasing number of daily COVID-19 confirmed cases in 120 cities of China

(Zhu et al., 2020), and the results in this study indicated a significant correlation of the number of

COVID-19 confirmed cases and the traffic intensity of Shenzhen, and which had positive correlation

with the air pollutants. Thus, the generalized social distancing measures, such as traffic control and

travel restriction, although being considered as bold measures facing of political, economic, social and

religious challenges (Yezli and Khan, 2020), could be an efficient way to control and prevent the

spread of the viruses as well as improving the air quality. On the other hand, the air pollution

conditions could be a possible indicator for people to plan the travel during the epidemic prevention

and control period.

Besides, previous study conducted by Zhu et al. in the 120 cities of China indicated that air

pollutants shown significant correlation with COVID-19 confirmed cases, however, the relationship of

the confirmed cases with PM_{10} and $PM_{2.5}$ was weak in this study. Possible reasons to explain were: first,

Shenzhen was one of the best air quality cities in China, the relatively low concentrations and little

fluctuation of the air pollutants in Shenzhen weakened the relationship of the variables statistically;

second, there might be a "scale effect" on the research results, which mean the different scale of the

study area and amount of data might result in different results; third, most of the studies focused on the

relationship between air pollution and respiratory diseases indicated that the health risk was positive

correlation with the exposure concentration of air pollutants and the exposure time (Krecl et al., 2020;

Phosri et al., 2020), the while the infection time of COVID-19 was relatively short (news reported that

a man with no mask was infected within 15 seconds while shopping in a food market,

https://www.thepaper.cn/newsDetail_forward_5844448.), thus the correlation of air pollutants and

COVID-19 infection should be evaluated more accurately considering the COVID-19 infection and

pathogenesis as well as the air quality conditions. In order to reveal this, further studies focused on the

nexus could be considered to follow two routes: (1) the nexus of these variables among different cities

and countries, especially the cities with wildly different air quality conditions, (2) a comprehensive

study in one city, regarding the differences of the source of the air pollutants, different travel restriction

and traffic control measures and COVID-19 infection and pathogenesis. Moreover, differentiated

epidemic control and prevention measures should be considered according to the different situation of

cities as well as countries.

Our study had several limitations. First, the travel intensity data was processed daily data which

could not reveal the variability of traffic and travel population more accurately, hourly or minutely

traffic volume data of the main artery of the city were more useful for further analysis. Second, this

study was conducted with statistical method, the meteorological conditions were not considered. Third,

this study was conducted at city scale owing to the lack of sufficient data of other cities and countries.

Further studies should overcome these limitations and challenges, and the authors hope this study could

bring the researcher's attention on the nexus of the generalized social distancing, air pollution and

COVID-19 infection, and conquer this novel disease.

5. Conclusion

Generalized social distancing, including traffic control and travel restriction has significant

correlation with the number of COVID-19 confirmed cases, which shows negative correlation with the

traffic intensity of the city, thus traffic control and travel restriction would slow down and prevent the

spread of the viruses at the outbreak period. NO_2, PM_{10} and $PM_{2.5}$ show significant positive correlation

with the traffic intensity, the research focused on the nexus of social distancing, air pollution and

COVID-19 should been enhanced in future.

Acknowledgements

The authors are grateful to Baidu qianxi (http://qianxi.baidu.com/) that provides the city travel

intensity data for free.

Conflicts of Interest: none.

References

Chen, H., Guo, J., Wang, C., Luo, F., Yu, X., Zhang,W., Li, J., Zhao, D., Xu, D., Gong, Q., 2020.
Clinical characteristics and intrauterine vertical transmission potential of COVID-19
infection in nine pregnant women: a retrospective review of medical records. Lancet 395,
809–815. https://doi.org/10.1016/S0140-6736(20)30360-3.
Chinazzi, M., Davis, J.T., Ajelli, M., et al. 2020. The effect of travel restrictions on the spread of
the 2019 novel coronavirus (COVID-19) outbreak. Science eaba9757.
https://doi.org/10.1126/science.aba9757.
Cui Y, Zhang Z, Froines J, Zhao J, Wang H, Yu S, Detels R. 2003. Air pollution and case fatality
of SARS in the People's Republic of China: an ecologic study. Environ Health. 2(1)15.
Grange, K., Lewis, A., Carslaw, David. 2016. Source apportionment advances using polar plots of
bivariate correlation and regression statistics. Atmos. Environ. 145:128-134.
https://doi.org/10.1016/j.atmosenv.2016.09.016.
Krecl, P., Cipoli, Y., Targino, A., Castro, L., Gidhagen, L., Malucelli, F., Wolf, A. 2020. Cyclists'
exposure to air pollution under different traffic management strategies. Sci. Total Environ.
723:138043. https://doi.org/10.1016/j.scitotenv.2020.138043.
Ma, Y., Zhao, Y., Liu, J., He, X., Wang, B., Fu S., Yan J., Niu J., Zhou, Ji., Luo, Bin. 2020. Effects
of temperature variation and humidity on the death of COVID-19 in Wuhan, China. Sci. Total
Environ. https://doi.org/10.1016/j.scitotenv.2020.138226.
Martelletti, L., Martelletti, P. 2019. Air Pollution and the Novel Covid-19 Disease: a Putative
Disease Risk Factor. SN Compr. Clin. Med. https://doi.org/10.1007/s42399-020-00274-4.
Pan A, Liu L, Wang C, Guo H, Hao X, wang Q, et al. 2020. Association of Public Health
Interventions With the Epidemiology of the COVID-19 Outbreak in Wuhan, China. J. Am.
Med. Assoc. https://doi.org/10.1001/jama.2020.6130.
Pasquier, A., André, M. 2017. Considering criteria related to spatial variabilities for the assessment
of air pollution from traffic. Transp. Res. Proc. 25:3354-3369.
https://doi.org/10.1016/j.trpro.2017.05.210.
Phosri, A., Ueda, K., Phung, V., Tawatsupa, B. , Honda, A., Takano H. 2020. Effects of ambient air
pollution on daily hospital admissions for respiratory and cardiovascular diseases in Bangkok,
Thailand. Sci. Total Environ. 651:1144-1153. https://doi.org/10.1016/j.scitotenv.2018.09.183.
Prin, M., Bartels, K. 2020. Social distancing: implications for the operating room in the faceof
COVID-19. Can. J. Anesth. https://doi.org/10.1007/s12630-020-01651-2.
Tian, H., Liu, Y., Li, Y., et al. An investigation of transmission controlmeasures during the first 50

days of the COVID-19 epidemic in China. 2020. Science. https://doi.org/10.1126/science.
abb6105.

Vellingiri, B., Jayaramayya, K., Iyer, M., et al. 2020. COVID-19: A promising cure for the global
panic. Sci. Total Environ. https://doi.org/10.1016/j.scitotenv.2020.138277.

Wang, Z., Lv, J., Tan, Y., Guo, M., Gu, Y. Xu, S., Zhou, Y. 2019. Temporospatial variations and
Spearman correlation analysis of ozone concentrations to nitrogen dioxide, sulfur dioxide,
particulate matters and carbon monoxide in ambient air, China. Atmo. Pollut. Res.
10(4):1203-1210. https://doi.org/10.1016/j.apr.2019.02.003.

World Health Organization. WHO Timeline - COVID-19. 8 April 2020.
https://www.who.int/news-room/detail/08-04-2020-who-timeline---covid-19.

World Health Organization. Coronavirus disease (COVID-19) Pandemic. 18 April 2020.
https://www.who.int/emergencies/diseases/novel-coronavirus-2019.

Wong E, Ho KF, Wong SY, Cheung AW & Yeoh E. Workplace safety and coronavirus disease
(COVID-19) pandemic: survey of employees. [Submitted]. Bull World Health Organ. E-pub:
20 March 2020. doi: http://dx.doi.org/10.2471/BLT.20.255893.

Xie, J., Zhu, Y. 2020. Association between ambient temperature and COVID-19 infection in 122
cities from China. 2020. Sci. Total Environ. 724, 138201
https://doi.org/10.1016/j.scitotenv.2020.138201.

Yezli, S., Khan, A. 2020. COVID-19 social distancing in the Kingdom of Saudi Arabia: Bold
measures in the face of political, economic, social and religious challenges. Travel Med.
Infect. Dis. In press. https://doi.org/10.1016/j.econlet.2020.109128.

Zhao, D., Bu, L., Alippi, C., Wei, Q. 2017.A Kolmogorov-Smirnov Test to Detect Changes in
Stationarity in Big Data. IFAC-PapersOnLine 50(1): 14260-14265.
https://doi.org/10.1016/j.ifacol.2017.08.1821.

Zhu, F., Ding, R., Lei, R., Cheng, H., Liu, J., Shen, C., Zhang, C., Xu, Y., Xiao, C., Li, X., Zhang,
J., Cao, J. 2019. The short-term effects of air pollution on respiratory diseases and lung
cancer mortality in Hefei: A time-series analysis. Respir. Med. 146:57-65.
https://doi.org/10.1016/j.rmed.2018.11.019.

Zhu, Y., Xie, J., Huang, F., Cao, L. 2020. Association between short-term exposure to air pollution
and COVID-19 infection: Evidence from China. Sci. Total. Environ.
https://doi.org/10.1016/j.scitotenv.2020.138704.

Figure list

Fig.1. Location of the study area and the air quality monitoring stations

Fig.2 Temporal variability of the confirmed cases and travel intensity

Fig.3 Social distancing, traffic control and travel restriction measures conducted in Shenzhen
(a) real-name registration and information collection
(b) registration and information collection on the subway
(c) registration and information collection on the texi>

Fig.4 The volume and variable ratio of monthly passenger population and public transport population in Shenzhen from January to March in 2020

Fig.5 The concentration variability of NO_2, PM_{10} and $PM_{2.5}$ ((a) the comparative results in 2020 and 2019; (b) the concentration of the air pollutants in different time period in 2020)

Fig.6 Cubic regression result of COVID-19 confirmed cased and travel intensity

Fig.7 Regression result of air pollutants, COVID-19 confirmed cased and travel intensity

Fig.1. Location of the study area and the air quality monitoring stations

Fig.2 Temporal variability of the confirmed cases and travel intensity

Fig.3 Social distancing, traffic control and travel restriction measures conducted in Shenzhen

(a) real-name registration and information collection

(b) registration and information collection on the subway

(c) registration and information collection on the taxi

Fig.4 The volume and variable ratio of monthly passenger population and public transport population in Shenzhen from January to March in 2020

Fig.5 The concentration variability of NO_2, PM_{10} and $PM_{2.5}$ ((a) the comparative results in 2020 and 2019; (b) the concentration of the air pollutants in different time period in 2020)

Fig.6 cubic regression result of COVID-19 confirmed cased and travel intensity

Fig.7 Regression result of air pollutants, COVID-19 confirmed cased and travel intensity

Table list

Table 1 Spearman's correlation test results of the variables

Table 1 Spearman's correlation test results of the variables

			travel intensity	confirmed cases	NO2	PM10	PM2.5
Spearman's rho	travel intensity	Correlation Coefficient	1.000	-.549[**]	.765[**]	.569[**]	.354[**]
		Sig. (2-tailed)		.000	.000	.000	.001
		N	91	73	91	91	91
	confirmed cases	Correlation Coefficient	-.549[**]	1.000	-.433[**]	-.115	.090
		Sig. (2-tailed)	.000		.000	.332	.451
		N	73	73	73	73	73
	NO2	Correlation Coefficient	.765[**]	-.433[**]	1.000	.556[**]	.411[**]
		Sig. (2-tailed)	.000	.000		.000	.000
		N	91	73	91	91	91
	PM10	Correlation Coefficient	.569[**]	-.115	.556[**]	1.000	.906[**]
		Sig. (2-tailed)	.000	.332	.000		.000
		N	91	73	91	91	91
	PM2.5	Correlation Coefficient	.354[**]	.090	.411[**]	.906[**]	1.000
		Sig. (2-tailed)	.001	.451	.000	.000	
		N	91	73	91	91	91

**. Correlation is significant at the 0.01 level (2-tailed).

*. Correlation is significant at the 0.05 level (2-tailed).