1

Title:

Impact of mitigating interventions and temperature on the instantaneous reproduction number in the COVID-19 epidemic among 30 US metropolitan areas

Author:

Xinhua Yu

Affiliation:

Division of Epidemiology, Biostatistics and Environmental Health

School of Public Health

University of Memphis

Email: xyu2@memphis.edu

Phone: 1-901-678-3433

Running title:

intervention & temperature on reproduction number

Keywords: virus transmissibility, reproduction number, mitigating intervention, temperature, epidemic, COVID-19

Impact of mitigating interventions and temperature on the instantaneous reproduction number in the COVID-19 epidemic among 30 US metropolitan areas

Abstract

Background: After more than three months into the coronavirus disease (COVID-19) epidemic, over 170,000 people had died worldwide. The current study aims to evaluate how mitigating interventions affected the epidemic process in the 30 largest metropolitan areas in the US and whether temperature played a role in the epidemic process.

Methods: Publicly available COVID-19 cases and deaths data and weather data were analyzed at the metropolitan level. The time-varying reproductive numbers were used to explore the trends.

Results: We found that virus transmissibility, measured by instantaneous reproduction number (R_t), had declined significantly since the end of March for all areas and almost all of them reached a R_t of 1 or below by April 15, 2020. Cities with warm temperature tended to have a lower peak R_t than that of cities with cold temperature. However, large geographic variations exist.

Conclusions: Though the end of epidemic of COVID-19 is near, temperature may have some weak effects on the virus transmission, and the return of the coronavirus outbreak is still possible.

Introduction

The coronavirus disease (COVID-19) epidemic caused by the novel severe acute respiratory

syndrome (SARS) associated coronavirus (SARS-CoV 2) infection [1] has not only affected

more than 2 million people and caused over 170,000 deaths worldwide

(https://coronavirus.jhu.edu/map.html, accessed April 20, 2020), but also induced significant

anxiety among the public [2]. Many people raised concerns about whether the stringent

interventions were over-reacted, and whether a second wave of outbreak was possible. In the

2003 SARS epidemic, the virus went away after June 2003 and never came back[3]. Will this

happen to SARS-CoV2?

There are several major differences between 2003 SARS coronavirus and 2019 SARS CoV2 [3,

4]. The 2003 coronavirus had much higher virulence, resulting in higher hospitalization and

mortality rates than the 2019 coronavirus. The transmission of the 2003 coronavirus almost

exclusively occurred among symptomatic cases[5], while the 2019 coronavirus can cause a large

percent of asymptomatic cases who can also transmit virus [6, 7]. Furthermore, the 2019

coronavirus is also circulating in the southern hemisphere where the current temperature is

warmer than that of northern hemisphere. By late 2020, it is possible the coronavirus may

circulate back to the northern hemisphere, leading to a second wave of epidemic[8]. Therefore, it

is important to evaluate the effects of mitigating interventions and examine whether temperature

may affect the virus transmissibility and virulence.

One key measure of virus transmissibility during an epidemic is effective reproduction number

(R), the average number of secondary cases infected by a primary case [9-11]. Based on the

susceptible-infectious-removed (SIR) model, the reproduction number can be conceptualized as

(number of contacts)*(infectivity per contact)*(generation interval), where generation interval

refers to the average duration between the time when a primary case becoming infectious and the

time when secondary cases being infected[10]. Clearly, interventions such as social distancing,

stay home rule, school or office closures, and prohibiting large gatherings will reduce the number

of contacts, thus reducing R. On the other hand, a lower virus infectivity can also reduce R,

assuming the number of contacts remains unchanged. Therefore, exploring the changes of R over

time and across different regions can shed new lights on the impact of interventions and

environmental factors during the epidemic.

A few studies have used time varying R_t to explore the effects of intervention on the epidemic

process[12-16]. For example, one recent study found significant effects of nonpharmaceutical

interventions on the transmissibility of SARS-CoV2, measured by R_t, in Hongkong[16].

However, few studies examined the impact of environmental factors on the COVID-19 epidemic.

A few unpublished manuscripts have examined the association between temperature and

COVID-19 case counts and found no or a negative but weak association[17-19]. However, their

studies were based on case counts among different countries, which subjects to myriads of

confounding effects due to different diagnostic criteria, availability of detection kits and

reporting biases.

In this study, we will compare the magnitude and changes of time-varying (instantaneous)

effective reproduction numbers (R_t) among 30 largest metropolitan areas in the US. We

hypothesize that stringent interventions are effective in curbing the epidemic, but temperature

may also facilitate the decline of the epidemic in some regions.

Methods

Data source

We obtained daily COVID-19 cases and deaths at the US county level from the data repository

provided by New York Times (https://github.com/nytimes/covid-19-data, accessed on April

17,2020). We further limited those counties to the 30 largest metropolitan areas (Table 1). All

cases and deaths were summarized at the metropolitan level. The sizes of total population and

people aged 65 or above for each metropolitan area were obtained from census bureau website.

Information about stay-home rule for each state was scraped from popular news media. The

historical daily average temperature was obtained from national climate data online

(https://www7.ncdc.noaa.gov/CDO), mostly based on temperature collected from stations at each

metropolitan's main airport.

Reproduction number and serial interval

The time varying reproduction number (R_t) was proposed by Cori A. et al. [11, 20]. This

approach assumes the occurrence of secondary cases follows a Poisson distribution, conditioning

on the time position in the whole infectious period of the primary case. The overall R_t at time t of

the epidemic is the average number of secondary cases for all prior infected cases who are still

infectious at a time window (t-s, t). The estimate is based on current secondary cases, not the

future infections. Therefore, it can be viewed as instantaneous R_t. To smooth the estimates, a

weighted average of R_t over a sliding window is used (e.g., one week window). Consequently,

we used a 7-day moving average of prior temperature when assessing the association between R_t

and temperature.

The estimation of R_t also depends on the estimation of virus infectivity profile that is

approximated from the distribution of generation interval. Since it is difficult to accurately

estimate generation interval, serial interval is often used in calculating R_t [11]. The serial interval

refers to the average duration between symptom onset of a primary case and symptom onset of

secondary cases. In this study, we assumed the serial interval has a gamma distribution with

mean=4.7 days and standard deviation = 2.9 days [21, 22], similar to that of recent studies[15].

This serial interval was applied to all regions throughout the whole study period to ensure the

compatibility of R_t across regions. However, different regions at different time might have

different serial intervals due to interventions and other environmental and social factors. We

also performed sensitivity analysis with a shorter serial interval (mean = 3.95, standard

deviation=4.75 [23] and a longer interval (mean = 7.5, standard deviation = 3.4) [4]. The patterns

of R_t were similar, except for different estimated R_t values (average peak R_t 1-4 for shorter

duration, and 2-9 for longer duration).

Statistical analysis

We adopted two time scales in the analysis. The first was calendar date to present the trends of

reproduction numbers for all metropolitan areas, starting from the date with at least 10 total

reported cases. Staggered entries into the outbreak were preserved. The second scale was the

time since the beginning of the outbreak, regardless what calendar date the outbreak happened.

This was to compare the declining patterns of R_t among metropolitan areas. We also realigned

the time scale from the peak of the outbreak. The first two weeks of R_t estimates were excluded,

as the first week R_t were zeros, and second week estimates were too variable due to small

number of cases.

Descriptive statistics and bivariate associations were reported. Pearson correlation coefficients

and student t-tests were used for comparisons. The sizes of total population and people aged 65

or older, and the percent of positive tests at each date were used for adjustment. R package

EpiEstim was used [11] to estimate the instantaneous reproduction numbers.

There were many statistical comparisons involved. Although we did not adjust for multiple

comparisons, we were cautious about over interpretations and conducted statistical tests only

between prior selected pairs (e.g., southern versus northern metropolitan areas).

In this study, the author has no financial and conflict of interest to disclose. The ethics approval

was exempted for this study, as no human subjects were involved, and all data were publicly

available. The statistical codes and data will be available online (*github address after blind*

review).

Results

The basic characteristics of metropolitan areas were presented in Table 1. All metropolitan areas

had at least 1.5 million people in 2019 and over 1,000 confirmed cases. The case-fatality rates

varied from 0.89 per 100 people in Salt Lake City to 7.28 per 100 people in Detroit. However,

since there were large variations in case ascertainment criteria and availability of detection kits

among different regions, comparing case-fatality rates was unreliable. Meanwhile, since majority

of deaths occurred among elderly people, we compared the ratios of deaths to the size of elderly

population among metropolitan areas. The ratios were generally lower in areas with warm

weather (mean 0.026, range 0.01 to 0.09 per 100 elderly), and higher in areas with cold

temperature (mean: 0.065, range 0.01 to 0.26 per 100 elderly) (p for difference = 0.046).

The trends of R_t for 30 metropolitan areas were shown in Figure 1a-1d, grouped by geographic

locations and temperature conditions. Overall, the instantaneous R_ts in all areas reached peaks or

some stable points after two to three weeks, decreased significantly since the end March, and

most areas reached a R_t of 1 or less by April 15. It is of note that around the week of March 25,

many schools were closed and many companies started offering employees working from home.

The US government has issued COVID-19 coping guideline to all US citizens, and many states

also issued stay-home rule.

Figure 1a compared R_t trends between typical northeastern cities and southern cities. Boston,

Chicago, New York and Philadelphia started the epidemic earlier, had higher peak R_ts than that

of Miami, Orlando, Houston, and Los Angeles. After some initial increases (though peaked at

different dates), all northern cities declined sharply after the mid-March. In addition, the

trajectories of Houston and Los Angeles were similar with initial peaks at around March 18,

somewhat decreased and then were stable around March 25. For Miami and Orland, the R_ts were

quite stable during the week of March 25, and declined sharply after about March 28. However,

the slopes of decline, when aligned by the time since the peak R_t, were similar except a few

spikes in Houston and Los Angeles (appendix Figure 1a).

On the other hand, the R_t curves were indistinguishable between upper midwestern cities and

other southern cities (Figure 1b). Upper midwestern cities except Pittsburg all had earlier

interventions in the mid-March (appendix Table 1 for dates stay-home rule issued). In addition,

the west coastal cities had an early start of the epidemic, and R_t curves were less volatile than

that of other cities during the study period (Figure 1c). The unusually high R_t in Salt Lake City

in the early epidemic may be due to small number of cases during that period (Figure 1d).

Furthermore, after realigning the starting time from their respective outbreak peaks for all

metropolitan areas, the overall declining patterns were similar among all regions (Appendix

Figure 1a-d).

To evaluate the association between R_t and temperature across regions, we compared the highest

R_t (occurred after the first two weeks) among them (Figure 2), most cities had a peak R_t between

1 and 5. At the low temperature, peak R_ts varied significantly. Cities with cold average

temperature generally had higher R_ts than those with warm temperature. However, upper

midwestern cities such as Minneapolis-St. Paul, Milwaukee, and Columbus had much lower

peak R_ts than the rest of cities. The average peak of R_ts in Boston, Chicago, New York, and

Philadelphia were marginally higher than that of Houston, Los Angeles, Orlando and Miami ($p =$

0.07). In addition, we also arbitrarily examined the R_t patterns on the 15^{th} day after the outbreak

and on March 24, 2020 when most interventions had not fully executed (Appendix Figure 2a and

2b). These cross-sectional analyses demonstrated similar patterns to that of peak R_t.

Discussion

Overall, since the end of March, the instantaneous reproduction number (R_t) declined over time

significantly and also similarly in 30 largest metropolitan areas, and by April 15, R_ts in almost all

areas reached 1 or below, suggesting stringent interventions were effective in halting the

epidemic. However, there were large geographic variations in R_t patterns, partly due to different

levels of interventions, and partly maybe due to temperature variations.

Without effective interventions, the reproduction number will not decline until a large proportion

of susceptible people are infected. For example, during the week of March 25, R_ts in Houston,

Miami, and Orlando were relatively stable (Figure 1a). After the end of March, due to national

effort in mitigating the epidemic, all R_t curves started declining. The state of Florida, however,

did not officially issue the stay-home rule until April 3, 2020, where the curves already declined

significantly. This posed some difficulties in assessing the intervention effects precisely. On the

other hand, a few cities demonstrated some significant impact of interventions on mitigating the

epidemic. For example, some upper midwestern cities (e.g., Minneapolis-St. Paul and

Milwaukee) implemented interventions earlier, had lower peak R_ts, and their R_ts started

declining early, while cities like Pittsburg and Detroit had much higher R_t at the beginning, and

R_t declined later. Even for cities like Chicago and New York, the intervention effects were

evident based on the sharp decline of R_t since the mid-March, despite they had much higher R_ts

in the beginning of epidemic.

It has been suggested that like many other respiratory virus infections, a seasonal pattern may

exist for SARS like coronavirus [24, 25]. However, as demonstrated in this study, the association

between reproduction number and temperature was deeply confounded by interventions and

other external factors. In this study, we found the peak R_ts in warm cities were lower on average

than those in cold cities, suggesting that the virus transmissibility might be lower in warm

temperature than cold temperature.

However, our analysis warned readers about interpreting selective findings. For emerging

epidemic like COVID-19, many things had been happening simultaneously. Effective

interventions such as travelling restriction, social distancing and stay-home rules will change the

epidemic process [26-28]. The availability of testing, diverse case ascertainment criteria, the

delay of diagnosis and case isolation, incomplete contact tracing, the percent of asymptomatic

cases, and the infectivity of asymptomatic cases will profoundly affect our ability to understand

the epidemic. In this study, we observed a possible negative association between temperature and

virus transmissibility (Figure 1a and 2). However, there were large variations in the peak R_ts

among regions with lower temperature, partly due to different intervention effects and also may

be due to cultural and social differences. It is also likely that other environmental factors such as

living conditions may affect virus transmission. For example, under cold weather, most people

will stay indoors and have close and more frequent contacts with other people. The indoor

environment such as air conditioning may be conducive for virus transmission.

Furthermore, the concurrent decline of R_t and increase of temperature over time are also

confounded by the epidemic process itself. As suggested before, a significant reduction of

susceptible people will lead to a decline of R_t. Longitudinally comparing the trend of R_t and

temperature is difficult if the epidemic evolves rapidly, as in the current COVID-19 epidemic.

For instance, from March to April, the epidemic had a sharp decline, while the temperature in the

whole US was still suitable for virus transmission.

There were some limitations in our study. The most important limitation was the inability to

account for the diverse detection capacities across regions (Appendix Table 1). In regions with

lower detection capacity, not only were there fewer cases detected (especially missing those with

no or mild symptoms), but also the eligibilities for detection were more stringent. Only those

with symptoms might be offered for virus detection. Thus, a sudden increase in case counts

might not be due to an actual increase of infected people, rather it reflected the increased

availability of detection kits. This was the main reason we had highly variable estimates of R_t in

the beginning of epidemic. Additionally, with more detection kits available, we will observe

more asymptomatic or mild symptomatic cases. Our methods assumed the same virus infectivity

between symptomatic and asymptomatic cases, which is likely not true.

Our estimation of R_t relied on many assumptions. R_t is determined by both the growth rate of

new cases and the distribution of generation interval or serial interval [10]. We assumed a

universal distribution of serial interval for all regions and over the whole time period. Serial

interval may change due to interventions, regional characteristics, and the stage of epidemic.

More stringent interventions and stay-home rule may result in shorter serial interval because the

transmission will likely occur inside the families.

We were also not able to rigorously evaluate the virulence of SARS -CoV 2. Although we briefly

compared death rates across regions, due to large and unknown delays between virus infection

and death in the US, most deaths would be diagnosed several weeks before. Additionally, most

died cases were elderly people or those with existing chronic conditions. Therefore, assessing the

virulence should untangle the confounding effects by health care resource capacities, evolving

treatments, and patient's characteristics. A possible measure of virulence is the pattern of

hospitalizations, as they are less likely affected by the availability of detection. After more

hospitalization data are available and more asymptomatic and mild symptomatic cases are

diagnosed, future research should focus on the impact of epidemic on the severity of disease and

health care resources instead of the magnitude of epidemic.

In addition, as mentioned before, secondary data analysis based on existing aggregated data

suffer many types of unmeasured confounding. Therefore, our analyses were mostly descriptive

and deliberately avoided over-interpretations through numerous comparisons between regions. A

careful exploration of individual level data may shed some lights on these issues.

Finally, our study has some unique strengths. First, we focus on large metropolitan areas to

ensure enough cases, have comparable societal structure, individual behaviors and health care

resources. Second, we explored the changes of effective reproduction number (R_t) over time and

across regions to understand the impact of interventions and temperature on virus transmissibility.

We presented and compared all regions to avoid selective reporting bias. To study the

association between temperature and the epidemic process, we focus on the comparisons of R_t at

the peak of outbreak and before interventions, presumably less confounded by interventions and

other factors.

In summary, we observed strong effects of interventions on mitigating the COVID-19 epidemic

in 30 US metropolitan areas. There was a possible negative association between instantaneous

reproduction number and temperature. However, whether this predicts the coronavirus will

disappear in the summer and never come back, like that of 2003 SARS coronavirus, is hard to

tell. Given that the virus is circulating in both northern and southern hemisphere, we need to be

vigilant about a possible second wave of outbreak later of the year.

References:

1.   Zhu, N., et al., *A Novel Coronavirus from Patients with Pneumonia in China, 2019.* N Engl J Med, 2020. **382**(8): p. 727-733.

2.   *Pew Research Center: Most Americans Say Coronavirus Outbreak Has Impacted Their Lives.* 2020, Pew Research Center.

3.   Peiris, J.S., et al., *The severe acute respiratory syndrome.* N Engl J Med, 2003. **349**(25): p. 2431-41.

4.   Li, Q., et al., *Early Transmission Dynamics in Wuhan, China, of Novel Coronavirus-Infected Pneumonia.* N Engl J Med, 2020. **382**(13): p. 1199-1207.

5.   Fraser, C., et al., *Factors that make an infectious disease outbreak controllable.* Proc Natl Acad Sci U S A, 2004. **101**(16): p. 6146-51.

6.   Li, C., et al., *Asymptomatic and Human-to-Human Transmission of SARS-CoV-2 in a 2-Family Cluster, Xuzhou, China.* Emerg Infect Dis, 2020. **26**(7).

7.   Bai, Y., et al., *Presumed Asymptomatic Carrier Transmission of COVID-19.* JAMA, 2020.

8.   Lipsitch, M., D.L. Swerdlow, and L. Finelli, *Defining the Epidemiology of Covid-19 - Studies Needed.* N Engl J Med, 2020. **382**(13): p. 1194-1196.

9.   Hethcote, H.W., *The Mathematics of Infectious Diseases.* SIAM Review 2000. **42**(4): p. 54.

10.   Wallinga, J. and M. Lipsitch, *How generation intervals shape the relationship between growth rates and reproductive numbers.* Proc Biol Sci, 2007. **274**(1609): p. 599-604.

11.   Cori, A., et al., *A new framework and software to estimate time-varying reproduction numbers during epidemics.* Am J Epidemiol, 2013. **178**(9): p. 1505-12.

12.   Cowling, B.J., L.M. Ho, and G.M. Leung, *Effectiveness of control measures during the*

SARS epidemic in Beijing: a comparison of the Rt curve and the epidemic curve.

Epidemiol Infect, 2008. **136**(4): p. 562-6.

13.   Cauchemez, S., et al., *Unraveling the drivers of MERS-CoV transmission.* Proc Natl Acad

Sci U S A, 2016. **113**(32): p. 9081-6.

14.   Gastanaduy, P.A., et al., *Impact of Public Health Responses During a Measles Outbreak*

in an Amish Community in Ohio: Modeling the Dynamics of Transmission. Am J

Epidemiol, 2018. **187**(9): p. 2002-2010.

15.   Zhang, J., et al., *Evolving epidemiology and transmission dynamics of coronavirus*

disease 2019 outside Hubei province, China: a descriptive and modelling study. Lancet

Infect Dis, 2020.

16.   Cowling, B.J., et al., *Impact assessment of non-pharmaceutical interventions against*

coronavirus disease 2019 and influenza in Hong Kong: an observational study. Lancet

Public Health, 2020.

17.   Pawar, S., et al., *Effects of temperature on COVID-19 transmission*, Y. University, Editor.

312   2020.

18.   Bannister-Tyrrell, M., et al., *Preliminary evidence that higher temperatures are*

associated with lower incidence of COVID-19, for cases reported globally up to 29th

February 2020. medRxiv, 2020: p. 2020.03.18.20036731.

19.   Troplett, M., *Evidence that higher temperatures are associated with lower incidence of*

COVID-19 in pandemic state, cumulative cases reported up to March 27, 2020. 2020.

20.   Thompson, R.N., et al., *Improved inference of time-varying reproduction numbers during*

infectious disease outbreaks. Epidemics, 2019. **29**: p. 100356.

21. Nishiura, H., N.M. Linton, and A.R. Akhmetzhanov, *Serial interval of novel coronavirus*

(COVID-19) infections. Int J Infect Dis, 2020. **93**: p. 284-286.

22. Lauer, S.A., et al., *The Incubation Period of Coronavirus Disease 2019 (COVID-19)*

From Publicly Reported Confirmed Cases: Estimation and Application. Ann Intern Med,

2020.

23. Du, Z., et al., *Serial Interval of COVID-19 among Publicly Reported Confirmed Cases.*

Emerg Infect Dis, 2020. **26**(6).

24. Lin, K., et al., *Environmental factors on the SARS epidemic: air temperature, passage of*

time and multiplicative effect of hospital infection. Epidemiol Infect, 2005. **134**(2): p. 7.

25. Shi, P., et al., *The impact of temperature and absolute humidity on the coronavirus*

disease 2019 (COVID-19) outbreak - evidence from China. medRxiv, 2020: p.

2020.03.22.20038919.

26. Chinazzi, M., et al., *The effect of travel restrictions on the spread of the 2019 novel*

coronavirus (COVID-19) outbreak. Science, 2020.

27. Hellewell, J., et al., *Feasibility of controlling COVID-19 outbreaks by isolation of cases*

and contacts. Lancet Glob Health, 2020. **8**(4): p. e488-e496.

28. Ferguson, N., et al., *Report 9: Impact of non-pharmaceutical interventions (NPIs) to*

reduce COVID-19 mortality and healthcare demand 2020, Imperial College: UK.

Table 1: Characteristics and death rates for 30 largest metropolitan areas

Metropolitan area	Total population	Age >=65	Cumulative cases	Cumulative deaths	Deaths per 100 Cases	Deaths per 100 old people
New York	16,669,277	2,789,981	103,943	4,358	4.19	0.16
Los Angeles	10,039,107	1,493,190	10,854	455	4.19	0.03
Washington DC - Baltimore	9,360,001	1,370,637	13,475	350	2.60	0.03
Houston	7,066,141	865,112	6,097	102	1.67	0.01
St. Francisco - St. Jose	6,463,637	991,147	5,364	173	3.23	0.02
Chicago	6,021,020	950,164	21,466	894	4.16	0.09
St Louis	5,485,267	797,415	3,936	140	3.56	0.02
Atlanta	5,261,067	741,175	8,165	274	3.36	0.04
Dallas-Fort Worth	5,081,942	615,560	3,295	80	2.43	0.01
Philadelphia	3,815,431	691,016	15,203	496	3.26	0.07
Minneapolis-St. Paul	3,654,908	580,638	1,276	71	5.56	0.01
Cleveland - Akron	3,149,448	669,206	2,587	132	5.10	0.02
Seattle	3,074,865	444,891	6,842	407	5.95	0.09
Boston	2,979,288	529,944	15,218	515	3.38	0.10
Denver	2,967,239	419,589	5,015	198	3.95	0.05
Miami - Fort Lauderdale	2,716,940	467,586	8,325	183	2.20	0.04
Charlotte	2,675,243	424,892	2,067	44	2.13	0.01
Orlando	2,608,147	430,027	1,989	33	1.66	0.01
Portland	2,492,412	424,694	1,289	57	4.42	0.01

Sacramento - Oakland	2,363,730	419,255	1,169	47	4.02	0.01
Pittsburg	2,317,600	542,666	1,652	81	4.90	0.01
Las Vegas	2,266,715	358,821	2,625	121	4.61	0.03
Cincinnati	2,198,450	394,794	1,249	59	4.72	0.01
Kansas	2,157,990	368,576	1,288	76	5.90	0.02
Columbus	2,122,271	325,706	1,776	35	1.97	0.01
Detroit	2,116,944	387,897	13,828	1,006	7.28	0.26
Indianapolis	2,074,537	337,623	5,373	284	5.29	0.08
Durham-Raleigh	1,974,709	289,249	1,340	19	1.42	0.01
Salt Lake City	1,880,948	207,100	1,804	16	0.89	0.01
Milwaukee	1,575,179	288,873	2,348	137	5.83	0.05

344 **Figure 1a-1d: Time trends of instantaneous reproduction number for 30 largest**
345 **metropolitan areas**

347   **(a)**

349   **(b)**

351 **(c)**

353 **(d)**

356   **Figure 2: Association between maximal reproduction number and 7-day average**
357   **temperature among 30 metropolitan areas**

Appendix:

Appendix Table 1: Positive detection rates at various time by state

State	Metro areas	%positive as of March 25, 2020	%positive as of April 15, 2020	Date of stay at home rule	% positive at rule mandating
CA	Los Angeles, St. Francisco-St. Jose, Sacramento-Oakland	12.9%	10.6%	19-Mar-20	9.5%
CO	Denver	11.8%	20.4%	11-Apr-20	19.9%
DC	Washington DC - Baltimore	11.4%	19.3%	1-Apr-20	15.2%
FL	Miami-Fort Lauderdale, Orlando	9.9%	10.4%	3-Apr-20	10.5%
GA	Atlanta	20.2%	23.1%	2-Apr-20	23.3%
IL	Chicago	13.1%	21.0%	21-Mar-20	12.1%
IN	Indianapolis	14.2%	18.7%	24-Mar-20	12.5%
KS	Kansas	5.1%	9.9%	30-Mar-20	7.5%
MA	Boston	9.3%	22.9%	24-Mar-20	8.4%
MI	Detroit	48.4%	31.0%	24-Mar-20	52.6%
MN	Minneapolis-St. Paul	2.5%	4.6%	27-Mar-20	2.8%
MO	St. Louis	49.1%	10.1%	3-Apr-20	9.8%
NC	Durham-Raleigh, Charlotte	4.8%	7.7%	30-Mar-20	6.3%
NV	Las Vegas	7.0%	11.7%	1-Apr-20	10.0%
NY	New York	29.8%	40.4%	22-Mar-20	24.7%
OH	Cleveland, Columbus, Cincinnati	4.8%	11.2%	22-Mar-20	71.5%
OR	Portland	4.6%	5.0%	23-Mar-20	5.0%
PA	Philadelphia, Pittsburg	9.1%	19.6%	1-Apr-20	12.0%
TX	Houston, Dallas-Fort Worth	7.2%	10.4%	2-Apr-20	9.2%
UT	Salt Lake City	5.1%	5.4%	27-Mar-20	5.2%
WA	Seattle	9.4%	8.7%	23-Mar-20	7.8%
WI	Milwaukee	5.5%	8.6%	25-Mar-20	5.5%

Appendix Figure 1a – d, Declining trend of instantaneous reproduction number over time after the peak of epidemic

(a)

(b)

373   **(c)**

375   **(d)**

Appendix Figure 2a-b: Variations of instantaneous reproduction numbers across metropolitan areas.

(a)

(b)