Title page

Correlation analysis of risk factors and GSI score of a medical team assisting

Wuhan city during the epidemic of COVID-19 in China -A cohort study

Academic titles of all authors:

Cheng Wang [1], Zhaohui Lu [2] ,Jingquan Wang [3] ,Xiaolei Jing [4] ,Yunyun Fang [5], Yanlin Wang [6], Jinlong

Zhang [*1,7]

First author: Cheng Wang [1] MA.Sc

Affiliations:

1. Department of Rehabilitation Medicine, The First Affiliated Hospital of University of Science

and Technology of China (Anhui Provincial Hospital), Hefei, Anhui Province, People's Republic

of China

Email: 2814836650@qq.com

Co-first author: Zhaohui Lu [2] MA.Sc

Affiliations:

2.Infectious Diseases Division, The First Affiliated Hospital of University of Science and

Technology of China (Anhui Provincial Hospital), Hefei, Anhui Province, People's Republic of

China

E-mail: lzh3290@sina.com

[*]**Corresponding Author :** Jinlong Zhang [1,6] MA.Sc

Affiliations:

1. Rehabilitation Department, The first affiliated Hospital of university of science and technology

of China (Anhui Provincial Hospital), Hefei, Anhui Province, People's Republic of China

7**.** Rehabilitation Department, Hefei BOE Hospital of BOE Technology Group, Hefei, Anhui

Province, People's Republic of China

Email: zhlyy520@tom.com

Address：

27   No.1 Swan Lake Road

28   Hefei City

29   Anhui Province

30   China

Code: 230001

Tel: 0086-51-62284092

Fax: 0086-51-62284091

E-mail: zhlyy520@tom.com

Other authors:

Jingquan Wang MA.Sc

Affiliations:

3. Intensive Care Unit, The First Affiliated Hospital of University of Science and Technology of China (Anhui Provincial Hospital), Hefei, Anhui Province, People's Republic of China

E-mail: jqwang604@163.com

Xiaolei Jing MA.Sc

Affiliations:

4.Department of Neurosurgery, The First Affiliated Hospital of University of Science and Technology of China (Anhui Provincial Hospital), Hefei, Anhui Province, People's Republic of China

E-mail: 3391791883@qq.com

Yunyun Fang[2] MA.Sc

Affiliations:

5.Traditional Chinese Medicine Department, Jiangning Hospital of Nanjing Medical University, Nanjing, Jiangsu Province, People's Republic of China

Email: cdfk22@tom.com

Yanlin Wang MA.Sc

Affiliations:

6.Department of ECG, The First Affiliated Hospital of University of Science and Technology of China (Anhui Provincial Hospital), Hefei, Anhui Province, People's Republic of China

E-mail:419939208@qq.com

Word count:3258

Key Points

Question: What is the correlation between risk factors and general symptom index

(GSI) score of medical team members who support Wuhan against COVID-19?

Findings: Dietary conditions (DC) had a positive and relationship between team

(RBT) had a negative correlation with GSI score of female, but for male was not

(P=0.59>0.05, P=0.08>0.05, respectively), lacking communication with teams

(LCWT) and afraid of being infected (AoBI) didn't had correlation with GSI score

between genders, a significant difference.

Meaning: Improving DC, RBT and decreasing LCWT between team members can

reduce the GSI score. Whether they are AoBI, didn't affect the psychological status,

male members have a more stable mood than female.

Abstract

Importance: There are few studies on the psychological status of medical staff during

the COVID-19 outbreak. This study is the first in the world about the psychological

status of the medical team during the COVID-19 outbreak.

Objective: To study the correlation between risk factors and general symptom index

(GSI) score of medical team members who support Wuhan against COVID-19.

Design: Cohort study.

Setting: Population-based.

Participants: Anhui Province sent a total of eight medical teams, including 1382

members, to support Hubei Province. We adopted a stratified sampling method and

selected the fourth team sent by Anhui Provincial Hospital, with a total of 137

members as our subjects.

Exposures: Four main exposures were collected, including basic information,

preparations before going to Wuhan, life issues and working issues after going to

Wuhan.

Main Outcomes and Measures: The GSI score of SCL-90 scale was used to reflect

the frequency and intensity of psychological symptoms. We made the hypothesis of

this study before data collection.

Results: 110(80.29%) members completed the questionnaire, of which, 77(70.00%)

female and 33(30.00%) male. When adjusted age, gender and covariates, DC, LCWT

had a positive correlations with GSI score(β was10.17, 95%CI was 3.30 to 17.04 for

DC, *P=0.00*<0.05; β was 11.55, 95%CI was 0.40 to 22.71 for LCWT, *P*

=0.04<0.05;respectively), RBT had a negative positive correlation with GSI score (β

was -28.09, 95%CI was -45.79 to -10.40, *P*=0.00<0.05), AoBI did not had a

correlation with GSI score (β was 11.55, 95%CI was 0.40 to 22.71, *P*=0.16>0.05).

When adjusted covariates, DC had a positive and RBT had a negative correlation with

GSI score of female (β was 13.20, 95%CI was 4.55 to 21.85, *P*=0.00<0.05; β was

-57.85, 95%CI was -94.52 to -21.18, *P*=0.00<0.05; respectively), but for male was not

(*P*=0.59>0.05, *P*=0.08>0.05, respectively), LCWT and AoBI didn't had correlation

with GSI score between genders (*P*>0.05).

Conclusions and Relevance: Improving DC, RBT and decreasing LCWT can reduce

the GSI score. AoBI didn't affect the psychological status; male members have a more

stable mood than female. Whether other countries medical team has the same result

still needs further research.

Correlation analysis of risk factors and GSI score of a medical team assisting

Wuhan city during the epidemic of COVID-19 in China -A cohort study

Abstract

Importance: There are few studies on the psychological status of medical staff during the COVID-19 outbreak. This study is the first in the world about the psychological status of the medical team during the COVID-19 outbreak.

Objective: To study the correlation between risk factors and general symptom index (GSI) score of medical team members who support Wuhan against COVID-19.

Design: Cohort study.

Setting: Population-based.

Participants: Anhui Province sent a total of eight medical teams, including 1382 members, to support Hubei Province. We adopted a stratified sampling method and selected the fourth team sent by Anhui Provincial Hospital, with a total of 137 members as our subjects.

Exposures: Four main exposures were collected, including basic information, preparations before going to Wuhan, life issues and working issues after going to Wuhan.

Main Outcomes and Measures: The GSI score of SCL-90 scale was used to reflect the frequency and intensity of psychological symptoms. We made the hypothesis of this study before data collection.

Results: 110(80.29%) members completed the questionnaire, of which, 77(70.00%) female and 33(30.00%) male. When adjusted age, gender and covariates, DC, LCWT had a positive correlations with GSI score(β was10.17, 95%CI was 3.30 to 17.04 for DC, *P=0.00*<0.05; β was 11.55, 95%CI was 0.40 to 22.71 for LCWT, *P* =0.04<0.05;respectively), RBT had a negative positive correlation with GSI score (β was -28.09, 95%CI was -45.79 to -10.40, *P*=0.00<0.05), AoBI did not had a correlation with GSI score (β was 11.55, 95%CI was 0.40 to 22.71, *P*=0.16>0.05). When adjusted covariates, DC had a positive and RBT had a negative correlation with GSI score of female (β was 13.20, 95%CI was 4.55 to 21.85, *P*=0.00<0.05; β was -57.85, 95%CI was -94.52 to -21.18, *P*=0.00<0.05; respectively), but for male was not (*P*=0.59>0.05, *P*=0.08>0.05, respectively), LCWT and AoBI didn't had correlation with GSI score between genders (*P*>0.05).

Conclusions and Relevance: Improving DC, RBT and decreasing LCWT can reduce the GSI score. AoBI didn't affect the psychological status; male members have a more stable mood than female. Whether other countries medical team have the same result still needs further research.

Keywords： Psychological status; Risk factors; COVID-19; SCL-90 scale; GSI score

1. Introduction

COVID-19 was endemic in China at the end of 2019. Data as received by WHO from national authorities by 10:00 CEST, 23 April 2020, there were 2 544 792 confirmed cases and 175 694 deaths[1], In this epidemic, China has quickly taken

various effective measures. Now the epidemic is basically under effective control in

China. These experiences are worth sharing with the world. One of the measures was

to recruit medical personnel from various provinces and cities across the country to

support the most severely affected province-Hubei, with a total of nearly 40,000

medical personnel. This type of disease has caused global panic, and medical staffs

are no exception，they may also feel panic[2-4]. The current research mainly focuses on

the psychological status of infected people[5-7]. Few people pay attention to the

psychological symptoms of medical staff，a study found that it is helpful to take

effective interventions to meet their needs if the needs of nurses caring for COVID-19

patients could be perceived well, the main needs was health and security[8]. At the time

of the COVID-19 epidemic outbreak in China, the heroic Chinese medical staffs were

under the tremendous pressure of fighting the COVID-19, whether they had a poor

psychological status? And what the risk factors were? So far, there was no research

report and also no similar literature in other countries around the world when we

search the database on line.

Symptom Checklist-90 (SCL-90) scale has been widely used in psychology

research in the world. The author used this scale to evaluate ordinary people

psychology status during COVID-19 pandemic in China, and found which also had a

significant adverse socio-psychological influence on ordinary citizens[9]. It has good

reliability and validity[10, 11], and can reflect the mental and psychological status of the

recipients from multiple dimensions. The GSI score of SCL-90 can reflect the

frequency and intensity of psychological symptoms[12, 13]. Therefore, this study

analyzed the GSI score of a medical team, we carried out systematic statistical

analysis to find out the risk factors that affected the GSI score, in order to provide a

theoretical basis for precise psychological intervention for medical staffs.

2. Methods

2.1 Design methods

We used a cohort study, using the single-blind method; the participants were not

clear about the purpose.

2.2 Data collecting methods

We used Questionnaire star APP (https://www.wjx.cn/) to collect the data, the

collected data set includes the basic information, preparations before going to Wuhan,

life issues and working issues after going to Wuhan. The basic information included

seven sub risk factors-age, working age, education background, only child of the

family, marriage status, childbirth status and family relationships. There were five sub

risk factors of preparations before going to Wuhan, including COVID-19 knowledge,

infection prevention and control knowledge, confidence to complete the task,

emergency aid experience and relationships with team (RBT). There were seven sub

risk factors of life issues after going to Wuhan, including dietary conditions (DC),

sleep quality (SQ), limit range of activities (LRoA), care about me (CAM), lavation

conditions (LC), residential conditions (RC) and surrounding conditions (SC).There

were nine sub risk factors of working issues after going to Wuhan, which were

unfamiliar with medical records system (UWMRS), unfamiliar with working

environment (UWWE), difficulty communicating with patients (DCWP), unfamiliar

with workflow (UWW), lacking medical equipments (LME), long working hours (LWH), lacking communication with teams (LCWT), difficulty communicating with local hospital (DCWLH) and fear of being infected (AoBI). The GSI score of SCL-90 scale was used to reflect the frequency and intensity of psychological symptoms

2.3 Statistical methods

In this survey study we took a purposive approach with no sample estimation. Data were analyzed using the statistical packages R (R Foundation; http://www.r-project.org;version3.4.3) and EmpowerStats (http:// www. empowerstats. com; X&Y Solutions Inc, Boston, MA). Multivariable logistic regression model, single factor logistic regression model, generalized estimated equation model, T-test and Pearson's chi-square tests were used to analyze the data, estimated change (β) and 95% confidence interval(CI) were used to establish the demographic and clinic characteristic of the sample, results were considered statistically significant with $P<0.05$.

2.4 Ethical considerations

The participants of this study filled out the questionnaire anonymously. Our research was approved by the Ethics Committee of the First Affiliated Hospital of University of Science and Technology of China (Anhui Provincial Hospital).

3. Results

3.1 Flow chart of the study

Anhui Province sent a total of eight medical teams to support Hubei, with a total of 1,362 medical members, including a total of 274 in the fourth medical team, which

were selected from the First Affiliated Hospital of Anhui Medical University and the First Affiliated Hospital of University of Science and Technology of China (Anhui Provincial Hospital). The first author of this study was one of the members of the latter medical constitution, so the object was the members of the latter as for the convenience of research. For details, see Fig1.

3.2 Demographic and risk factors of members

110(80.29%) members completed the questionnaire, of which, 77(70.00%) female and 33(30.00%) male. For basic information, 76.6% female and 69.7% male had a bachelor degree, the difference of education background ratio between the genders was statistically significant(X^2=9.77, df=3, P=0.02<0.05). 41.6% female and 21.2% male unmarried, the marriage ratio between the genders was also statistically significant(X^2=4.18, df=1, P=0.04<0.05). For preparations before going to Wuhan, the relationships with team (RBT) factor had a statistically significant between genders, 98.7% female and 90.9% male felt a good interpersonal relationship with the team(X^2=4.00, df=1, P=0.04<0.05).For life issues after going to Wuhan, 96.1% female and 84.8% male felt care about me(CAM) from the team, and the differences also had a statistically significant(X^2=4.34, df=1, P=0.04<0.05).The other factors of genders had no statistically significant(P>0.05).

3.3 Crude correlation associations of exposure risk factors and GSI score of the members

As seen in table2,we analyzed the exposure risk factors, the single factor analysis showed risk factors of basic information had no correlation with GSI score(P>0.05).

For preparations before going to Wuhan, there were five risk factors, RBT had a negative correlation with GSI score, estimated change was -20.63, 95%CI was -39.65 to -1.60, the difference had a statistically significant(P=0.04<0.05). For life issues after going to Wuhan, there were seven risk factors, DC had a positive correlation with GSI score, estimated change was 11.08, 95% CI was 4.11 to 18.05, the difference had a statistically significant(P=0.00<0.05). For working issues after going to Wuhan, there were nine risk factors, LCWT and AoBI had a positive correlation with GSI score, the estimated change was 17.11 and 16.32 respectively, 95%CI was 6.32 to 27.90 and 5.08 to 27.57 respectively, the difference had a statistically significant (P=0.00<0.05,P=0.01<0.05,respectively).

3.4 Multivariate logistic regression model for DC, LCWT, AoBI, RBT and GSI score of the members

Multivariate logistic regression analysis showed that, for DC, LCWT, AoBI and RBT, when unadjusted covariates, DC,LCWT and AoBI had a positive correlations with GSI score(P<0.05), RBT had a negative correlation with GSI score(P<0.05); When adjusted age and gender, DC,LCWT and AoBI also had a positive correlations with GSI score(P<0.05), RBT   also had a negative correlation with GSI; For DC, when adjusted LCWT, AoBI and RBT, DC also had a positive correlation with GSI score(P=0.00<0.05), estimated change was 10.17, 95%CI was 3.30 to 17.04; For LCWT, when adjusted DC, AoBI and RBT, LCWT also had a positive correlation with GSI score(P=0.04<0.05), estimated change was 11.55, 95%CI was 0.40 to 22.71; For AoBI, when adjusted DC, LCWT and RBT, AOBI did not had a correlation with

GSI score(P=0.16>0.05), estimated change was 11.55, 95%CI was 0.40 to 22.71, this

result showed that if control other covariates, AOBI will not influence GSI score. For

RBT, when adjusted DC, LCWT and AoBI, RBT also had a negative positive

correlation with GSI score (P=0.00<0.05), estimated change was -28.09, 95%OR was

-45.79 to -10.40.(Table3)

3.5 Stratification analysis of DC, LCWT, AoBI, RBT and GSI score of the members between genders

Multivariate logistic regression stratification analysis showed that DC，LCWT

and AoBI of female had a positive correlation with GSI score for non- adjusted model

and adjusted I model, RBT of female had a negative correlation with GSI score for

non- adjusted model and adjusted I model, the differences had statistically

significant(P<0.05). But for male, above factors had no correlation with GSI score for

non-adjusted model, the differences did not had statistically significant (P>0.05), DC

and RBT had no correlation with GSI score for adjusted I model, the differences also

did not had statistically significant (P>0.05), LCWT and AoBI had correlation with

GSI score for adjusted I model, the differences had statistically significant (P<0.05).

This results showed that when adjusted age, for female, the relationships of DC,

LCWT,AoBI, RBT and GSI score didn't changed. But for male, when adjusted age,

LCWT and AoBI correlations with GSI score started to appear, which showed that age

may was an effect modifier. The adjusted II model showed when adjusted covariates

of LCWT, AoBI and RBT, DC had a positive correlation with GSI score of female,

estimated change was 13.20, 95%CI was 4.55 to 21.85, the difference was statistically

significant(P=0.00<0.05), but for male was not(P=0.59>0.05). RBT had a negative

correlation with GSI score of female, when adjusted covariates of DC, LCWT and

AoBI, the estimated change was -57.85,95%CI was -94.52 to -21.18, the difference

was statistically significant(P=0.00<0.05), for male was also not(P=0.08>0.05).

LCWT and AoBI didn't had correlation with GSI score between genders, the

differences were not statistically significant (P>0.05), when adjusted covariates of DC

and RBT in adjusted II model (table4).

4. Discussion

The 2019-nCoV(2019 new coronavirus) is officially called SARS-CoV-2 (severe

acute respiratory syndrome-coronavirus-2) and the disease is named

COVID-19(Corona virus disease-2019), which is currently causing a pandemic in the

world. More than 1 million people have been infected with coronavirus, and tens of

thousands have died, causing panic in the whole society. People who are not infected

are worried and afraid of being infected[7, 14, 15]. Patients who are already infected are

worried about whether they can get timely and effective treatment. The pandemic of

SARS-CoV-2 has had significant social, psychological and economic consequences

worldwide[16], especially for low-income patients, this is a serious concern when linked

to the pandemic[17].

The outbreak of the SARS-CoV-2 is undesirable by people and needs to be

faced with all over the world, but unfortunately, there are too many news reports that

SARS-CoV-2 has caused racial hatred and violence in some countries. The occurrence

of these incidents is not conducive to fight the epidemic. Medical staffs are the main

force to fight against COVID-19. They directly face the patients who have been

infected and the chance of being infected is relatively large at work. At present, the

psychological research objects of COVID-19 are mainly patients or the general public,

but few people pay attention to and study the psychological status of medical staff.

A study found that the sudden increase in confirmed cases had brought tremendous

stress and anxiety to frontline surgical staff, and researchers believe that early

psychological intervention were essential[18]. Researchers from Italy found that at least

2,629 health workers have been infected with coronavirus since the outbreak onset in

February, representing 8.3% of total cases[19]. If the medical staffs also feel panic, it

may affect their medical behavior, such as avoiding, reducing communication with

infected people, or even leaving the job. Therefore, improving the enthusiasm of the

medical staff and reducing their panic will play a decisive role in improving the

clinical efficacy and even achieving the successful control of the epidemic.

Preliminary evidence suggests that symptoms of anxiety and depression

(16–28%) and self-reported stress (8%) are common psychological reactions to the

COVID-19 pandemic, and may be associated with disturbed sleep[20]. The outbreak of

COVID-19 infection had an effect on the psychology of the elderly people, resulting

in anxiety and depression[6]. Our research found that the GSI scores of medical team

members were not affected by age, as well as working age, educational background,

family relationship, marital status, and fertility status. Before going to Wuhan,

medical team members need to receive hospital infection knowledge prevention and

control training, learn COVID-19 knowledge, and build confidence in defeating the

1 disease. Knowledge of COVID-19, confidence to complete the task and emergency

2 aid experience also did not have correlations with GSI score. We found that after

3 training, 99.09% members are confident to fight against COVID-19, but just 42.73%

4 members had emergency aid experience and 4 (3.64%) felt that the relationship with

5 other team members was not in harmony. Such research results showed that in order

6 to control the outbreak of COVID-19, China quickly established medical teams, the

7 team members were from various clinical disciplines.

8  Before going to Wuhan, they did not have time to receive systematic emergency

9 first aid training, and even some team members may not know each other. We have

10 reviewed the literature database in detail, and found no literature about the medical

11 team against COVID-19. A study found that communication had achieved satisfactory

12 results and may play a role in the protection on the psychological condition on the

13 people with close contact with influenza A(H1N1)[21]. Another study showed that

14 patients with difficulty in communicating significant aspects of their complaints to the

15 physician or in understanding his instructions for treatment , the authors believed

16 SCL-90 was a quickly administered and easily scored test that can screen for both

17 psychopathology and communication problems[21]. Our experience is worth sharing

18 with the world, but if there is enough time, the members should best to receive

19 systematic emergency first aid training and to increase communication between them,

20 because our research found that RBT and GSI score were negatively correlated, the

21 more harmonious the relationship between the members, the lower the GSI score.

22  After going to Wuhan, the team members faced life and work issues. Our study

showed that DC and AoBI had a positive correlation with GSI score, LCWT had a

negative correlation with GSI score for non- adjusted model, but when we adjusted

age, gender and covariates, there was no correlation between AoBI and GSI score, but

correlations of DC,RBT and LCWT with GSI score didn't changed. This result

showed that the influence of AoBI on GSI score was influenced by the three

covariates of DC, RBT and LCWT. A literature published in 1990 found that refined

sucrose and caffeine free diet can significantly decline in depression on all depression

measures, including SCL-90, Beck Depression Inventory (BDI), and Interpersonal

Style Inventory (ISI)[22].A study found that adding certain substances to food can

improve sleep and depression[23]. Our research found that DC was related to GSI scores.

Good diet can improve the psychological status of medical team members, but

standards for DC are different for different people. This risk factor is greatly

influenced by personal subjective attitudes. Our medical team members came from all

corners of China. Everyone has different preferences for food. Individuals in one

study having poor diets were more likely to suffer from depression than those eating

good diets[24] .There is a Chinese proverb called "difficult to adjust for

mouth" .However, we still recommend the logistics support department of medical

team, if time and material conditions permit, it would better try to enrich and diversify

the catering, and customize the catering according to each person's different hobbies,

so that can alleviate the psychological anxiety and depression.

If the RBT was not good, it will be easy result in LCWT, there was a correlation

between the two, and some researchers had already confirmed that lacking

1. communication with others could increase anxiety or depression[25]. As said above , our

2. study also found if the members had bad RBT and LCWT, the GSI Score will

3. increased. Fighting the SARS-CoV-2 epidemic requires joint efforts of medical and

4. nursing members. Due to the short time, members were transferred from various

5. departments to form a team. Many of them did not know each other and understand

6. each other, which will increase their psychological pressure. Some studies have found

7. that recreational activities can improve psychological conditions [26, 27], so we suggest

8. that if there is time between the members, they can participate in some collective

9. recreational activities while ensuring protection, which helps increase mutual

10. understanding and can reduce the chance of bad mood.

11.   This study has a shortcoming ,due to lack of time , we did not follow up the team

12. members. We will observe the changes of their psychological status in the later period,

13. to further understand the long-term effects of RBT, LCWT and AoBI on GSI score.

14. **Declaration of competing interest**

15.   The author(s) declared no potential conflicts of interest with respect to the

16. research, authorship, and/or publication of this article.

17. **Funding source**

18.   This research received no specific grant from any funding agency in the public,

19. commercial, or not-for-profit sectors.

20. **Disclosure statement**

21.   No potential conflict of interest was reported by the authors.

22. **Author contributions**

Cheng Wang conducted analyses and drafted the manuscript in consultation with Jinglong Zhang and Yunyun Fang. All authors provided critical revisions and approved a final version of the manuscript prior to publication.

Acknowledgements

We sincerely thank Jian Tian, Yan Ma, Jun Wang, Shiyang Zhangand Kai Liu for assorting the data, we thank Changcheng Zheng, Yuyou Zhu, Xuhan Zhang, and Hongzhi Ji for devising and issuing the questionnaire. We also like to thank all the members for their cooperation in our study.

5. References

1.https://www.who.int/docs/default-source/coronaviruse/situation-reports/20200423-sitrep-94-covid-19.pdf?sfvrsn=b8304bf0_4.

2.   Xu J, Xu Q-h, Wang C-m and Wang J. Psychological status of surgical staff during the COVID-19 outbreak. *Psychiatry Research*. 2020; 288: 112955.

3.   Lu W, Wang H, Lin Y and Li L. Psychological status of medical workforce during the COVID-19 pandemic: A cross-sectional study. *Psychiatry Research*. 2020; 288: 112936.

4.   Chen Y, Zhou H, Zhou Y and Zhou F. Prevalence of self-reported depression and anxiety among pediatric medical staff members during the COVID-19 outbreak in Guiyang, China. *Psychiatry Research*. 2020; 288: 113005.

5.   Sun N, Wei L, Shi S, et al. A qualitative study on the psychological experience of caregivers of COVID-19 patients. *American Journal of Infection Control*. 2020.

6.   Meng H, Xu Y, Dai J, Zhang Y, Liu B and Yang H. The Psychological effect of COVID-19 on the Elderly in China. *Psychiatry Research*. 2020: 112983.

7.   The L. COVID-19: fighting panic with information. *The Lancet*. 2020; 395: 537.

8.   Yin X and Zeng L. A study on the psychological needs of nurses caring for patients with coronavirus disease 2019 from the perspective of the existence, relatedness, and growth theory. *International Journal of Nursing Sciences*. 2020.

9.   Tian F, Li H, Tian S, Yang J, Shao J and Tian C. Psychological symptoms of ordinary Chinese citizens based on SCL-90 during the level I emergency response to COVID-19. *Psychiatry Research*. 2020; 288: 112992.

10.   Luo W, Guo CX, Han DL and Li ZJ. Reliability and Validity of Chinese Version of the Addiction Severity Index among Drug Users in the Community. *Biomedical and Environmental Sciences*. 2012; 25: 684-689.

11.   Ransom D, Ashton K, Windover A and Heinberg L. Internal consistency and validity assessment of

SCL-90-R for bariatric surgery candidates. *Surgery for Obesity and Related Diseases*. 2010; 6: 622-627.

12. Tunçel ÖK, Kayıkçıoğlu M, Pırıldar Ş, et al. Mental Status And Physical Activity In Patients With Homozygous Familial Hypercholesterolemia: A Subgroup Analysis Of A Nation-Wide Survey (A-Hit1 Registry). *Journal of Clinical Lipidology*. 2020.

13. Meng C, Wang W, Hao Z and Liu H. Investigation on the influence of isolated environment on human psychological and physiological health. *Science of The Total Environment*. 2020; 716: 136972.

14. Vellingiri B, Jayaramayya K, Iyer M, et al. COVID-19: A promising cure for the global panic. *Science of The Total Environment*. 2020; 725: 138277.

15. Parry NMA. COVID-19 and pets: When pandemic meets panic. *Forensic Science International: Reports*. 2020; 2: 100090.

16. Yang Y, Peng F, Wang R, et al. The deadly coronaviruses: The 2003 SARS pandemic and the 2020 novel coronavirus epidemic in China. *Journal of Autoimmunity*. 2020; 109: 102434.

17. Neto MLR, de Souza RI, Quezado RMM, et al. WHEN BASIC SUPPLIES ARE MISSING, WHAT TO DO? SPECIFIC DEMANDS OF THE LOCAL STREET POPULATION IN TIMES OF CORONAVIRUS – A CONCERN OF SOCIAL PSYCHIATRY. *Psychiatry Research*. 2020: 112939.

18. Xu J, Xu Q-h, Wang C-m and Wang J. Psychological status of surgical staff during the COVID-19 outbreak. *Psychiatry Research*. 2020: 112955.

19. Neto MLR, Almeida HG, Esmeraldo JDa, et al. When health professionals look death in the eye: the mental health of professionals who deal daily with the 2019 coronavirus outbreak. *Psychiatry Research*. 2020: 112972.

20. Rajkumar RP. COVID-19 and mental health: A review of the existing literature. *Asian Journal of Psychiatry*. 2020: 102066.

21. Jian-Yun LU, Mei-Xia LI, Feng J, et al. Investigation on the Influence of Risk Communication to the Psychological Status of People in the Influenza A(H1N1) Quarantine Center. *Journal of Tropical Medicine*. 2010.

22. Christensen L and Burrows R. Dietary treatment of depression. *Behavior Therapy*. 21: 183-193.

23. Cauffield JS and Forbes HJ. Dietary supplements used in the treatment of depression, anxiety, and sleep disorders. *Lippincotts Primary Care Practice*. 1999; 3: 290-304.

24. Liu Y, Ozodiegwu ID, Yu Y, Hess R and Bie R. An association of health behaviors with depression and metabolic risks: Data from 2007 to 2014 U.S. National Health and Nutrition Examination Survey. *Journal of Affective Disorders*. 2017; 217: 190-196.

25. . !!! INVALID CITATION !!!

26. Furuhashi Y and Furuhashi S. Group Therapy for University Students with Autism Spectrum Disorders. *European Psychiatry*. 2015; 30: 440.

27. Furuhashi Y and Shusuke F. Group therapy for university students with autism spectrum disorders. *European Psychiatry*. 2016; 33: S631.

2   *Figure1 Flow chart of the study*

3   Table 1 Demographic and risk factors of the sample: 77 women and 33 men

Risk factors	Mean± SD/ N%		P Value
	Women（77）	Men（33）	
Basic information			
Age(years)			0.24
20-29	36.4	21.2	
30-39	50.6	57.6	
>39	13.0	21.2	
Working age（years）			0.21
<5	24.7	15.2	
6-10	33.8	39.4	
10-20	32.5	30.3	
21-30	9.1	9.1	
>30	0.0	6.1	
Education background			0.02[*]
College	11.7	0.0	
Bachelor	76.6	69.7	
Master	11.7	27.3	
Doctor	0.0	3.0	
Only child of the Family			0.22
Yes	19.5	30.3	
No	80.5	69.7	
Marital Status			0.04[*]
Married	58.4	78.8	
Unmarried	41.6	21.2	
Childbirth status			0.17
Yes	55.8	69.7	
No	44.2	30.3	
Family relationships			0.86
Good	94.8	93.9	
Poor	5.2	6.1	
Preparations before going to Wuhan city			
Infection prevention and control knowledge			0.17
Familiar	84.4	93.9	
Unfamiliar	15.6	6.1	
COVID-19 knowledge			0.47
Familiar	89.6	93.9	
Unfamiliar	10.4	6.1	
Confidence to complete the task			0.12

Yes	100.0	96.8	
No	0.0	3.2	
Emergency aid experience			0.97
Yes	42.9	42.4	
No	57.1	57.6	
RBT			0.04[*]
Good	98.7	90.9	
Poor	1.3	9.1	
Life issues after going to Wuhan city			
DC			0.97
Good	48.1	48.5	
Poor	51.9	51.5	
SQ			0.11
Good	35.1	51.5	
Poor	64.9	48.5	
LRoA			0.45
Adapted	44.2	36.4	
Unadapted	55.8	63.6	
CAM			0.04[*]
Yes	96.1	84.8	
No	3.9	15.2	
LC			0.53
Good	57.1	63.6	
Poor	42.9	36.4	
RC			0.62
Good	96.1	93.9	
Poor	3.9	6.1	
SEC			0.07
Good	90.9	78.8	
Poor	9.1	21.2	
Working issues after going to Wuhan city			
UWMRS			0.92
Yes	77.9	78.8	
No	22.1	21.2	
UWWE			0.40
Yes	48.1	39.4	
No	51.9	60.6	
DCWP			0.93
Yes	59.7	60.6	
No	40.3	39.4	
UWW			0.28
Yes	50.6	39.4	
No	49.4	60.6	
LME			0.82

Yes	7.8	9.1	
No	92.2	90.9	
LWH			0.86
Yes	5.2	6.1	
No	94.8	93.9	
LCWT			0.32
Yes	14.3	6.1	
No	85.7	87.9	
DCWLH			0.20
Yes	5.2	12.1	
No	94.8	90.3	
AoBI			0.29
Yes	13.0	6.1	
No	87.0	93.9	
SCL-90 Score			
GSI	1.20±0.24	1.17±0.16	0.56

Note: GSI=general symptom index, RBT= relationship between team, DC=dietary conditions, SQ=Sleep quality,

LRoA=limit range of activities, CAM=care about me, LC=lavation conditions, RC=residential conditions, SEC=

surrounding environmental conditions, UWMRS=unfamiliar with medical records system, UWWE=unfamiliar

with working environment, DCWP=difficulty communicating with patients, UWW=unfamiliar with workflow,

LME=lacking medical equipments, LWH=long working hours, LCWT=lacking communication with teams,

DCWLH=difficulty communicating with local hospital, AoBI= afraid of being infected.

Table2.Crude correlation associations of exposure risk factors and GSI of the

members

Variables	Statistics	GSI	
		β95%CI	*P Value*
Basic information			
Gender			
Female	77(70.00%)	0	
Male	33(30.00%)	-2.35(-10.27, 5.57)	0.56
Age(years)			
20-29	35(31.82%)	0	
30-39	58 (52.73%)	-4.07 (-12.23, 4.09)	0.33
>39	17 (15.45%)	-1.45 (-12.72, 9.82)	0.80
Working age（years）			
<5	24(21.82%)	0	
6-10	39(35.45%)	-3.39(-13.15, 6.38)	0.50
11-20	35(31.82%)	-7.31(-17.29, 2.66)	0.15
21-30	10(9.09%)	6.96(-7.21, 21.12)	0.34

>30	2(1.82%)	-15.04(-42.74, 12.66)	0.29
Education background			
Bachelor	82(74.55%)	0	
College	9(8.18%)	-5.92(-19.28, 7.44)	0.39
Master	18(16.36%)	3.97(-5.94, 13.87)	0.44
Doctor	1(0.91%)	-17.37(-55.65, 20.92)	0.38
Only Child of the Family			
No	85(77.27%)	0	
Yes	25(22.73%)	-0.53(-9.21, 8.14)	0.90
Marital Status			
No	39(35.45%)	0	
Yes	71(64.55%)	-1.29(-8.88, 6.31)	0.74
Childbirth status			
No	44(40.00%)	0	
Yes	66(60.00%)	-1.23(-8.65, 6.18)	0.75
Family relationships			
Poor	6(5.45%)	0	
Good	104(94.55%)	-13.88(-29.68, 1.91)	0.09
Preparations before going to Wuhan city			
Infection prevention and control knowledge			
Unfamiliar	14(12.73%)	0	
Familiar	96(87.27%)	-2.85(-13.74, 8.05)	0.61
COVID-19 knowledge			
Unfamiliar	10(9.09%)	0	
Familiar	100(90.91%)	-3.22(-15.85, 9.41)	0.62
Confidence to complete the task			
No	1(0.91%)	0	
Yes	(99.09%)	6.43(-31.85, 44.71)	0.74
Emergency aid experience			
No	63(57.27%)	0	
Yes	47(42.73%)	5.26(-2.03, 12.54)	0.16
RBT			
Poor	4(3.64%)	0	
Good	106(96.36%)	-20.63(-39.65, -1.60)	0.04[*]
Life issues after going to Wuhan city			
DC			
Poor	57(51.82%)	0	
Good	53(48.18%)	11.08(4.11, 18.05)	0.00[*]
SQ			
Poor	66(60.00%)	0	
Good	44(40.00%)	-1.83(-9.25, 5.58)	0.63
LRoA			
Unadapted	64(58.18%)	0	
Adapted	46(41.82%)	3.28(-4.06, 10.63)	0.38

CAM			
NO	8(7.27%)	0	
Yes	102(92.73%)	-4.05(-18.02, 9.93)	0.57
LC			
Poor	45 (40.91%)	0	
Good	65 (59.09%)	-0.20 (-7.59, 7.20)	0.96
RC			
Good	105 (95.45%)	0	
Poor	5 (4.55%)	3.17 (-14.27, 20.61)	0.72
SEC			
Good	96 (87.27%)	0	
Poor	14 (12.73%)	5.06 (-5.81, 15.92)	0.36
Working issues after going to Wuhan city			
UWMRS			
No	24 (21.82%)	0	
Yes	86 (78.18%)	5.70 (-3.04, 14.43)	0.20
UWWE			
No	60 (54.55%)	0	
Yes	50 (45.45%)	5.33 (-1.90, 12.56)	0.15
DCWP			
No	44 (40.00%)	0	
Yes	66 (60.00%)	6.87 (-0.43, 14.18)	0.07
UWW			
No	58(52.73%)	0	
Yes	52 (47.27%)	2.61(-4.65, 9.88)	0.48
LME			
No	101(91.82%)	0	
Yes	9(8.18%)	0.93(-12.34, 14.19)	0.89
LWH			
No	104 (94.55%)	0	
Yes	6 (5.45%)	7.36 (-8.58, 23.31)	0.37
LCWT			
No	97 (88.18%)	0	
Yes	13 (11.82%)	17.11 (6.32, 27.90)	0.00[*]
DCWLH			
No	102 (92.73%)	0	
Yes	8(7.27%)	0.54(-13.46, 14.54)	0.94
AoBI			
No	98 (89.09%)	0	
Yes	12(10.91%)	16.32(5.08, 27.57)	0.01[*]

Note: CI indicates confidence interval, [*]$P<0.05$.

6   Table3 Multivariate logistic regression model for DC, LCWT, AoBI, RBT and GSI of

7   the members

Variables in model	Non-adjusted		Adjust I		Adjust II	
	β (95%CI)	P Value	β (95%CI)	P Value	β (95%CI)	P Value
DC						
No	0		0		0	
Yes	11.08(4.11,18.05)	0.00[*]	10.85(3.61,18.09)	0.00[*]	10.17(3.30,17.04)	0.00[*]
LCWT						
No	0		0		0	
Yes	17.11(6.32,27.90)	0.00[*]	16.54(5.28,27.81)	0.01[*]	11.55(0.40,22.71)	0.04[*]
AoBI						
No	0		0		0	
Yes	16.32(5.08,27.57)	0.01[*]	15.66(4.11,27.21)	0.01[*]	8.51(-3.14,20.15)	0.16
RBT						
Good	0		0		0	
Poor	-20.63(-39.65,-1.60)	0.04[*]	-22.40(-41.89,-2.90)	0.03[*]	-28.09(-45.79,-10.40)	0.00[*]

8   Note: Non-adjusted model adjust for: None; Adjust I model adjust for: age and gender; [*]$P<0.05$.DC: Adjust II

9   model adjust for: LCWT, AoBI, RBT. LCWT: Adjust II model adjust for: DC, AoBI, RBT. AoBI: Adjust II model

10   adjust for: DC, LCWT, RBT. RBT: Adjust II model adjust for: DC, LCWT, AoBI.

13   Table4 Stratification analysis of DC, LCWT, AoBI, RBT and GSI score of the

14   members between genders

Exposure	Female		Male		Total	
	β (95%CI)	P Value	β (95%CI)	P Value	β (95%CI)	P Value
Non-adjusted						
DC						
No	0		0		0	
Yes	14.89 (5.94, 23.83)	0.00[*]	2.23 (-7.60, 12.06)	0.66	11.09 (4.10, 18.08)	0.00[*]
LCWT						
No	0		0		0	
Yes	17.20 (4.12, 30.27)	0.01[*]	15.73 (-4.17, 35.62)	0.13	16.95 (6.04, 27.86)	0.00[*]
AoBI						
No	0		0		0	

Yes	16.23 (2.51, 29.95) 0.02[*]	15.73 (-4.17, 35.62)0.13	16.14 (4.79, 27.49) 0.01[*]
RBT			
Good	0	0	0
Poor	-48.55 (-89.27, -7.84)0.02[*]	-13.13 (-29.64, 3.37)0.13	-22.40 (-41.89, -2.90) 0.03[*]

Adjusted Ⅰ

DC			
No	0	0	0
Yes	14.43 (5.08, 23.78) 0.00[*]	2.79 (-7.35, 12.92) 0.59	10.85 (3.61, 18.09) 0.00[*]
LCWT			
No	0	0	0
Yes	16.62 (3.10, 30.15) 0.02[*]	26.30 (4.08, 48.52) 0.03[*]	16.54 (5.28, 27.81) 0.01[*]
AoBI			
No	0	0	0
Yes	16.61 (2.55, 30.66) 0.02[*]	26.30 (4.08, 48.52) 0.03[*]	15.66 (4.11, 27.21 0.01[*]
RBT			
Good	0	0	0
Poor	-47.89 (-90.74, -5.04)0.03[*]	-12.40 (-29.60, 4.79)0.17	-28.09(-45.79,-10.40) 0.00[*]

Adjusted Ⅱ

DC			
No	0	0	0
Yes	13.20 (4.55, 21.85) 0.00[*]	3.07 (-7.02, 13.16) 0.56	10.40 (3.50, 17.29) 0.00[*]
LCWT			
No	0	0	0
Yes	11.89 (-0.53, 24.31) 0.06	15.57 (-4.61, 35.76) 0.14	11.14 (-0.06, 22.35) 0.054
AoBI			
No	0	0	0
Yes	8.73 (-4.34, 21.80 0.19	15.57 (-4.61, 35.76)0.14	8.17 (-3.51, 19.86) 0.17
RBT			
Good	0	0	0
Poor	-57.85 (-94.52, -21.18)0.00[*]	-15.81 (-32.80, 1.18)0.08	-29.62 (-47.67, -11.57) 0.00[*]

1   Note: Non-adjusted model adjust for: None; Adjust I model adjust for: age; [*]$P<0.05$. DC: Adjust II model adjust

2   for: LCWT, AoBI, RBT. LCWT: Adjust II model adjust for: DC, AoBI, RBT. AoBI: Adjust II model adjust for: DC,

3   LCWT, RBT. RBT: Adjust II model adjust for: DC, LCWT, AoBI.