

1 **Distinguishing between COVID-19 and the common cold in a primary care setting**  
2 **- comparison of patients with positive and negative SARS-CoV-2 PCR results.**

3  
4 Johannes Just<sup>1\*</sup>, Marie-Therese Puth<sup>2</sup>, Felix Regenold<sup>1</sup>, Klaus Weckbecker<sup>1#</sup>, Markus  
5 Bleckwenn<sup>3#</sup>

6  
7 1:  
8 Institute of General Practice and Interprofessional Care  
9 Witten/Herdecke University  
10 Alfred-Herrhausen-Straße 50  
11 58453 Witten

12  
13  
14 2:  
15 Department of Medical Biometry, Informatics and Epidemiology (IMBIE)  
16 University Hospital Bonn  
17 Sigmund-Freud-Str. 25,  
18 53127 Bonn

19  
20 3:  
21 Department of General Practice  
22 Leipzig University  
23 Medical Faculty  
24 Ph.-Rosenthal-Str. 55  
25 04103 Leipzig

26  
27  
28 \* Corresponding author: [johannes.just@uni-wh.de](mailto:johannes.just@uni-wh.de)  
29 # joined last authorship

30  
31  
32  
33  
34  
35  
36  
37

38 **Background:** Combating the COVID-19 pandemic is a major challenge for health  
39 systems, citizens and policy makers worldwide. Early detection of affected patients  
40 within the huge population of patients with common cold symptoms is an important  
41 element of this effort but often hindered by limited testing resources. We aimed to  
42 identify predictive risk profiles for a positive PCR result in primary care.

43 **Methods:** Multi-center cross-sectional cohort study on predictive characteristics over a  
44 period of 4 weeks in primary care patients in Germany. We evaluated age, sex, reason  
45 for testing, risk factors, symptoms, and expected PCR result for their impact on the test  
46 result.

47 **Results:** In total, 374 patients in 14 primary care centers received SARS-CoV-2 PCR  
48 swab testing and were included in this analysis. A fraction of 10.7% (n=40) tested  
49 positive for COVID-19. Patients who reported anosmia had a higher odds ratio (OR:  
50 4.54; 95%-CI: 1.51-13.67) for a positive test result while patients with a sore throat had  
51 a lower OR (OR: 0.33; 95%-CI: 0.11-0.97). Patients who had a first grade contact with  
52 an infected persons and showed symptoms themselves also had an increased OR for  
53 positive testing (OR: 5.16; 95% CI: 1.72-15.51). This correlation was also present when  
54 they themselves were still asymptomatic (OR: 12.55; 95% CI: 3.97-39.67).

55 **Conclusion:** The reported contact to an infected person is the most important factor for  
56 a positive PCR result, independent of any symptoms of illness in the tested patient.  
57 Those persons with contact to an infected person should always get a PCR test. If no  
58 contact is reported and testing material is scarce, anosmia should increase the  
59 likelihood of performing a test, while a sore throat should decrease it.

60  
61  
62  
63  
64  
65  
66

## 67 **Introduction**

68  
69 The COVID-19 pandemic is a major challenge for health systems, citizens and policy  
70 makers worldwide (1). The early political implementation of individual distancing has  
71 slowed down the reproduction rate of the virus in Germany significantly (2). In addition,  
72 compared to other countries, Germany has implemented an intensive testing strategy  
73 for new infections using PCR swab testing, funded by the national social health  
74 insurance system (statutory health insurance). According to date from “Robert-Koch  
75 Institut” (RKI), roughly an equivalent to the CDC in the US, 176 participating laboratories  
76 reported a number of 348,619 PCR tests carried out in calendar week 12/2020 (419  
77 tests/100.000 inhabitants). The tests were mostly carried out in doctors' (GP's)  
78 practices, newly formed test centers and hospitals (3).

79  
80 In view of the potentially exponential increase in the number of new cases and finite  
81 laboratory resources (test stations, reagents, etc.), there is an urgent need to collect  
82 and evaluate clinical data on the clinical features of tested patients. Of particular interest  
83 is the question of how far positive and negative patients differ in their initial clinical  
84 presentation. The more precisely physicians use SARS-CoV-2 PCR testing, the more  
85 efficient it is. Statistically, this is reflected in Baye's theorem: The quality of a test is not  
86 only determined by specificity and sensitivity, but also depends on the pre-test  
87 probability of the event to be tested. Thus, if the pre-test probability for COVID-19 is  
88 higher, the probability that a SARS-CoV-2 PCR-positive patient is actually ill (positive-  
89 predictive value = PPV) is increased (4).

90  
91 In this paper we aimed to identify predictive risk profiles for a positive PCR result in  
92 primary care.

93  
94  
95  
96

## 97 **Method**

98 In this multi-center, cross-sectional study, the characteristics of patients who tested  
99 positive and negative for COVID-19 were assessed. A total of 26 office-based  
100 specialists for internal and/or general medicine with a full primary care mandate from 14  
101 different locations participated in the study.

102 All locations collected patient-related data based on a uniform quality standard in the  
103 documentation of COVID-19 suspect cases that we provided.

104 Each site reported anonymous data on all SARS-CoV-2 PCR swabs taken. These  
105 included: Age, sex, reason for the test, risk factors, symptoms leading to the test, result  
106 expected by the GP and actual PCR result. All data were noted on a structured  
107 documentation sheet and passed on anonymously for evaluation. The documentation  
108 form was prepared on the basis of the recommendations of the RKI as well as the  
109 available evidence on symptoms and risk factors for COVID-19 and was subjected to  
110 pre-testing with experienced family doctors (5,6,7,8). The test reason "Patient is contact  
111 person without symptoms" was included in addition to the test reasons recommended  
112 by the RKI, as it had already been used frequently in everyday practice.

113

114 The survey period was from 24.03.2020 to 17.04.2020 - this period was chosen  
115 because the RKI criteria for carrying out a PCR remained stable.

116

117 Tests that had been carried out for procedural reasons and did not correspond to a  
118 specific clinical indication were excluded (e.g. testing of recovered patients after end of  
119 quarantine). There were no other exclusion criteria.

120

121 The participating GP's practices represented a "convenience sample". Selection criteria  
122 were the performance of smear tests, structured documentation and willingness to  
123 participate.

124

125 In order to ensure that there was no excessive distortion in the patient collective due to  
126 the convenience sampling of practices, age and sex of the tested persons were  
127 compared with demographic data from a large national sample.

128

129 Data entry was performed twice and the data sets were compared digitally, any  
130 deviations were checked and a plausibility check of the data entered was performed.  
131 The statistical evaluation was carried out with the statistics program R. In addition to the  
132 descriptive statistics, initial chi-square tests were performed for different potential  
133 influencing factors (symptoms, test reason, etc.) on the test result (PCR positive  
134 yes/no). Statistically significant influencing factors were then transferred to a logistic  
135 regression model to check their effect strength. Missing data were reported in the  
136 descriptive part, for the regression model missing values were assigned to the more  
137 frequent result (deviation to the middle), a proportion of missing values >5% was not  
138 exceeded.

139  
140 The study design was reviewed by the ethics committee of the University of Leipzig,  
141 Germany under the procedure number 184/20-ek, there were no ethical concerns.

142  
143 This study was conducted without external financial support.

144  
145

## 146 **Results**

147  
148 The participating practices tested n=374 patients per swab test for SARS-CoV-2 in the  
149 investigated period. Of these, 10.7% (n=40) tested positive. The symptom anosmia, as  
150 well as contact with infected persons, was associated with a positive PCR result.


151  
152 A total of 26 specialists with a general practitioner mandate at 14 different locations  
153 (individual and group practices) from North Rhine-Westphalia, Rhineland-Palatinate,  
154 Hesse and Saxony-Anhalt (four of 16 federal German states) took part in the study. All  
155 practices are located in urban and rural districts with moderate to increased COVID-19  
156 activity (cumulative incidence 50-250 cases per 100,000 inhabitants) (2).

157

158 Parallel to the number of new infections reported throughout Germany, the number of  
 159 tests performed decreased in the course of the study, as did the rate of positively tested  
 160 patients.

161

162 Figure 1: Number of tests separated by test reason per calendar week



163

164

165 The examined patients were 58.5% female and the median age was 44 years. Further  
 166 characteristics of the investigated cohort can be found in table 1.

167

168

169 Table 1: Patient characteristics

	Total	COVID-19 negative	COVID-19 positive	p-value
	(n = 374)	(n = 334)	(n = 40)	

Age				0.147*
MV (SD)	47.0 (20.2)	46.4 (19.7)	52.4 (23.5)	
median (Q1-Q3)	44.0 (31.0-59.0)	43.5 (31.0-58.0)	52.0 (31.2-73.2)	
missing, n (%)	4 (1.1)	4 (1.2)	0 (0.0)	
gender, n (%)				0.475
male	154 (41.2)	140 (41.9)	14 (35.0)	
female	217 (58.0)	191 (57.2)	26 (65.0)	
missing	3 (0.8)	3 (0.9)	0 (0.0)	
reason for testing, n (%)				<0.001
First grade contact (has symptoms)	72 (19.3)	58 (17.4)	14 (35.0)	
First grade contact (has no symptoms)	40 (10.7)	27 (8.1)	13 (32.5)	
Indication of viral pneumonia	13 (3.5)	12 (3.6)	1 (2.5)	
Works in medical profession	51 (13.6)	47 (14.1)	4 (10.0)	
belongs to risk group	60 (16.0)	59 (17.7)	1 (2.5)	
free capacity/other	134 (35.8)	127 (38.0)	7 (17.5)	

missing	4 (1.1)	4 (12.0)	0 (0.0)	
GP expects positive result, n (%)				0.010
no	245 (65.5)	227 (68.0)	18 (45.0)	
yes	119 (31.8)	99 (29.6)	20 (50.0)	
missing	10 (2.7)	8 (2.4)	2 (5.0)	
Number of risk factors, n (%)				N/A
none	187 (50.0)	168 (50.3)	19 (47.5)	
one or two	138 (36.9)	125 (37.4)	13 (32.5)	
more than two	49 (13.1)	41 (12.3)	8 (20.0)	

170 \*Mann-Whitney U Test

171 MV=mean value; SD=Standard deviation; Q1 = 25%-quantile; Q3 = 75%-quantile

172

173

174 The most frequent symptoms at the presentation of patients who later tested positive for  
175 COVID-19 were cough, fever, anosmia and muscle pain.

176 The most common symptoms at presentation of patients who later tested negative for  
177 COVID-19 were cough, sore throat, fatigue and fever.

178

179 Table 2: Symptoms of patients tested negative and positive for COVID-19

Variable	COVID-19 negative (n = 307)	COVID-19 positive (n = 27)	p-value
----------	--------------------------------	-------------------------------	---------



cough (n, %)	214	69.7	19	70.4	1.000
Sore throat (n, %)	120	39.1	5	18.5	0.038
fatigue (n, %)	89	29.0	5	18.5	0.371
fever (n, %)	84	27.4	9	33.3	0.507
nasal congestion (n, %)	84	27.4	5	18.5	0.373
muscle pain (n, %)	59	19.2	7	25.9	0.448
dyspnea (n, %)	56	18.2	4	14.8	0.798
headache (n, %)	47	15.3	3	11.1	0.780
anorexia (n, %)	28	9.1	2	7.4	1.000
diarrhea (n, %)	23	7.5	1	3.7	0.707
anosmia (n, %)	22	7.2	7	25.9	0.005
chills (n, %)	20	6.5	5	18.5	0.040
nausea (n, %)	11	3.6	0	0.0	1.000
vomiting (n, %)	4	1.3	0	0.0	1.000
other symptoms (n, %)	37	12.1	3	11.1	1.000

180 Asymptomatic patients (n=40) were excluded in this table.

181

182 The regression model used to determine the individual effect sizes included all patient  
 183 characteristics that showed a statistically significant influence on the test result in the  
 184 Chi-square test. The odds ratio (OR) for a positive COVID-19 PCR was increased in  
 185 patients who had contact with an infected person who were older and in patients with  
 186 anosmia and decreased in patients suffering from a sore throat. Table 3 shows all  
 187 results of the regression model.

188

189 Table 3: Logistic regression model: factors influencing positive COVID-19 PCR

190

Variable	COVID-19 negative (n=330)  n (%)	COVID-19 positive (n=40)  n (%)	adjusted  OR  (95% CI)	p-value
First grade contact (has symptoms)	58 (17.5)	14 (35.0)	5.16 (1.72-15.51)	0.002
First grade contact (has no symptoms)	27 (11.2)	13 (32.5)	12.55 (3.97-39.67)	<0.001
Free capacity/other	125 (37.9)	7 (17.5)	1.50 (0.46-4.87)	0,497
age*	46,4 (19.7)	52,4 (23.5)	1.02 (1.00-1.04)	0,019
GP expects positive result	97 (29.4)	20 (50.0)	1.98 (0.90-4.38)	0.092
chills	20 (6.1)	5 (12.5)	2.80 (0.83-9.43)	0.117

Anosmia	22 (6.7)	7 (17.5)	4.54 (1.51-13.67)	0.011
sore throat	118 (35.8)	5 (12.5)	0.33 (0.11-0.97)	0.029

191 OR = odds ratio; CI = confidence intervals  
192 \* mean value (standard deviation); Nagelkerke R<sup>2</sup> 0.265

193  
194

195

196

## 197 Discussion

198

### 199 Main result

200 The main result of this work is the shown correlation between a positive COVID-19 PCR  
201 and the symptom anosmia as well as a confirmed contact with an infected person,  
202 regardless of whether the tested person itself had symptoms or not.

203

### 204 Strengths and limitations

205 In this work, predictive factors for a positive COVID-19 PCR were investigated for the  
206 first time. The evaluation was performed with a multicentric approach and was based on  
207 a common documentation standard. The selection of the practices was non-  
208 randomized, as a rapid implementation of the study was preferred to a lengthier  
209 randomized selection in respect to the pandemics' rapid evolution. This may have  
210 resulted in a distortion of the investigated patient collective, although, no conspicuous  
211 deviations were found in comparison to the demographic data of all patients tested in  
212 Germany as published by the RKI: the age median in our population was 44 years (IQR:  
213 31-59) compared to 42 years (IQR: 29-56) in a sample of +1.000.000 patients tested in  
214 Germany (3). Therefore, the results could be considered representative for patients in  
215 Germany.

216

### 217 Number of patients tested and rates of positives

218 The proportion of test-positive patients in our population was 10.7%, which is slightly  
219 above the average of positive tests performed in Germany during this period (North  
220 Rhine-Westphalia (9.3%), Rhineland-Palatinate (9.6%), Hesse (11.8%)) (3). The  
221 number of tests decreased towards the end of the investigation period, which parallels  
222 the decreasing number of new COVID-19 infections in Germany as well as the end of  
223 the pandemic cold season in early spring.

224

225

### 226 **Factors that were associated with a negative test result**

227 Negative tests were associated with the symptom "sore throat". This result is consistent  
228 with the low prevalence of the symptom "sore throat" in confirmed COVID-19 cases (6).  
229 The correlation found between age and test results is marginal (younger people were  
230 tested negative more often) and may be explained by local outbreaks in nursing homes.  
231 This effect will most likely dissolve due to diffusion processes during the course of the  
232 pandemic and may not be present in countries with a more homogenous spreading  
233 pattern.

234

### 235 **Factors that were associated with a positive test result**

236 Positive tests were associated with confirmed contact. This was to be expected, but it  
237 was surprising that this also applied to asymptomatic patients. He et al. concluded that  
238 45% of infections occur in the pre-symptomatic phase of the patient, i.e. the patient is  
239 still asymptomatic but already has large amounts of virus replicated in the throat and is  
240 highly contagious (9). This result underlines the particular relevance of contact  
241 anamnesis for the test decision, but also the relevance of early isolation of potentially  
242 affected contacts.

243 Of particular clinical importance is the observed association between the symptom  
244 anosmia and a positive PCR result. Anosmia has already been described as a relevant  
245 COVID-19 symptom in other studies, but this is the first study to show that anosmia may  
246 be a useful discriminator between COVID-19 and other, endemic respiratory tract  
247 infections (8,10,11). Since the examined patients mainly presented with fever, cough,  
248 muscle pain and other cold symptoms, which are common with endemic respiratory

249 tract infections, anosmia can be an important indication of COVID-19. Although anosmia  
250 is a secondary symptom of a variety of viral respiratory infections, the incidence of  
251 anosmia without acute or chronic rhinitis in COVID-19 is striking (8,10-12). Reporting  
252 from our clinical experience with COVID-19, anosmia was sometimes so pronounced in  
253 COVID-19 patients that it was presented as the sole symptom upon presentation.  
254 Therefore, we recommend that patients are always asked for fever, cough, dyspnea and  
255 anosmia as part of the initial stratification process. In case of isolated anosmia in the  
256 absence of chronic or acute rhinosinusitis, we recommend a testing for COVID-19  
257 infection, as already suggested in a French case series by Villalba et al. and others  
258 (8,11,13)

259

## 260 **Conclusion**

261 Rapid testing of first degree contacts, consistent tracing of infection chains and  
262 quarantine will remain essential parts of the COVID-19 containment strategy. Our data  
263 show that the testing of all first-degree contacts, including asymptomatic ones can be of  
264 merit and that anosmia, even as an isolated symptom is a clear indication of COVID-19  
265 infection.

266 General practitioners are an essential part of the testing strategy in most countries.  
267 Based on their intensive experience with respiratory tract infections, they are able to  
268 clinically assess a large number of patients with cold symptoms in a short period of time  
269 and, if necessary, perform PCR testing. Further GP research should pay particular  
270 attention to the three areas of diagnostics, therapy and advance care planning for  
271 patients with COVID-19.

272

273

274

275

- 276 1. Statement on the second meeting of the International Health Regulations (2005)  
277 Emergency Committee regarding the outbreak of novel coronavirus (2019-nCoV) [Internet].  
278 [cited 2020 Apr 1]. Available from: [https://www.who.int/news-room/detail/30-01-2020-](https://www.who.int/news-room/detail/30-01-2020-statement-on-the-second-meeting-of-the-international-health-regulations-(2005)-emergency-committee-regarding-the-outbreak-of-novel-coronavirus-(2019-ncov))  
279 [statement-on-the-second-meeting-of-the-international-health-regulations-\(2005\)-](https://www.who.int/news-room/detail/30-01-2020-statement-on-the-second-meeting-of-the-international-health-regulations-(2005)-emergency-committee-regarding-the-outbreak-of-novel-coronavirus-(2019-ncov))  
280 [emergency-committee-regarding-the-outbreak-of-novel-coronavirus-\(2019-ncov\)](https://www.who.int/news-room/detail/30-01-2020-statement-on-the-second-meeting-of-the-international-health-regulations-(2005)-emergency-committee-regarding-the-outbreak-of-novel-coronavirus-(2019-ncov))

- 281 2. RKI: Daily situation report on COVID-19. Available from:  
282 [https://www.rki.de/DE/Content/InfAZ/N/Neuartiges\\_Coronavirus/Situationsberichte/2020-04-19-de.pdf?\\_\\_blob=publicationFile](https://www.rki.de/DE/Content/InfAZ/N/Neuartiges_Coronavirus/Situationsberichte/2020-04-19-de.pdf?__blob=publicationFile)  
283
- 284 3. RKI: epidemiological update 15/2020. SARS-CoV-2-testing numbers. [cited 2020 Apr 15].  
285 Available from:  
286 [https://www.rki.de/DE/Content/Infekt/EpidBull/Archiv/2020/Ausgaben/15\\_20.pdf?\\_\\_blob=pu](https://www.rki.de/DE/Content/Infekt/EpidBull/Archiv/2020/Ausgaben/15_20.pdf?__blob=publicationFile)  
287 [blicationFile](https://www.rki.de/DE/Content/Infekt/EpidBull/Archiv/2020/Ausgaben/15_20.pdf?__blob=publicationFile)
- 288 4. Schneider A, Dinant G-J, Szecsenyi J. Stepwise diagnostic workup in general practice as a  
289 consequence of the Bayesian reasoning. *Z Arztl Fortbild Qualitatssich.* 2006;100(2):121–7.
- 290 5. RKI: Testing guidelines for doctors [Internet]. [cited 2020 Apr 1]. Available from:  
291 [https://www.rki.de/DE/Content/InfAZ/N/Neuartiges\\_Coronavirus/Massnahmen\\_Verdachtsfal](https://www.rki.de/DE/Content/InfAZ/N/Neuartiges_Coronavirus/Massnahmen_Verdachtsfall_Infografik_DINA3.pdf?__blob=publicationFile)  
292 [l\\_Infografik\\_DINA3.pdf?\\_\\_blob=publicationFile](https://www.rki.de/DE/Content/InfAZ/N/Neuartiges_Coronavirus/Massnahmen_Verdachtsfall_Infografik_DINA3.pdf?__blob=publicationFile)
- 293 6. Huang C, Wang Y, Li X, Ren L, Zhao J, Hu Y, et al. Clinical features of patients infected  
294 with 2019 novel coronavirus in Wuhan, China. *Lancet.* 2020 Feb 15;395(10223):497–506.
- 295 7. Lechien JR, Chiesa-Estomba CM, De Siati DR, Horoi M, Le Bon SD, Rodriguez A, et al.  
296 Olfactory and gustatory dysfunctions as a clinical presentation of mild-to-moderate forms of  
297 the coronavirus disease (COVID-19): a multicenter European study. *Eur Arch*  
298 *Otorhinolaryngol* [Internet]. 2020 Apr 6; Available from: [http://dx.doi.org/10.1007/s00405-](http://dx.doi.org/10.1007/s00405-020-05965-1)  
299 [020-05965-1](http://dx.doi.org/10.1007/s00405-020-05965-1)
- 300 8. Spinato G, Fabbris C, Polesel J, Cazzador D, Borsetto D, Hopkins C, et al. Alterations in  
301 Smell or Taste in Mildly Symptomatic Outpatients With SARS-CoV-2 Infection. *JAMA*  
302 [Internet]. 2020 Apr 22; Available from: <http://dx.doi.org/10.1001/jama.2020.6771>
- 303 9. He X, Lau EHY, Wu P, Deng X, Wang J, Hao X, et al. Temporal dynamics in viral shedding  
304 and transmissibility of COVID-19. *Nat Med* [Internet]. 2020 Apr 15; Available from:  
305 <http://dx.doi.org/10.1038/s41591-020-0869-5>
- 306 10. Brann DH, Tsukahara T, Weinreb C, Lipovsek M, Van den Berge K, Gong B, et al. Non-  
307 neuronal expression of SARS-CoV-2 entry genes in the olfactory system suggests  
308 mechanisms underlying COVID-19-associated anosmia [Internet]. Available from:  
309 <http://dx.doi.org/10.1101/2020.03.25.009084>
- 310 11. Giacomelli A, Pezzati L, Conti F, Bernacchia D, Siano M, Oreni L, et al. Self-reported  
311 Olfactory and Taste Disorders in Patients With Severe Acute Respiratory Coronavirus 2  
312 Infection: A Cross-sectional Study [Internet]. *Clinical Infectious Diseases.* 2020. Available  
313 from: <http://dx.doi.org/10.1093/cid/ciaa330>
- 314 12. van Riel D, Verdijk R, Kuiken T. The olfactory nerve: a shortcut for influenza and other viral  
315 diseases into the central nervous system [Internet]. Vol. 235, *The Journal of Pathology.*  
316 2015. p. 277–87. Available from: <http://dx.doi.org/10.1002/path.4461>
- 317 13. Villalba NL, Maouche Y, Ortiz MBA, Sosa ZC, Chahbazian JB, Syrovatkova A, et al.  
318 Anosmia and Dysgeusia in the Absence of Other Respiratory Diseases: Should COVID-19  
319 Infection Be Considered? *Eur J Case Rep Intern Med.* 2020 Apr 3;7(4):001641.