A model showing the relative risk of viral aerosol infection from breathing and the benefit of wearing masks in different settings with implications for Covid-19

.

Gerald D Barr MBChB , FRCS (Ed) ,Cert. Nat. Sci. (Open), Cert. HE (Open)

Independent Health Research , Peat Inn , UK

Abstract

Background . Widespread use of masks in the general population is being used in many countries for Covid-19 . There has been reluctance on the part of the WHO and some governments to recommend this .

Methodology . A basic model has been constructed to show the relative risk of aerosol from normal breathing in various situations together with the benefit from use of masks which is multiplicative .

Results . Social distancing at 2 metres appears validated but in confined areas is time limited and the use of masks together with efficient ventilation is important. Where social distancing is not possible at all times or a potentially infectious person is in a confined area for a prolonged period there is a higher risk of infection requiring protection .

Conclusions . The use of masks should be factored into models and used at an early stage as widespread use of more efficient masks could have a large impact on control and spread of infection . Public health planning requires stockpiling of masks and encouraging everyone to have suitable masks in their household when supplies are normalised . The use of a cloth mask will be better than no protection at all .

Introduction

Concerning Covid-19 the WHO advice from January 2020 [1] is that masks for the general population are not required except for those with respiratory symptoms. In April this changed to medical masks should be reserved for health care professionals [2] , citing that mask wearing by the public might lead to complacency . However , it could be argued that informing the public that masks are not needed is potentially more damaging leading some individuals to believe that Covid-19 is only contracted by poor hand hygiene . The advice appears to be contradictory to the 2019 document for influenza which states that the approach should be multifaceted including ,'Face masks worn by asymptomatic people are conditionally recommended in severe epidemics or pandemics, to reduce transmission in the community.' , although it does mention the possibility of reduced supply to health professionals and increase in prices as a consequence. [3] Even basic masks can help to prevent droplet spread which the WHO states is the primary spread of Covid-19 together with touching fomites , while acknowledging aerosol spread is an important consideration in hospitals .[4] Given the relatively high R number [5] and findings of aerosol spread in other respiratory viruses such as SARS and influenza this route must be a consideration in the community unless proven otherwise . The UK has adopted a similar policy that a decision cannot be made regarding general use of masks in the population based on having an insufficient evidence base . A rapid Cochrane review commissioned by the WHO assessing quarantine alone or with other public health measures to control Covid-19 does not mention masks . [6] A past Cochrane review [7] for influenza showed low level evidence in favour of general mask

wearing but due to the lack of good quality studies could not make a positive recommendation pointing to compliance issues , but these studies do not translate to a pandemic with significant morbidity . A further review by Public Health England concluded that low level evidence suggested there was benefit of masks for health care workers and in the community but community use could not be recommended [8] . It seems illogical in a pandemic with significant morbidity that only health care workers need protected when the driving force for their need is the numbers of people from the community being admitted to hospital. Greenhalgh referred to the precautionary principle when there is no evidence for doing something but where the risk of doing nothing could cause considerable harm in relation to recommending the use of masks in the general population [9] and Cowling asked for a rational approach to the use of masks . [10]

Aerosol spread will be much more in coughing and sneezing , than in speaking or normal breathing , however some individuals produce excessively high amounts of aerosol particles > 10,000 per litre in normal breathing [11] which may be one mechanism for superspreading . From an observation in influenza [12] causing 100 per cent infection in a confined space it was postulated that there must be a relationship between the amount of virus expelled into the air , the ventilation of the area and the time exposed . From this a simple model has been constructed to show the relative benefit of mask wearing in relation to the infectious dose of a virus in aerosol . A Health and Safety review found a six times reduction in bioaerosol using a surgical mask with and estimated 100 times reduction for an FFP3 mask , with much of the inefficiency of surgical masks being due to leakage or poor fit [13] .

Cloth masks are acknowledged to give some protection which may only have an improvement factor of two .[14]

Method

Infectious dose (ID) is the amount of virus needed to be inhaled to cause infection . Many factors could affect this and the bioavailability of the virus once inhaled together with mucosal defences . A review by Nikitin suggested a figure of 1.9×10^3 for influenza [15] . Regarding fine aerosol that can penetrate more deeply into the lungs it is quite possible that a much lower dose is required . For this comparative study an ID at 1000 was used especially as this fitted best with scenario 2 (Table 2) and the original observation . Not all aerosol particles contain virus which may only be 10 % of aerosol but can be more .[16] For this study 30% of particles being infectious was taken at $av = 0.3$.

For an infected person breathing out aerosol particles $P = 500$ / litre is used which is the cut off value between a low and high producer of aerosol .[11]

Ev is the volume of air expired by the infectious person per minute [tidal volume x breaths/min] for this study taken as normal breathing 500ml x12 breaths per minute = 6 litres /min .

Bv is the volume of air inspired by a non infected person per minute [tidal volume x breaths/min] for this study taken as normal breathing 500ml x12 = 6 litres /min .

F_1 is the inspiratory filtration factor of a mask for an infectious person breathing out , F_2 is the filtration factor for non infected person breathing in . For no mask F is 1 , for x 6 reduction is 0.167 , x100 reduction 0.01 . Using this data and that from a study [17] on the reduction factor , inspiratory and expiratory at 30 , 50 and 80

l/min extrapolating to 6 l/min : cloth mask F_1 is 0.75 F_2 0.5 , surgical mask F_1 is 0.333 F_2 is 0.167 , FFP3 F_1 is 0.333 F_2 0.01 .

At a distance under one metre the volume is not applicable due to risk of directly breathing the same air   e.g. one person breathes in as the other breathes out .

Infectious Dose ID = $P(av)$ EvF_1 (Bv /Ev) F_2 t where t is time in minutes . (Bv cannot exceed Ev i.e. max for Bv/Ev is 1)

Ve is the amount of ventilation given by the percentage of air replaced with fresh air per minute   Ve is % /100

Considering the exhaled infectious air dispersed into a volume V .

The time evolution of the infectious particle concentration C in the volume V is given by:

$$\frac{dC}{dt} = \left(\frac{Pa_v E_v F_1}{V}\right) - \lambda C$$

The time evolution of the inspired dose N is given by:

$$\frac{dN}{dt} = (B_v F_2)C$$

For zero decay constant λ the solutions are (where we consider that the infectious person has been present for a time T_{start} before the uninfected person arrives):

$$C(t) = C_{start} + \left(\frac{Pa_vE_vF_1}{V}\right)t$$

$$N(t) = (B_vF_2)\left[C_{start}t + \left(\frac{Pa_vE_vF_1}{V}\right)\left(\frac{1}{2}t^2\right)\right]$$

$$C_{start} = \left(\frac{Pa_vE_vF_1}{V}\right)T_{start}$$

For nonzero λ the solutions are (where we allow a different decay constant λ_{pre} during the warm-up phase before the uninfected person arrives):

$$C(t) = C_{final} + (C_{start} - C_{final})\exp(-\lambda t)$$

$$N(t) = (B_vF_2)\left[C_{final}t + (C_{start} - C_{final})\left(\frac{1 - \exp(-\lambda t)}{\lambda}\right)\right]$$

$$C_{final} = \left(\frac{Pa_vE_vF_1}{V\lambda}\right)$$

$$C_{start} = C_{pre-sat}\left(1 - \exp\left(-\lambda_{pre}T_{start}\right)\right)$$

$$C_{pre-sat} = \left(\frac{Pa_vE_vF_1}{V\lambda_{pre}}\right)$$

$$t_{1/2} = \frac{\ln(2)}{\lambda}$$

The aerosol absorbed by each person inhaling would reduce the amount of infectious particles also increased by use of a mask affecting exhalation too . Fx is the inhalation filtration factor for the infectious person and Fy is the filtration factor for the non infected person exhaling . Using the calculated loss from a medical aerosol study of absorption giving an estimated 45% of particles absorbed . [18] The inhalation absorption loss for inhalation

$$\lambda_{insp/exp} = (BvF_{loss})/V$$

Where F_{loss} is the effective fraction of particles lost in one complete breath and accounts for the effect of any masks during breathing in or out. The contributions of multiple individuals could be added to this value.

F_{abs} is the fraction of particles absorbed with no mask then the effect of a mask can be estimated as:

$F_{loss} Bv = 1-(Fy\ F2)\ (1-Fabs)$

$F_{loss} Ev = 1-(FxF1)\ (1-Fabs)$

The exponential decay constant λ represents all loss processes which can be combined as:

$\lambda = \lambda_1 + \lambda_2 + \cdots$ $\lambda = -ln(2)/(half\text{-}life\ for\ aerosol\ decay)$

$$\lambda = -ln(Volume\ retained\ after\ 1\ min/V\ )$$

$$\lambda = -ln(\ proportion\ inspiratory\ and\ expiratory\ loss\ )$$

General comments and limitation

This is a comparative analysis not an actual situation for any particular virus and estimated values from the literature are used . Any of the chosen parameters could be less than the actual for instance the aerosol may only have 10% infectious particles , ID could be 3000 but equally particles exhaled by an individual could be 10 times more.

The model assumes that the particles expired are instantly diluted to give a uniform concentration in the volume. In reality the distribution in space would be complex and so V must be regarded as an effective interaction volume rather than the precise volume of any particular region. Additionally, ventilation and convection or diffusion would create a complex spatial and temporal dependence of the concentration so the contribution to λ by these processes must be considered as an effective value to model the loss of the virus from the system and not a precise value. It does not take account of localised directional air flow which in theory could increase the infectious dose i.e. directing higher concentration aerosol towards a person .

Decay of aerosol will vary for virus , particle size and humidity .   Adams et al. found a sharp decay at the start for Reo virus , although this was considerably reduced by humidity , with recovery over longer periods [19] which may be due to smaller particles surviving longer . Cowling found that 87% of particles [20] emitted are less than 1 micron .   At this size there is less potential space for a virus , for instance an aerosol particle 1 micron in diameter is much more likely to have a

maximum of one SARS-CoVid-2 particle measuring 0.12 microns in diameter . Smaller particles can stay airborne for hours in a humid environment with only a 10% loss of infectivity . [21] Yang found enough influenza virus in the air to cause infection in different public places and that a steady state will be reached [22] . For this study it was assumed one infectious particle releases only one viral particle and in the calculations the median aerosol half life of 1.2 hours for SARS-CoVid-2 found by van Doremalen was used . [23]

It is not certain what the minimum cumulative dose for infection is and over what period i.e. the minimum number of virus particles that non specific defences can cope with per hour or minute . It is assumed the ID is cumulative at least over a relatively short period for this study minutes to a few hours .

Results

The model shows for two people wearing masks the protection is multiplicative not additive . Scenario 1 (table I) shows a high risk being close to someone breathing normally, potentially only for a short time . Wearing a mask will reduce the number of infectious particles which could mean reducing the chance of infection or even having a milder infection . Combining a surgical mask in the infectious and non infected person gives a 2.8 times improvement from a surgical mask only on the non infected person , and a 17 times improvement from no mask at all . The more people wearing higher quality masks will give more improvement and less risk of infection . Scenario 2 (table II) with two people on a half hour car journey reaching the ID by the end . There is a greatly reduced infectious dose with a non infected person

wearing an FFP3 mask but all the masks reduce the ID . Both wearing a surgical mask is of considerable benefit although not as good as FFP3 . Scenario 3 (table III) shows that an infectious person in a confined area for 3 hours with no ventilation will give a high risk of infection . Such a situation could be a taxi where the driver is not separated from the passengers , or an infectious person in a cubicle with little ventilation . When someone enters the benefit of FFP3 is high but also there is a marked improvement by both wearing surgical masks . As there is only a short time benefit from a surgical mask alone or a cloth mask then ventilation with clean air is needed for further improvement . Figure1 shows in a restricted volume using cloth masks the difference with 5% of the air being replaced per minute. Scenario 4 (table IV) shows a person in an enclosed area with a larger volume for 3 hours and a non infected person entering . This gives a low risk situation but a cloth mask can give an even larger improvement . Scenario 5 (table V) shows two people socially distanced in a confined area such as a work place . It shows how social distancing can work but it is time limited which needs to be extended with masks and ventilation over a longer period . Interestingly the effect of both wearing surgical masks was greater than 200 litres per minute of air replacement but both combined made an extremely large difference .

Discussion

Wearing a mask is beneficial at every aspect of infection modality . It can reduce aerosol inhalation with varying degrees of efficiency , from cloth to FFP3, all masks will reduce large droplets and reducing both will reduce fomite contamination ; the wide distribution of virus in CoVid-19 infected patients' rooms shows that aerosol deposition is important [24] in addition to droplets and direct touch . Any

reduction in the ID could be significant by preventing infection or reducing viral load and degree of infection . [25] High risk situations can be identified where greater protection is needed especially where a potentially infectious person is in one place for a long period or people are in close contact even for a short time , then use of masks is important . In addition to masks good ventilation and possibly air filtration could be important in work places or anywhere social distancing is time limited by a confined area . It should be noted that ventilation can be beneficial or harmful   depending on air flow patterns [26] and input from health and safety and occupational health agencies would be useful in minimising risk in such places . Better quality protection is needed in environments such as public transport where air borne infections have been identified   [27,28] , inside shops where social distancing cannot be fully maintained at all times , or anywhere there is poor ventilation   . The model suggests validation of strategies already in use such as putting a mask on a patient and ventilation of rooms by opening windows .

In a pandemic early intervention is essential using good quality masks as part of a multifaceted strategy . Reductions in infections   early on will have considerable benefit due to the number of people each infected person infects .   Tracht et al produced a model for influenza that showed even with 50% compliance in the population wearing N95 masks , there would be more than a 36% reduction in the number of cases. [29] They found less benefit for surgical masks but may have underestimated the combined efficiency of wearing surgical masks . Reducing the numbers of infected people early on will give contact tracing and isolation a much better chance to work . Those returning from infected areas should use masks

immediately together with isolation . Models in pandemics should include aerosol spread and mask use , noting there was no mention of either in the models published by the UK government .[30,31] If there is an epidemic with rapid spread in a population with high morbidity a warning is sounding that in that country, or for other countries as soon as any spread out with that country occurs , plans for implementing the general use of masks should be engaged . The evidence for this is in the countries that have good control of the outbreak even with an initial exponential rise in cases , such as China , Japan , Korea , and Taiwan , although it is confounded with other measures such as contact tracing and social distancing . However , to suggest a deviation from an effective policy which includes the use of masks requires strong evidence that masks are of no benefit . Anytime lockdown is likely masks should already be in general use and would be cost effective even if it is only shortened by a week .

For Covid-19 the lack of resources is all too evident but a lack of resources should not dictate public health policy , public health policy should ensure resources are available when needed . With regard to masks national stock piles are necessary , high grade protection for the care sector and at least surgical masks for the general population . The more people that can use higher grade protection such as N95 especially on public transport and crowded places the better but if there is a national supply of surgical masks there would be a fall back for those unable to tolerate the higher grade masks or unable to procure them . Manufacturers increasing the shelf life from 5 years to 10 years would make this much more feasible . In addition to a national endeavour once supplies are normalised at a domestic level all households should be encouraged through public health information to keep a supply of good quality masks that can be used for dusty activities and DIY , then replaced when

dirty . Alternatively there are reusable masks such as sports masks which are usually available with PM2.5 2.5 micron pollen or dust filters but could be supplied with additional PM1 1 micron filters for pandemics , balancing comfort with reduced efficiency compared to N95 masks . Exhalation valves are designed to reduce moisture and CO_2 in theory but it appears with the current design most valves do not work at low levels of breathing [32] which allows a protective effect in exhaling ; masks without exhalation valves still seem preferable for general use . Improving the exhalation filtering and fit would improve the combined efficacy of wearing masks .

Most of the issues postulated that masks in the community would be detrimental are hand hygiene issues, such as touching the mask with unclean hands , meaning public educational information is needed and similarly advice on fitting could be addressed . As viruses cannot multiply on a mask there is no situation where wearing a mask can be worse than not wearing one . Compliance is not an issue when there is a national emergency with significant risk to individual health and the wider economy provided the correct information is given [32] , and if necessary relatively easy to enforce on public transport , at work or in shops providing supplies are available . Protection against eye risk is not a case against masks , which is more likely when close to someone from droplets compared to aerosol (taking the study results and the surface area of the lungs compared to the eyes) . In this situation a minimum of everyone using a basic mask and hand hygiene , similar to precautions needed in commercial food preparation , will help to reduce eye contamination risk .

Conclusion

The model shows the relative risk of aerosol in various situations together with the benefit from use of masks which is multiplicative . Social distancing at 2 metres appears validated but in confined areas is time limited and the use of masks in addition to efficient ventilation is important. Where social distancing is not possible at all times or a potentially infectious person is in a confined area for a long time there is a higher risk of infection requiring protection . The use of masks should be factored into models and used at an early stage as widespread use of efficient masks could have a large impact on control and spread of infection . Public health planning requires stockpiling of masks and encouraging everyone to have suitable masks in their household when supplies are normalised . The use of a cloth mask is better than no protection at all .

Acknowledgement to Timothy Heelis for providing assistance with the numerical calculations .

References

1.WHO. Advice on the use of masks the community, during home care and in health care settings in the context of the novel coronavirus (2019-nCoV) outbreak . Interim guidance 29 January 2020.

https://www.who.int/docs/defaultsource/documents/advice-on-the-use-of-masks-2019-ncov.pdf

2. WHO. Advice on the use of masks in the context of COVID-19 . Interim guidance 6 April 2020. https://apps.who.int/iris/rest/bitstreams/1274280/retrieve

3. WHO Non-pharmaceutical public health measures for mitigating the risk and impact of epidemic and pandemic influenza . 2019. Licence: CC BY-NC-SA 3.0 IGO. https://apps.who.int/iris/bitstream/handle/10665/329438/9789241516839-eng.pdf

4. WHO . Modes of transmission of virus causing COVID-19: implications for IPC precaution recommendations Scientific brief . 29 March 2020 .

https://apps.who.int/iris/rest/bitstreams/1273450/retrieve.

5. Choi S, Moran Ki M Estimating the reproductive number and the outbreak size of COVID-19 in Korea Epidemiol Health . 2020; 42: e2020011.

2020 DOI: https://doi.org/10.4178/epih.e2020011

6. Nussbaumer-Streit B , Mayr V , Dobrescu AI et al Quarantine alone or in combination with other public health measures to control COVID-19: a rapid review

Cochrane Database of Systematic Reviews 2020 NO: 4 PB: John Wiley & Sons, Ltd

SN:1465-1858 DOI:10.1002/14651858.CD013574

https://doi.org//10.1002/14651858.CD013574

7. Jefferson T, Del Mar C B, Dooley L, et al. Physical interventions to interrupt or reduce the spread of respiratory viruses. Cochrane Systematic Review - Intervention Version published: 2011. https://doi.org/10.1002/14651858.CD006207.pub4

8. Dixon M , Phin N. The Use of Facemasks and Respirators during an Influenza Pandemic Scientific Evidence Base Review. Department of Health and produced by Public Health England . 2014.

https://assets.publishing.service.gov.uk/government/uploads/system/uploads/attachment_data/file/316198/Masks_and_Respirators_Science_Review.pdf

9. Greenhalgh G , Schmid M B , Czypionka T, Face masks for the public during the covid-19 crisis BMJ 2020; 369 doi: https://doi.org/10.1136/bmj.m1435 (April 2020) Cited as: BMJ 2020;369:m1435

10. Feng S, Shen C, Xia N, Song W, Fan M, Cowling BJ. Rational use of face masks in the COVID-19 pandemic [published online ahead of print, 2020 Mar 20]. Lancet Respir Med. 2020;S2213-2600(20)30134-X. doi:10.1016/S2213-2600(20)30134-X

11. Edwards D A, Man J C, Brand P et al .Inhaling to mitigate exhaled bioaerosols Proc Natl Acad Sci USA . 2004 ;101(50) : 17383-17388 .

https://doi.org/10.1073/pnas.0408159101

12. Barr G D . The Covid-19 Crisis and the need for suitable face masks for the general population . Chinese J Med Res . 2020; 3:28-31 doi:10.37515/cjmr.091X.3108

13. Gawn J, Clayton M, Makison C , Crook B . Evaluating the protection afforded by surgical masks against influenza bioaerosols . Gross protection of surgical masks compared to filtering facepiece respirators . 2008 Health Improvement and Human Factors Groups Health and Safety Laboratory Health and Safety Executive . https://www.hse.gov.uk/research/rrpdf/rr619.pdf

14. C Davies, Anna & Thompson, Katy-Anne & Giri, Karthika & Kafatos, George & Walker, James & Bennett, Allan. Testing the Efficacy of Homemade Masks: Would They Protect in an Influenza Pandemic? Disaster medicine and public health preparedness. 2013 ; 7: 413-418. 10.1017/dmp.2013.43.

15. Nikitin N , Petrova E, Trifonova E,   Karpova O Influenza Virus Aerosols in the Air and Their Infectiousness. Adv. Virol. 2014 . doi: 10.1155/2014/859090 https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4147198/#__ffn_sectitle

16. Leung , N H L , Chu, D K W, Shiu, E.Y.C. et al. Respiratory virus shedding in exhaled breath and efficacy of face masks. Nat Med . 2020 ; 26: 676–680 . https://doi.org/10.1038/s41591-020-0843-2

17. Van der Sande M , Teunis P, Sabel R . Professional and home-made face masks reduce exposure to respiratory infections among the general population. PLoS One. 2008 ; 3(7):e2618. doi:10.1371/journal.pone.0002618

18. Ari A, Fink JB, Pilbeam S. Secondhand aerosol exposure during mechanical ventilation with and without expiratory filters: An in-vitro study. Ind J Resp Care 2016; 5 : 677-82.

19. Adams D J , Spendlove J C , Spendlove R S , et al . Aerosol Stability of Infectious and Potentially Infectious Reovirus Particles.   Applied and Environmental Microbiology . 1982. 198: 44 ; 903-908

20. Fabian P, McDevitt J J, DeHaan W H, et al . Influenza Virus in Human Exhaled Breath: An Observational Study Plos One 2008 ; 3 (7)
doi:10.1371/journal.pone.0002691

21. Yang W , Elankumaran S , Marr .L C .Concentrations and size distributions of airborne influenza A viruses measured indoors at a health centre, a day-care centre and on aeroplanesYJ R Soc Interface. 2011 ; 8(61): 1176–1184.

22. Zhili Zuo , Thomas H. Kuehn , Harsha Verma , et al . Association of Airborne Virus Infectivity and Survivability with its Carrier Particle Size . Aerosol Science and Technology . 2013 ; 47: (4) 373-382 .

23 . Van Doremalen N , Bushmaker T, Morris D , et al . Aerosol and Surface Stability of SARS-CoV-2 as Compared with SARS-CoV-1. New England Journal of Medicine 2020; 382:1564-1567 . doi:10.1056/NEJMc2004973

24. Santarpia1 J L, Rivera D N, Hererra V, et al. Transmission Potential of SARS-CoV-2 in Viral Shedding Observed at the University of Nebraska Medical Center. accessed 2020. preprint from the University of Nebraska Medical Center. https://www.medrxiv.org/content/10.1101/2020.03.23.20039446v1.full.pdf .   doi: https://doi.org/1.1101/2020.03.23.20039446

25. PauloA C , Correia-Neves M , Domingos T , et al   Influenza Infectious Dose May Explain the High Mortality of the Second and Third Wave of 1918–1919 Influenza Pandemic . Plos One . 2010 ;5(7): e11655 . https://doi.org/10.1371/journal.pone.0011655

26. Villafruela JM, Olmedo I, Berlanga FA, Ruiz de Adana M . Assessment of displacement ventilation systems in airborne infection risk in hospital rooms. PLoS ONE 2019 ;14(1): e0211390. https://doi.org/10.1371/journal.pone.0211390

27. Zhang L, Peng Z, Ou J, Zeng G, Fontaine RE, Liu M, et al. Protection by face masks against influenza A(H1N1)pdm09 virus on trans-Pacific passenger aircraft, 2009. Emerg Infect Dis. 2013;19 (9) :1403-1410. doi:10.3201/eid1909.121765

28. Xiaopeng L , Zhang S , COVID-19: Face masks and human-to-human transmission. Influenza and other respiratory Viruses . 2020 .

https://doi.org/10.1111/irv.12740

https://onlinelibrary.wiley.com/doi/full/10.1111/irv.12740

29. Tracht S M , Del Valle S Y , Hyman J M . Mathematical Modeling of the Effectiveness of Facemasks in Reducing the Spread of Novel Influenza A (H1N1) PLoS One. 2010; 5(2): e9018. doi: 10.1371/journal.pone.0009018

30. Petra Klepac P, Kucharski A J ,Conlan A J K, Kissler S . Contacts in context: large-scale setting-specific social mixing matrices from the BBC Pandemic project .

https://www.medrxiv.org/content/10.1101/2020.02.16.20023754v2.full.pdf+html

31. Joël Mossong , Niel Hens, Mark Jit, Philippe Beutels Social Contacts and Mixing Patterns Relevant to the Spread of Infectious Diseases Social Contacts and Mixing Patterns Relevant to the Spread of Infectious Diseases . Plos One. 2008 PLoS Med 5(3): e74. https://doi.org/10.1371/journal.pmed.0050074

https://journals.plos.org/plosmedicine/article?id=10.1371/journal.pmed.0050074

32. Roberge RJ Are exhalation valves on N95 filtering facepiece respirators beneficial at low-moderate work rates: an overview.J Occup Environ Hyg. 2012; 9(11):617-23

33. Sim S W, Moey K S, Tan N C. The use of facemasks to prevent respiratory infection: a literature review in the context of the Health Belief Model. Singapore Med J. 2014;55(3):160–167. doi:10.11622/smedj.2014037

Table I   Scenario 1

Close by , breathing the exhaled air of an infectious person with no dilution , av is taken at 0.3 , both breathing normally so Ev and Bv are each 6 l/min	
ID Infectious dose is 1000 particles	
ID 1000 = 500 x 0.3 x 1 x 6 x (6 / 6) x 1 x 1 x t , t = 1000/900	**t = 1.1 mins**
Non infected wearing surgical mask (F$_2$ 0.167)	**t = 6.7 mins**
Both wearing surgical mask (F$_1$ 0.333 F$_2$= 0.167)	**t = 19 mins**
Non infected surgical mask (F$_2$ = 0.167;Infected FFP3 mask F$_1$= 0.333)	**t = 19 mins**
Non infected FFP3 (F$_2$ = 0.01 ; infected surgical mask F$_1$=0.333)	**t= 334 mins**
Both wearing cloth mask (F2= 0.5 F1=0.75)	**t=2.96 mins**

Table II   Scenario 2

2 people in a car one infected , 2 m apart , Volume 2 m^3 no ventilation , t=30 mins Fresh uncontaminated air at the start		
No masks	ID = 1078 p	t 30 mins
Non infected wearing surgical mask	ID = 177 p	t 30 mins
Non infected wearing FFP3	ID = 11 p	t 30 mins
For both using surgical mask	ID = 58 p	t 30 mins
Both using cloth mask	ID = 396 p	t 30 mins

Table III   Scenario 3

2 people in confined space such as a car 2 m^3 , one infected person already present for 3 hours and no ventilation .	
Starting concentration at 180 minutes is 35 p/l	
No masks	ID is 1060 p at 5 mins
Non infected FFP3	ID is 1011 p at 511mins
Non infected Surgical mask	ID is 1011 p at 29 mins
Non infected Cloth mask	ID is 1056 at 10 mins
Both wearing surgical masks from the start ID is 1001 at 95 mins	
Relatively short time so ventilation with fresh air is needed unless using FFP3 or both using surgical masks	
Introducing ventilation with 5 % of air replaced / min from the start of the three hours	
Starting concentration at 180 minutes is 7 p/l	
Non infected with no mask	ID is 1008 at 24 mins
Non infected with cloth mask	ID is 1003 at 48 mins

Table IV Scenario 4

Infected person in a room or working in a small shop 30 m^3 for 3 hours , no ventilation
Starting concentration at 180 minutes is 2.6 p/l
Non infected with no mask ID 1004 at 63 minutes
Non infected with cloth mask ID 1003 at 122 minutes

Table IV   Scenario 5

2 people socially distanced at 2m in a confined area with a 4m^3 corridor of air between (assuming the air is static and no diffusion to a wider volume) , one infected and no ventilation , time to reach ID of 1000		
No masks	ID of 1027	reached at 42 mins
Both wearing surgical masks	ID of 1000	reached at 243 mins
With ventilation 5% air replaced per minute (200 litres/min)		
No masks	ID of 1003	reached at 62 minutes
Both wearing surgical masks	ID of 1001	reached at 866 mins

Figure 1   Scenario 3 Cloth Mask

Scenario 3. (a) Entering car   wearing   a Cloth Mask with infected person present for 3 hours

Scenario 3 .   (b) Entering car wearing   Cloth   Mask   with infected person present for 3 hours   and 5% of the air replaced per minute