

1 **BRAZIL IS PROJECTED TO BE THE NEXT GLOBAL COVID-19 PANDEMIC**
2 **EPICENTER**

3 **Running title: Brazil the next pandemic epicenter**

4
5 Pedro de Lemos Menezes¹, David M. Garner², Vitor E. Valenti^{3*}

6
7 ¹University of Health Sciences of Alagoas, AL, Brazil. ²Cardiorespiratory Research
8 Group, Department of Biological and Medical Sciences, Faculty of Health and Life
9 Sciences, Oxford Brookes University, Headington Campus, Oxford, OX3 0BP, United
10 Kingdom. ³Autonomic Nervous System Center (CESNA), São Paulo State University,
11 UNESP, Marília, Brazil.

12
13 * **Corresponding author:** Vitor E. Valenti
14 São Paulo State University, UNESP.
15 Av. HyginoMuzzi Filho, 737
16 Marília, SP, Brazil - 17.525-000
17 E-mail: vitor.valenti@unesp.br

18
19
20

21 **ABSTRACT**

22 Coronavirus disease 2019 (COVID-19) is a disease triggered by SARS-CoV-2 infection,
23 which is related in the most recent pandemic situation, significantly affecting health and
24 economic systems. In this study we assessed the death rate associated to COVID-19 in
25 Brazil and the United States of America (USA) to estimate the probability of Brazil
26 becoming the next pandemic epicenter. We equated data between Brazil and USA
27 obtained through the Worldometer website (www.worldometer.info). Epidemic curves
28 from Brazil and USA were associated and regression analysis was undertaken to predict
29 the Brazilian death rate regarding COVID-19 in June. In view of data from April 9th
30 2020, death rates in Brazil follow a similar exponential increase to USA ($r=0.999$;
31 $p<0.001$), estimating 64,310 deaths by June 9th 2020. In brief, our results demonstrated
32 that Brazil follows an analogous progression of COVID-19 deaths cases when compared
33 to USA, signifying that Brazil could be the next global epicenter of COVID-19. We
34 highlight public strategies to decrease the COVID-19 outbreak.

35

36 **Keywords:** Beta-coronavirus; Centers for Disease Control and Prevention (U.S.A.);
37 Coronavirus; Coronavirus Infections; Death; SARS Virus.

38

39 INTRODUCTION

40 The Coronavirus disease 2019 (COVID-19) is a new disease that has developed
41 into a global public health concern as it has rapidly extended worldwide. The
42 transmission of severe acute respiratory syndrome Coronavirus 2 (SARS-CoV-2),
43 initiating COVID-19, has led to this urgent pandemic status (Angelos et al, 2020).

44 This disease initially appeared in Wuhan, China, which became the first epicenter
45 (Nature, 2020). Next, when the cases were partially controlled in China, Europe became
46 affected and was an added epicenter of COVID-19 (Nature, 2020). At present, the
47 United States of America (USA) is the current pandemic epicenter with in excess of
48 700,000 COVID-19 cases, consistent with the Worldometer website
49 (www.worldometer.info).

50 In this scenario, we understand that Brazil has the potential for a considerable
51 increase in deaths related to COVID-19 (www.worldometer.info). We comprehend that
52 Brazil has the likelihood to be the next pandemic epicenter of the disease. So, in this
53 report, we aim to evaluate deaths associated to COVID-19 in Brazil and USA, so as to
54 authenticate any likenesses between the countries.

55

56 METHODS

57 The mortality data for COVID-19, which occurred in Brazil and the USA, was
58 obtained from the Worldometer website (www.worldometer.info).

59 Worldometer is managed by an international team of developers, researchers, and
60 volunteers' with the objective of providing global health statistics available in a
61 challenging and time relevant format, to an audience world-wide. It is published by a
62 small and independent digital media company located in the USA. They have no
63 political, governmental, or corporate affiliations.

64 Deaths were estimated until the 31st day after the 5th death in both countries.

65 At first, a link was achieved between COVID-19 mortality in the USA, which is
66 related to the Pandemic development process and is considered as the current pandemic
67 epicenter. Temporal modifications were completed, enabling the comparison of data on
68 the Cartesian x-axis, and the adjustment of the number of deaths for each 10,000,000
69 inhabitants, which allowed the procurement of data on the Cartesian y-axis.

70 To relate the mortality results between the two countries, the Bivariate correlation
71 test was enforced and the degree of linear relationship was considered by the Spearman
72 coefficient.

73 Formerly, there were some regressions to discover the best mathematical model
74 which would allow the representation of the mortality estimation curve for COVID-19
75 in Brazil and USA. Whilst searching for the mathematical model that best fits the
76 mortality estimation curves, the following models were evaluated: Linear, Logarithmic,
77 Inverse, Quadratic, Cubic, Composite, S, Growth, Exponential and Logistic. For this
78 estimate, a significance of alpha, (α) less than 0.01 ($p < 0.01$, $< 1\%$) and $R^2 > 0.95$ was
79 adopted. Furthermore, the one-way analysis of variance (ANOVA1) test was applied.

80 The data were managed via Microsoft Excel version 16.36, for MacOS Catalina
81 version 10.15.4. Statistical assessments were computed by IBM SPSS Statistic
82 Subscription application 1.0.0.1347 64Bits for MacOS. These values were considered
83 statistically significant when ($p < 0.05$, $< 5\%$) and the established beta, (β) value was 0.1.

84

85 RESULTS

86 Until the 31st day after the 5th death, the increase in mortality in Brazil and USA
87 follow a similar pattern. The bivariate correlation test was significant with $p < 0.001$ and
88 degree of linear relationship was $R^2 = 0.99$ (Figure 1).

89 In Brazil, the mathematical models attempted to estimate the mortality curve in
90 absolute values, from the 5th to the 30th death (April 17th 2020) and the models that came
91 nearest were the cubic and the quadratic, as illustrated in Figure 2.

92 Nevertheless, as illustrated in Table 1, the cubic model fits best, $R^2 = 0.998$.

93 Consequently, the cubic model, thus far, best fits an approximation of the growth
94 curve of mortality owing to COVID-19 in Brazil and is in the following manner:

95
96 $x(t) = 11,422 + 0,269 * t + 0,265 * t^2 + 0,069 * t^3$
97

98 Where t is the number of days from the 5th death.
99

100 With regards this mathematical model, it was demonstrated in Figure 3 that the
101 estimation of total deaths for Brazil, assuming no behavioral changes amongst the
102 population for the 50th day after the 5th death is 9,312 deaths by COVID-19.
103

104 **DISCUSSION**

105 Our brief report was commenced to assess the level of COVID-19 related deaths
106 in Brazil and USA. This intention was explained so as to estimate the possibility of
107 Brazil becoming the next pandemic global epicenter. Our results described significant
108 connections between Brazil and USA about total mortality related to COVID-19 up
109 until the 31st day after the 5th death.

110 The main concern regarding this mathematical forecast is to enable data to permit
111 the Brazilian government to format policies to protect its own population.

112 Our study highlights some problems which need addressing. Some Brazilian
113 states, including Sao Paulo, reported only a few COVID-19 cases (Sao Paulo sanctions
114 warnings only in severe cases of COVID-19 and declares that it maintains advice from
115 the Brazilian Ministry of Health), we emphasize the potential of the under reporting of
116 cases.

117 In summary, the mathematical model enforced in Brazil and USA advocates that
118 Brazil may be the next global pandemic epicenter. We encourage adherence to Brazilian
119 public guidelines to reduce the impact of COVID-19 in Brazil.
120

121 **ACKNOWLEDGMENTS**

122 Authors thank the technical staff from Sao Paulo State University,
123 UNESP/Marilia, SP, Brazil and University of Health Sciences of Alagoas, AL, Brazil.
124 Dr. Vitor receives financial support from FAPESP (Process number 2012/01366-6)
125 financial support from the National Council for Scientific and Technological
126 Development, an entity linked to the Ministry of Science, Technology, Innovations and
127 Communications from Brazil (Process number 302197/2018-4).
128

129 **AUTHOR CONTRIBUTIONS**

130 Pedro de Lemos Menezes draft the manuscript, wrote introduction and discussion
131 section gave final approval for the version submitted for publication.

132 David M. Garner draft the manuscript, performed statistical analysis and improved
133 interpretation analysis, reviewed English Grammar and Spelling.

134 Vitor E. Valenti draft the manuscript, wrote introduction and discussion section
135 and gave final approval for the version submitted for publication.
136

137 **COMPETING INTERESTS**

138 The authors declare absence of financial and non-financial interests.

139

140 **REFERENCES**

141 Angelos, G., Dockter, A.G., Gachabayov, M., Latifi, R., Bergamaschi, R. (2020).
142 Emergency Colorectal Surgery in a COVID-19 Pandemic Epicenter. *Surg.*
143 *Technol. Int.* 36. pii: sti36/1297.

144 No authors listed. (2020). Stop the Wuhan virus. *Nature* 577, 450. doi: 10.1038/d41586-
145 020-00153-x

146 No author listed. (2020). Europe now centre of pandemic, says WHO. *Nature*. 13 March
147 22:00 GMT.

148 SP keeps notification only for severe cases of coronavirus and says it follows Brazilian
149 Ministry of Health. G1. Assessed on April 13th 2020:
150 [https://g1.globo.com/sp/sao-paulo/noticia/2020/04/08/sp-mantem-notificacao-
153 apenas-para-casos-graves-de-coronavirus-e-diz-que-segue-ministerio-da-
saude.ghtml](https://g1.globo.com/sp/sao-paulo/noticia/2020/04/08/sp-mantem-notificacao-
151 apenas-para-casos-graves-de-coronavirus-e-diz-que-segue-ministerio-da-
152 saude.ghtml)

153

154 **FIGURE LEGENDS**

155 **Figure 1:** Deaths associated to COVID-19 in Brazil and USA.

156 **Figure 2:** Total deaths related to COVID-19 in Brazil.

157 **Figure 3:** Comparison between total deaths related to COVID-19 in Brazil and
158 projected via mathematical model.

159

160 **Table 1:** Model summary and parameter estimates.

Dependent variable: Deaths									
Equation	R ²	Model summary				Parameter estimates			
		Z	df1	df2	Sig.	Constant	b1	b2	b3
Linear	.859	170.682	1	28	.000	-450.108	67.532		
Logarithmic	.545	33.587	1	28	.000	-789.488	556.986		
Invers	.188	6.495	1	28	.017	788.676	-1442.125		
Quadratic	.995	2654.798	2	27	.000	124.413	-40.191	3.475	
Cubic	.998	4350.630	3	26	.000	11.422	.269	.265	.069
Composed	.963	732.622	1	28	.000	13.567	1.204		
Potency	.947	497.163	1	28	.000	2.110	1.902		
S	.565	36.326	1	28	.000	6.343	-6.475		
Growth	.963	732.622	1	28	.000	2.608	.185		
Exponential	.963	732.622	1	28	.000	13.567	.185		
Logistic	.963	732.622	1	28	.000	.074	.831		


The independent variable is Days from the 5th Death.

161

162

163 **FIGURE 1**

164




165

166

167

168 **FIGURE 2**

169




170

171

172

173 **FIGURE 3**

174



175