Effects of High Intensity Interval Exercise on Cerebrovascular Function: A Systematic

Review

Alicen A. Whitaker, DPT[1], Mohammed Alwatban, PhD[1], Andrea Freemyer, PhD[1], Jaime

Perales-Puchalt, PhD, MPH[2,3], and Sandra A. Billinger, PT, PhD[1,3,4]

[1] Department of Physical Therapy and Rehabilitation Science, University of Kansas Medical

Center, Kansas City, KS

[2] University of Kansas Alzheimer's Disease Center, Fairway, KS

[3] Department of Neurology, University of Kansas Medical Center, Kansas City, KS

[4] Department of Physical Medicine and Rehabilitation, University of Kansas Medical Center,

Kansas City, KS

Corresponding Author:

Sandra A. Billinger, PT, PhD

Address: 3901 Rainbow Blvd, Mail Stop 2002, Kansas City, KS 66160

Phone number: 913-945-6685

Fax: 913-588-9428

Email: sbillinger@kumc.edu

Conflicts of Interest: None

Funding: The Georgia Holland Research in Exercise and Cardiovascular Health (REACH)

laboratory space was supported by the Georgia Holland Endowment Fund. This publication was

partly supported by the National Institute on Aging of the National Institutes of Health under

Award Number P30AG035982.

Abstract

High intensity interval exercise (HIIE) has emerged as an effective method to improve health and aerobic fitness with the benefit of decreased exercise time when compared to conventional moderate continuous exercise. However, a gap in knowledge exists regarding the effects of HIIE on contributors to overall cerebrovascular function such as cerebral blood velocity and autoregulation. The objective of this review was to ascertain the effect of HIIE on cerebrovascular function in healthy individuals. We searched PubMed and the Cumulative Index to Nursing and Allied Health Literature (CINAHL) databases for articles pertaining to the apriori chosen key words. The review followed guidelines for Preferred Reporting Items for Systematic Reviews. Six articles were reviewed via the modified Sackett's quality evaluation. HIIE outcomes included middle cerebral artery blood velocity (MCAv) (n=4), dynamic cerebral autoregulation (dCA) (n=2), cerebral de/oxygenated hemoglobin (n=2), cerebrovascular conductance index (n=1), cerebrovascular resistance index (n=1), and cerebrovascular reactivity to carbon dioxide (CO_2) (n=1). Overall quality review was moderately poor ranging from 5/7 to 3/7 quality criteria met. Compared to moderate intensity exercise, HIIE resulted in a lower exercise MCAv and HIIE decreased dCA phase following exercise compared to rest. However, we report HIIE increased oxygenated and deoxygenated hemoglobin compared to rest. This systematic review outlines the cerebrovascular response during and following HIIE in healthy individuals. Future research is critically needed to better understand the effects of HIIE on cerebrovascular function in healthy individuals and individuals with chronic conditions. In order to conduct rigorous systematic reviews in the future, we recommend studies assess MCAv, dCA, and CO_2 reactivity during and post HIIE.

Introduction

High intensity interval exercise (HIIE) has emerged at the forefront of exercise regimens due to

the shorter activity time needed to benefit[1-3]. HIIE confers similar or significant increased

aerobic fitness compared to conventional moderate intensity continuous exercise[1, 4-7]. While

aerobic fitness is a measure of increased cardiovascular health, the entire vascular system

(including the cerebral vascular system) may be improved following increased aerobic fitness[8].

A review and meta-analysis of HIIE in healthy adults has shown significant increases in aerobic

fitness [1, 5, 6, 9]. However, the effects of HIIE on cerebrovascular function have not been

systematically reviewed.

Cerebrovascular function is the ability of the cerebral blood vessels to deliver oxygen and

nutrients for neuronal metabolism and maintain cerebral blood flow through dynamic

autoregulation (dCA). dCA is the ability of the brain to sustain a constant cerebral blood flow

despite large fluctuations in peripheral blood pressure[10] [11]. During resting conditions cerebral

blood flow responds to arterial blood pressure fluctuations, neuronal metabolism, cortical

activation, arterial blood gases, and cardiac output [12]. Cerebral blood flow can be measured at

rest using magnetic resonance imaging or transcranial Doppler ultrasound (TCD). Middle

cerebral artery blood velocity (MCAv) measured by TCD is the only technique to measure

cerebral blood flow during exercise, with high temporal resolution [13]. MCAv is linearly related to

cerebral blood flow with the caveat that the MCA diameter remains unchanged[14].

Cerebrovascular response to submaximal exercise results in increased cerebral blood flow [15-

17], increased cerebral oxygenation [18, 19], and sustained dCA[20, 21]. MCAv has been shown to

increase in parallel with exercise up to moderate intensity [15, 22-25]. During continuous high

intensity exercise and hyperventilation, MCAv is decreased due to a reduction in arterial carbon

dioxide (CO_2)[26, 27] causing downstream arteriole constriction[12, 28]. Cerebrovascular reactivity is

the ability of the small vessels in the brain to vasodilate and vasoconstrict in response to

fluctuating CO_2 levels [29, 30].

Therefore, cerebrovascular function measures provide real-time dynamic responses within the

cerebrovascular system during HIIE. Previous scientific statements and narrative reviews have

recounted the latent physiological effects of HIIE on cerebrovascular function[31, 32]. However,

studies involving cerebrovascular function during HIIE have not been systematically reviewed.

The purpose of this systematic review was to address this gap in knowledge and report the

effects of HIIE on cerebrovascular function compared to moderate continuous exercise or rest

conditions. We systematically examined the results of HIIE studies in healthy individuals based

on the operationalization of cerebrovascular outcomes.

Methods

This review follows the guidelines for Preferred Reporting Items for Systematic Reviews [33].

Literature searches and reviews were performed using PubMed and the Cumulative Index to

Nursing and Allied Health Literature (CINAHL) databases. The University of Kansas Medical

Center Online Library system was used to access these databases in February - March 2020. In

this systematic review, we included peer-reviewed manuscripts written in English from January

2010 to March 2020. We also incorporated inclusion of literature via recommendation emails in

March 2020 from experts in the field.

Key words used to search the databases included "high intensity interval training", "HIIT", "high

intensity interval exercise" "HIIE" AND "cerebral blood flow", "cerebral blood velocity", "dynamic

autoregulation". We believe these key words primarily reflect high intensity interval intervention

and cerebrovascular function outcome measures. The main outcomes of this systematic review

were MCAv and dCA. However, additional cerebrovascular measures were also included such

as oxygenated hemoglobin, cerebrovascular reactivity, cerebrovascular conductance index, and

cerebrovascular resistance index. Cerebrovascular conductance index is a measure of the

conductance of peripheral blood pressure to cerebral blood velocity and is calculated as

MCAv/mean arterial pressure (MAP). Cerebrovascular resistance index (MAP/MCAv) measures

the resistance of cerebral perfusion pressure to cerebral blood velocity.

The identified abstracts from PubMed and CINAHL were screened using the following inclusion

criteria: 1) experimental or quasi-experimental, 2) aerobic exercise identified as the primary

means of performing HIIE, 3) cerebrovascular measures were primary or secondary outcomes,

and 4) human subjects across the lifespan with no current disease. After the removal of

duplicates, 2 researchers screened titles/abstracts for inclusion criteria (A.W. and M.A.). The full

texts were examined, and data extracted (A.W. and M.A.). If the authors were unable to come to

an agreement, a third author moderated incongruity (A.F.).

A quality review was performed for each article using the modified version of Sackett's 1981

criteria[34]. We critically analyzed each article's study design, population, HIIE protocols,

cerebrovascular outcomes, and results. If an article did not report enough information to

determine sufficient quality criteria a "No" rating was given. Articles were rated based on the

level of evidence including level 1 for large randomized control trials, level II for small

randomized trial, level III for nonrandomized design, Level IV for case series, and Level V for

case reports [35].

Results

The search methods resulted in 64 articles. After removal of duplicates, 12[36-47] articles were

identified in PubMed and 5[48-52] new articles in CINAHL. One recent article (published March

2020) was not recovered through key word searches. However, experts in the field

recommended the recently published article to the authors[53]. During the initial screening of

titles/abstracts, 11 articles were excluded due to HIIE not being the primary experimental

protocol performed (n=6), studies not measuring cerebral arteries (n=4) and an animal study

(n=1). Studies that combined other interventions with HIIE were excluded due to the

confounding variables that could affect cerebrovascular outcomes. After the full text

assessment, 1 article was excluded due to not meeting experimental or quasi experimental

criteria (n=1). *[See Figure 1 at end of document]*

We included 6 articles describing cerebrovascular outcomes following HIIE within this review[37,

40, 43, 50, 51, 53]. The full texts are described in Table 1. *[See Table 1 at end of document]*. Of the

articles reviewed 5 were small, randomized trials and 1 nonrandomized cross-over trial. All the

studies involved healthy individuals although some only included men (n=1), women (n=1), or

children (n= 1). Prior activity levels of participants ranged from inactive, recreationally active and

endurance trained.

High Intensity Interval Protocols

Methods of prescribing HIIE varied greatly between studies. HIIE protocols included 6 to 16-

week exercise interventions (n = 3) or 1 single bout of exercise(n=3). By examining 6 to 16

weeks of HIIE, the long-term or chronic effects of this intervention were studied. By examining a

single bout of HIIE, the immediate or acute effects of the exercise were reported. In addition to

the duration variability, we found that the mode of HIIE also differed across the included studies.

One study used a treadmill as the mode of exercise with 4-minute intervals of 90-95% maximal

heart rate for 30 minutes. The remaining 5 studies used cycling as the mode of exercise but

differed in parameters ranging from 30 seconds to 7-minute intervals at 85% to 115% of

maximal watts or ~ 85% to 90% maximal heart rate. A constant between all studies included an

active (rather than passive) recovery interval between sprints. However, the intensity and

duration of recovery intervals differed greatly.

Cerebrovascular Outcome Measures

The results of this review can be operationalized based on the outcome variables measured

during HIIE such as MCAv (n=4), dCA (n=2), cerebral de/oxygenated hemoglobin (n=2),

cerebrovascular reactivity to CO_2 (n=1) and cerebrovascular conductance/resistance index

(n=1). Table 3 describes whether HIIE increased, decreased or had no influence on the

operationalized cerebrovascular measures. *[See Table 3 at end of document]* A meta-analysis

was not performed due to low number of studies (≤ 2) reporting each operationalized

cerebrovascular measure.

<u>*MCAv*</u>

Of the studies reporting MCAv outcomes, resting MCAv (n = 2), exercise MCAv (n = 2) and

MCAv immediately post exercise (n = 1) were used. No significant differences were found for

resting MCAv after 6 or 12 weeks of HIIE when compared to moderate continuous exercise or

control[37, 40]. During HIIE, exercise MCAv was significantly decreased compared to moderate

continuous exercise[50, 53]. Conflicting results were found between two studies comparing

exercise MCAv to rest. Burma et al[53] reported no significant difference between average

exercise MCAv and rest in adults. However, rather than reporting average exercise MCAv of the

entire HIIE bout, Tallon et al[50] reported exercise MCAv for each 1-minute sprint interval of HIIE.

During the 6th sprint interval of HIIE, Tallon et al[50] reported significantly decreased exercise

MCAv compared to rest which remained immediately following and up to 30 minutes post

exercise[50].

<u>*dCA*</u>

Transfer function analysis of dCA was reported in the very low and low frequency bands (n = 2).

Drapeau et al[37] conducted a 6-week intervention of HIIE and reported a significant decrease in

phase compared to rest with no significant change in coherence or gain. Burma et al[53]

conducted a single bout of HIIE and reported decreased MCAv systolic phase immediately

following exercise that extended up to 4 hours later.

De/Oxygenated Hemoglobin

Oxygenated and deoxygenated hemoglobin were reported during a single bout of HIIE (n=1)

and during a 16-week HIIE intervention (n=1). Monroe et al[51] conducted a single bout of HIIE

and reported an increase in oxygenated and deoxygenated hemoglobin during HIIE compared

to moderate continuous exercise. Coetsee et al[43] conducted a 16-week intervention of HIIE and

reported no significant lasting changes in oxygenated or deoxygenated hemoglobin during

cortical activation.

Cerebrovascular Reactivity, Conductance, and Resistance

The other outcomes measures such as cerebrovascular reactivity to CO_2, cerebrovascular

conductance index, and cerebrovascular resistance index were only reported in a single study

each. Cerebrovascular reactivity to CO_2 was not significantly different following 12 weeks of

HIIE[40]. A 6-week HIIE intervention reported no significant changes in cerebrovascular

conductance index or cerebrovascular resistance index[37].

Quality Review

The quality review of each study is presented in Table 2. *[See Table 2 at end of document].* Out

of 7 total quality criteria, (n=1) study met 5 criteria, (n=1) study met 4 criteria, and the remaining

(n=4) studies met 3 criteria. Therefore, the overall quality criteria results were moderately poor.

All studies accounted for subjects and monitored the HIIE protocol parameters. No studies

reported avoidance of contamination or co-intervention. No studies reported blinding of the

outcome assessments. Only Burma et al[53] and Monroe et al[51] reported their reliability via

coefficient of reproducibility and intraclass coefficients of their measures. And only Burma et

al[53] and Drapeau et al[37] reported validity of their respective cerebrovascular outcomes.

Discussion

This review met the objective of reporting various HIIE effects on operationalized

cerebrovascular function in healthy individuals. This review is the first to systematically report

the effects HIIE on cerebrovascular function compared to moderate continuous exercise and

rest.

MCAv

In these studies, 6 to 12-week HIIE interventions had no effect on resting MCAv in healthy

individuals. A ceiling effect may be observed for young, healthy individuals and could explain no

changes in resting MCAv[16]. During a single bout of HIIE, hyperventilation and downstream

arteriole vasoconstriction may explain the acute decreases in exercise MCAv compared to

moderate continuous exercise[12, 28, 54, 55]. Vasoconstriction may also play a protective role during

HIIE due to heightened peripheral blood pressure potentially causing hyper-perfusion or

damage to the blood brain barrier[10, 32, 56]. During a single bout of HIIE, there is contradictory

evidence comparing exercise MCAv to rest. One study reported no change in average exercise

MCAv compared to resting. Another study reported decreased exercise MCAv after 6 intervals

of HIIE and remained decreased compared to rest immediately following and up to 30-minutes

after HIIE[50]. The differences reported in exercise MCAv compared to rest could be due to age

(adults versus prepubertal children) or due to the analysis of MCAv during HIIE (average over

entire exercise versus separate sprint intervals). Decreases in exercise MCAv compared to rest

may only occur in the late intervals of HIIE, during hyperventilation. Therefore, exercise MCAv

should be reported for each interval of HIIE rather than an average of the entire exercise bout.

dCA

After a 6-week intervention and single bout of HIIE, dCA phase was decreased compared to

rest. In healthy individuals, increased frequency within MCAv and MAP waveforms (that can

occur with HIIE) caused a reduction in phase due to dCA being a high-pass filter[57, 58]. Burma et

al[53] also suggests that systolic phase may reveal greater changes than both diastolic and mean

phase.

Although not included in this review due to the observational study design, contradictory

evidence of sustained dCA during HIIE has been reported[41]. Differences in exercise

parameters between HIIE may be the cause to contradictory findings due to exhaustive exercise

showing decreased dCA [59, 60]. More studies are needed to confirm the acute and chronic

decreases in dCA following HIIE.

De/Oxygenated Hemoglobin

After a 16-week HIIE intervention, oxygenated and deoxygenated hemoglobin during cortical

activation did not change. However, the 16-week HIIE intervention decreased reaction time

during cortical activation and therefore may have increased efficiency of cortical oxygen use.

Also, during a single bout of HIIE oxygenated and deoxygenated hemoglobin increased

compared to moderate continuous exercise. The acute and chronic effects of HIIE on

oxygenated and deoxygenated hemoglobin still needs further investigation due to each only

being reported in a single study.

Cerebrovascular Reactivity, Conductance, Resistance

Due to cerebrovascular reactivity, conductance index or resistance index being reported in only

a single study, no conclusive effects of HIIE can be determined. However, 6 to 12 week HIIE

interventions may not produce changes in cerebrovascular reactivity due to vascular

desensitization from chronic exposure to CO_2 during HIIE[61].

Article Limitations

The individual study limitations have been previously acknowledged within the published articles

or have been identified by authors (AW and MA) during the quality review. Most of the studies

were small, randomized trials and included only healthy individuals. The small sample size

available in males (n = 17), females (n = 17), or children (n = 8) limit the generalizability of

results presented in this systematic review to a larger population. All studies implemented HIIE

protocols that differed in time, repetitions and intensity levels, therefore making comparisons

between studies difficult. These studies report potential limitations to the methods of measuring

cerebrovascular function. TCD is the preeminent method of measuring MCAv during exercise

but MCAv during HIIE may be underestimated[62]. However, these studies report the minor

changes in expired CO_2 during HIIE were not likely to induce changes in MCA diameter[14, 50, 51].

The studies also report underestimation of cerebral oxygenation due to the two-channel near

infrared spectrometer not measuring the motor, occipital, or parietal cortex [43, 51].

Review Limitations

The authors acknowledge a risk of publication bias by only including peer-reviewed articles

written in English and did not include grey literature. The cerebrovascular function measures

included within this review vary greatly and have vast heterogeneity. While HIIE is not a new

mode of exercise, studying cerebrovascular measures during HIIE is novel. Therefore, authors

could only identify 6 small studies with the oldest article dating back to 2015. The primary

outcome of MCAv (n=4) and dCA (n=2) were reported in few studies with low power. Therefore,

a meta-analysis could not be performed due to insufficient mathematical combination.

Conclusion

This review has provided preliminary information studying cerebrovascular function with HIIE.

We propose that a single bout of HIIE may decrease MCAv, decrease dCA, and increase

oxygenated hemoglobin during or immediately following HIIE compared to moderate continuous

exercise or rest conditions. However, no significant effects on resting MCAv, cerebrovascular

reactivity to CO_2, cerebrovascular conductance index or cerebrovascular resistance index were

found after weeks of HIIE interventions. With increased interest in healthy brain aging and

implementing interventions to maintain or improve brain health, studying the effects of HIIE

interventions are critically needed. While this review only included healthy individuals, we

provide an early reference to understanding "normal" physiological effects of HIIE on

cerebrovascular function and the need to compare to clinical populations.

Future HIIE research is needed to evaluate cerebrovascular function during and immediately

following exercise. More studies such as randomized controlled trials with large sample sizes

are needed to conduct a meta-analysis to combine and statistically analyze the summary results

of HIIE on cerebrovascular function. Additionally, more studies are needed to determine the

optimal parameters for HIIE and cerebrovascular function for humans.

References

1. Batacan, R.B., Jr., et al., *Effects of high-intensity interval training on cardiometabolic health: a systematic review and meta-analysis of intervention studies.* Br J Sports Med, 2017. **51**(6): p. 494-503.
2. Costa, E.C., et al., *Effects of High-Intensity Interval Training Versus Moderate-Intensity Continuous Training On Blood Pressure in Adults with Pre- to Established Hypertension: A Systematic Review and Meta-Analysis of Randomized Trials.* Sports Med, 2018. **48**(9): p. 2127-2142.
3. Ramos, J.S., et al., *The impact of high-intensity interval training versus moderate-intensity continuous training on vascular function: a systematic review and meta-analysis.* Sports Med, 2015. **45**(5): p. 679-92.
4. Rakobowchuk, M., et al., *Sprint interval and traditional endurance training induce similar improvements in peripheral arterial stiffness and flow-mediated dilation in healthy humans.* Am J Physiol Regul Integr Comp Physiol, 2008. **295**(1): p. R236-42.
5. Tabata, I., et al., *Effects of moderate-intensity endurance and high-intensity intermittent training on anaerobic capacity and VO2max.* Med Sci Sports Exerc, 1996. **28**(10): p. 1327-30.
6. Cunningham, D.A., D. McCrimmon, and L.F. Vlach, *Cardiovascular response to interval and continuous training in women.* Eur J Appl Physiol Occup Physiol, 1979. **41**(3): p. 187-97.
7. Edge, J., D. Bishop, and C. Goodman, *The effects of training intensity on muscle buffer capacity in females.* Eur J Appl Physiol, 2006. **96**(1): p. 97-105.
8. Ainslie, P.N., et al., *Elevation in cerebral blood flow velocity with aerobic fitness throughout healthy human ageing.* J Physiol, 2008. **586**(16): p. 4005-10.
9. Bayati, M., et al., *A practical model of low-volume high-intensity interval training induces performance and metabolic adaptations that resemble 'all-out' sprint interval training.* J Sports Sci Med, 2011. **10**(3): p. 571-6.
10. Aaslid, R., et al., *Cerebral autoregulation dynamics in humans.* Stroke, 1989. **20**(1): p. 45-52.
11. Li, Z., et al., *Assessment of cerebral oxygenation oscillations in subjects with hypertension.* Microvasc Res, 2013. **88**: p. 32-41.
12. Smith, K.J. and P.N. Ainslie, *Regulation of cerebral blood flow and metabolism during exercise.* Exp Physiol, 2017. **102**(11): p. 1356-1371.
13. Brassard, P., P.N. Ainslie, and N.H. Secher, *Cerebral oxygenation in health and disease.* Front Physiol, 2014. **5**: p. 458.
14. Serrador, J.M., et al., *MRI measures of middle cerebral artery diameter in conscious humans during simulated orthostasis.* Stroke, 2000. **31**(7): p. 1672-8.
15. Witte, E., et al., *Exercise intensity and middle cerebral artery dynamics in humans.* Respir Physiol Neurobiol, 2019. **262**: p. 32-39.
16. Ward, J.L., et al., *Effect of healthy aging and sex on middle cerebral artery blood velocity dynamics during moderate-intensity exercise.* Am J Physiol Heart Circ Physiol, 2018. **315**(3): p. H492-H501.
17. Brugniaux, J.V., et al., *Acute exercise stress reveals cerebrovascular benefits associated with moderate gains in cardiorespiratory fitness.* J Cereb Blood Flow Metab, 2014. **34**(12): p. 1873-6.
18. Davenport, M.H., et al., *Cerebrovascular reserve: the link between fitness and cognitive function?* Exerc Sport Sci Rev, 2012. **40**(3): p. 153-8.
19. Fisher, J.P., et al., *Cerebral perfusion, oxygenation and metabolism during exercise in young and elderly individuals.* J Physiol, 2013. **591**(7): p. 1859-70.
20. Brys, M., et al., *Dynamic cerebral autoregulation remains stable during physical challenge in healthy persons.* Am J Physiol Heart Circ Physiol, 2003. **285**(3): p. H1048-54.

348   21.   Ogoh, S., et al., *Middle cerebral artery flow velocity and pulse pressure during dynamic exercise*
349   *in humans.* Am J Physiol Heart Circ Physiol, 2005. **288**(4): p. H1526-31.
350   22.   Steventon, J.J., et al., *Cerebrovascular Function in the Large Arteries Is Maintained Following*
351   *Moderate Intensity Exercise.* Front Physiol, 2018. **9**: p. 1657.
352   23.   Herholz, K., et al., *Regional cerebral blood flow in man at rest and during exercise.* J Neurol,
353   1987. **234**(1): p. 9-13.
354   24.   Jorgensen, L.G., et al., *Middle cerebral artery flow velocity and blood flow during exercise and*
355   *muscle ischemia in humans.* J Appl Physiol (1985), 1992. **72**(3): p. 1123-32.
356   25.   Thomas, S.N., et al., *Cerebral blood flow during submaximal and maximal dynamic exercise in*
357   *humans.* J Appl Physiol (1985), 1989. **67**(2): p. 744-8.
358   26.   Hellstrom, G. and N.G. Wahlgren, *Physical exercise increases middle cerebral artery blood flow*
359   *velocity.* Neurosurg Rev, 1993. **16**(2): p. 151-6.
360   27.   Moraine, J.J., et al., *Relationship of middle cerebral artery blood flow velocity to intensity during*
361   *dynamic exercise in normal subjects.* Eur J Appl Physiol Occup Physiol, 1993. **67**(1): p. 35-8.
362   28.   Ogoh, S. and P.N. Ainslie, *Regulatory mechanisms of cerebral blood flow during exercise: new*
363   *concepts.* Exerc Sport Sci Rev, 2009. **37**(3): p. 123-9.
364   29.   Maeda, H., et al., *Reactivity of cerebral blood flow to carbon dioxide in various types of ischemic*
365   *cerebrovascular disease: evaluation by the transcranial Doppler method.* Stroke, 1993. **24**(5): p.
366   670-5.
367   30.   Willie, C.K., et al., *Regional brain blood flow in man during acute changes in arterial blood gases.*
368   J Physiol, 2012. **590**(14): p. 3261-75.
369   31.   Franklin, B.A., et al., *Exercise-Related Acute Cardiovascular Events and Potential Deleterious*
370   *Adaptations Following Long-Term Exercise Training: Placing the Risks Into Perspective-An*
371   *Update: A Scientific Statement From the American Heart Association.* Circulation, 2020. **141**(13):
372   p. e705-e736.
373   32.   Lucas, S.J., et al., *High-intensity interval exercise and cerebrovascular health: curiosity, cause,*
374   *and consequence.* J Cereb Blood Flow Metab, 2015. **35**(6): p. 902-11.
375   33.   Moher, D., et al., *Preferred reporting items for systematic reviews and meta-analyses: the*
376   *PRISMA statement.* BMJ, 2009. **339**: p. b2535.
377   34.   Sackett, D.L., *How to read clinical journals.* Can Med Assoc J, 1982. **126**(12): p. 1373.
378   35.   Sackett, D.L., *Rules of evidence and clinical recommendations for the management of patients.*
379   Can J Cardiol, 1993. **9**(6): p. 487-9.
380   36.   Dupuy, O. and J. Tremblay, *Impact of Carbohydrate Ingestion on Cognitive Flexibility and*
381   *Cerebral Oxygenation during High-Intensity Intermittent Exercise: A Comparison between Maple*
382   *Products and Usual Carbohydrate Solutions.* Nutrients, 2019. **11**(9).
383   37.   Drapeau, A., et al., *Six weeks of high-intensity interval training to exhaustion attenuates dynamic*
384   *cerebral autoregulation without influencing resting cerebral blood velocity in young fit men.*
385   Physiol Rep, 2019. **7**(15): p. e14185.
386   38.   Willis, S.J., F. Borrani, and G.P. Millet, *Leg- vs arm-cycling repeated sprints with blood flow*
387   *restriction and systemic hypoxia.* Eur J Appl Physiol, 2019. **119**(8): p. 1819-1828.
388   39.   Klein, T., et al., *Cerebral Blood Flow during Interval and Continuous Exercise in Young and Old*
389   *Men.* Med Sci Sports Exerc, 2019. **51**(7): p. 1523-1531.
390   40.   Northey, J.M., et al., *Cognition in breast cancer survivors: A pilot study of interval and continuous*
391   *exercise.* J Sci Med Sport, 2019. **22**(5): p. 580-585.
392   41.   Tsukamoto, H., et al., *Dynamic Cerebral Autoregulation Is Maintained during High-Intensity*
393   *Interval Exercise.* Med Sci Sports Exerc, 2019. **51**(2): p. 372-378.

42. Hanssen, H., et al., *Effects of different endurance exercise modalities on migraine days and cerebrovascular health in episodic migraineurs: A randomized controlled trial.* Scand J Med Sci Sports, 2018. **28**(3): p. 1103-1112.

43. Coetsee, C. and E. Terblanche, *Cerebral oxygenation during cortical activation: the differential influence of three exercise training modalities. A randomized controlled trial.* Eur J Appl Physiol, 2017. **117**(8): p. 1617-1627.

44. Kirkham, A.A., et al., *Protective effects of acute exercise prior to doxorubicin on cardiac function of breast cancer patients: A proof-of-concept RCT.* Int J Cardiol, 2017. **245**: p. 263-270.

45. Gjellesvik, T.I., et al., *Effects of High-Intensity Interval Training after Stroke (The HIIT-Stroke study) - A Multicenter Randomized Controlled Trial.* Arch Phys Med Rehabil, 2020.

46. Scott, J.M., et al., *Association of Exercise and Swimming Goggles With Modulation of Cerebro-ocular Hemodynamics and Pressures in a Model of Spaceflight-Associated Neuro-ocular Syndrome.* JAMA Ophthalmol, 2019. **137**(6): p. 652-659.

47. Saoi, M., et al., *Characterization of the Human Skeletal Muscle Metabolome for Elucidating the Mechanisms of Bicarbonate Ingestion on Strenuous Interval Exercise.* Anal Chem, 2019. **91**(7): p. 4709-4718.

48. Peyravi, A., et al., *The effect of endurance training with crocin consumption on the levels of MFN2 and DRP1 gene expression and glucose and insulin indices in the muscle tissue of diabetic rats.* J Food Biochem, 2020. **44**(2): p. e13125.

49. Komiyama, T., et al., *Cognitive Impairment during High-Intensity Exercise: Influence of Cerebral Blood Flow.* Med Sci Sports Exerc, 2020. **52**(3): p. 561-568.

50. Tallon, C.M., et al., *Intracranial Vascular Responses to High-Intensity Interval Exercise and Moderate-Intensity Steady-State Exercise in Children.* Pediatr Exerc Sci, 2019. **31**(3): p. 290-295.

51. Monroe, D.C., et al., *Effects of Sprint Interval Cycling on Fatigue, Energy, and Cerebral Oxygenation.* Med Sci Sports Exerc, 2016. **48**(4): p. 615-24.

52. Hansen, R.K., et al., *CO2 supplementation dissociates cerebral oxygenation and middle cerebral artery blood velocity during maximal cycling.* Scand J Med Sci Sports, 2020. **30**(3): p. 399-407.

53. Burma, J.S., et al., *Dynamic cerebral autoregulation across the cardiac cycle during 8 hr of recovery from acute exercise.* Physiol Rep, 2020. **8**(5): p. e14367.

54. Gonzalez-Alonso, J., et al., *Brain and central haemodynamics and oxygenation during maximal exercise in humans.* J Physiol, 2004. **557**(Pt 1): p. 331-42.

55. Linkis, P., et al., *Dynamic exercise enhances regional cerebral artery mean flow velocity.* J Appl Physiol (1985), 1995. **78**(1): p. 12-6.

56. Brassard, P., et al., *Endotoxemia reduces cerebral perfusion but enhances dynamic cerebrovascular autoregulation at reduced arterial carbon dioxide tension.* Crit Care Med, 2012. **40**(6): p. 1873-8.

57. van Beek, A.H., et al., *Cerebral autoregulation: an overview of current concepts and methodology with special focus on the elderly.* J Cereb Blood Flow Metab, 2008. **28**(6): p. 1071-85.

58. Hamner, J.W., et al., *Spectral indices of human cerebral blood flow control: responses to augmented blood pressure oscillations.* J Physiol, 2004. **559**(Pt 3): p. 965-73.

59. Ogoh, S., et al., *Dynamic cerebral autoregulation during exhaustive exercise in humans.* Am J Physiol Heart Circ Physiol, 2005. **288**(3): p. H1461-7.

60. Bailey, D.M., et al., *Exercise-induced oxidative-nitrosative stress is associated with impaired dynamic cerebral autoregulation and blood-brain barrier leakage.* Exp Physiol, 2011. **96**(11): p. 1196-207.

61. Thomas, B.P., et al., *Life-long aerobic exercise preserved baseline cerebral blood flow but reduced vascular reactivity to CO2.* J Magn Reson Imaging, 2013. **38**(5): p. 1177-83.

62.   Coverdale, N.S., et al., *Cerebral blood flow velocity underestimates cerebral blood flow during*
443   *modest hypercapnia and hypocapnia.* J Appl Physiol (1985), 2014. **117**(10): p. 1090-6.

445 **Figure 1: Flow Diagram of Study Selection**

469 HIIE = high intensity interval exercise

Table 1: Summary of Reviewed Articles

Study	Design	Level of Evidence	Subjects	Intervention	Outcome measures	Results
Burma et al, 2020	Small Randomized Cross-Over Trial	II	9 Young Adults	3 cycling conditions: - High intensity interval training (HIIT, 1-min interval at 85-90% predicted heart rate reserve with 1-min active recovery 15% power output for 10 intervals) - Moderate intensity continuous exercise (MICT, 50-60% predicted heart rate reserve for 45 min) - No-exercise control	Average Exercise MCAv dCA via Transfer Function Analysis during forced MAP oscillations (repeated squat-stand maneuver)	Significant increase in exercise MCAv during MICT (69 ± 5.6) compared to HIIT (58 ± 11.0, p<.05) and baseline (63 ± 7.2, p<.05). No significant change in exercise MCAv during HIIT compared to baseline (p>.05). Significantly higher systolic gain/phase compared to diastolic/mean gain/phase at 0.05 and 0.10 Hz during control (p<.05). Decreased systolic phase in 0.05 Hz immediately following HIIT (p>.102) and MICT until hour 4 (p>.079). Decreased systolic phase in 0.10 Hz immediately following HIIT until hour 2 (p>.11) and immediately following MICT until hour 4 (p>.079). No change in gain or coherence in 0.05 Hz or 0.10 Hz during HIIT or MICT.

Coetsee et al, 2017	Small Randomized Controlled Trial	II	67 Inactive Adults	16-week intervention Treadmill 30 min, 3x/week 4 groups: - High intensity Interval training (HIIT, 4 min interval at 90-95% HRmax with 3 min active recovery 70% HRmax) - Moderate continuous training (MCT, 70-75% HRmax) - Resistance training (RT, 50,75,100% 10 RM) - No-exercise control (CON).	Cerebrovascular measures taken during cortical activation (Stroop test): - Oxygenated Hemoglobin - Deoxygenated Hemoglobin - Total Hemoglobin Index	No significant differences in oxygenated (effect size=.45, p=.3), deoxygenated (effect size=.67, p=.14), or total hemoglobin (effect size<.6, p>.18) after HIIT. Significant decrease in deoxygenated hemoglobin (effect size=1.14, p=.01) and total hemoglobin index (effect size=1.49, p<.01) after MCT. Significant increase in oxygenated hemoglobin in CON (effect size=.76, p=.03).
Drapeau et al, 2019	Small Randomized Clinical Trial	II	17 Endurance Trained Males	6-week intervention. Cycled until exhaustion, 3x/week 2 groups: - $HIIT_{85}$ (1-7 min interval at 85% of maximal aerobic power, with active recovery of 50% of maximal aerobic power) - $HIIT_{115}$ (30sec – 1min interval at 115% of maximal aerobic power, with active recovery of 50% of maximal aerobic power).	Resting MCAv Resting CVCi Resting CVRi dCA via Transfer Function Analysis during forced MAP oscillations (repeated squat-stand maneuver)	Significant decrease in phase at 0.10 Hz in $HIIT_{85}$ and $HIIT_{115}$ (p = .048) with no differences between intensity groups. No significant difference in power spectral density (p > .39), gain (p > .05), or coherence (p>.05) between time or intensity. No significant differences in MCAv (p = .4), CVCi (p = .87), or CVRi (p = .87).

Northey et al, 2019	Small Randomized Controlled Trial	II	17 Female Breast Cancer Survivors	12-week intervention Cycled 20-30 min 3x/week 3 groups: - High intensity interval training (HIIT, 30 sec intervals at ~90% maximal heart rate or ~105% peak power with 2 min active recovery) - Moderate intensity continuous exercise (MOD, 55-65% peak power) - No-exercise control (CON)	Resting MCAv, Cerebrovascular Reactivity to CO_2	No significant differences in resting MCAv (p=.24) or cerebrovascular reactivity (p=.54) after HIIT compared to MOD. No significant differences in resting MCAv (p=.86) or cerebrovascular reactivity (p=.72) after HIIT compared to CON.
Tallon et al, 2019	Small Randomized Cross-over Trial	II	8 Prepubertal Children	2 Cycling conditions: - High intensity interval exercise (HIIE, 1 min interval at 90%max watt with 1 min active recovery at 20%max watt for 6 intervals) - Moderate-intensity steady-state exercise (MISS, 15 min at 44%max watt)	Exercise MCAv during each interval Immediate post-exercise MCAv 30-minutes post-exercise MCAv	Significant decrease in exercise MCAv during the 6th interval of HIIE compared to baseline (10.7%, p=.004). Significant decrease in exercise MCAv during the 3rd and 4th intervals of HIIE compared to MISS (p=.001). Significant decrease in MCAv immediately post-exercise following HIIE and MISS (p<.001). No significant difference in MCAv at 30-minutes post-exercise following HIIE and MISS compared to baseline (p>.05). Significant increase in exercise MCAv during the 2nd minute of MISS compared to baseline (5.8%, p=.004).

| Monroe et al, 2015 | Nonrandomized Cross-Over Trial | III | 15 Recreationally Active Adults | 2 cycling conditions:
- Sprint Interval Cycling (SIC, 30 second all-out sprint interval with 4 min active recovery for 4 intervals)
- Constant Resistance Cycling (CRC, 18 min at 70rpm with resistance set by matching total work performed during SIC) | Oxygenated Hemoglobin (HbO2)

Deoxygenated Hemoglobin (HHb) | Significant increase in average HbO2(effect size=.536, p=.001), minimum HbO2 during recovery (effect size=.392, p<.001), and maximum HbO2 during recovery (effect size=.588, p=.001) in SIC compared to CRC.

Significant increase in average HHb during SIC compared to CRC (effect size=.386, p=.003).

Significant increase in average HbO2 (effect size=.373, p=.001), minimum HbO2 during recovery (effect size=.223, p=.038), and maximum HbO2 during recovery (effect size=.333, p=.006) in men compared to women.

Significant increase in average HHb (effect size=.198, p=.034), minimum HHb during recovery (effect size=.264, p=.012), and maximum HHb during recovery (effect size=.358, p=.001) in men compared to women. |

MCAv = middle cerebral artery blood velocity, dCA = dynamic cerebral autoregulation, min = minute, HRmax = maximum heart rate, CVCi = cerebrovascular conductance index, CVRi = cerebrovascular resistance index, CO_2 = carbon dioxide

Table 2: Summary of Quality Review

	Avoided Contamination and Co-Intervention	Random Assignment to Conditions	Blinded Assessment	Monitored Intervention	Accounted for All Subjects	Reported Reliability of Measures Used	Reported Validity of Measures Used	Total Number of Criteria Met
Burma et al, 2020	No	Yes	No	Yes	Yes	Yes	Yes	5
Coetsee et al, 2017	No	Yes	No	Yes	Yes	No	No	3
Drapeau et al, 2019	No	Yes	No	Yes	Yes	No	Yes	4
Northey et al, 2019	No	Yes	No	Yes	Yes	No	No	3
Tallon et al, 2019	No	Yes	No	Yes	Yes	No	No	3
Monroe et al, 2015	No	No	No	Yes	Yes	Yes	No	3

Table 3: Summary of Effects of HIIE on Operationalized Cerebrovascular Measures

	Resting MCAv	Exercise MCAv	Post-Exercise MCAv	dCA phase	dCA Gain	dCA Coherence	De/Oxygenated Hemoglobin during Cognition	De/Oxygenated Hemoglobin during Exercise	Resting Cerebrovascular Conductance/ Resistance Index	Cerebrovascular Reactivity to CO$_2$
Burma et al, 2020		↓ to moderate		↓	no effect	no effect				
Coetsee et al, 2017							no effect			
Drapeau et al, 2019	no effect			↓	no effect	no effect			no effect	
Northey et al, 2019	no effect									no effect
Tallon et al, 2019		↓ to moderate and rest	↓							
Monroe et al, 2015								↑		

↓ = Decreased effect, ↑ = Increased effect, ⦸ = no effect