A SURPRISING FORMULA FOR THE SPREAD OF COVID-19 UNDER AGGRESSIVE MANAGEMENT

IVAN CHEREDNIK [†]

ABSTRACT. We propose an algebraic-type formula that describes with high accuracy the spread of Covid-19 pandemic under aggressive management for the periods of the intensive growth of the total number of infections. The formula can be used as a powerful forecasting tool. The parameters of the theory are the transmission rate, reflecting the viral fitness and "normal" frequency of contacts in the infected areas, and the intensity of prevention measures. The duration of the period of intensive growth is essentially inversely proportional to the square root of the intensity of hard measures. A more precise formula is based on Bessel functions. The data for the USA, UK, Sweden, Israel are provided.

1. Our approach and main findings. A formula describing almost perfectly the spread of Covid-19 during the periods of intensive growth is the main result. We model the effects of *momentum management of epidemics*, which can be defined as a system of measures of any nature aimed at reducing the epidemic spread by regulating the intensity of these measures on the basis of the latest absolute or average *total* numbers of infections. The usage of the absolute numbers combined with "hard measures", such as detection and prompt isolation of infected people and closing the places where the spread is the most likely, ensures a relatively fast saturation. This is not the end of the spread of the virus, but then the growth of the total number of infections becomes essentially linear and mild assuming that the measures are in place. According to the complete theory, the relatively short first period of exponential growth is followed by the period of essentially quadratic growth for *Covid-19*, which reasonably quickly becomes linear upon the aggressive protection measures, and then evolves toward the saturation. Our approach is of behavioral and sociological nature; the fact that it provides such a surprising accuracy seems a real discovery.

____

[†] Department of Mathematics, UNC Chapel Hill, NC, USA.
Email: chered@email.unc.edu .

Power law of epidemics. The simplest equation for the spread of communicable diseases results in exponential growth of the number of infections, which is mostly applicable to the initial stages of epidemics. See e.g. [1, 2, 3, 4] here and below and [5] about some perspectives with *Covid-19*. We focus on the middle stages, where the growth is no greater than some power functions in time, which requires a different approach and different equations.

The equally classical *logistic* models of the spread, as well as the SIR, SID models an generalizations, assume that the number of infections is comparable with the whole population, which we do not impose. Major new epidemics were not really of this kind during the last 100 years, which is obviously due to better disease control worldwide.

The reality now is the power-type growth of *total* number of infections $U(t)$ after a possible short period of exponential growth, *Covid-19* included. Our approach is based on this assumption. Generally, the rate of change of the total number of infections $dU(t)/dt$ is mostly related to $U(t) - U(t-p)$, where p is the period when the infected people spread the virus in the most intensive way. Assuming that $U(t) = t^\alpha$: $dU(t)/dt = \alpha t^{\alpha-1}$ and the leading term of $U(t) - U(t-p)$ is $p\alpha t^{\alpha-1} = p\lambda U(t)/t$, i.e. it is essentially proportional to $dU(t)/dt$. However, if $\alpha > 1$, there will be other terms in the expansion of $U(t) - U(t-p)$ and the proportionality with $U(t)/t$ can be only if either the virus transmission strength diminishes over time or we reduce our contacts over time *when the total number of cases growths faster than linearly*. With *Covid-19*, we attribute it mostly to the latter, to the reduction of the contacts over time of infected individuals with the rest of population, i.e. to *behavioral* and *sociological* factors. This is different from other power laws for infectious diseases; compare e.g. with [3].

Generally, if someone wants to "see" the *trend* of the epidemic using *only* the total number of infections to date, then $U(t)/t$ is the best way of course. This is qualitative, but the corresponding differential equation $dU(t)/dt = c\,U(t)/t$ immediately gives that $U(t) = Ct^c$ for some constant C, i.e. results in a power growth. This can be really seen for *Covid-19* and other epidemics before the active management begins. The coefficient c is approximately $c \approx 2$ for *Covid-19* initially; when the management begins, c drops and then the Bessel functions describe the increase of $U(t)$ till the "saturation", which is a technical end of the epidemic, though not its real end of course.

Some details. The full theory is presented in [6]. If the number of new infections is proportional to the current number of those infected, then the exponential growth of the spread is granted. By analogy with *news impact over time* from [7], and similar sociological-type processes, it is quite likely that the number of such contacts is proportional to

the current *total* number of the infected individuals to date *divided by the time to date*. The coefficient of proportionality is essentially the intensity of the spread. Sociologically, it is related to the intensity of the discussion of the epidemic by the authorities in charge and everywhere, which directly influences our understanding the gravity of the situation and results in the reduction of our contacts. This can happen even *before* any active prevention measures begin.

Starting with the equation $dU(t)/dt = c\,U(t)/t$ for the total number of infections $U(t)$, the coefficient of proportionality, the *basic transmission rate* c, is therefore a combination of the transmission strength of the virus and the "normal" frequency of the contacts in the infected place. There can be some other mechanisms for the "power law", including the biological ones. Self-isolation of infected species is common not only for humans ... unless for *rabies*; it can grow over time and when the intensity of the spread increases. Another possible mechanism can be due to the replication processes for viruses, but this is well beyond this article.

Anyway a sociological approach to the spread, which "explains" under some assumptions the power growth of the number of total cases, is quite natural in our work, because the active managements of epidemics is clearly of sociological nature, applicable only to humans. The coefficient c is one of the two main mathematical parameters of our theory. It can be practically seen as follows.

Before the prevention measures are implemented, approximately it reveals itself in the growth $\sim t^c$ of the total number of detected infections, where t is time. For *Covid-19*, c is around 2, which results in the quadratic growth of total number of infections after a short period of its exponential growth. Upon the active management of *Covid-19*, the growth quickly becomes $t^{c/2}$, i.e. essentially linear, which is part of our theory in [6]. This can be observed in many countries.

Ending epidemics. The "power law" is only a starting point of our analysis. The main problem is of course to "add" here some mechanisms ending epidemics and then prevent their possible recurrence. These are major challenges, biologically, psychologically, sociologically and mathematically. One can expect this *megaproblem* to be well beyond the power law itself, but we demonstrate that mathematically there is a path based on Bessel functions. The power growth alone obviously cannot lead to any saturation.

We are of course fully aware of the statistical nature of the problem, but the formula for the growth of the total number of infections we propose works almost with an accuracy of fundamental physics laws. This is very surprising for such stochastic processes as epidemics.

An important outcome of our modeling is that the measures of "hard type", like detecting and isolating infected people and closing the places where the spread is almost inevitable, are the key for ending an epidemic. Moreover, such measures must be employed strictly proportionally to the current numbers of infections, not to its derivative of any kind, which is the most aggressive "momentum" way to react to them, the "hardest" possible as we will explain now.

For instance, if the number of infections doubled during some period, then the *operation formula* from [6] requires that *testing* must be increased 4–fold. Assuming that we approach the saturation, where the number of new infections becomes almost zero, the "hard way" is to keep *testing* at the same level as before, i.e. performing the same number of tests every day. This is in spite of almost zero number of new infections. Practically, this is not always the case.

If we react to the *average* number instead of the *absolute* number of infections, then we are supposed to stop testing and other measures when there are no new "cases". This is the "soft way". If this is coupled with "soft" measures (see below), then mathematically the epidemic will never reach the saturation point.

The u–formula. The *intensity* of "hard" measures will be denoted by a in this article. Upon composing the corresponding differential equations and integrating them we obtain the following. The function $u(t) = C \, t^{c/2} J_{c/2-1/2}(\sqrt{a}t)$ for the Bessel function $J_\alpha(x)$ of the first kind models the growth *total* number of infections, where C is the scaling parameter necessary to adjust this formula to the actual numbers. Here and below time t is normalized *days*/10 for the number of days from the beginning of the period of the intense growth of the epidemic. This functions matches the total number of infections with high accuracy.

The parameter a is 0.2 for the USA and UK; it can be about 0.3 for countries with somewhat more proactive approach, like Austria and Israel, and it becomes 0.15 and even 0.1 for "the world" and Sweden. The basic transmission rates are: $c = 2.2$ for the USA, $c = 2.4$ for UK, Sweden, and it is currently 2.8 for "the world". Recall that c approximately corresponds to the growth $\sim t^c$ of total infections in the beginning of the epidemic, when no protection measures are in place.

Limitations. Importantly, the function $u(t)$ is proportional to $t^{c/2+1}$ for t near $t = 0$, which is very close to $\sim t^c$, assuming that $c \approx 2$. So the phase of "parabolic growth" is generally covered well by the u–formula too, though formally this does not follow from the theory in [6].

Of course there can be other reasons for $u(t)$ to "serve epidemics"; it is not impossible that there are connections to the replication process of viruses, but this we do not touch upon. As always, there are limitations, which we will address now.

First and foremost, the available infection numbers are for the *detected* cases, which are mostly *symptomatic*. However this is not of much concern for us. We understand managing epidemics sociologically. Those who are detected mostly have symptoms, but the management is mostly focused on them too. So all our arguments fully applicable within this group; we just restrict ourselves with symptomatic cases. No assumptions on asymptomatic cases are necessary for the u–formula. We of course understand that when the number of new *reported* infections drops to zero, there can be many non-detected asymptomatic cases, which can potentially lead to the recurrence of the epidemic. Such "saturation" is only a *technical* end of the epidemic.

The second reservation concerns *newly emerged clusters of infections* and the countries where the spread is on the rise. The u–formula can be used in spite of such fluctuations, but it is now a statistical tool; see Figure 5. The predictions must be regularly updated.

The third reservation is related to the management of the epidemic. Not all countries employ the protection measures in similar ways, but this is not a problem for us. The problem is if the intensity of the measures and the criteria are changed in some unpredictable ways. Diminishing the "hard" measures too early or even dropping them completely at later stages is quite possible.

Last but not the least is the *data quality*. Changing the ways they are collected and the criteria frequently makes such data useless for us. Though, if the number of detected infections is underreported in some regular ways, whatever the reasons, such data can be generally used. Thus this is sufficiently relaxed, but the data for several countries, not too many, are not suitable for the usage of our "forecasting tool".

Forecasting the spread. With these reservations, the first point of maximum t_{top} of $u(t)$ is a good estimate for the duration of the epidemic and the corresponding estimate for the top value of the total number of infections. This gives an important "forecasting tool", assuming that "hard measures" are the key in practical management of the epidemic.

We note that the approximate reflection symmetry of $du(t)/dt$ for the $u(t)$ in the range from $t = 0$ to t_{top} can be interpreted as *Farr's law of epidemics under aggressive management*. Generally, the portions of the corresponding graph before and after the *turning point* are supposed to be essentially symmetric to each other. This is not exactly true for $du(t)/dt$, but close enough. See Figure 2 and the other ones; the turning point is at $\max\{du(t)/dt\}$.

As any model, our one is based on various simplifications. We assume that the number of people perceptive to the virus is unlimited, i.e. we do not consider epidemics with the number infections comparable with the whole population, as well as *herd immunity*. Also, we disregard

the average duration of the disease and that for the quarantine periods imposed. The total number of infections regardless of the output is what we are going to model, which is commonly used. In spite of all these assumptions, the u–formula works unexpectedly well.

We mention here a strong connections with *behavioral finance*; see [7]. For instance, practically the same $u(t)$ in terms of Bessel function as above models "profit taking" in stock markets. The polynomial growth of $u(t)$ is parallel to the "power law" for share prices.

2. Two kinds of management. There is a long history and many aspects of mathematical modeling epidemic spread; see e.g. [1] for a review. We restrict ourselves only with the dynamic of *momentum managing epidemics*, naturally mostly focusing on the middle stages, when our actions must be as precise as possible. The two basic modes we consider are essentially as follows:

(A) aggressive enforcement of the measures of immediate impact reacting to the current *absolute* numbers of infections and equally aggressive reduction of these measures when these numbers decrease;

(B) a more balanced and more defensive approach when mathematically we react to the *average* numbers of infections to date and the employed measures are of more indirect and palliative nature.

Hard and soft measures. The main examples of (A)–type measures are: prompt detection and isolation of infected people and those of high risk to be infected, and closing places where the spread is the most likely. Actually the primary measure here is *testing*; the number of tests is what we can really implement and control. The *detection* of infected people is its main purpose, but the number of tests is obviously not directly related to the number of detections, i.e. to the number of *positive tests*. The efficiency of testing requires solid priorities, focus on the groups with main risks, and solving quite a few problems.

Even simple mentioning problems with testing, detection and isolation is well beyond our article. However, numerically we can use the following. During the epidemics, essentially during the stages of linear growth, which are the key for us, the number of positive tests can be mostly assumed a stable fraction of the total number of tests. This is demonstrated in Figure 1. Such proportionality can be seen approximately from March 16. However, at the end of this chart, testing was reduced below the levels required for mode (A); it must remain at least constant until the saturation of the number of total cases.

The measures like wearing protective masks, social distancing, recommended self-isolation, restricting the size of events are typical for (B). This distinction heavily impact the differential equations we obtained in [6]. However the main difference between the modes, (A) vs.

FIGURE 1. Daily testing vs. new daily cases for USA

(B), is the way the number of infections is treated: the *absolute* number of infections is the trigger for (A), whereas the *average* number of infections to date is what we monitor under (B).

Hard measures are the key. Generally, the (A)–type approach provides the fastest possible and "hard" response to the changes with the number of infections, whereas we somewhat postpone with our actions until the averages reach proper levels under (B), and the measures we implement are "softer". Mathematically, the latter way is better protected against stochastic fluctuations, but (B) is slower and cannot alone lead to the termination of the epidemic, which we justify mathematically within our approach.

The main objective of any managing epidemics is to quickly end them. However the excessive usage of hard measures can lead to the recurrence of the epidemic, some kind of "cost" of our aggressive interference in a natural process. This can be avoided only if we continue to stick to the prevention measures as much as possible even when the number of new infections goes down significantly. Reducing them too much on the first signs of improvement is a way to the recurrence of the epidemic, which we see mathematically within our approach.

Some biological aspects. The *viral fitness* is an obvious component of the transmission rate c. Its diminishing over time can be expected, but this is involved. This can happen because of the virus replication errors. The *RNA* viruses, *Covid-19* included, replicate with fidelity that is close to error catastrophe. See e.g. [8] for some review and predictions. Such matters are well beyond this paper, but one biological aspect must be mentioned, concerning the *asymptomatic cases*.

The viruses mutate at very high rates. They can "soften" over time to better coexist with the hosts, though fast and efficient spread is of course the "prime objective" of any virus. Such softening can result in an increase of asymptomatic cases, difficult to detect. So this can

contribute to diminishing c we observe, though this is not because of the decrease of the spread of the disease. We model the available (posted) numbers of total infections, which mostly reflect the symptomatic cases.

To summarize, it is not impossible that the replication errors and "softening the virus" may result in diminishing c at later stages of the epidemic, but we think that the reduction of the contacts of infected people with the others dominates here, which is directly linked to behavioral science, sociology and psychology.

3. The formula for the growth.. We will need the definition of the *Bessel functions* of the first kind:

$$J_\alpha(x) = \sum_{m=0}^{\infty} \frac{(-1)^m (x/2)^{2m+\alpha}}{m! \Gamma(m+\alpha+1)}.$$

See [9] (Ch.3, S 3.1). Practically, use $\mathtt{BesselJ}[\alpha, \mathtt{x}]$ in Mathematica.

The key point is that measures of type (A) have "ramified" consequences, in some contrast to mode (B). Namely, an isolated infected individual will *not* transmit the virus to many people and the number of those protected due to this isolation grows over time. Combining this with our understanding of the power law of epidemics, we arrive at the differential equation for the total number of infections $u(t)$:

$$\frac{du(t)}{dt} = c\,\frac{u(t)}{t} - \frac{1}{a}\int_0^t u(s)ds,$$

which can be solved in terms of Bessel functions. The dominant solution is $u(t) = Ct^{c/2}J_{c/2-1/2}(\sqrt{a}t)$ for some constant C.

There is a surprisingly perfect match of the *total* number of infections for *Covid-19* in the USA and UK till April 14 with our solutions $u(t)$ above. This is from the moment when these numbers begin to grow "significantly", approximately around March 16 for the USA and UK. Epidemics are very stochastic processes, so such a precision is surprising. The site https://ourworldindata.org/coronavirus is mostly used for data, updated at 11:30 London time. *We take $x = days/10$.*

The USA data. The scaling coefficient 1.7 in Figure 2 is adjusted to match the real numbers. For the USA, we set $y = \text{infections}/100K$, and take March 17 the beginning of the period of "significant growth". The parameters are $c = 2.2$, $a = 0.2$.

The *red dots* show the corresponding actual *total* numbers of infections. They *perfectly* match $u(t) = 1.7t^{1.1}J_{0.6}(t\sqrt{0.2})$ in Figure 2, which results in the following: the number of cases in the USA can be expected to reach its preliminary saturation at $t_{top} = 4.85$ (48.5 days

from 03/17 to May 5) with $u_{top} = 10.3482$, i.e. with 1034820 infections (it was 609516 at 04/15). This is of course a *projection*: about $1M$ of total cases at the saturation point t_{top} near May 5.

The *black dots* show the test period, till April 29. The predictions are of course based on the assumption that the intensity of hard measures continues to be proportional to the total number of detected infections to date, as it was clearly the case for the *red dots*. The jumps like the one at about $x = 4.1$ can be because of various reasons impossible to forecast, as well as some period of linear growth after it. However the general trend for black dots matches our $u(t)$ well enough.

FIGURE 2. $u(t) = 1.7\, t^{c/2} J_{(c-1)/2}(\sqrt{a}t)$ for $c = 2.2, a = 0.2$

Obviously $t = t_{top}$ cannot be the end of the epidemic. The data from South Korea and those from other countries that went through the "saturation", demonstrate that *a linear growth of the total number of cases can be expected around and after t_{top}, with periodic fluctuations.* The obvious reasons are *(a)* reducing the "hard" and "soft" measures, *(b)* no country is isolated from infections from other places, *(c)* continued testing can result in finding more asymptomatic cases, *(d)* new clusters of disease are always possible. Anyway, fluctuations closer to the "saturation" and after it are quite likely. Resuming the corresponding measures is possible, but this is not always the case. Our formula is just a forecasting tool, which is mostly applicable to the period when the protection measures, especially "hard" ones, are applied in a regular manner. Then the match with the real data is good.

Covid-19 in UK. One of the reasons of the strong match of *red dots* and our $u(t)$ can be that the USA consists of 50 states, and therefore the total number of infections is quite an average. However this is

no worse for UK. The data will be from 03/16 till 04/15; add 18 to
our "red dots", the initial number at 03/16, to match the actual total
numbers. The *black dots* constitute the control period: 04/16-04/29.

FIGURE 3. $u(t) = 2.2\,t^{c/2}J_{(c-1)/2}(\sqrt{a}t)$ for $c = 2.4, a = 0.2$

Now $c = 2.4, a = 0.2$ work fine, and the scaling coefficient is 2.2. The
total number of cases is divided by $10K$, not by $100K$ as for the USA.
The expectation is now as follows: the "saturation moment" can be
5.17, i.e about 51 days after March 16, somewhere around May 6. The
estimate for corresponding number of infections is about 170000, with
all ifs. This is assuming that the "hard" measures will be employed at
the same pace as before April 15. The black dots confirm the trend.

Sweden 03/07-04/23. This is an example of the country that remains
essentially "open". Actually, they actively do testing followed by the
isolation of infected people, the key "hard" measure from our perspec-
tive. Also, the strength of the health-care in this country must be taken
into consideration, and that it is surrounded by the countries that fight
Covid-17 aggressively. The growth of the total number of cases was
essentially quadratic for a relatively long period, which is what "power
law" states for the epidemics with minimal "intervention". By now,
the growth is linear, which means that what the measures they use
appeared working. Interestingly, our $u-$ formula is applicable, but for
record law $a = 0.1$; it is $a = 0.15$ for "the world". Here y is the to-
tal number of cases divided by 1000 and add 137, the initial value, to
our "dots". The projected saturation is around May 17. As always,
it is with a reservation about their future policies, which seem suffi-
ciently stable by now. In spite of their "soft" approach, our $u-$formula
obviously can be used; see Figure 4.

Sweden: 03/07-04/23
top=7.32 (at May 19)
u(top)=23.5612 (23.5K)
(a=0.1; c=2.4; C=1.7)

Black dots:
control period
04/24-04/29

du(t)/ dt

FIGURE 4. $u(t) = 1.7\, t^{c/2} J_{(c-1)/2}(\sqrt{a}t)$ for $c = 2.4, a = 0.1$

Israel: "saturation". The last example we provide is what can be expected when the country went through the "saturation". Israeli population is diverse, which has a potential of significant fluctuations of the number of cases and various clusters of infection. However its solid response to *Covid-19* and good overall health-system, made the growth of the spread sufficiently predictable. Generally, for small countries the fluctuations can be expected higher than for countries like the USA, UK, though it depends on how "homogeneous" they are. We divide the total number of cases by 1000, as for Sweden; see Figure 5.

The *red dots* began March 13, when the *total* number of detected infections was 96, and stopped April 17; the remaining period till April 29, shown by the black dots, was the "control one". The saturation forecast went through almost perfectly, but there were significant fluctuations in process. After April 26, the predicted moment of the saturation, the growth of the total number of (known) infections is supposed to be mild linear. The parameters are: $a = 0.3$, i.e. the intensity of hard measures is better than with the USA, UK, and $c = 2.6$. The latter means that the initial transmission coefficient was somewhat worse than $c = 2.4$ in the USA, UK, possibly due to the greater number of "normal" contacts. Recall that a, c are parameters of our theory, related to but not immediately connected with the real factors.

Some discussion. Recall that we consider only "total cases", the numbers of all detected infections, and begin at the moment when the "significant" growth begins, which is also essentially when the active measures start. According to our theory, the match with $u(t)$ above can be expected in some interval around the turning point. However

FIGURE 5. $u(t) = 2.2\, t^{c/2} J_{(c-1)/2}(\sqrt{a}t)$ for $c = 2.6, a = 0.3$

practically it holds better than this: an almost perfect match is for about 80% of the whole periods of intensive growth.

Our restriction to the period till April 15, 2020 with fixing the parameters a, c, except for Sweden, is not accidental. It was sufficient for the practical confirmations of the theory presented in Figures 2,3, and with other countries. The latest data, *the black dots*, provide the "real-time checks"; they were obtained *after* the parameters were fixed. The pandemic is far from over, but within the scope of this paper, the data we needed for the "forecasting tool" were essentially present by 04/15.

When the epidemic approaches the saturation, which is the first maximum of $u(t)$ in our model, its management can be expected to evolve toward reducing and abandoning "hard" measure. This can result in a growth like $Ct^{c/2}$ with some "mild" C and $c/2 \approx 1$ and significant fluctuations, which really happens. Such growth at the end is generally covered by our theory; it is not connected with Bessel functions.

The superb match of our $u(t)$ during the significant part of the period of intensive growth of the spread can be of real importance for practical managing epidemics. On the basis of what we see, the best ways to use the u–curves seem as follows: *(1):* determine a, c using about the first 30-40% after the intensive growth begin, *(2):* update them constantly till the turning point and somewhat beyond, *(3):* try to the adjust the measures at later stages "to stay close to the curve". With *(3)*, the constant response is needed to new clusters of the disease, some jumps with the new cases due to the reductions of the measures and so on. Generally, our forecasting tool can serve the best if the data and the measures are as uniform and "stable" as possible. Then underreporting, focusing on symptomatic cases, and inevitable fluctuations with

the data may not influence too much the match with the u–curves; this is of course statistically and with usual reservations. Also, *(4):* testing the population and some "soft" measure must be continued well after the "saturation" to prevent the recurrence of the epidemic.

Acknowledgements. I'd like to thank ETH-ITS for outstanding hospitality. My special thanks are to Giovanni Felder, Rahul Pandharipande. I also thank very much David Kazhdan for his valuable comments and suggestions. Funding: partially supported by NSF grant DMS–1901796 and the Simons Foundation.

References.

[1] H. Hethcote, *The mathematics of infectious diseases*, SIAM Review, 42:4. (2000), 599–653. 2, 6

[2] H. Hethcote, and S. Levin, *Periodicity in Epidemiological Models*, In: Applied Mathematical Ecology. Biomathematics, 18, 193–211, Springer, Berlin, Heidelberg, S. Levin, T. Hallam, L. Gross (eds), 1989. 2

[3] S. Meyer, and L. Held, *Power-Law models for infectious disease spread*, The Annals of Applied Statistics 8:3 (2014), 1612–1639. 2

[4] Ph. Strong, *Epidemic psychology: a model*, Sociology of Health & Illness 12:3 (1990), 249–259. 2

[5] S. Cobey, *Modeling infectious disease dynamics*, Science, 24 Apr 2020; DOI: 10.1126/science.abb5659. 2

[6] I. Cherednik, *Momentum managing epidemic spread and Bessel functions*, Preprint: arxiv 2004.06021v3 (q-bio), 2020. 2, 3, 4, 6

[7] I. Cherednik, *Artificial intelligence approach to momentum risk-taking*, Preprint: arxiv 1911.08448v4 (q-fin), 2019. 2, 6

[8] R. Carrasco-Hernandez, and R. Jácome, and Y. López Vidal, and S. Ponce de León, *Are RNA viruses candidate agents for the next global pandemic? A review*, ILAR Journal, 58:3 (2017), 343–358. 7

[9] G.N. Watson, *A Treatise on the Theory of Bessel Functions*, 2nd Edition, Cambridge University Press, Cambridge, 1944. 8