The impact of social distancing on COVID19 spread: State of Georgia

case study

Pinar Keskinocak[1,2¶ *], Buse Eylul Oruc[1¶], Arden Baxter[1¶], John Asplund[1,3¶], Nicoleta

Serban[1¶]

[1] H. Milton Stewart School of Industrial and Systems Engineering, Georgia Institute of

Technology, Atlanta, Georgia, United States

[2] Department of Environmental Health, Rollins School of Public Health, Emory

University, Atlanta, Georgia, United States

[3] Metron Inc., Reston, VA, United States

* Corresponding Author

Email: pinar@isye.gatech.edu (PK)

[¶] These authors contributed equally to this work.

Abstract

As the spread of COVID19 in the US continues to grow, local and state officials face difficult decisions about when and how to transition to a "new normal." The goal of this study is to project the number of COVID19 infections and resulting severe outcomes, and the need for hospital capacity under social distancing, particularly, shelter-in-place and voluntary quarantine for the State of Georgia. We developed an agent-based simulation model to project the infection spread. The model utilizes COVID19-specific parameters and data from Georgia on population interactions and demographics. The simulation study covered a seven and a half-month period, testing different social distancing scenarios, including baselines (no-intervention or school closure only) and combinations of shelter-in-place and voluntary quarantine with different timelines and compliance levels. The following outcomes are compared at the state and community levels: the number and percentage of cumulative and daily new symptomatic and asymptomatic infections, hospitalizations, and deaths; COVID19-related demand for hospital beds, ICU beds, and ventilators. The results suggest that shelter-in-place followed by voluntary quarantine reduced peak infections from approximately 180K under no intervention and 113K under school closure, respectively, to below 53K, and delayed the peak from April to July or later. Increasing shelter-in-place duration from four to five weeks yielded 2-9% and 3-11% decrease in cumulative infection and deaths, respectively. Regardless of the shelter-in-place duration, increasing voluntary quarantine compliance decreased daily new infections from almost 53K to 25K, and decreased cumulative infections by about 50%. The cumulative number of deaths ranged from 6,660 to 19,430 under different scenarios. Peak infection date varied

38   across scenarios and counties; on average, increasing shelter-in-place duration delayed

39   the peak day by 6 days. Overall, shelter-in-place followed by voluntary quarantine

40   substantially reduced COVID19 infections, healthcare resource needs, and severe

41   outcomes.

Introduction

The novel coronavirus SARS-CoV-2 causes a rapidly spreading respiratory illness, Coronavirus Disease 2019 (COVID19), which has become a pandemic [1]. During the early stages of a pandemic, medical interventions, such as vaccines or antiviral treatments, are either non-existent or extremely limited [2]. Hence, local, national, and global governments and public officials wrestle with the difficult decisions of how, when, and where to implement non-medical interventions [3]. The decision-makers also need to understand how the type and duration of interventions, as well as the public's compliance levels, impact their effectiveness [4].

In this study, we developed an agent-based simulation model to predict the spread of COVID19 geographically and over time. The model captures both the natural history of the disease and interactions in households, workplaces, schools, and communities [5-9]. The model was populated with COVID19 parameters from the literature and population-related data from the State of Georgia, including demographic information, household sizes, and travel patterns, and validated using data regarding COVID19 confirmed infections and deaths in Georgia. The model's outputs include new daily infections (symptomatic and asymptomatic by age group), hospitalizations, and deaths at the census tract level.

We utilized the model to evaluate the effectiveness and impact of non-medical social-distancing interventions, including school closure, shelter-in-place (SIP), and voluntary quarantine (VQ) [6, 10-16]. We tested various scenarios with different durations and time-varying compliance levels for interventions to inform decision-makers about

potential social distancing recommendations to be shared with the public. We also

developed a hospital resource estimation decision-support tool, which takes as input the

model's daily COVID19-related hospitalization estimates, and predicts the number of

hospital beds, ICU beds, and ventilators needed geographically (at the county level) and

over time. We then aggregated these estimates across the fourteen coordinating

hospital regions in Georgia, to provide insights about potential capacity shortages in the

healthcare system [17].

Methods

Study population

73 Population in Georgia stratified by age groups 0-4, 5-9, 10-19, 20-64, 65+. In Georgia

74 there are 1,336,490, 1,418,910, 6,685,870, and 1,356,730 people in age groups 0-9,

75 10-19, 20-64, and 65 or over, respectively, with a total population of approximately

76 10,519,000 [18, 19].

Infection projection model

We adapted an agent-based simulation model with heterogeneous population mixing to

predict the spread of the disease geographically during the study period of February 18,

2020 to September 30, 2020 [5-8]. The model captures the natural history of the

disease at the individual level, by age group, as well as the infection spread via a

contact network consisting of interactions in households, peer groups (workplaces,

schools), and communities, with different rates of transmission [13, 20-25].

The model was populated with COVID19-specific parameters [20-22, 26-40] and data

from Georgia, including household type [18], household size [18], children status [18,

41], workflow [42], and population demographics [18] at the census tract level. To seed

the model, we utilized the confirmed infection data for Georgia, at the county level [43].

The main assumptions in the model were (1) every individual is in one of the following

states at any given time (see Fig 1): susceptible (S), exposed (E), transition (IP),

asymptomatic (IA), symptomatic (IS), hospitalized (H), recovered (R), or dead (D) and

(2) three levels of mixing in the population: (i) community (day and night), (ii) peer

groups (day), and (iii) household (night). During the exposed state, an infected

individual shows no symptoms and does not infect others. During the transition state an

infected individual shows no symptoms but could infect others. From the transition state,

an infected individual moves to the symptomatic or the asymptomatic state.

Fig 1. Model descriptions. Agent-based model incorporates the natural history of the

disease for each individual agent, by age group, and the interactions at the household,

peer group, and community, across different geographic areas. Outcome measures

reported are averages of 30 replications ran for each scenario.

S1 Appendix provides additional details on the model implementation. Table 1 provides

the input model parameters, and Fig A in S1 Appendix provides model validation using

COVID19 confirmed cases and deaths reported in Georgia.

Table 1. Model parameters. Descriptions and references for the model input

parameters.

Parameters	Estimates	References
Probability of Symptomatic	0.63	[26, 27, 38-40] [44]
Probability of Hospitalization	0.016 for age 0-19, 0.18 for age 20-64, 0.30 for age 65+	[21]
Probability of Death	0 for age 0-19, 0.103 for age 20-64, 0.375 for age 65+	[21, 44]
Reproductive Number (R_0)	2.4	[28-30]
Transmission Rate (β)	1.02	[29]
Exposed Duration	Weibull with mean 4.6 days	[20, 26, 32, 45]
Transition Duration	0.5 days	[20]
Hospitalized Duration	Exponential with mean 7 days	[20, 34, 44]
Symptomatic Duration	Exponential with mean 2.9 days	[22]
Symptomatic-Asymptomatic Duration Ratio	1.5	[20]
Proportion of Transmission that Occurs at the Transition or Asymptomatic Stage (θ)	0.48	[35]
Proportion of Infections Generated by those who are Asymptomatic (ω)	0.24	[35]
Percentage of hospitalizations that require ICU	0% for age 0-19, 29.9% for age 20-64, 35.1% for age 65+	[21, 44]
Percentage of ICU patients that require ventilation	64%	[46]
ICU Duration (days)	8	[36]
Ventilation Duration (days)	6	[47]

Intervention analysis

The following baseline scenarios and social distancing interventions are analyzed in our study:

1. _No intervention (NI)_ – the population interacts with each other normally;

2. _School Closure (SC)_ – no peer group interactions among children (i.e., no K-12 school interactions);

3. _Voluntary Quarantine (VQ)_ – All household members stay home if a household member experiences symptoms, until the entire household is symptom-free.

4. _Shelter-in-Place (SIP)_ – Household members stay home complying with a state order.

5. _Voluntary Shelter-in-Place (VSIP)_ – Household members choose to follow SIP voluntarily.

Household members complying with SIP, VSIP, or VQ do not engage in peer group or community interactions. Compliance levels (<100%) under SIP, VSIP, and VQ probabilistically determine individual compliance and corresponding community interactions.

NI and SC were considered as _baselines_ for comparison. In Scenarios 1-9, SIP durations (4, 5, and 6 weeks) and gradually decreasing post-SIP VQ compliance levels (low, medium, high) were tested (Fig 2_); s_helter-in-place was in effect for 4 weeks (April 3-April 30) in Georgia [48]. Because all K-12 schools in Georgia were closed starting

March 16, 2020 until the end of July, Scenarios 1-9 assumed school closure.

Additionally, after the end of SIP in Scenarios 1-9, decreasing compliance with voluntary

shelter-in-place was considered, chosen to be in line with social mobility indicators [49].

Further details on the choice of VSIP compliance levels can be found in S1 Appendix.

Fig 2. Intervention scenarios. Description of the intervention scenarios considered in

this study.

Healthcare resource needs projection model

The hospitalization output from the simulation model was used to estimate the daily

demand for hospital beds (general inpatient beds and intensive care unit (ICU) beds)

and ventilators for COVID19 patients. Daily hospital bed demand was calculated by

aggregating the number of hospital beds needed in the previous day with the number of

new hospitalizations, minus the proportion of the population that was discharged from

the hospital based on the average hospitalization duration. ICU bed and ventilator

demand were estimated using the percentage of hospitalized patients that require ICU

(by age), average ICU duration, percentage of ICU patients that require ventilation, and

the average ventilation duration.

We derived county-level risk factors (see Fig B in S1 Appendix) by applying the principal

component analysis [50] on several factors known to impact a higher risk of

complications and severe outcomes for COVID19 infections, including prevalence of

asthma, diabetes, obesity, smoking, cardiovascular disease and chronic conditions [51].

We then adjusted the estimated demand using these risk factors.

S1 Appendix and Table 1 provide additional details on the estimation approach and the

input model parameters along with their references.

Outcome measures

The outcome measures considered for the study period include:

- *New Infection Count (NIC):* number of daily new symptomatic and asymptomatic

infections.

- *Infection attack rate (IAR):* cumulative percentage of the population infected.

- *Peak infection (PI):* maximum percentage of the population infected on a given

day.

• *Peak day*: The day when NIC is highest.

• *Clinical attack rate (CAR):* percentage of symptomatic infections among the total

population.

• *Hospital Bed Demand (HB):* number of hospital beds (general inpatient and ICU

beds) needed due to severe outcomes among the infected.

• *Intensive Care Unit Bed Demand (ICUB):* number of ICU beds needed due to

severe outcomes among the infected.

• *Ventilator Demand (V):* number of ventilators needed due to severe outcomes

among the infected.

169 Results

170 State-level outcome measure analysis

Fig 3 shows the NIC outcome for all scenarios. Table 2 includes summaries across all

scenarios and outcomes. Fig A in S1 Appendix and S1, S2 Figs provide state-level

outcomes for Scenarios 1-9.

Fig 3. State level outcomes: NIC under all scenarios; 4-week, 5-week and 6-week

SIP scenarios. NIC under all scenarios including baseline (top left plot), NIC under 4-

week SIP followed by Low (Scenario 1), Medium (Scenario 2), High (Scenario 3) VQ

compliance (top right plot), NIC under 5-week SIP followed by Low (Scenario 4),

Medium (Scenario 5), High (Scenario 6) VQ compliance (bottom left plot), NIC under 6-

week SIP followed by Low (Scenario 7), Medium (Scenario 8), High (Scenario 9) VQ

compliance (bottom right plot).

Table 2. Statewide outcome measures.

	Peak Infection (%)	Peak Day	IAR %	CAR %	Deaths	Cumulative Hospitalizations	Infections	Peak HB	ICUB	V
NI	1.67	18-Apr	59.15	37.26	30646	193089	6387094	33979	11356	5702
SC	1.05	26-Apr	50.31	31.70	27356	173196	5432630	23638	7995	3970
1	0.49	28-Jul	37.81	23.74	19431	127589	4082885	11166	3804	1861
2	0.38	22-Aug	31.47	19.60	15209	104050	3398305	8753	2993	1454
3	0.25	28-Sep	19.45	11.95	8509	61535	2100760	5638	1922	940
4	0.48	5-Aug	37.03	23.20	18850	124248	3998923	11197	3823	1861
5	0.37	2-Sep	29.62	18.36	13566	95984	3197887	8728	2980	1452
6	0.25	29-Sep	17.71	10.85	7734	55673	1912008	5484	1858	914
7	0.47	15-Aug	35.65	22.27	17481	118664	3849091	10911	3717	1816
8	0.37	8-Sep	27.45	16.93	12205	88083	2963912	8720	2984	1448

| 9 | 0.23 | 29-Sep | 15.36 | 9.40 | 6662 | 47910 | 1658130 | 4659 | 1573 | 781 |

Statistical summaries that compare baseline and intervention scenarios with respect to

Peak Infection (%), Peak Day, IAR (%), CAR (%), Cumulative Deaths, Cumulative

Hospitalization, Cumulative Infections, Peak HB, Peak ICUB, and Peak V.

The maximum NIC was approximately 180K (April 18th) and 113K (April 26th) under NI

and SC, respectively. Under Scenarios 1-9, the maximum NIC was below 53K, with the

earliest peak in July. Compared to NI and SC, in Scenarios 1-9 NIC was at least 36%

and 25% lower, and approximately 2.3 million and 1.4 million fewer people infected,

respectively. Extending SIP by 1-2 weeks or following SIP by high VQ further reduced

NIC and delayed the peak day. Similar trends were observed for CAR, with the number

of symptomatic infections reducing by more than 36% and 25%, compared to NI and

SC, respectively.

Increasing the SIP duration from four to five weeks (i.e., extending by one week) yielded

a decrease of approximately 2-9% in IAR and CAR, approximately 3-10% in cumulative

hospitalizations, and 3-11% in cumulative deaths. Increasing the SIP duration from five

to six weeks (i.e., extending by two weeks) yielded a decrease of approximately 4-13%

in IAR, 4-14% in cumulative hospitalizations, and 7-14% in cumulative deaths.

Higher VQ compliance after SIP, regardless of the SIP duration, decreased the peak

NIC from approximately 53K to 25K and decreased IAR by up to 57%.

SC had a lower IAR outcome than NI, but led to similar severe outcomes (cumulative

deaths and hospitalizations). Healthcare resource demand estimates (HB, ICUB, and

V) were about 30% lower under SC versus NI. In the absence of social distancing

interventions, approximately 30,640 and 27,350 people in Georgia were projected to

die, and approximately 193,080 and 173,190 people were projected to be hospitalized

under NI and SC, respectively.

The total number of deaths for Scenarios 1-9 ranged from 6,662 (Scenario 9) to 19,431

(Scenario 1); the number hospitalized was between 47,910 (Scenario 9) and 127,589

(Scenario 1); peak hospital bed needs ranged from 4,659 (Scenario 9) to 11,166

(Scenario 1). Peak ICU bed and ventilator needs ranged from 1,573 (Scenario 9) to

3,804 (Scenario 1) and from 781 (Scenario 9) to 1,861 (Scenario 1), respectively.

Scenario 9, where SIP extended until mid-May followed by high VQ, provided the lowest

cumulative infections, hospitalizations, and deaths.

Infection spread outcome measure analysis by county

Tables A and B in S1 Appendix include the estimated peak day and percentage by

county across all scenarios. S3 Fig includes the maps of the NIC by county for different

dates.

Peak day varied across scenarios and counties; however, on average, increasing SIP

duration by one week moved the peak day by 6 days across counties. Each week of

SIP extension delayed the peak day by an average of 6-7 days in the most populated

counties (e.g., Fulton, Gwinnett, Cobb, DeKalb, Chatham) and 3-6 days in smaller

counties (e.g., Glascock, Clay, Webster, Quitman, Taliaferro), respectively. Increasing

VQ compliance from low to medium and from medium to high delayed the peak day by

an average of 24 and 19 days, respectively. In most scenarios, rural counties peaked on

average 5 days later compared to urban counties. Differences in the peak day observed

in rural and urban counties were mostly consistent across the scenarios.

PI fluctuated depending on SIP duration. Rural counties and urban counties did not

differ much in terms of PI. See S1 Appendix for examples of an analysis of urban and

rural counties in Georgia.

The NIC was highest in the densely populated Fulton county and other surrounding

counties in the Atlanta metropolitan area across all scenarios.

246 Healthcare resource needs analysis by coordinating hospital

247 region

Fig 4 presents the healthcare resource peak demand under Scenario 2, by hospital

region. (A map of the 14 coordination hospital regions of Georgia can be found in [52].)

Fig 5 shows the hospital and ICU bed needs over time for region N under Scenario 2.

Similar patterns were observed for other scenarios.

Fig 4. COVID19-related peak healthcare resource demand for 4-week SIP, Medium

VQ compliance post-SIP. COVID19-related hospital bed (top left plot), ICU bed (top

right plot), and ventilator (bottom center plot) peak needs for 4-week SIP, Medium VQ

compliance post-SIP (Scenario 2).

Fig 5. COVID19-related healthcare resource demand over time for region N for 4-

week SIP, Medium VQ compliance post-SIP. COVID19-related hospital bed (left plot)

and ICU bed (right plot) needs over time for region N under 4-week SIP, Medium VQ

compliance post-SIP (Scenario 2).

The highest need for COVID19-related healthcare resources was in region D, with

3,117 hospital beds, 1,058 ICU beds, and 516 ventilators. Region D includes three of

the top four populous counties in Georgia. The four regions with the highest need

across all outcomes were regions D, F, J, and N, which are regions that include

populous counties.

There is a gap between HB and ICUB demand and availability around the peak day in

many regions. For example, consider region N under Scenario 2. The peak hospital bed

need for COVID19 patients is projected to occur around September 2nd with a demand

of 1,164 beds, yet the estimated staffed hospital bed availability for all (including non-

COVID19) patients is 1,607 [53]. Peak ICU bed demand for COVID19 patients was

projected as 396, also occurring around September 2nd whereas estimated availability

for all patients is 134.

Discussion

During the early phases of a pandemic, in the absence of a vaccine or effective

treatments, non-medical interventions are of utmost importance. During COVID19, the

majority of schools in the US closed around mid-March for the remainder of the school

year [54], and governors issued shelter-in-place orders [55] during March or April. Many

states ended their shelter-in-place orders towards the end of April or early May, given

the financial, social, and psychological impacts of shelter-in-place. For example, in

Georgia, schools closed on March 16, 2020 [56], shelter-in-place orders were issued on

April 3, 2020 ,transitioning out of shelter-in-place started on April 24, 2020 [57], and

shelter-in-place officially ended on April 30, 2020. The number of new COVID19

confirmed infections in Georgia have rapidly increased since early June [46]. In this

paper, we analyzed the impact of shelter-in-place duration and social distancing

compliance levels, particularly voluntary quarantine, using data from Georgia.

For baseline comparisons, we ran two scenarios: no intervention and school closure

only. We tested nine intervention scenarios, assuming school closures starting on

March 16, 2020, followed by shelter-in-place on April 3, 2020. In these nine scenarios,

we modeled a slowly increasing social distancing compliance prior to school closures,

shelter-in-place durations of 4-6 weeks, voluntary shelter-in-place and voluntary

quarantine compliance levels of low, medium, and high after the end of shelter-in-place.

Our study period extended through the end of September.

Compared to Scenarios 1-9, infections are higher in the baseline scenarios, with peak

infections occurring around mid-to-late April. Scenarios 1-9 show that shelter-in-place

could significantly slow down the disease spread, protecting public health, and offering

the opportunity for better preparedness of healthcare resource capacity. Each week

extension of shelter-in-place (beyond 4 weeks) could delay the peak day by about 6

days.

Social distancing (modeled by voluntary quarantine compliance) showed a significant

impact on all outcome metrics, particularly, peak day and peak infections. Depending on

social distancing compliance levels, the peak infections under low compliance levels

could exceed 50K. State level peak percentage could also decrease significantly as the

voluntary quarantine compliance increases, observing the same effect of post shelter-in-

place compliance at the county level as well.

After the end of shelter-in-place, voluntary quarantine compliance had a significant

impact on COVID19-related deaths, which could approach 20K by the end of

September in the case of low voluntary quarantine compliance. The death numbers

could be even higher because the demand would significantly exceed healthcare

capacity, especially around the peak.

Infection spread across counties varied over time, with earlier peak days in some

counties compared to others, which impacts resource allocation decisions across the

state. For example, in Scenario 1 (shelter-in-place 4 weeks, low voluntary quarantine

compliance after shelter-in-place), several of the larger counties could reach their peak

around late July, with the peak days of other counties extending to late July or mid-

August. Similarly, there was a variation among counties in terms of the peak infection

percentage under different voluntary quarantine compliance levels.

Estimates for COVID19-related needs for hospital beds, ICU beds, and ventilators

suggested shortages in all scenarios across the 14 coordinating hospital regions in

Georgia. Even if all available hospital resources were used for COVID19 patients, at the

peak, these resources would still not be sufficient for some of the regions. In some

regions, the shortage would continue for several weeks. These results further

emphasize the importance of voluntary shelter-in-place and high compliance levels for

social distancing after the end of state-mandated shelter-in-place.

During shelter-in-place, the withdrawal of a large percentage of the population (e.g.,

80%, depending on compliance level) from interactions raises social and economic

concerns. Hence, voluntary quarantine is an effective intervention if widely adopted, but

significantly less disruptive than shelter-in-place.

Another important advantage of voluntary quarantine is the ease of communication and

implementation. When COVID19 diagnostic testing capacity is limited, voluntary

quarantine can be implemented based on symptoms. Household members are advised

to stay home if there is a person with cold- or flu-like symptoms in the household (even

in the absence of testing or confirmation of COVID19), until the entire household is

symptom-free. High compliance with voluntary quarantine would reduce not only the

spread of COVID19 but would have the side benefit of also reducing the spread of the

cold or flu.

350   Limitations

Most of the limitations of this study lie in the limited data available regarding COVID19

infection and transmission, and the related parameters (which were drawn from the

literature), which impact the natural history and severe outcomes for the study period.

While we incorporated Georgia data on household types and sizes, children status,

workflow, and population demographics, our model made assumptions about the peer-

to-peer interactions between different age groups, which impact the virus transmission

under various intervention strategies. Testing of a wide range of scenarios enabled a

better understanding of the impact of social distancing compliance on COVID19

outcomes. This study did not consider the usage of face coverings; the projected

infections and deaths would decrease, but the relative effectiveness of interventions

would not change with the use of face coverings.

362   Conclusions

As states continuously evaluate the benefits versus social and economic costs of non-

pharmaceutical interventions such as school closures and shelter-in-place, our results

suggest that there needs to be a very strong messaging to the public about social

distancing. It is important to re-emphasize that some people might be infected with little

or no symptoms and infect others [58]. Voluntary quarantine is one form of social

distancing that is easy to communicate; it reduces infection spread (both from

symptomatic and asymptomatic individuals), but does not entirely prevent the spread.

There may be households with COVID19 infection, and yet no household member might

be experiencing symptoms – these households would not be impacted by voluntary

quarantine. Therefore, while it is essential to promote voluntary quarantine, strongly

encouraging households to continue voluntary shelter-in-place, to the extent possible, or

other forms of social distancing would help slow the spread of COVID19. It is also

important to enact policies and guidelines for promoting voluntary quarantine at the local

and national levels. Without such policies, compliance will be low, and hence, such

interventions will become quickly ineffective.

Acknowledgement

The authors of this paper are thankful to state of Georgia representatives, including Garry McGiboney from Georgia Department of Education, Laura Edison from Georgia Department of Public Health, and Susan Miller and Natalie Lee from Georgia Geospatial Information Office for their support, guidance, or sharing data sources. The authors would like to thank StreetLight Data, Inc. for providing data that motivated our scenario choices. The authors are also thankful to Melody Shellman, Hannah Lin, Ethan Channel, Pravara Harati, April Zhuoting Yu, Gabriel Siewert, and Christopher Stone for supporting various parts of the projects, and to the anonymous PLoS One reviewers and editors whose constructive comments and suggestions helped significantly improve the content and exposition.

REFERENCES

1.   WHO. WHO Timeline - COVID-19 2020 [27 April 2020]. Available from: https://www.who.int/news-room/detail/27-04-2020-who-timeline---covid-19.
2.   Boyle P. Here's why we can't rush a COVID-19 vaccine: Association of American Medical Colleges; 2020 [25 April 2020]. Available from: https://www.aamc.org/news-insights/here-s-why-we-can-t-rush-covid-19-vaccine.
3.   Hartley DM, Perencevich EN. Public Health Interventions for COVID-19: Emerging Evidence and Implications for an Evolving Public Health Crisis. JAMA. 2020. doi: 10.1001/jama.2020.5910.
4.   Giordano G, Blanchini F, Bruno R, Colaneri P, Di Filippo A, Di Matteo A, et al. Modelling the COVID-19 epidemic and implementation of population-wide interventions in Italy. Nature Medicine. 2020. doi: 10.1038/s41591-020-0883-7.
5.   Wu JT, Riley S, Fraser C, Leung GM. Reducing the Impact of the Next Influenza Pandemic Using Household-Based Public Health Interventions. PLOS Medicine. 2006;3(9):e361. doi: 10.1371/journal.pmed.0030361.
6.   Ekici A, Keskinocak P, Swann JL. Modeling Influenza Pandemic and Planning Food Distribution. Manufacturing & Service Operations Management. 2014;16(1):11-27. doi: 10.1287/msom.2013.0460.
7.   Shi P, Keskinocak P, Swann JL, Lee BY. The impact of mass gatherings and holiday traveling on the course of an influenza pandemic: a computational model. BMC Public Health. 2010;10(1):778. Epub 2010/12/24. doi: 10.1186/1471-2458-10-778. PubMed PMID: 21176155; PubMed Central PMCID: PMCPMC3022852.
8.   Li Z, Swann JL, Keskinocak P. Value of inventory information in allocating a limited supply of influenza vaccine during a pandemic. PLoS One. 2018;13(10):e0206293. Epub 2018/10/26. doi: 10.1371/journal.pone.0206293. PubMed PMID: 30359445; PubMed Central PMCID: PMCPMC6201932.
9.   Ekici A, Keskinocak P, Swann JL, editors. Pandemic influenza response. 2008 Winter Simulation Conference; 2008 7-10 Dec. 2008.
10.   Valdez LD, Macri PA, Braunstein LA. Intermittent social distancing strategy for epidemic control. Physical review E: Statistical, nonlinear, and soft matter physics. 2012;85(3 Pt 2):036108. Epub 2012/05/17. doi: 10.1103/PhysRevE.85.036108. PubMed PMID: 22587150.
11.   Towers S, Feng Z. Social contact patterns and control strategies for influenza in the elderly. Mathematical Biosciences. 2012;240(2):241-9. Epub 2012/08/11. doi: 10.1016/j.mbs.2012.07.007. PubMed PMID: 22877728.
12.   Mossong J, Hens N, Jit M, Beutels P, Auranen K, Mikolajczyk R, et al. Social contacts and mixing patterns relevant to the spread of infectious diseases. PLOS Medicine. 2008;5(3):e74. Epub 2008/03/28. doi: 10.1371/journal.pmed.0050074. PubMed PMID: 18366252; PubMed Central PMCID: PMCPMC2270306.
13.   Ferguson NM, Cummings DAT, Cauchemez S, Fraser C, Riley S, Meeyai A, et al. Strategies for containing an emerging influenza pandemic in Southeast Asia. Nature. 2005;437(7056):209-14. doi: 10.1038/nature04017.
14.   Jackson C, Mangtani P, Hawker J, Olowokure B, Vynnycky E. The effects of school closures on influenza outbreaks and pandemics: systematic review of simulation studies. PLoS One. 2014;9(5):e97297. Epub 2014/05/17. doi: 10.1371/journal.pone.0097297. PubMed PMID: 24830407; PubMed Central PMCID: PMCPMC4022492.
15.   Prem K, Liu Y, Russell T, Kucharski AJ, Eggo RM, Davies N, et al. The effect of control strategies that reduce social mixing on outcomes of the COVID-19 epidemic in Wuhan, China. medRxiv. 2020:2020.03.09.20033050. doi: 10.1101/2020.03.09.20033050.

16.   Lai S, Ruktanonchai NW, Zhou L, Prosper O, Luo W, Floyd JR, et al. Effect of non-pharmaceutical interventions for containing the COVID-19 outbreak in China. medRxiv. 2020:2020.03.03.20029843. doi: 10.1101/2020.03.03.20029843.

17.   Greenblatt Library. Regional Maps: Augusta University; 2020 [24 April 2020]. Available from: https://www.augusta.edu/library/greenblatt/disaster/map-ga.php/.

18.   U.S. Census Bureau. American Community Survey, 2017 American Community Survey 5-year Estimates. 2017.

19.   U.S. Census Bureau. American Community Survey, 2018 American Community Survey 1-year Estimates. U.S. Census Bureau; 2018.

20.   Ferguson NM, Laydon D, Nedjati-Gilani G, Imai N, Ainslie K, Baguelin M, et al. Impact of non-pharmaceutical interventions (NPIs) to reduce COVID19 mortality and healthcare demand. Imperial College London, Team ICC-R; 2020 16 March 2020. Report No.

21.   CDC COVID-19 Response Team. Severe Outcomes Among Patients with Coronavirus Disease 2019 (COVID-19) — United States, February 12–March 16, 2020. Morbidity and Mortality Weekly Report (MMWR): CDC, 18 March 2020. Report No.

22.   Riou J, Hauser A, Counotte MJ, Althaus CL. Adjusted age-specific case fatality ratio during the COVID-19 epidemic in Hubei, China, January and February 2020. medRxiv. 2020:2020.03.04.20031104. doi: 10.1101/2020.03.04.20031104.

23.   The Novel Coronavirus Pneumonia Emergency Response Epidemiology Team. The Epidemiological Characteristics of an Outbreak of 2019 Novel Coronavirus Diseases (COVID-19) — China, 2020. China CDCWeekly: 2020 14 February 2020. Report No.: Contract No.: x.

24.   Epicentro Epidemiology for Public Health. COVID-19 integrated surveillance: key national data 2020 [15 April 2020]. Available from: https://www.epicentro.iss.it/en/coronavirus/sars-cov-2-integrated-surveillance-data.

25.   Maharaj S, Kleczkowski A. Controlling epidemic spread by social distancing: do it well or not at all. BMC Public Health. 2012;12(1):679. Epub 2012/08/22. doi: 10.1186/1471-2458-12-679. PubMed PMID: 22905965; PubMed Central PMCID: PMCPMC3563464.

26.   Mizumoto K, Kagaya K, Zarebski A, Chowell G. Estimating the asymptomatic proportion of coronavirus disease 2019 (COVID-19) cases on board the Diamond Princess cruise ship, Yokohama, Japan, 2020. Eurosurveillance. 2020;25(10):2000180. doi: doi:https://doi.org/10.2807/1560-7917.ES.2020.25.10.2000180.

27.   Mandavilli A. Infected but Feeling Fine: The Unwitting Coronavirus Spreaders: The New York Times; 2020 [31 March 2020].

28.   Walker PG, Whittaker C, Watson O, Baguelin M, Ainslie KEC, Bhatia S, et al. The Global Impact of COVID-19 and Strategies for Mitigation and Suppression. Imperial College COVID-19 Response Team, 2020.

29.   Li R, Pei S, Chen B, Song Y, Zhang T, Yang W, et al. Substantial undocumented infection facilitates the rapid dissemination of novel coronavirus (SARS-CoV2). Science. 2020:eabb3221. Epub 2020/03/18. doi: 10.1126/science.abb3221. PubMed PMID: 32179701.

30.   WHO. Report of the WHO-China Joint Mission on Coronavirus Disease 2019 (COVID-19) World Health Organization, 2020 16-24 February 2020. Report No.

31.   Chen TM, Rui J, Wang QP, Zhao ZY, Cui JA, Yin L. A mathematical model for simulating the phase-based transmissibility of a novel coronavirus. Infectious Diseases of Poverty. 2020;9(1):24. Epub 2020/03/01. doi: 10.1186/s40249-020-00640-3. PubMed PMID: 32111262; PubMed Central PMCID: PMCPMC7047374.

32.   Linton NM, Kobayashi T, Yang Y, Hayashi K, Akhmetzhanov AR, Jung S-m, et al. Incubation Period and Other Epidemiological Characteristics of 2019 Novel Coronavirus Infections with Right Truncation: A Statistical Analysis of Publicly Available Case Data. medRxiv. 2020:2020.01.26.20018754. doi: 10.1101/2020.01.26.20018754.

33.   Li Q, Guan X, Wu P, Wang X, Zhou L, Tong Y, et al. Early Transmission Dynamics in
Wuhan, China, of Novel Coronavirus–Infected Pneumonia. New England Journal of Medicine.
2020;382(13):1199-207. doi: 10.1056/NEJMoa2001316.
34.   Weitz J. Intervention Serology and Interaction Substitution: Exploring the Role of
`Immune Shielding' in Reducing COVID-19 Epidemic Spread 2020.
35.   Ganyani T, Kremer C, Chen D, Torneri A, Faes C, Wallinga J, et al. Estimating the
generation interval for COVID-19 based on symptom onset data. medRxiv.
2020:2020.03.05.20031815. doi: 10.1101/2020.03.05.20031815.
36.   Zhou F, Yu T, Du R, Fan G, Liu Y, Liu Z, et al. Clinical course and risk factors for
mortality of adult inpatients with COVID-19 in Wuhan, China: a retrospective cohort study. The
Lancet. 2020;395(10229):1054-62. doi: 10.1016/s0140-6736(20)30566-3.
37.   Xie J, Tong Z, Guan X, Du B, Qiu H, Slutsky AS. Critical care crisis and some
recommendations during the COVID-19 epidemic in China. Intensive Care Medicine. 2020. doi:
10.1007/s00134-020-05979-7.
38.   Andrei M. Iceland's testing suggests 50% of COVID-19 cases are asymptomatic ZME
Science2020 [updated 26 March 202014 April 2020]. Available from:
https://www.zmescience.com/medicine/iceland-testing-covid-19-0523/.
39.   Day M. Covid-19: four fifths of cases are asymptomatic, China figures indicate. BMJ.
2020;369:m1375. doi: 10.1136/bmj.m1375.
40.   Nishiura H, Kobayashi T, Suzuki A, Jung SM, Hayashi K, Kinoshita R, et al. Estimation
of the asymptomatic ratio of novel coronavirus infections (COVID-19). International Journal of
Infectious Diseases. 2020. Epub 2020/03/18. doi: 10.1016/j.ijid.2020.03.020. PubMed PMID:
32179137.
41.   U.S. Census Bureau. Census Summary File 1. U.S. Census Bureau; 2010.
42.   U.S. Census Bureau. Census Transportation Planning Products, 5-year data. U.S.
Census Bureau; 2016.
43.   The New York Times. Coronavirus (Covid-19) Data in the United States. The New York
Times; 2020.
44.   CDC. COVID-19 Pandemic Planning Scenarios 2020 [5 June 2020]. Available from:
https://www.cdc.gov/coronavirus/2019-ncov/hcp/planning-scenarios.html.
45.   Backer JA, Klinkenberg D, Wallinga J. Incubation period of 2019 novel coronavirus
(2019-nCoV) infections among travellers from Wuhan, China, 20–28 January 2020.
Eurosurveillance.   2020;25(5):2000062.   doi: doi:https://doi.org/10.2807/1560-
7917.ES.2020.25.5.2000062.
46.   Georgia Department of Public Health. Georgia Department of Public Health COVID-19
Daily Status Report 2020 [updated 14 April 2020; cited 2020 14 April 2020]. Available from:
https://dph.georgia.gov/covid-19-daily-status-report.
47.   Xie J, Tong Z, Guan X, Du B, Qiu H, Slutsky AS. Critical care crisis and some
recommendations during the COVID-19 epidemic in China. Intensive care medicine. 2020:1-4.
doi: 10.1007/s00134-020-05979-7. PubMed PMID: 32123994.
48.   CBS Atlanta. BREAKING: COVID-19: Gov. Kemp Extends Statewide Shelter In Place
Through April 30. CBS Atlanta. 2020.
49.   StreetLight Data Inc. Understand the impact of COVID-19 on traffic, travel patterns, toll
revenues and more   [cited 2020]. Available from: https://www.streetlightdata.com/covid-
transportation-metrics/.
50.   Lever J, Krzywinski M, Altman N. Principal component analysis. Nature Methods.
2017;14(7):641-2. doi: 10.1038/nmeth.4346.
51.   CDC. People Who Are at Higher Risk for Severe Illness 2020 [25 April 2020]. Available
from: https://www.cdc.gov/coronavirus/2019-ncov/need-extra-precautions/groups-at-higher-
risk.html.

52. Georgia Department of Public Health. Regional healthcare coalitions 2018 [24 July
2020]. Available from: https://northcentralhealthdistrict.org/regional-healthcare-coalitions/.
53. Esri. COVID-19 GIS Hub: Esri; 2020 [29 April 2020]. Available from: http://coronavirus-
resources.esri.com/.
54. Chavez N, Moshtaghian A. 40 states and Washington, DC have ordered or
recommended that schools don't reopen this academic year: CNN; 2020 [24 April 2020].
Available from: https://www.cnn.com/2020/04/18/us/schools-closed-coronavirus/index.html.
55. Mervosh S, Lu D, Swales V. See Which States and Cities Have Told Residents to Stay
at Home: The New York Times; 2020 [24 April 2020]. Available from:
https://www.nytimes.com/interactive/2020/us/coronavirus-stay-at-home-order.html.
56. Walker M, Broady A, McCray V. Kemp orders schools closed through end of school
year: The Atlantic Journal-Constitution; 2020 [24 April 2020]. Available from:
https://www.ajc.com/news/state--regional-education/schools-closed-until-
fall/r7QgK2idaQ0681UafbW3XP/.
57. WTVC / The Associated Press. Kemp: Some Georgia businesses allowed to reopen
April 24; shelter-in-place to end April 30: WTVC / The Associated Press; 2020 [24 April 2020].
Available from: https://newschannel9.com/news/local/gov-kemp-certain-georgia-businesses-
allowed-to-reopen-april-24.
58. Bai Y, Yao L, Wei T, Tian F, Jin D-Y, Chen L, et al. Presumed Asymptomatic Carrier
Transmission of COVID-19. JAMA. 2020;323(14):1406–7. doi: 10.1001/jama.2020.2565.
59. Weinberger DM, Chen J, Cohen T, Crawford FW, Mostashari F, Olson D, et al.
Estimation of Excess Deaths Associated With the COVID-19 Pandemic in the United States,
March to May 2020. JAMA Internal Medicine. 2020. doi: 10.1001/jamainternmed.2020.3391.
60. CDC. Commercial Laboratory Seroprevalence Survey Data [Online]. 2020 [updated 26
June 2020]. Available from: https://www.cdc.gov/coronavirus/2019-ncov/cases-
updates/commercial-lab-surveys.html.
61. U.S. Census Bureau. Cartographic Boundary Files - Shapefile.
https://www.census.gov/geographies/mapping-files/time-series/geo/carto-boundary-file.html
2018.
566

Supporting information

 S1 Appendix. Data sources, model description and model inputs.

 S1 Fig. State level outcomes: IAR, CAR, HB, ICUB, V across all scenarios. State

 Level Outcomes: IAR (first row left plot), CAR (first row right plot), HB (second row left

 plot), ICUB (second row right plot), V (third row center plot) across all scenarios

 (including the baseline scenarios).

S2 Fig. State level outcomes: NIC for low, medium, and high levels of VQ after 4,

5, and 6 weeks of SIP. Daily new COVID19 infections when Low VQ is combined with

4 week (Scenario 1), 5 week (Scenario 4), 6 week (Scenario 7) SIP (top left plot),

Medium VQ is combined with 4 week (Scenario 2), 5 week (Scenario 5), 6 week

(Scenario 8) SIP (top right plot), High VQ is combined with 4 week (Scenario 3), 5 week

(Scenario 6), 6 week (Scenario 9) SIP (bottom center plot).

S3 Fig. Number of new infections per 100,000 people. Four maps of Georgia at the

county level recording the number of new infections per 100,000 people for June 23,

2020 (using the actual number of infections), July 15, 2020 (simulated data from our

model), August 15, 2020 (simulated data), and September 15, 2020 (simulated data),

respectively [61].

588