Lack of Evidence for a Reduced Late Positive Potential in Major Depressive Disorder

Stevan Nikolin[1,2], Nicholas Chand[2], Donel Martin[1,2], Jacqueline Rushby[3], Colleen K. Loo[1,2,4], Tjeerd W. Boonstra[1,5]

[1] School of Psychiatry, University of New South Wales, Sydney, Australia

[2] Black Dog Institute, Sydney, Australia

[3] School of Psychology, University of New South Wales, Sydney, Australia

[4] St. George Hospital, Sydney, Australia

[5] Department of Neuropsychology and Psychopharmacology, Faculty of Psychology and Neuroscience, Maastricht University, Maastricht, the Netherlands

Keywords: Depression, Emotional Regulation, Electroencephalography, Event-Related Potentials, Late Positive Potential

Running title: Affective Processing in Depression

Corresponding Author:

Tjeerd W. Boonstra;

tjeerd.boonstra@maastrichtuniversity.nl

ABSTRACT

Background: Individuals with major depressive disorder (MDD) present with deficits in emotional reactivity. Conflicting models have been proposed to explain this effect. We sought to test the *emotional context insensitivity* hypothesis, which suggests that reactivity to positive and negatively-valenced emotional stimuli is blunted in depression, in a preregistered study.

Methods: Forty-one depressed participants and 41 age- and gender-matched healthy controls were presented a series of unpleasant and neutrally-valenced pictures in a passive view paradigm while acquiring electroencephalography (EEG). The late positive potential (LPP), an EEG correlate of emotional reactivity, was compared between groups using mixed-effects repeated-measures models and exploratory cluster-based permutation tests. A sensitivity analysis was performed to assess the robustness of LPP findings by reanalysing the LPPs using 22 EEG pipelines from studies identified in the literature.

Results: We found no difference in LPP amplitudes between MDD and healthy individuals using the preregistered analysis pipeline. The sensitivity analysis revealed that the magnitude and direction of LPP effect sizes were affected by the analysis pipeline. Exploratory permutation analyses revealed an electrode cluster that showed a significant reduction in the LPP for MDD participants while viewing unpleasant pictures.

Conclusions: These results do not provide evidence in support of the emotional context insensitivity hypothesis, except for the exploratory data-driven approach. Methodological differences, in particular in the analysis pipeline, contribute to the heterogeneity of LPP

modulation in depression. A standardised approach to quantify EEG correlates of emotional

reactivity is needed to evaluate alternative models of emotional reactivity in depression.

INTRODUCTION

Major depressive disorder (MDD) is a mental illness that presents as persistent low mood and anhedonia, loss of interest in pleasurable activities, or both, in conjunction with other symptoms (American Psychiatric Association, 2013). Consistent with the core symptom of disturbed mood, depressed individuals report altered reactivity to emotion-eliciting contexts of positive or negative affect (Rottenberg et al., 2005). An improved understanding of the emotional deficits in MDD may thus provide further insight into mood symptoms (Rottenberg, 2005). Although the terms mood and emotion are sometimes used interchangeably, mood typically refers to slow-moving affective states that are weakly influenced by acute external stimuli, whereas emotions are considered to be rapid reactions to meaningful affective stimuli (Watson, 2000). Notwithstanding, the two are deeply intertwined and an improved understanding of the emotional deficits in MDD may thus provide further insight into mood symptoms (Rottenberg, 2005).

Three competing hypotheses have been proposed to describe the observed abnormalities in emotional reactivity in depressed individuals: 1) The *positive attenuation* model suggests that the physiological response is diminished to positive stimuli and unchanged for unpleasant, negatively-valenced stimuli (Clark and Watson, 1991), 2) The *negative potentiation* hypothesis states that depression increases reactivity towards negative contexts (Browning et al., 2010), and 3) The *emotional context insensitivity* model argues that depression results in blunted emotional reactivity to positive stimuli as well as to negative emotional contexts (Rottenberg et al., 2005). These theories disagree with regards to the physiological response in MDD to negative stimuli: *positive attenuation* predicts no difference in reactivity compared to healthy individuals, *negative potentiation* predicts

increased reactivity, and the *emotional context insensitivity* predicts decreased reactivity in depression.

Emotional reactivity is commonly assessed using electroencephalography (EEG). Specifically, the late positive potential (LPP) may provide a reliable measure of affective cortical processing (Codispoti et al., 2006, 2007). The LPP is characterized as the mean amplitude of the event-related potential (ERP) within a time window approximately 400-1000ms following the visual presentation of emotionally laden images (Cuthbert et al., 2000; Schupp et al., 2000). Self-reported depressive symptoms have been associated with reduced LPP amplitudes to positive and unpleasant emotional stimuli in a non-clinical sample (Hill et al., 2019). Further, studies of depressed individuals have demonstrated attenuated LPP amplitudes in response to unpleasant stimuli (Benau et al., 2019; Foti et al., 2010; Grunewald et al., 2019; MacNamara et al., 2016), and pleasant pictures (Klawohn et al., 2020), in agreement with a summary of the LPP literature in a recent review by Hajcak and Foti (2020). However, these findings are not consistently observed, with some research reporting a significant increase in LPP amplitudes while passively observing unpleasant pictures (Auerbach et al., 2015; Burkhouse et al., 2017; Zhang et al., 2016), or no statistically significant difference (Weinberg et al., 2016). Although a meta-analysis found that MDD was characterized by reduced emotional reactivity to both positively and negatively valenced stimuli (Bylsma et al., 2008), in agreement with the *emotional context insensitivity* hypothesis, the results also demonstrated substantial heterogeneity between studies that could not be attributed to a specific moderator.

One of the potential contributing factors of heterogeneity between studies examining LPP amplitudes in response to emotional images is the role of emotion regulation in depression.

In healthy individuals, the LPP is sensitive to the use of emotion regulation strategies, which refers to conscious or subconscious efforts to exert control over one's emotional experience (Gross and John, 2003). Differences in the frequency or type of emotion regulation strategy use could partly explain the observed differences in the LPP to unpleasant images in MDD participants compared to healthy controls (Foti and Hajcak, 2008; Hajcak and Nieuwenhuis, 2006; Moser et al., 2009). In addition to emotion regulation strategy, anxiety, a common comorbidity of depression (Brown et al., 1998), has also been found to modulate the LPP. Individuals with high self-reported anxiety showed enhanced LPP amplitudes to aversive pictures compared to neutral pictures (MacNamara and Hajcak, 2009; MacNamara et al., 2016). Anxiety in depressed individuals may therefore contribute to the conflicting evidence for alternative models of emotional reactivity in MDD.

The primary aim of this study was therefore to verify previous reports in support of the *emotional context insensitivity* hypothesis by comparing the contrast in LPP amplitudes for unpleasant *vs* neutral pictures between MDD participants and healthy controls. We hypothesised that the LPP amplitude difference (i.e., unpleasant *vs* neutral) would be attenuated in MDD participants in support of *emotional context insensitivity*. We performed additional secondary analyses to examine whether emotion regulation strategy use and comorbid anxiety in MDD participants modulate the LPP amplitude in response to unpleasant pictures. We anticipated that MDD participants would use emotion regulation strategies less frequently than healthy controls, and that LPP amplitude would be moderated by emotion regulation strategy use and anxiety comorbidity in MDD participants. We based the research protocol used to test our hypotheses – the required sample size, use of affective picture stimuli, time interval, and region of interest to quantify

the LPP amplitude – on previous literature, and documented this in a preregistration form (Nikolin et al., 2020). Preregistration provides a clear distinction between confirmatory and exploratory research and improves effect-size estimation, and may therefore help to evaluate the LPP as a biomarker for depression (Clayson et al., 2020). In addition, we identified 22 EEG processing pipelines reported in previous studies on the LPP in depression and reanalysed our data using these alternative processing pipelines to assess the robustness of the attenuation of the LPP amplitude difference to variations in processing parameters.

MATERIALS AND METHODS

Participants

A total of 82 participants were recruited in this study, of whom 41 were currently depressed and 41 were healthy age- and gender-matched controls. Individual participant data could only be obtained from Foti et al. (2010) to estimate the effect size for the contrast in LPP amplitudes between MDD participants and healthy controls for unpleasant *vs* neutral pictures. This showed a large effect size (Cohen's d = 0.89), which would require 21 participants per diagnostic group to detect. We approximately doubled this sample size to allow for the detection of a more conservative, moderate effect size of Cohen's d = 0.6 (using β = 80% and α = 0.05).

Inclusion criteria shared by both MDD participants and healthy controls were: aged 18 years and above, able to provide informed consent, normal or corrected-to-normal vision, no recent serious head injury in the last 12 months (e.g. loss of consciousness of more than 30

minutes), no current or past neurological condition (e.g. stroke, epilepsy), no history of drug or alcohol dependence in the last three months, not pregnant or possibly pregnant, no past history of distress reaction to affective images, and an ability to cooperate with the EEG procedure (e.g. no tremor).

Depressed participants were recruited from those seeking entry into clinical treatment trials at the Black Dog Institute, Sydney Australia. They were assessed by an experienced study Psychiatrist and confirmed to meet DSM-5 criteria for a current MDD diagnosis using the Structured Clinical Interview for DSM-5 Research Version (First et al., 2015), as well as score ≥ 20 on the Montgomery–Åsberg Depression Rating Scale for depressive symptoms (Montgomery and Asberg, 1979). Healthy controls were recruited via the University of New South Wales (UNSW) School of Psychology Paid Sign-Up System, Black Dog Institute Volunteer Research Register, or flyers distributed in the local area. Healthy control participants were eligible to participate if they had no current or past psychiatric or neurological disorders and were selected to be the same gender and within five years of age of the matched MDD participants since these factors have previously been shown to influence the latency, amplitude, and topography of ERP components (Kayser et al., 2000; Oliver-Rodriguez et al., 1999; Renfroe et al., 2016). The study protocol was approved by the UNSW Human Research Ethics Committee (HC16617).

Procedure

Participants were instructed to passively view and attend to picture stimuli, selected from the International Affective Picture System (IAPS; Lang et al. (2008)). Pictures were chosen to be consistent with those used in previous EEG studies investigating LPP effects (Hajcak and Nieuwenhuis, 2006; MacNamara et al., 2016; Moser et al., 2006; Weinberg et al., 2016).

Participants viewed 56 images once each, of which 28 were unpleasant (e.g., threatening scenes, violence, weapons) and 28 were neutral (e.g., neutral faces, household objects, landscapes). The normative valence ratings were lower for unpleasant pictures (2.79 ± 0.59) than for neutral pictures (5.02 ± 0.20), indicating that they elicited more negative emotional affect. Normative arousal ratings were higher for unpleasant pictures (5.74 ± 0.84) than for neutral pictures (2.98 ± 0.59; see Supplementary Materials for further details).

The stimuli were presented on a 58.4 cm (23-inch) computer screen in a dimly lit, sound-attenuated room using Inquisit 4 software (Version 4, Millisecond Software). Digitised pictures sized 20×30 cm were presented on a black background, spanning a visual angle of 20-30 degrees, consistent with previous EEG studies (Dunning and Hajcak, 2009; MacNamara et al., 2016). The pictures were displayed for 5 s in a pre-determined randomised order, so participants were unable to anticipate the valence of each picture. A black screen with a centered white fixation cross was presented for 2 s between stimuli (Fig. 1). Pictures of the same valence were not presented more than three times consecutively to prevent participants from generating expectations of upcoming stimuli and to avoid over-saturation of emotional salience. Participants were not informed of any emotional regulation strategies prior to viewing the IAPS images.

Figure 1. Study protocol. Participants passively observed a total of 56 picture stimuli. These were presented in a randomised order and consisted of 28 unpleasant images with a negative valence, and 28 neutral images with no emotional valence, sourced from the International Affective Picture System (IAPS). Pictures were displayed for 5 s, followed by a 2 s fixation cross. A sample of two trials is displayed, showing a neutral image (a towel) and an unpleasant image (a gun).

Behavioural measures

Prior to the experimental session, MDD participants had their mood assessed by a trained rater using the ten-item Montgomery-Åsberg Depression Rating Scale (MADRS) for depressive symptoms (Montgomery and Asberg, 1979). All participants completed the 21-item self-reported Depression, Anxiety and Stress Scale (DASS; Lovibond and Lovibond (1995)). The DASS was chosen as it reliably discriminates between depression, anxiety, and stress states (Brown et al., 1997; Crawford and Henry, 2003; Henry and Crawford, 2005). Upon completion of the experimental session, we used an emotion regulation questionnaire adapted from Gross and John (2003), provided in Supplementary Materials. Participants

were able to select from five emotion regulation strategies, including no strategy, reinterpretation, distraction, detachment, and 'other' in which participants could specify their strategy. Strategies specified in the 'other' category were re-assigned to one of the former strategy options by two study personnel who were blind to participant group allocation.

Electroencephalography data acquisition and processing

EEG data were acquired using a 64-channel setup and a TMSi Refa amplifier (TMS International, Oldenzaal, Netherlands) with a sampling rate of 1024 Hz. Offline EEG processing and analyses were conducted using custom-developed MATLAB scripts (v.R2019a; MathWorks) and the Fieldtrip toolbox (Oostenveld et al., 2011). All scripts used for EEG processing and calculation of neurophysiological measures are available at the following link: https://github.com/snikolin/MDDvsCTRL_IAPS.

EEG data were pre-processed using a Butterworth IIR digital filter to remove 50 Hz electrical line noise. Following this, a second-order band-pass filter with cut-offs of 0.5 and 70 Hz was used to remove low and high frequency noise generated from head movements and muscle activity. The EEG data were then epoched into 6 s intervals, beginning 1 s before each picture onset and continuing for 5 s after picture onset. Epochs were rejected firstly using a semi-automated procedure, and then on visual inspection to identify any remaining trials with large artefacts. Smaller, non-cortical physiological activity (e.g., cardiac, muscle, or ocular) and non-physiological activity (e.g., environmental noise, or movement) were removed using an Independent Components Analysis (ICA) algorithm (Delorme et al., 2007;

Hyvärinen et al., 2001; Makeig et al., 1996). Finally, EEG data was re-referenced to the

common average reference, which reduces the distortion of measured EEG potentials as

compared to a linked mastoids reference (Hu et al., 2018).

Event-related potentials

ERPs were constructed by separately averaging trials for neutral and unpleasant pictures

time-locked to picture presentation. Each averaged ERP was baseline corrected to the mean

activity occurring in the 200 ms time interval prior to stimulus onset. The late positive

potential (LPP) is a slow, positive wave occurring maximally at centroparietal recording sites,

beginning approximately 400 ms after stimulus onset and lasting several seconds (Cuthbert

et al., 2000; Schupp et al., 2000). We quantified the LPP component as the mean amplitude

of the ERP from 400 to 1000 ms following stimulus onset at a centroparietal cluster of EEG

channels (Cz, CPz, Pz, CP1, and CP2), where it has previously been shown to be maximal

(Foti et al., 2010; MacNamara et al., 2016). Participants with LPP component amplitudes

greater than three standard deviations from the mean were considered outliers and

excluded from statistical analyses.

Statistical analyses

For the primary analysis, LPPs were investigated using mixed-effects repeated-measures

model (MRMM), performed using SPSS software (IBM SPSS Statistics 25, SPSS Inc.), and

were in accordance with our preregistered protocol (Nikolin et al., 2020). The model

included a repeated factor of Valence (unpleasant vs neutral) using an unstructured

covariance matrix, as well as the between-subjects factor of Group (MDD vs healthy

controls), and the Valence × Group interaction. Subject was included as a random effect.

Residuals were visually inspected to ensure an adequate fit of the model. Cohen's d was calculated to estimate the effect size for the LPP amplitude interaction contrast between unpleasant and neutral images comparing MDD participants to healthy controls:

$$[\text{MDD LPP}_{\text{unpleasant}} - \text{MDD LPP}_{\text{neutral}}] \; vs \; [\text{Healthy LPP}_{\text{unpleasant}} - \text{Healthy LPP}_{\text{neutral}}]$$

A follow-up exploratory non-parametric cluster-based permutation test was performed using the Fieldtrip toolbox in MATLAB to identify any significant differences in emotional stimulus processing between MDD participants and healthy controls beyond the *a priori* time interval and channels operationalised for the LPP. This method controls for multiple comparisons while comparing differences across a large spatiotemporal parameter space (Maris and Oostenveld, 2007). Participant ERPs, incorporating all EEG channels within the time interval 0 – 1000 ms following presentation of IAPS stimuli, were randomly assigned to one of two group conditions in the same proportion as the original data for 3000 permutations. The permuted data were compared using independent samples t-tests, producing a data-driven distribution of the test statistics. The t-test statistic for the original, non-permuted data was then compared against the permuted distributions to identify any significant clusters. A value of $\alpha < 0.05$ was adopted as the two-tailed significance threshold. Statistically significant clusters were required to comprise at least two neighbouring channels.

For secondary analyses, a Fisher's Exact Test was used to investigate whether there were differences in emotion regulation strategy use between diagnostic groups. Emotion regulation strategies were entered into a 2 x 2 contingency table using the binary variables

of Strategy (no strategy vs strategy) and Group (MDD vs healthy controls). Emotional

regulation strategies consisted of reinterpretation, distraction, or detachment.

The effects of depression on LPP amplitude may become more apparent when accounting

for variability in emotion regulation strategy use or comorbid anxiety symptoms across

diagnostic groups. To this end, we repeated MRMM analyses with emotion regulation

strategy or comorbid anxiety entered separately as covariates. Strategy was represented as

a binary categorical variable (no strategy vs strategy). Anxiety was included as a continuous

variable using the anxiety subscore of the DASS. Additional confirmatory simple univariate

linear regression analyses were performed with LPP amplitude as the dependent variable,

restricted to amplitudes obtained from MDD participants while viewing unpleasant images.

Scripts for EEG analyses in MATLAB and statistical analyses using SPSS are provided at:

https://github.com/snikolin/MDDvsCTRL_IAPS.

Sensitivity analysis

Given the variability in the methodology used in previous studies to quantify the LPP, we

performed additional analyses to assess the robustness of our findings and recomputed the

LPP using pipelines identified in the depression literature. We performed a PubMed review

on the 8th of May 2021 for manuscripts containing the terms: depression AND

(electroencephalography OR EEG) AND "late positive potential". Studies were included if

they reported LPP results in participants with depression or depressive tendencies and

provided a comprehensive description of their EEG processing pipeline. We supplemented

the review with additional studies identified while reading this literature. We also assessed

alternative variants of our preregistered pipeline that were not observed in the literature

but may be of interest. These included: Repeating our preregistered pipeline analysis using linked mastoid re-referencing (Alternative #1) and reducing the high-pass filter to 0.1Hz (Alternative #2). We used a collapsed localiser approach (Luck and Gaspelin, 2017), which entails calculating the grand-average ERP for all participants (MDD and controls combined) for unpleasant relative to neutral IAPS images and using the resultant topography to select a sample-specific region of interest for the LPP (Alternative #3). Lastly, we repeated Alternative #2 using an average common reference (Alternative #4), and similarly repeated Alternative #3 using an average common reference (Alternative #5). We estimated effect sizes for each pipeline using Cohen's d, as described previously. We additionally report significance p-values for simple independent samples t-tests for each pipeline's contrast comparison (i.e., MDD $LPP_{unpleasant} -$ MDD $LPP_{neutral}$ vs Healthy $LPP_{unpleasant} -$ Healthy $LPP_{neutral}$). In total, we assessed 22 unique EEG processing pipelines and MRMMs were repeated using LPP amplitudes obtained from each pipeline.

Lastly, we repeated exploratory non-parametric cluster-based permutation tests using 1) a 0.5Hz high-pass filter with referencing to linked mastoids; 2) a 0.1Hz high-pass filter and common average reference; and lastly 3) a 0.1Hz high-pass filter with referencing to linked mastoids.

RESULTS

Behavioural and EEG measures were assessed in 82 participants (41 with MDD and 41 aged-
and gender-matched controls). Table 1 shows baseline demographic and clinical information
for MDD participants and healthy controls. One participant was excluded from EEG analyses
due to values greater than three standard deviations from the group mean, leaving a total
of 40 MDD participants and 41 healthy controls. Following rejection of epochs with EEG
artefacts, MDD participants had an average of 25.5 unpleasant trials (SD=2.1; range=20-28)
and 26.1 neutral trials (SD=1.7; range=21-28). Healthy controls had an average of 25.8
unpleasant trials (SD=1.8; range=20-28) and 26.4 neutral trials (SD=1.5; range=21-28).

Table 1. Baseline demographic and clinical information. Means, standard deviations (SD),
and ranges [minimum-maximum] are shown for continuous measures, and frequency
counts for categorical measures.

	MDD	Controls
n	41	41
Age (years)	46.3 (13.2) [21-76]	46.0 (13.9) [21-77]
Education (years)	15.8 (3.6) [9-30]	17.3 (3.6) [10-28]
Gender (females/males)	15/26	15/26
TRD (yes/no)	32/9	-
Concurrent antidepressants (yes/no)	31/9	-
SSRI (yes/no)	6/25	-
SNRI (yes/no)	7/24	-
TCA (yes/no)	8/23	-
MAOI (yes/no)	4/27	-
Other (yes/no)	6/25	-
MADRS	29.6 (5.0) [20-43]	-
DASS Depression	33.7 (7.1) [16-42]	4.2 (5.2) [0-24]
DASS Anxiety	11.0 (9.7) [0-32]	3.3 (4.5) [0-18]
DASS Stress	21.3 (9.5) [0-40]	6.8 (5.5) [0-20]
DASS Total	66.5 (19.7) [34-108]	14.2 (12.9) [0-50]

MDD: Major Depressive Disorder; TRD: Treatment-Resistant Depression – Defined as failure to respond to at least two adequate courses of antidepressant medication; SSRI: selective serotonin reuptake inhibitor; SNRI: serotonin-norepinephrine reuptake inhibitor; TCA: tricyclic antidepressant; MAOI: monoamine oxidase inhibitor; MADRS: Montgomery-Asperg Depression Rating Scale (Montgomery and Asberg, 1979); DASS: 21-item Depression, Anxiety and Stress Scale (Lovibond and Lovibond, 1995). DASS scores were not available for two MDD participants.

Primary Analysis

The LPP amplitude was compared across groups (MDD and controls) and valence conditions (unpleasant and neutral). MRMM results showed a significant main effect of Valence (F = 19.62, $p < 0.001$), indicating increased LPP amplitude for unpleasant compared to neutral images (Fig. 2). The main effect of Group (F = 0.39, $p = 0.54$) and the Group $\times$ Valence interaction (F = 0.26, $p = 0.61$) were not significant. Comparison of the valence difference between MDD participants and healthy controls showed a small effect size ($d = -0.11$).

Figure 2. Preregistered EEG measures of affective processing. The LPP was quantified as

the mean EEG amplitude in the time interval from 400 to 1000 ms following stimulus onset

for a cluster of centroparietal EEG channels (highlighted in red). ERPs, spatial topographies,

and scatter plots of LPP amplitude are presented for MDD participants and controls while

viewing unpleasant and neutral images. ERPs are displayed with bootstrapped 95%

confidence intervals and with the time interval used to calculate LPPs shaded in light grey.

Spatial topographies are of average ERP amplitude from 400 to 1000 ms. Scatter plots show

the mean (black line), median (red line), and the 95% confidence interval (dark grey region).

A) ERPs and topographies for MDD participants. **B)** ERPs and topographies for healthy

controls. **C)** ERPs and spatial topographies of the valence difference (unpleasant – neutral)

for MDD participants and healthy controls. **D)** Scatter plots of LPP amplitude for unpleasant

(U) and neutral (N) images at the centroparietal cluster. **E)** Spatial topography of the valence

difference (unpleasant – neutral) contrasting MDD participants with healthy controls.

Exploratory non-parametric cluster-based permutation testing compared the difference

scores (unpleasant – neutral stimuli) between MDD participants and healthy controls. These

revealed a cluster showing a significant difference between MDD participants and controls

(p = 0.01; Fig. 3). The time interval for the cluster fell within the preregistered LPP window

(662 – 755 ms). EEG channels identified in this cluster were slightly more posterior than

those for our preregistered LPP analysis and included CPz, CP1, Pz, POz, P1, and P2. Three of

these channels are also found in the preregistered cluster (CPz, CP1, and Pz). See

Supplementary Material for further details regarding cluster channels.

Figure 3. Exploratory EEG measures of affective processing. Exploratory cluster-based permutation testing revealed a significant cluster from 662 to 755 ms in EEG channels CPz, CP1, Pz, POz, P1, and P2 (highlighted in red). ERPs, spatial topographies, and scatter plots of LPP amplitude are presented for MDD participants and controls while viewing unpleasant and neutral images. ERPs are displayed with bootstrapped 95% confidence intervals and with the time interval used to calculate the mean amplitude within the exploratory cluster

shaded in light grey. Spatial topographies are of average ERP amplitude from 662 to 755 ms. Scatter plots show the mean (black line), median (red line), and the 95% confidence interval (dark grey region). **A)** ERPs and topographies for MDD participants. **B)** ERPs and topographies for healthy controls. **C)** ERPs and spatial topographies of the valence difference (unpleasant – neutral) for MDD participants and healthy controls. **D)** Scatter plots of LPP amplitude for unpleasant (U) and neutral (N) images at the exploratory cluster. Comparison of the valence differences between MDD participants and healthy controls showed a large effect size of Cohen's d = -0.88. **E)** Spatial topography of the valence difference (unpleasant – neutral) for the time interval 662 – 755 ms contrasting MDD participants with healthy controls.

Secondary Analyses

As part of our secondary confirmatory analyses, we also investigated the association between strategy use and anxiety on the LPP amplitude. We compared the emotion regulation strategies used by MDD participants and control. Most participants reported that they did not use a strategy (Table 2). A Fisher's Exact Test revealed that strategy use was not significantly different between both groups (p = 0.34).

Table 2. Emotion regulation strategy use.

	MDD (*n* = 40)	Controls (*n* = 41)
No strategy	30	26
Strategy	10	15
Reinterpretation	1	5
Distraction	2	5
Detachment	7	5

MDD: Major Depressive Disorder

We performed MRMMs with the inclusion of either Anxiety or Strategy as covariates.

Neither the model including Strategy (F = 0.2, p = 0.67), nor the model including Anxiety (F =

0.2, p =0.63) were substantively changed (see Supplementary Materials). Next, we

performed simple univariate linear regression analyses, restricting the dataset to LPP values

obtained from MDD participants while viewing unpleasant images. Strategy was not

significantly associated with LPP amplitudes (r = 0.13, r^2 = 0.02, β = -0.18, p = 0.44; Fig. S1A).

Similarly, Anxiety was not significantly associated with the LPP (r = 0.21, r^2 = 0.05, β = 0.01, p

= 0.20; Fig. S1B).

Sensitivity analysis

We reanalysed our data using 22 alternative EEG processing pipelines (Table 3). A literature

review identified 18 studies and the most used parameters were a high-pass filter of 0.1Hz

(k=9), low-pass filter of 30Hz (k=16), correction for ocular artefacts using a regression

method such as that proposed by Gratton et al., (1983) (k=14), and re-referencing to linked

mastoids (k=12). The LPP time of interest was most often operationalised as starting at

either 400 or 600 ms (k=7), and ending at 1000 ms (k=9). The region of interest was typically

centro-parietal, mostly consisting of channels Pz (k=11), Cz (k=9), CP1 (k=9), and CP2 (k=9).

1 **Table 3. EEG pipeline processing steps.**

Study	Bandpass Filter	Semi-automated trial rejection	Ocular artefact removal	Referencing channels	Trial TOI	LPP TOI	LPP ROI	Sample
Originally submitted pipeline & analyses								
Preregistered	0.5 – 70Hz	Range > 3 SD of trials Absolute maximum > 12 SD trials	ICA	Average reference	-200 – 1000ms	400 – 1000ms	Cz, CP1, CPz, CP2, Pz	MDD: 41 HC: 41
Exploratory cluster	0.5 – 70Hz	Range > 3 SD of trials Absolute maximum > 12 SD trials	ICA	Average reference	-200 – 1000ms	662 – 755ms	CP1, P1, Pz, P2, POz	MDD: 41 HC: 41
Reviewer-suggested pipeline & analyses								
Alternative #1	0.5 – 70Hz	Range > 3 SD of trials Absolute maximum > 12 SD trials	ICA	Linked mastoids	-200 – 1000ms	400 – 1000ms	Cz, CP1, CPz, CP2, Pz	MDD: 41 HC: 41
Alternative #2	0.1 – 70Hz	Range > 3 SD of trials Absolute maximum > 12 SD trials	ICA	Linked mastoids	-200 – 1000ms	400 – 1000ms	Cz, CP1, CPz, CP2, Pz	MDD: 41 HC: 41
Alternative #3	0.1 – 70Hz	Range > 3 SD of trials Absolute maximum > 12 SD trials	ICA	Linked mastoids	-200 – 1000ms	400 – 1000ms	CP1, P5, P3, P1, Pz, PO3	MDD: 41 HC: 41

Alternative #4	0.1 – 70Hz	Range > 3 SD of trials Absolute maximum > 12 SD trials	ICA	Average reference	-200 – 1000ms	400 – 1000ms	Cz, CP1, CPz, CP2, Pz	MDD: 41 HC: 41
Alternative #5	0.1 – 70Hz	Range > 3 SD of trials Absolute maximum > 12 SD trials	ICA	Average reference	-200 – 1000ms	400 – 1000ms	CP1, P5, P3, P1, Pz, PO3	MDD: 41 HC: 41
Representative LPP studies in depression								
Foti et al. (2010); MacNamara et al. (2016)	0.1 – 30Hz	>50μV step between samples >300μV range within trial <0.5μV range within 100ms interval	Regression Method	Linked mastoids	-200 – 1200ms	400 – 1000ms	Cz, CP1, CP2, Pz	MDD: 22; HC: 25 MDD: 36; HC: 23
Shestyuk and Deldin (2010)	0.01 – 30Hz	-	Regression Method[1]	Linked mastoids	-250 – 1200ms	600 – 800ms	FC4	MDD: 17 RMDD: 18 HC: 17
Hilimire et al. (2015)	0.1 – 30Hz	>50μV step between samples >300μV range within trial <0.5μV range within 100ms interval	Regression Method	Average reference	-200 – 1700ms	600 – 800ms	P3, Pz, P4	TRD: 7
Auerbach et al. (2015)	0.1 – 30Hz	>50μV step between samples >300μV range within trial <0.5μV range within 100ms interval	ICA	Average reference	-200 – 1200ms	600 – 1200ms	Fz, FCz, Cz	MDD: 22[*] HC: 28

[1] Proprietary regression method developed by James Long Company software.
[*] Study participants consisted of adolescents.

Weinberg et al. (2016)	0.01 – 30Hz	>50µV step between samples >300µV range within trial <0.5µV range within 100ms interval	Regression Method	Linked mastoids	-200 – 1700ms	400 – 1000ms	CP1, CP2, Pz	MDD: 94 HC: 32
Zhang et al. (2016)	0.01 – 30Hz	> 150µV or < -150µV	Regression Method[2]	Linked mastoids	-200 – 1200ms	430 – 650ms	CP1, CP2, Pz	MDD: 26 HC: 26
Stange et al. (2017b); Stange et al. (2017a)	0.01 – 30Hz	>50µV step between samples >300µV range within trial <0.5µV range within 100ms interval	Regression Method	Linked mastoids	-200 – 1000ms	400 – 1000ms	Cz, CP1, CP2, Pz	ANX/DEP: 39 ANX/DEP: 32
Burkhouse et al. (2017)	0.1 – 30Hz	>50µV step between samples >300µV range within trial <0.5µV range within 100ms interval	Regression Method	Linked mastoids	-100 – 2000ms	400 – 1000ms	P3, Pz, P4, PO3, PO4, O1, Oz, O2	MDD: 18[*] RMDD: 27 HC: 26
Webb et al. (2017)	0.1 – 30Hz	>50µV step between samples >300µV range within trial <0.5µV range within 100ms interval	ICA	Linked mastoids	-200 – 1000ms	600 – 1000ms	Fz, FCz, Cz	MDD: 26 HC: 25
Xie et al. (2018)	0.01 – 30Hz	Range > 200µV	Regression Method[3]	Linked mastoids	-200 – 1200ms	500 – 900ms	P1, Pz, P2	DT: 30 NDT: 30
Buchheim et al. (2018)	0.5 – 90Hz	-	Regression Method	Average reference	-100 – 1000ms	600 – 1000ms	Fz	MDD: 17 HC: 13

[2] Proprietary regression method developed by NeuroScan software.
[*] Study participants consisted of adolescents.

Grunewald et al. (2019)	0.1 – 30Hz	> 100μV or < -100μV <0.5μV range within 100ms interval	ICA	Average reference	-200 – 1500ms	450 – 1500ms	Cz, C4, CP1, CP2, P3, Pz, P4	MDD: 26[*] HC: 26
Benau et al. (2019)	0.01 – 30Hz	>50μV step between samples >300μV range within trial <0.5μV range within 100ms interval	Regression Method	Linked mastoids	-500 – 2000ms	600 – 800ms	CP1, CP2	MDD: 36 HC: 34
MacNamara et al. (2019)	0.01 – 30Hz	>50μV step between samples >300μV range within trial <0.5μV range within 100ms interval	Regression Method	Linked mastoids	-200 – 1000ms	1000 – 2000ms	Cz, CP1, CP2, Pz	MDD: 36 HC: 23
Trotti et al. (2020)	0.1 – 50Hz	> 120μV or < -120μV	ICA	Average reference	-200 – 1000ms	400 – 900ms	FCz, C1, Cz, C2, CPz	BDP: 130 BDNP: 75 HC: 181
Whalen et al. (2020)	0.1 – 30Hz	>50μV step between samples >175μV range within trial <0.5μV range within 100ms interval	Regression Method	Average reference	-200 – 1000ms	600 – 1000ms	O1, Oz, O2	MDD: 99[*] HC: 65

[*] Study participants consisted of adolescents.

3 We repeated our primary MRMM analyses using the pipelines described in Table 3 with the

4 results provided in Supplementary Table S2. In all pipelines where a regression was used to

5 remove the ocular artefact we implemented the procedure as outlined in Gratton et al.

6 (1983), as this was used most often and contained sufficient information for replication.

7 Statistical findings remained qualitatively similar for these alternative analysis approaches.

8 Briefly, all pipelines showed a significant effect of Valence, with the exception of Whalen et

9 al. (2020). Two pipelines resulted in a significant effect of Group: in one pipeline the MDD

10 LPPs were more negative than controls (Hilimire et al., 2015), the other pipeline resulted in

11 the opposite effect of MDD LPPs being more positive than controls (Buchheim et al., 2018),

12 although in both cases depressed LPP amplitude was attenuated. No processing pipeline

13 resulted in a significant Group × Valence interaction effect (all $p > 0.05$). Bootstrapped

14 Cohen's d effect sizes ranged from 0.3 (in the direction of *negative potentiation* in

15 depression) to -0.4 (in the direction of *emotional context insensitivity*); however, these were

16 not statistically significant (Fig. 4).

Figure 4. Effect sizes of 22 alternative EEG processing pipelines. Bootstrapped Cohen's *d*

effect sizes with 95% confidence intervals were calculated for the interaction contrast

between unpleasant and neutral images comparing MDD participants to healthy controls:

[MDD $LPP_{unpleasant}$ − MDD $LPP_{neutral}$] *vs* [Healthy $LPP_{unpleasant}$ − Healthy $LPP_{neutral}$]. Numbered

text to the right of the effect size bar graphs indicates *p*-values from independent samples *t*-

tests comparing the LPP difference score (i.e. $LPP_{unpleasant}$ − $LPP_{neutral}$) between MDD and

control participants. Results are formulated such that a negative Cohen's *d* indicates

support for *emotional context insensitivity*, and a positive Cohen's *d* indicates support for

negative potentiation. AR: Average reference; Localiser: collapsed localiser (Luck and

Gaspelin, 2017); ROI: Region of interest (preregistered LPP channels Cz, CPz, Pz, CP1, and

CP2);

Three additional exploratory non-parametric cluster-based permutation tests compared the

difference scores ($LPP_{unpleasant}$ − $LPP_{neutral}$) between MDD participants and healthy controls

using alternative pipelines (0.5Hz or 0.1Hz high-pass filters and referencing to linked

mastoids; and 0.1Hz high-pass filter using the common average reference). No significant

clusters were identified using these data-driven comparisons (all $ps > 0.27$)

DISCUSSION

We performed a preregistered study to verify the presence of *emotional context insensitivity* in depression by comparing the LPP amplitude for unpleasant and neutral stimuli in MDD participants with age- and gender-matched healthy controls. Our preregistered data analysis plan found no difference in emotional reactivity between MDD participants and controls within the preregistered centroparietal region of interest and time interval. Further re-analysis of the data using 16 different EEG processing pipelines reported in the literature and five additional alternative pipelines also found no evidence of a difference in emotional reactivity between groups. Exploratory permutation analyses only revealed a significant difference between MDD participants and healthy individuals using the preregistered settings. This effect was in the hypothesised direction. The difference in amplitudes between unpleasant pictures relative to neutral pictures was lower for MDD participants compared to healthy controls, in agreement with the *emotional context insensitivity* model of depression. The exploratory results consisted of parietal EEG channels slightly posterior to the preregistered region of interest within a narrower post-stimulus time interval (662-755 ms). We found no evidence that emotion regulation strategy use differed between diagnostic groups, nor that strategy use or anxiety influenced the LPP in a passive view paradigm. Overall, our results provide no or weak evidence in support of *emotional context insensitivity* in our depressed sample and showed that quantification of the LPP in depression strongly depended on EEG processing pipeline parameters.

Evidence in support of emotional context insensitivity in depression arises from attenuated LPP responses to emotionally evocative stimuli (Hajcak and Foti, 2020), and from broader physiological research demonstrating reduced responses to both pleasant and unpleasant

stimuli (Bylsma et al., 2008; Rottenberg et al., 2005; Rottenberg and Hindash, 2015).

However, mixed effects have been reported in some EEG studies, including suggestions of

reduced emotional reactivity to unpleasant stimuli in MDD relative to a healthy cohort (Foti

et al., 2010; MacNamara et al., 2016), increased reactivity (Zhang et al., 2016), and no

difference between unpleasant and neutral stimuli (Weinberg et al., 2016). The aim of the

present study was therefore to obtain additional evidence for *emotional context*

insensitivity in depression through a registered study (Nosek et al., 2018), replicating

observations of reduced LPP to unpleasant stimuli in a large sample of adult MDD

participants. We did not find evidence of a significant reduction in LPP amplitude for

negatively-valenced stimuli using either our preregistered analysis plan or processing

pipelines reported in literature. Similarly, no differences between groups were observed

using a data-driven approach (i.e., cluster permutation testing), except for a statistically

significant reduction in MDD LPP amplitudes using our preregistered EEG processing

parameters (i.e., 0.5Hz high-pass filter and common average reference). This cluster

significantly differentiated diagnostic groups within a narrower time window of the LPP

(662-755 ms) and a slightly more posterior region of interest. Although these findings

matched our *a priori* directional prediction (i.e., low LPP in the MDD group while viewing

unpleasant images compared to healthy controls), they should be interpreted with caution.

Only one exploratory analysis, out of a total of 26 variations reported in the present study,

detected a significant LPP effect for the interaction contrast between groups for unpleasant

and neutral images. While cluster-based permutation tests control the family-wise error

rate (Maris and Oostenveld, 2007; Pernet et al., 2015), simulations conducted using similarly

multidimensional neuroimaging data (obtained using fMRI) note that such procedures may

have low sensitivity (i.e. true positive rate) (Noble et al., 2020). Furthermore, they should

not be used to make inferences about the latency or location of effects (Sassenhagen and

Draschkow, 2019). As such, these exploratory findings, which only marginally exceeded the

significance threshold (p = 0.01), should not be used as conclusive evidence in favour of

emotional context insensitivity in depression.

It is unclear why we did not observe more convincing evidence of attenuated LPP

amplitudes in MDD sample. Potential reasons include the stimuli used to elicit the LPP, and

the passive viewing study design. Stimuli selection differs markedly between studies,

specifically with regards to the selection of aversive, unpleasant stimuli. These have

included negatively-valenced scenes (MacNamara et al., 2016; Weinberg et al., 2016),

affective and self-referential adjectives (Auerbach et al., 2015; Hilimire et al., 2015; Shestyuk

and Deldin, 2010; Xie et al., 2018), and pictures of faces depicting sadness, anger, or fearful

expressions (Foti et al., 2010; Zhang et al., 2016). The latter have been shown to produce a

stronger amygdala response in comparison to matched fearful or threatening scenes (Hariri

et al., 2002), which has important implications for the expected neurophysiological

response measured using EEG (Mavratzakis et al., 2016). In addition to the type of stimulus,

an important consideration is the context in which participants are instructed to interact

with stimuli, either as part of a task, through active viewing of images with directed

attention (e.g. using preselected emotional regulation strategies), or in a passive view

paradigm. Hajcak et al. (2009) showed that instructing participants to attend to either

neutral or arousing features in an image resulted in an attenuated LPP as compared to

passive viewing. This agrees with research suggesting emotion regulation strategy use

reduces LPP amplitudes (Dunning and Hajcak, 2009; Foti and Hajcak, 2008; Hajcak et al.,

2007; Hajcak et al., 2006; Hajcak and Nieuwenhuis, 2006; Moser et al., 2006; Thiruchselvam

et al., 2011; Uusberg et al., 2014; Zhang et al., 2014). Interestingly, in the present study we

did not find supporting evidence that viewing strategies moderated outcomes in our

dataset. An important caveat, however, is that we did not systematically instruct

participants to adopt a particular strategy. Most participants in the current study reported

no emotion regulation strategy use, thus limiting the ability to statistically infer the effects

of strategy on LPP amplitudes.

A key outcome of the present study is that LPP findings may not necessarily generalise

across stimulus processing steps and measurement contexts (Yarkoni, 2019). Sensitivity

analyses performed by re-processing and operationalising the LPP using multiple parameter

variations empirically demonstrate that relatively minor differences can impact effect

estimates. Effect sizes were observed to change both in magnitude and direction depending

on the EEG pipeline used to generate the LPP (Fig. 3). However, we note that there was

considerable overlap in effect size intervals and that statistical results remained qualitatively

unchanged regardless of processing methodology, i.e., none of the pipelines found a

significant group by valence interaction effect. Based on these results it is plausible that

mixed findings of emotional reactivity in the EEG studies described earlier may in part be

caused by variations in pipeline processing steps and definitions of the LPP (see Table 3).

Interpretation of findings within the context of the broader literature is therefore

complicated by differences in the methodology, i.e., researcher degrees of freedom

(Simmons et al., 2011), used to obtain the LPP. This provides a compelling argument for the

development of a standardized EEG measure of emotional reactivity, encompassing the

stimuli set, task protocol, EEG processing pipeline, and statistical analysis procedure used to

assess the LPP. Though beyond the scope of the current manuscript, construction of such a

tool could follow established methods to develop psychometric instruments of behaviour

(Boateng et al., 2018).

Limitations

A potential limitation to our study design is that depressed participants were allowed to be

on stable ongoing antidepressant medications, which may have influenced LPP amplitudes

(Kerestes et al., 2009). The impact of antidepressants on ERP components is unclear, having

been found to both reduce (d'Ardhuy et al., 1999) or increase amplitudes (Ozgocmen et al.,

2003; Sanz et al., 2001). We note that previous studies in depression have used similar

recruitment strategies and have reported no differences in LPP component amplitudes

between medicated and unmedicated participants (Auerbach et al., 2015; MacNamara et

al., 2016; Webb et al., 2017).

Our stimuli set was restricted to negative and neutral pictures, omitting positive images. The

reason for this was to reduce the total duration of the experiment session and minimise

participant burden, as this study was part of a larger battery of cognitive tests. Given the

weight of the evidence in the literature, we sought to replicate research in support of

emotional context insensitivity, which uniquely predicts decreased reactivity in response to

unpleasant stimuli.

Conclusion

We investigated emotional reactivity in a large sample of MDD and age- and gender-

matched controls to obtain additional evidence in support of *emotional context insensitivity*

in depression. Contrary to our hypothesis, we did not find a statistical difference in

emotional reactivity between groups using our preregistered analysis plan or the processing

pipelines of several representative LPP studies in depression. Exploratory cluster analyses

identified a single cluster, which was marginally more posterior and subtended a narrower

time interval (660-755ms) than our preregistered LPP, that revealed a significant difference

between MDD and control participants. However, as this cluster was the only significant

finding of a total of 26 analysis variations, this only provides weak evidence in support of

emotional context insensitivity in depression. Interestingly, EEG processing pipelines were

found to affect both the magnitude and direction of the LPP comparison between MDD and

control participants. These results suggest that methodological differences between studies

(e.g., choice of affective stimuli, EEG processing pipeline, and analysis methods) may play an

important role in observed heterogeneity of the cortical response to emotional stimuli in

depression. This heterogeneity in EEG correlates of affective processing strongly indicates

the need for a standardised EEG test and analysis pipeline for emotional reactivity.

ACKNOWLEDGEMENTS

162 We would like to thank William Flannery, Liyi Tan and Yi Yin Tan for their contributions to

163 EEG acquisitions.

FUNDING

Donel Martin and Tjeerd Boonstra were both funded by a NARSAD Young Investigator

Award (grant numbers 24015 and 26060, respectively) from the

Brain and Behavior Research Foundation.

REFERENCES

American Psychiatric Association, 2013. Diagnostic and statistical manual of mental
disorders (DSM-5®). American Psychiatric Pub.

Auerbach, R.P., Stanton, C.H., Proudfit, G.H., Pizzagalli, D.A., 2015. Self-referential
processing in depressed adolescents: A high-density event-related potential study. Journal
of abnormal psychology 124, 233.

Benau, E.M., Hill, K.E., Atchley, R.A., O'Hare, A.J., Gibson, L.J., Hajcak, G., Ilardi, S.S., Foti, D.,
2019. Increased neural sensitivity to self-relevant stimuli in major depressive disorder.
Psychophysiology 56, e13345.

Boateng, G.O., Neilands, T.B., Frongillo, E.A., Melgar-Quiñonez, H.R., Young, S.L., 2018. Best
practices for developing and validating scales for health, social, and behavioral research: a
primer. Frontiers in public health 6, 149.

Brown, T.A., Chorpita, B.F., Barlow, D.H., 1998. Structural relationships among dimensions
of the DSM-IV anxiety and mood disorders and dimensions of negative affect, positive
affect, and autonomic arousal. Journal of abnormal psychology 107, 179.

Brown, T.A., Chorpita, B.F., Korotitsch, W., Barlow, D.H., 1997. Psychometric properties of
the Depression Anxiety Stress Scales (DASS) in clinical samples. Behaviour research and
therapy 35, 79-89.

Browning, M., Holmes, E.A., Harmer, C.J., 2010. The modification of attentional bias to
emotional information: A review of the techniques, mechanisms, and relevance to
emotional disorders. Cognitive, Affective, & Behavioral Neuroscience 10, 8-20.

Buchheim, A., Labek, K., Taubner, S., Kessler, H., Pokorny, D., Kachele, H., Cierpka, M., Roth,
G., Pogarell, O., Karch, S., 2018. Modulation of gamma band activity and late positive
potential in patients with chronic depression after psychodynamic psychotherapy.
Psychotherapy and psychosomatics 87, 252-255.

Burkhouse, K.L., Owens, M., Feurer, C., Sosoo, E., Kudinova, A., Gibb, B.E., 2017. Increased
neural and pupillary reactivity to emotional faces in adolescents with current and remitted
major depressive disorder. Social Cognitive and Affective Neuroscience 12, 783-792.

Bylsma, L.M., Morris, B.H., Rottenberg, J., 2008. A meta-analysis of emotional reactivity in
major depressive disorder. Clinical psychology review 28, 676-691.

Clark, L.A., Watson, D., 1991. Tripartite model of anxiety and depression: psychometric
evidence and taxonomic implications. Journal of abnormal psychology 100, 316.

Clayson, P.E., Carbine, K.A., Larson, M.J., 2020. A registered report of error-related
negativity and reward positivity as biomarkers of depression: P-Curving the evidence.
International Journal of Psychophysiology 150, 50-72.

Codispoti, M., Ferrari, V., Bradley, M.M., 2006. Repetitive picture processing: autonomic and
cortical correlates. Brain research 1068, 213-220.

Codispoti, M., Ferrari, V., Bradley, M.M., 2007. Repetition and event-related potentials:
distinguishing early and late processes in affective picture perception. J Cogn Neurosci 19,
577-586.

Crawford, J.R., Henry, J.D., 2003. The Depression Anxiety Stress Scales (DASS): Normative
data and latent structure in a large non-clinical sample. British journal of clinical psychology
42, 111-131.

Cuthbert, B.N., Schupp, H.T., Bradley, M.M., Birbaumer, N., Lang, P.J., 2000. Brain potentials
in affective picture processing: covariation with autonomic arousal and affective report.
Biological psychology 52, 95-111.

d'Ardhuy, X.L., Boeijinga, P., Renault, B., Luthringer, R., Rinaudo, G., Soufflet, L., Toussaint,
M., Macher, J.-P., 1999. Effects of serotonin-selective and classical antidepressants on the
auditory P300 cognitive potential. Neuropsychobiology 40, 207-213.

Delorme, A., Sejnowski, T., Makeig, S., 2007. Enhanced detection of artifacts in EEG data
using higher-order statistics and independent component analysis. Neuroimage 34, 1443-
1449.

Dunning, J.P., Hajcak, G., 2009. See no evil: Directing visual attention within unpleasant
images modulates the electrocortical response. Psychophysiology 46, 28-33.

First, M., Williams, J., Karg, R., Spitzer, R., 2015. Structured clinical interview for DSM-5—
Research version (SCID-5 for DSM-5, research version; SCID-5-RV). Arlington, VA: American
Psychiatric Association, 1-94.

Foti, D., Hajcak, G., 2008. Deconstructing reappraisal: descriptions preceding arousing
pictures modulate the subsequent neural response. J Cogn Neurosci 20, 977-988.

Foti, D., Olvet, D.M., Klein, D.N., Hajcak, G., 2010. Reduced electrocortical response to
threatening faces in major depressive disorder. Depression and anxiety 27, 813-820.

Gratton, G., Coles, M.G., Donchin, E., 1983. A new method for off-line removal of ocular
artifact. Electroencephalography and clinical Neurophysiology 55, 468-484.

Gross, J.J., John, O.P., 2003. Individual differences in two emotion regulation processes:
implications for affect, relationships, and well-being. Journal of personality and social
psychology 85, 348.

Grunewald, M., Döhnert, M., Brandeis, D., Klein, A.M., von Klitzing, K., Matuschek, T.,
Stadelmann, S., 2019. Attenuated LPP to emotional face stimuli associated with parent-and
self-reported depression in children and adolescents. Journal of abnormal child psychology
47, 109-118.

Hajcak, G., Dunning, J.P., Foti, D., 2007. Neural response to emotional pictures is unaffected
by concurrent task difficulty: an event-related potential study. Behavioral neuroscience 121,
1156.

Hajcak, G., Dunning, J.P., Foti, D., 2009. Motivated and controlled attention to emotion:
time-course of the late positive potential. Clinical Neurophysiology 120, 505-510.

Hajcak, G., Foti, D., 2020. Significance?... Significance! Empirical, methodological, and
theoretical connections between the late positive potential and P300 as neural responses to
stimulus significance: An integrative review. Psychophysiology, e13570.

Hajcak, G., Moser, J.S., Simons, R.F., 2006. Attending to affect: appraisal strategies modulate
the electrocortical response to arousing pictures. Emotion 6, 517.

Hajcak, G., Nieuwenhuis, S., 2006. Reappraisal modulates the electrocortical response to
unpleasant pictures. Cognitive, Affective, & Behavioral Neuroscience 6, 291-297.

Hariri, A.R., Tessitore, A., Mattay, V.S., Fera, F., Weinberger, D.R., 2002. The amygdala
response to emotional stimuli: a comparison of faces and scenes. Neuroimage 17, 317-323.

Henry, J.D., Crawford, J.R., 2005. The short-form version of the Depression Anxiety Stress
Scales (DASS-21): Construct validity and normative data in a large non-clinical sample.
British journal of clinical psychology 44, 227-239.

Hilimire, M.R., Mayberg, H.S., Holtzheimer, P.E., Broadway, J.M., Parks, N.A., DeVylder, J.E.,
Corballis, P.M., 2015. Effects of subcallosal cingulate deep brain stimulation on negative
self-bias in patients with treatment-resistant depression. Brain Stimulation 8, 185-191.

Hill, K.E., South, S.C., Egan, R.P., Foti, D., 2019. Abnormal emotional reactivity in depression:
Contrasting theoretical models using neurophysiological data. Biological psychology 141, 35-
43.

Hu, S., Lai, Y., Valdes-Sosa, P.A., Bringas-Vega, M.L., Yao, D., 2018. How do reference
montage and electrodes setup affect the measured scalp EEG potentials? Journal of neural
engineering 15, 026013.

Hyvärinen, A., Hoyer, P.O., Inki, M., 2001. Topographic independent component analysis.
Neural Comput 13, 1527-1558.

Kayser, J., Bruder, G.E., Tenke, C.E., Stewart, J.E., Quitkin, F.M., 2000. Event-related
potentials (ERPs) to hemifield presentations of emotional stimuli: differences between
depressed patients and healthy adults in P3 amplitude and asymmetry. Int J Psychophysiol
36, 211-236.

Kerestes, R., Labuschagne, I., Croft, R.J., O'Neill, B.V., Bhagwagar, Z., Phan, K.L., Nathan, P.J.,
2009. Evidence for modulation of facial emotional processing bias during emotional
expression decoding by serotonergic and noradrenergic antidepressants: an event-related
potential (ERP) study. Psychopharmacology 202, 621-634.

Klawohn, J., Burani, K., Bruchnak, A., Santopetro, N., Hajcak, G., 2020. Reduced neural
response to reward and pleasant pictures independently relate to depression. Psychological
Medicine, 1-9.

Lang, P., Bradley, M., Cuthbert, B., 2008. International affective picture system (IAPS):
affective ratings of pictures and instruction manual. University of Florida, Gainesville. Tech
Rep A-8.

Lovibond, P.F., Lovibond, S.H., 1995. The structure of negative emotional states:
Comparison of the Depression Anxiety Stress Scales (DASS) with the Beck Depression and
Anxiety Inventories. Behav Res Ther 33, 335-343.

Luck, S.J., Gaspelin, N., 2017. How to get statistically significant effects in any ERP
experiment (and why you shouldn't). Psychophysiology 54, 146-157.

MacNamara, A., Hajcak, G., 2009. Anxiety and spatial attention moderate the electrocortical
response to aversive pictures. Neuropsychologia 47, 2975-2980.

MacNamara, A., Jackson, T.B., Fitzgerald, J.M., Hajcak, G., Phan, K.L., 2019. Working memory
load and negative picture processing: neural and behavioral associations with panic, social

anxiety, and positive affect. Biological Psychiatry: Cognitive Neuroscience and Neuroimaging
4, 151-159.

MacNamara, A., Kotov, R., Hajcak, G., 2016. Diagnostic and symptom-based predictors of
emotional processing in generalized anxiety disorder and major depressive disorder: An
event-related potential study. Cognitive Therapy and Research 40, 275-289.

Makeig, S., Bell, A.J., Jung, T.-P., Sejnowski, T.J., 1996. Independent component analysis of
electroencephalographic data. Adv Neural Inf Process Syst, 145-151.

Maris, E., Oostenveld, R., 2007. Nonparametric statistical testing of EEG-and MEG-data. J
Neurosci Methods 164, 177-190.

Mavratzakis, A., Herbert, C., Walla, P., 2016. Emotional facial expressions evoke faster
orienting responses, but weaker emotional responses at neural and behavioural levels
compared to scenes: A simultaneous EEG and facial EMG study. Neuroimage 124, 931-946.

Montgomery, S.A., Asberg, M., 1979. A new depression scale designed to be sensitive to
change. Br J Psychiatry 134, 382-389.

Moser, J.S., Hajcak, G., Bukay, E., Simons, R.F., 2006. Intentional modulation of emotional
responding to unpleasant pictures: an ERP study. Psychophysiology 43, 292-296.

Moser, J.S., Krompinger, J.W., Dietz, J., Simons, R.F., 2009. Electrophysiological correlates of
decreasing and increasing emotional responses to unpleasant pictures. Psychophysiology
46, 17-27.

Nikolin, S., Chand, N., Martin, D., Rushby, J., Loo, C.K., Boonstra, T., 2020. EEG correlates of
affective processing in major depressive disorder. OSF.

Noble, S., Scheinost, D., Constable, R.T., 2020. Cluster failure or power failure? Evaluating
sensitivity in cluster-level inference. Neuroimage 209, 116468.

Nosek, B.A., Ebersole, C.R., DeHaven, A.C., Mellor, D.T., 2018. The preregistration
revolution. Proceedings of the National Academy of Sciences 115, 2600-2606.

Oliver-Rodriguez, J.C., Guan, Z., Johnston, V.S., 1999. Gender differences in late positive
components evoked by human faces. Psychophysiology 36, 176-185.

Oostenveld, R., Fries, P., Maris, E., Schoffelen, J.-M., 2011. FieldTrip: open source software
for advanced analysis of MEG, EEG, and invasive electrophysiological data. Comput Intell
Neurosci 2011, 1.

Ozgocmen, S., Yoldas, T., Kamanli, A., Yildizhan, H., Yigiter, R., Ardicoglu, O., 2003. Auditory
P300 event related potentials and serotonin reuptake inhibitor treatment in patients with
fibromyalgia. Annals of the rheumatic diseases 62, 551-555.

Pernet, C., Latinus, M., Nichols, T., Rousselet, G., 2015. Cluster-based computational
methods for mass univariate analyses of event-related brain potentials/fields: A simulation
study. J Neurosci Methods 250, 85-93.

Renfroe, J.B., Bradley, M.M., Sege, C.T., Bowers, D., 2016. Emotional modulation of the late
positive potential during picture free viewing in older and young adults. PloS one 11.

Rottenberg, J., 2005. Mood and emotion in major depression. Current Directions in
Psychological Science 14, 167-170.

Rottenberg, J., Gross, J.J., Gotlib, I.H., 2005. Emotion context insensitivity in major
depressive disorder. Journal of abnormal psychology 114, 627.

Rottenberg, J., Hindash, A.C., 2015. Emerging evidence for emotion context insensitivity in
depression. Current Opinion in Psychology 4, 1-5.

Sanz, M., Molina, V., Martin-Loeches, M., Calcedo, A., Rubia, F.J., 2001. Auditory P300 event
related potential and serotonin reuptake inhibitor treatment in obsessive-compulsive
disorder patients. Psychiatry research 101, 75-81.

Sassenhagen, J., Draschkow, D., 2019. Cluster-based permutation tests of MEG/EEG data do
not establish significance of effect latency or location. Psychophysiology 56, e13335.

Schupp, H.T., Cuthbert, B.N., Bradley, M.M., Cacioppo, J.T., Ito, T., Lang, P.J., 2000. Affective
picture processing: the late positive potential is modulated by motivational relevance.
Psychophysiology 37, 257-261.

Shestyuk, A.Y., Deldin, P.J., 2010. Automatic and strategic representation of the self in major
depression: trait and state abnormalities. American Journal of Psychiatry 167, 536-544.

Simmons, J.P., Nelson, L.D., Simonsohn, U., 2011. False-positive psychology: Undisclosed
flexibility in data collection and analysis allows presenting anything as significant.
Psychological science 22, 1359-1366.

Stange, J.P., MacNamara, A., Barnas, O., Kennedy, A.E., Hajcak, G., Phan, K.L., Klumpp, H.,
2017a. Neural markers of attention to aversive pictures predict response to cognitive
behavioral therapy in anxiety and depression. Biological psychology 123, 269-277.

Stange, J.P., MacNamara, A., Kennedy, A.E., Hajcak, G., Phan, K.L., Klumpp, H., 2017b. Brain-
behavioral adaptability predicts response to cognitive behavioral therapy for emotional
disorders: A person-centered event-related potential study. Neuropsychologia.

Thiruchselvam, R., Blechert, J., Sheppes, G., Rydstrom, A., Gross, J.J., 2011. The temporal
dynamics of emotion regulation: An EEG study of distraction and reappraisal. Biological
psychology 87, 84-92.

Trotti, R.L., Parker, D.A., Sabatinelli, D., Tamminga, C.A., Gershon, E.S., Keedy, S.K.,
Keshavan, M.S., Pearlson, G.D., Sweeney, J.A., McDowell, J.E., 2020. Electrophysiological
correlates of emotional scene processing in bipolar disorder. Journal of psychiatric research
120, 83-90.

Uusberg, A., Thiruchselvam, R., Gross, J.J., 2014. Using distraction to regulate emotion:
Insights from EEG theta dynamics. International Journal of Psychophysiology 91, 254-260.

Watson, D., 2000. Mood and temperament. Guilford Press.

Webb, C.A., Auerbach, R.P., Bondy, E., Stanton, C.H., Foti, D., Pizzagalli, D.A., 2017.
Abnormal neural responses to feedback in depressed adolescents. Journal of abnormal
psychology 126, 19.

Weinberg, A., Perlman, G., Kotov, R., Hajcak, G., 2016. Depression and reduced neural
response to emotional images: Distinction from anxiety, and importance of symptom
dimensions and age of onset. Journal of abnormal psychology 125, 26.

Whalen, D.J., Gilbert, K.E., Kelly, D., Hajcak, G., Kappenman, E.S., Luby, J.L., Barch, D.M.,
2020. Preschool-onset major depressive disorder is characterized by electrocortical deficits
in processing pleasant emotional pictures. Journal of abnormal child psychology 48, 91-108.

Xie, H., Jiang, D., Zhang, D., 2018. Individuals with depressive tendencies experience
difficulty in forgetting negative material: two mechanisms revealed by ERP data in the
directed forgetting paradigm. Scientific reports 8, 1-14.

Yarkoni, T., 2019. The generalizability crisis.

Zhang, D., He, Z., Chen, Y., Wei, Z., 2016. Deficits of unconscious emotional processing in
patients with major depression: an ERP study. Journal of affective disorders 199, 13-20.

Zhang, W., Li, H., Chen, J., Chen, N., Liu, X., Wang, D., Shen, J., 2014. Posterior P1 and early
frontal negativity reflect developmental changes in attentional distraction during
adolescence. Brain and cognition 87, 30-38.