EEG Correlates of Affective Processing in Major Depressive Disorder

Stevan Nikolin[1,2], Nicholas Chand[2], Donel Martin[1,2], Jacqueline Rushby[3], Colleen K. Loo[1,2,4], Tjeerd W. Boonstra[1,5]

[1] School of Psychiatry, University of New South Wales, Sydney, Australia

[2] Black Dog Institute, Sydney, Australia

[3] School of Psychology, University of New South Wales, Sydney, Australia

[4] St. George Hospital, Sydney, Australia

[5] Faculty of Psychology and Neuroscience, Maastricht University, Maastricht, Netherlands

Acknowledgements: We would like to thank William Flannery, Liyi Tan and Yi Yin Tan for their contributions to EEG acquisitions.

Keywords: Depression, Emotional Regulation, Electroencephalography, Event-Related Potentials, Late Positive Potential

Running title: Affective Processing in Depression

ABSTRACT

Background: Individuals with major depressive disorder present with deficits in emotional reactivity. Conflicting models have been proposed to characterise the direction of this effect. Previous studies mostly support the *emotional context insensitivity* model, which suggests that reactivity to positive and negatively-valenced emotional stimuli is blunted in depression. We sought to test the *emotional context insensitivity* hypothesis in a preregistered study.

Methods: Forty-one depressed participants and 41 age- and gender-matched healthy controls were shown a series of unpleasant and neutrally-valenced pictures in a passive view paradigm while acquiring electroencephalography (EEG). The late positive potential (LPP), an EEG marker of emotional reactivity, was compared between groups using mixed-effects repeated-measures models and exploratory cluster-based permutation tests.

Results: We found no difference in LPP amplitudes between MDD and healthy individuals using the preregistered analysis plan. However, exploratory permutation analysis revealed a significant reduction in the LPP for MDD participants while viewing unpleasant pictures in EEG electrodes marginally more posterior and within a narrower time interval than specified in our original analysis plan. Secondary analyses found that emotion regulation strategy use and anxiety comorbidity did not influence the LPP.

Conclusions: We found significantly reduced LPP amplitudes to unpleasant pictures in MDD, in support of the *emotional context insensitivity* hypothesis. However, the parameters used to estimate the LPP deviated from the preregistered analysis plan, suggesting that methodological differences between studies may result in variability in the cortical response to emotional stimuli.

INTRODUCTION

Major depressive disorder (MDD) is a mental illness that presents as persistent low mood, loss of interest in pleasurable activities, or both, in conjunction with other symptoms (American Psychiatric Association, 2013). Consistent with the core symptom of disturbed mood, depressed individuals report altered reactivity to emotion-eliciting contexts of positive or negative affect (Rottenberg et al., 2005). Although the terms mood and emotion are sometimes used interchangeably, mood typically refers to slow-moving affective states that are weakly influenced by external stimuli, whereas emotions are considered to be rapid reactions to meaningful affective stimuli (Watson, 2000). The two are deeply intertwined and an improved understanding of the emotional deficits in MDD may thus provide further insight into mood symptoms (Rottenberg, 2005). Currently, three competing models have been proposed to describe the observed abnormalities in emotional reactivity in depressed individuals. The oldest of these, the *positive attenuation* model, suggests that the physiological response is diminished to positive stimuli and unchanged for unpleasant, negatively-valenced stimuli (Clark & Watson, 1991). The *emotional context insensitivity* model argues that depression results in blunted emotional reactivity to positive stimuli as well as to negative emotional contexts (Rottenberg et al., 2005). The most recently proposed *negative potentiation* hypothesis states that depression increases reactivity towards negative contexts (Browning et al., 2010). These theories disagree with regards to the physiological response in MDD to negative stimuli: *positive attenuation* predicts no difference in reactivity compared to healthy individuals, *negative potentiation* predicts increased reactivity, and the *emotional context insensitivity* predicts decreased reactivity in depression.

Emotional reactivity is commonly assessed using electroencephalography (EEG). Specifically,

the late positive potential (LPP) may provide a reliable measure of affective cortical

processing (Codispoti et al., 2006, 2007). The LPP is characterized as the mean amplitude of

the event-related potential (ERP) within a time window approximately 400-1000ms

following the visual presentation of emotionally laden images (Cuthbert et al., 2000; Schupp

et al., 2000). EEG studies in depressed individuals have shown mixed findings of the LPP

amplitude and do not provide conclusive support in favour of any current model of

emotional reactivity in depression. Two studies have demonstrated attenuated LPP

amplitudes in response to unpleasant stimuli, lending support to the *emotional context*

insensitivity hypothesis (Foti et al., 2010; MacNamara et al., 2016). In contrast, Zhang et al.

(2016) found a significant increase in LPP amplitudes while passively observing pictures of

faces showing sad emotional expressions as compared to neutral expressions, in agreement

with the *negative potentiation* hypothesis. Lastly, Weinberg et al. (2016) did not find a

significant difference in LPP amplitudes comparing healthy controls to depressed

participants. Although most studies on emotional reactivity reveal attenuation of LPP

amplitudes in depressed individuals, in agreement with the *emotional context insensitivity*

hypothesis, the evidence is not conclusive.

The primary aim of this study was therefore to test the *emotional context insensitivity*

hypothesis by comparing the contrast in LPP amplitudes for unpleasant *vs* neutral pictures

between MDD participants and healthy controls. As previous studies (Foti et al., 2010;

MacNamara et al., 2016; Weinberg et al., 2016; Zhang et al., 2016) did not preregister their

study protocols and should hence be considered preliminary, we sought to test the

difference in emotional reactivity in MDD participants in a preregistered study. We

hypothesised that the LPP amplitude difference (i.e. unpleasant *vs* neutral) would be

attenuated in MDD participants in support of the *emotional context insensitivity* model. In

addition, studies examining LPP amplitudes in response to emotional images did not

investigate the role of emotion regulation in depression. In healthy individuals, the LPP is

sensitive to the use of emotion regulation strategies, which refers to conscious or

subconscious efforts to exert control over one's emotional experience (Gross & John, 2003).

Differences in the frequency or type of emotion regulation strategy use could partly explain

the observed differences in the LPP to unpleasant images in MDD participants compared to

healthy controls (Foti & Hajcak, 2008; Hajcak & Nieuwenhuis, 2006; Moser et al., 2009). In

addition to emotion regulation strategy, anxiety, a common comorbidity of depression

(Brown et al., 1998), has also been found to modulate the LPP. Compared to neutral

pictures, MacNamara and Hajcak (2009) showed enhanced LPP amplitudes to aversive

pictures in individuals with high self-reported anxiety. Anxiety in depressed individuals may

therefore explain the conflicting evidence supporting hypotheses of emotional reactivity in

MDD. Therefore, the secondary aims of this study were to determine whether the

frequency of emotion regulation strategy use differs between MDD and healthy participants

and whether emotion regulation strategy use and comorbid anxiety in MDD participants

modulates the LPP amplitude in response to unpleasant pictures. We based the research

protocol that we used to test our hypotheses – the required sample size and the EEG pre-

processing steps to quantify the LPP amplitude – on previous literature, and documented

this in a preregistration form (Nikolin et al., 2020). We anticipated that MDD participants

would use emotion regulation strategies less frequently than healthy controls, and that LPP

amplitude would be moderated by emotion regulation strategy use and anxiety comorbidity

in MDD participants.

MATERIALS AND METHODS

Participants

A total of 82 participants were recruited in this study, of whom 41 were currently depressed

and 41 were healthy adults with no current or past diagnosis of a psychiatric illness. Power

analysis shows that this sample is sufficient to detect moderate effect sizes of Cohen's d =

0.6 (using β = 80% and α = 0.05).

Inclusion criteria shared by both MDD participants and healthy controls were: aged 18 years

and above, able to provide informed consent, normal or corrected-to-normal vision, no

recent serious head injury in the last 12 months (e.g. loss of consciousness of more than 30

minutes), no current or past neurological condition (e.g. stroke, epilepsy), no history of drug

or alcohol dependence in the last three months, not pregnant or possibly pregnant, no past

history of distress reaction to affective images, and an ability to cooperate with the EEG

procedure (e.g. no tremor).

Depressed participants were recruited from those seeking entry into clinical treatment trials

at the Black Dog Institute, Sydney Australia. Healthy controls were recruited via the

University of New South Wales (UNSW) School of Psychology Paid Sign-Up System, Black

Dog Institute Volunteer Research Register, or from flyers distributed in the local area.

Healthy controls were age- and gender-matched to MDD participants since these factors

have previously been shown to influence the latency, amplitude, and topography of ERP

components (Kayser et al., 2000; Oliver-Rodriguez et al., 1999; Renfroe et al., 2016). The

study protocol was approved by the UNSW Human Research Ethics Committee (HC16617).

Procedure

Participants were instructed to passively view and attend to picture stimuli. Pictures were selected to have valence and arousal ratings consistent with those used in previous EEG studies investigating LPP effects (Hajcak & Nieuwenhuis, 2006; MacNamara et al., 2016; Moser et al., 2006). A total of 168 pictures were selected from the International Affective Picture System (IAPS; Lang et al. (2008)). Of these, 84 were unpleasant (e.g. threatening scenes, violence, weapons) and 84 were neutral (e.g. neutral faces, household objects, landscapes). These pictures were chosen from the IAPS based on published valence and arousal ratings. Valence ratings were lower for unpleasant pictures (2.78 ± 0.54) compared to neutral pictures (5.01 ± 0.21), indicating that they elicited more negative emotional affect. Arousal ratings were higher for unpleasant pictures (5.80 ± 0.61) compared to neutral pictures (3.04 ± 0.52). The pictures were divided into three parallel lists consisting of 56 images each (28 unpleasant and 28 neutral) with similar valence and arousal ratings (see Supplementary Materials for further details).

The stimuli were presented on a 58.4 cm (23-inch) computer screen in a dimly lit, sound-attenuated room using Inquisit 4 software (Version 4, Millisecond Software). Digitised pictures sized 20×30 cm were presented on a black background, consistent with previous EEG studies (Dunning & Hajcak, 2009; MacNamara et al., 2016). The pictures were displayed for 5 s in a pre-determined block-randomised order so participants were unable to anticipate the valence of each picture. A black screen with a centered white fixation cross was presented for 2 s between stimuli (Fig. 1). Pictures of the same valence were not presented more than three times consecutively to prevent participants from generating expectations of upcoming stimuli and to avoid over-saturation of emotional salience.

Participants were not informed of any emotional regulation strategies prior to viewing the

IAPS images.

Figure 1. Study protocol. Participants passively observed a total of 56 picture stimuli. These

were presented in a block-randomised order and consisted of 28 unpleasant images with a

negative valence, and 28 neutral images with no emotional valence, sourced from the

International Affective Picture System (IAPS). Pictures were displayed for 5 s, followed by a 2

s fixation cross. A sample of two trials, showing a neutral image (a towel) and an unpleasant

image (a gun), is displayed.

Behavioural measures

Prior to the experimental session, MDD participants had their mood assessed by a trained

rater using the ten-item Montgomery-Asberg Depression Rating Scale (MADRS) for

depressive symptoms (Montgomery & Asberg, 1979). All participants completed the 21-item

self-reported Depression, Anxiety and Stress Scale (DASS; Lovibond and Lovibond (1995)).

The DASS was chosen as it reliably discriminates between depression, anxiety, and stress

states (Crawford & Henry, 2003). Upon completion of the experimental session, we used an

emotion regulation questionnaire adapted from Gross et al. (2003), provided in

Supplementary Materials. Participants were able to select from five emotion regulation

strategies, including no strategy, reinterpretation, distraction, detachment, and 'other' in

which participants could specify their strategy. Strategies specified in the 'other' category

were re-assigned to one of the former strategy options by study personnel.

Electroencephalography data acquisition and processing

EEG data were acquired using a 64-channel setup and a TMSi Refa amplifier (TMS

International, Oldenzaal, Netherlands) with a sampling rate of 1024 Hz. Offline EEG

processing and analysis was conducted using custom-developed MATLAB scripts (v.R2019a;

MathWorks) and the Fieldtrip toolbox (Oostenveld et al., 2011). All scripts used for EEG

processing and calculation of neurophysiological measures are available at the following

link: https://github.com/snikolin/MDDvsCTRL_IAPS.

EEG data was pre-processed using a Butterworth IIR digital filter to remove 50 Hz electrical

line noise. Following this, a second-order band-pass filter with cut-offs of 0.5 and 70 Hz was

used to remove low and high frequency noise generated from head movements and muscle

activity. The EEG data were then epoched into 6 s intervals, beginning 1 s before each

picture onset and continuing for 5 s after picture onset. Epochs were rejected firstly using a

semi-automated procedure, and then on visual inspection to identify any remaining trials

with large artefacts. Smaller, non-cortical physiological activity (e.g. cardiac, muscle, or

ocular) and non-physiological activity (e.g. environmental noise, or movement) were

removed using an Independent Components Analysis (ICA) algorithm (Delorme et al., 2007;

Hyvärinen et al., 2001; Makeig et al., 1996). Finally, EEG data was re-referenced to the

common average reference.

Event-related potentials

ERPs were constructed by separately averaging trials for neutral and unpleasant pictures

time-locked to picture presentation. Each averaged ERP was baseline corrected to the mean

activity occurring in the 200 ms time interval prior to stimulus onset. The late positive

potential (LPP) is a slow, positive wave occurring maximally at centroparietal recording sites,

beginning approximately 400 ms after stimulus onset and lasting several seconds (Cuthbert

et al., 2000; Schupp et al., 2000). We quantified the LPP component as the mean amplitude

of the ERP from 400 to 1000 ms following stimulus onset at a centroparietal cluster of EEG

channels (Cz, CPz, Pz, CP1, and CP2), where it has previously been shown to be maximal

(Foti et al., 2010; MacNamara et al., 2016). Participants with LPP component amplitudes

greater than three standard deviations from the mean, per diagnostic group, were

considered outliers and excluded from statistical analyses.

Statistical Analyses

All analyses were performed using SPSS software (IBM SPSS Statistics 25, SPSS Inc.), and

were in accordance with our preregistered protocol (Nikolin et al., 2020). For the primary

analysis, LPPs were investigated using a mixed-effects repeated-measures model (MRMM).

The model included a repeated factor of Valence (unpleasant vs neutral) using an

unstructured covariance matrix, as well as the between-subjects factor of Group (MDD vs

healthy controls), and the Valence × Group interaction. Subject was included as a random

effect. Residuals were visually inspected to ensure an adequate fit of the model. Cohen's *d*

was calculated to estimate the effect size for the LPP amplitude contrast between

unpleasant and neutral images comparing MDD participants to healthy controls:

[MDD LPP$_{unpleasant}$ − MDD LPP$_{neutral}$] *vs* [Healthy LPP$_{unpleasant}$ − Healthy LPP$_{neutral}$]

A follow-up exploratory non-parametric cluster-based permutation test was performed to

identify any significant differences between MDD participants and healthy controls beyond

the *a priori* time interval and channels operationalised for the LPP. This method controls for

multiple comparisons while comparing differences across a large spatiotemporal parameter

space (Maris & Oostenveld, 2007). Permutation testing was performed across all EEG

channels within the time interval 0 – 1000 ms following presentation of IAPS stimuli.

Participants' ERPs were permuted 3000 times, and the resulting distributions were

compared using independent samples *t*-tests. A value of α < 0.05 was adopted as the two-

tailed significance threshold. Statistically significant clusters were required to comprise at

least two neighbouring channels.

For secondary analyses, a Fisher's Exact Test was used to investigate whether there were

differences in emotion strategy regulation use between diagnostic groups. Emotion

regulation strategies were entered into a 2 x 2 contingency table using the binary variables

of Strategy (no strategy vs strategy) and Group (MDD vs healthy controls). Acceptable

strategies consisted of reinterpretation, distraction, or detachment.

We performed simple univariate linear regression analyses to confirm whether emotion

regulation strategy or comorbid anxiety had an effect on LPP amplitudes. LPP amplitude was

included as the dependent variable, with emotion regulation strategy or comorbid anxiety

entered separately as independent variables. Strategy was included as a binary categorical

variable (no strategy vs strategy). Anxiety was included as a continuous variable using the

anxiety subscore of the DASS. These analyses were initially restricted to LPP amplitudes

obtained from MDD participants while viewing unpleasant images. Additional confirmatory

multivariate linear regression analyses were then performed using the full dataset of MDD

and healthy participants, including both unpleasant and neutral pictures. Independent

variables in these regressions included Valence (unpleasant vs neutral), Group (MDD vs

healthy controls), as well as Strategy or Anxiety as described previously for univariate

analyses.

Scripts for EEG analyses in MATLAB and statistical analyses using SPSS are provided at:

https://github.com/snikolin/MDDvsCTRL_IAPS.

RESULTS

Behavioural and EEG measures were assessed in 82 participants (41 with MDD and 41 aged-
and gender-matched controls). Table 1 shows baseline demographic and clinical information
for MDD participants and healthy controls. One participant was excluded from EEG analyses
due to values greater than three standard deviations from the group mean, leaving a total
of 40 MDD participants and 41 healthy controls.

Table 1. Baseline demographic and clinical information. Means and standard deviations
(SD) are shown for continuous measures, and frequency counts for categorical measures.

	MDD	Controls
n	41	41
Age (years)	46.3 (13.2)	46.0 (13.9)
Education (years)	15.8 (3.6)	17.3 (3.6)
Gender (females/males)	15/26	15/26
TRD (yes/no)	32/9	-
Concurrent antidepressants (yes/no)	31/9	-
MADRS	29.6 (5.0)	-
DASS Depression	33.7 (7.1)	4.2 (5.2)
DASS Anxiety	11.0 (9.7)	3.3 (4.5)
DASS Stress	21.3 (9.5)	6.8 (5.5)
DASS Total	66.5 (19.7)	14.2 (12.9)

MDD: Major Depressive Disorder; TRD: Treatment-Resistant Depression – Defined as failure to respond to at least two
adequate courses of antidepressant medication; MADRS: Montgomery-Asperg Depression Rating Scale (Montgomery et
al., 1979); DASS: 21-item Depression, Anxiety and Stress Scale (Lovibond et al., 1995). DASS scores were not available for
two MDD participants.

Primary Analysis

The LPP amplitude was compared across groups (MDD and controls) and valence conditions

(unpleasant and neutral). MRMM results showed a significant main effect of Valence ($F =$

19.62, $p < 0.001$), indicating increased LPP amplitude for unpleasant compared to neutral

images (Fig. 2). The main effect of Group ($F = 0.39$, $p = 0.54$) and the Group $\times$ Valence

interaction ($F = 0.26$, $p = 0.61$) were not significant. Comparison of the valence difference

between MDD participants and healthy controls showed a small effect size ($d = 0.11$).

Figure 2. Preregistered EEG measures of affective processing. The LPP was quantified as

the mean EEG amplitude in the time interval from 400 to 1000 ms following stimulus onset

for a cluster of centroparietal EEG channels (highlighted in red). ERPs, spatial topographies,

and scatter plots of LPP amplitude are presented for MDD participants and controls while

viewing unpleasant and neutral images. ERPs are displayed with bootstrapped 95%

confidence intervals and with the time interval used to calculate LPPs shaded in light grey.

Spatial topographies are of average ERP amplitude from 400 to 1000 ms. Scatter plots show

the mean (black line), median (red line), and the 95% confidence interval (dark grey region).

A) ERPs and topographies for MDD participants. **B)** ERPs and topographies for healthy

controls. **C)** ERPs and spatial topographies of the valence difference (unpleasant – neutral)

for MDD participants and healthy controls. **D)** Scatter plots of LPP amplitude for unpleasant

(U) and neutral (N) images at the centroparietal cluster. **E)** Spatial topography of the valence

difference (unpleasant – neutral) contrasting MDD participants with healthy controls.

Exploratory non-parametric cluster-based permutation testing compared the difference

scores (unpleasant – neutral stimuli) between MDD participants and healthy controls. These

revealed a significant cluster (Fig. 3). The time interval for the significant cluster fell within

the preregistered LPP window (662 – 755 ms). EEG channels identified in this cluster were

slightly more posterior than those for our preregistered LPP analysis and included CPz, CP1,

Pz, POz, P1, and P2. Three of these channels are also found in the preregistered cluster (CPz,

CP1, and Pz). See Supplementary Material for further details regarding cluster channels.

Repeating the MRMM using the EEG channels and time interval identified in the exploratory

cluster revealed a main effect of Valence (F = 9.55, p = 0.003) and a significant Group ×

Valence interaction effect (F = 15.59, p < 0.001). The main effect of Group was not

significant (F = 0.56, p = 0.46). Post-hoc independent *t*-tests found that the MDD group had

lower LPPs while viewing unpleasant images compared to healthy controls (p = 0.02), but

there was no difference for neutral images (p = 0.23). Comparison of the valence differences

between MDD participants and healthy controls showed a large effect size of Cohen's *d* =

0.88.

Figure 3. Exploratory EEG measures of affective processing. Exploratory cluster-based permutation testing revealed a significant cluster from 662 to 755 ms in EEG channels CPz, CP1, Pz, POz, P1, and P2 (highlighted in red). ERPs, spatial topographies, and scatter plots of LPP amplitude are presented for MDD participants and controls while viewing unpleasant and neutral images. ERPs are displayed with bootstrapped 95% confidence intervals and with the time interval used to calculate the mean amplitude within the exploratory cluster

shaded in light grey. Spatial topographies are of average ERP amplitude from 662 to 755 ms.

Scatter plots show the mean (black line), median (red line), and the 95% confidence interval

(dark grey region). **A)** ERPs and topographies for MDD participants. **B)** ERPs and

topographies for healthy controls. **C)** ERPs and spatial topographies of the valence

difference (unpleasant – neutral) for MDD participants and healthy controls. **D)** Scatter plots

of LPP amplitude for unpleasant (U) and neutral (N) images at the exploratory cluster. **E)**

Spatial topography of the valence difference (unpleasant – neutral) for the time interval 662

– 755 ms contrasting MDD participants with healthy controls.

Secondary Analyses

As a secondary analysis, we compared the emotion regulation strategies used by MDD

participants and control. The majority of participants reported that they did not use a

strategy (Table 2). A Fisher's Exact Test revealed that strategy use was not significantly

different between both groups (p = 0.34).

Table 2. Emotion regulation strategy use.

	MDD (*n* = 40)	Controls (*n* = 41)
No strategy	30	26
Strategy	10	15
Reinterpretation	1	5
Distraction	2	5
Detachment	7	5

MDD: Major Depressive Disorder

We also investigated the association between strategy use and anxiety on the LPP

amplitude. We performed simple univariate linear regression analyses, restricting the

dataset to LPP values obtained from MDD participants while viewing unpleasant images.

Strategy was not significantly associated with LPP amplitudes ($r = 0.13$, $r^2 = 0.02$, $\beta = -0.18$, p

$= 0.44$; Supp Fig. 1A). Similarly, Anxiety was not significantly associated with the LPP ($r =$

0.21, $r^2 = 0.05$, $\beta = 0.01$, $p = 0.20$; Supp Fig. 1B). Next, we performed multivariate linear

regression analyses using the full dataset (incorporating healthy controls as well as neutral

images), including Group and Valence as independent variables. Neither the regression

model including Strategy ($r = 0.18$, $r^2 = 0.03$, $p = 0.17$), nor the model including Anxiety ($r =$

0.17, $r^2 = 0.03$, $p = 0.20$), met the significance threshold.

Univariate regression analyses were repeated for the exploratory cluster identified during

permutation testing with similar results. Neither Strategy ($r = 0.02$, $r^2 = 0.00$, $\beta = -0.04$, $p =$

0.90) nor Anxiety ($r = 0.05$, $r^2 = 0.00$, $\beta = -0.00$, $p = 0.76$) were significantly associated with

LPP amplitudes. After expanding the dataset to incorporate healthy controls and neutral

images, and including Group and Valence as independent variables, neither the model

including Strategy ($r = 0.14$, $r^2 = 0.02$, $p = 0.39$), nor Anxiety ($r = 0.21$, $r^2 = 0.04$, $p = 0.08$),

were significant.

DISCUSSION

We performed a preregistered study to test the *emotional context insensitivity* model by comparing the LPP amplitude for unpleasant and neutral stimuli in depressed participants and age- and gender-matched healthy controls. Our preregistered data analysis plan found no difference in emotional reactivity between MDD participants and controls within the preregistered centroparietal region of interest and time interval, which were based on previous studies. However, exploratory permutation analysis revealed a significant difference between MDD participants and healthy individuals in the hypothesised direction. The difference in amplitudes between unpleasant pictures relative to neutral pictures was lower for MDD participants compared to healthy controls, in agreement with the *emotional context insensitivity* model of depression. The exploratory results revealed a significant cluster of parietal EEG channels slightly posterior to the preregistered region of interest and within a narrower post-stimulus time interval (662-755 ms). We also investigated the effect of strategy use and anxiety comorbidity but found no evidence that emotion regulation strategy use differed between diagnostic groups, nor that strategy use or anxiety influenced the LPP in a passive view paradigm.

The primary aim of the study was to obtain stronger evidence for the *emotional context insensitivity* model through registered predictions (Nosek et al., 2018), and to replicate the reduced LPP to unpleasant stimuli in depression. We found a large reduction (Cohen's *d* = 0.88) in LPP amplitude in the predicted direction for negatively-valenced stimuli. However, the reduction was not statistically significant when using the preregistered analysis plan to quantify the LPP (i.e. mean ERP amplitude in a time interval of 400-1000ms following simulation presentation in a cluster of predefined EEG electrodes). The predicted reduction

in LPP amplitude was only observed when using a narrower time window (662-755 ms) and

slightly more posterior EEG electrodes, which were determined using a data-driven

approach. Ledgerwood (2018) made a distinction between two types of preregistration:

preregistering a theoretical, *a priori*, directional prediction to ensure a study tests (rather

than informs) a theory and preregistering an analysis plan to help control type I error. The

current findings matched the *a priori* directional prediction (i.e. lower LPP in the MDD group

while viewing unpleasant images compared to healthy controls) and as such provides

evidence supporting the *emotional context insensitivity* model of depression. While the

predicted difference was not observed with the preregistered analysis plan, we used a

cluster-based permutation test that controls the family-wise error rate (Maris et al., 2007;

Pernet et al., 2015). Cluster-based permutation tests should not be used to make inferences

about the latency or location of effects (Sassenhagen & Draschkow, 2019), but the current

results show that the temporal and spatial extent of the LPP – as outlined in the

preregistered analysis plan and based on previous literature – may not necessarily

generalise across stimulus steps and measurement contexts (Yarkoni, 2019).

Relatively few EEG studies have investigated the emotional reactivity of depressed

individuals (Foti et al., 2010; MacNamara et al., 2016; Weinberg et al., 2016; Zhang et al.,

2016). These have reported mixed effects, including studies suggesting reduced reactivity to

unpleasant stimuli in MDD relative to a healthy cohort (Foti et al., 2010; MacNamara et al.,

2016), increased reactivity (Zhang et al., 2016), and no difference between unpleasant and

neutral stimuli (Weinberg et al., 2016). The discrepancy observed in the present study

between preregistered and exploratory analyses, as well as in outcomes between previously

mentioned studies, suggests that relatively minor differences in the protocol used to

generate and operationalise the LPP may contribute to mixed findings. These include the

selection of EEG electrodes for analysis, the LPP time interval, as well as other

methodological choices such as the set of images used to evoke a physiological response.

The set of images used to generate the LPP differs markedly between studies, specifically

with regards to the selection of aversive, unpleasant stimuli. Zhang et al. (2016) and Foti et

al. (2010) used pictures of faces bearing emotional expressions. The former using faces

depicting sadness, whereas the latter displayed angry and fearful expressions. In contrast,

Weinberg et al. (2016) and MacNamara et al. (2016) chose their stimuli from the IAPS,

which contains a broader array of picture options (e.g. photographs of people, faces,

everyday objects, and scenes). However, even working from the same image set, the choice

of unpleasant stimuli was substantially different between these studies. Weinberg et al.

(2016) selected threatening images, defined as "*images of mutilation, death, and animal*

and human threat", whereas MacNamara et al. (2016) used less restrictive criteria to pick

their negatively-valenced content. Although there is overlap in the brain regions used to

process faces and emotionally evocative scenes, specialised regions of the brain activate in

response to human faces, i.e. both image types activate the visual cortex, amygdala,

posterior hippocampus, and ventromedial prefrontal cortex, whereas only images of

expressive faces activated the superior temporal gyrus, insula and anterior cingulate

(Britton et al., 2006). Furthermore, angry or fearful facial expressions have been shown to

produce a stronger amygdala response in comparison to matched fearful or threatening

scenes (Hariri et al., 2002). Consequently, this has important implications for the expected

neurophysiological response measured using EEG, which may differ between these image

types (Mavratzakis et al., 2016).

Regarding the EEG electrodes preselected for measuring the LPP, the exploratory cluster

identified in the present study was marginally more posterior than the centroparietal region

of interest used in previous studies of depression, but in agreement with studies of healthy

participants reporting that a maximum effect occurred parietally (Hajcak, Moser, et al.,

2006; Kessel et al., 2017; Liu et al., 2012). Interestingly, there is evidence to suggest that the

LPP is initially largest parietally, and then shifts to a more centroparietal location in the

latter stages of affective processing (Hajcak et al., 2012).

Studies of affective processing have operationalised the LPP using a diverse range of time

intervals. These report it as commencing as early as 300 ms (Liu et al., 2012; Thiruchselvam

et al., 2011), and as late as 600 ms (Mehmood & Lee, 2016). The LPP has been suggested to

terminate as early as 500 ms (Schupp et al., 2003; Schupp et al., 2004), and as late as 1700

ms (MacNamara, Foti, et al., 2009; Thiruchselvam et al., 2011). Lastly, the duration of the

LPP interval has been defined to be as narrow as 50 ms (Schupp et al., 2003), and as wide as

1400 ms (Liu et al., 2012). Broadly, the cluster identified in our exploratory analysis falls

within the time interval of previous LPP studies.

Our secondary hypotheses stated that LPP amplitude would be moderated by emotion

regulation strategy use and anxiety comorbidity in MDD participants. Contrary to these

hypotheses, emotion regulation strategy use did not differ between diagnostic groups.

Further, we found no evidence to suggest that emotion regulation strategy use attenuated

LPP amplitudes to unpleasant pictures, in contrast to previous EEG studies. Extensive

research has shown reduced LPP amplitude to unpleasant pictures when

reinterpretation/reappraisal (Foti et al., 2008; Hajcak, Moser, et al., 2006; Hajcak &

Nieuwenhuis, 2006; Moser et al., 2006) or distraction were used to regulate negative

emotions (Dunning et al., 2009; Hajcak et al., 2007; Thiruchselvam et al., 2011; Uusberg et

al., 2014; Zhang et al., 2014). A possible reason for this discrepancy may be that the majority

of healthy and depressed participants in the current study reported no emotion regulation

strategy use (MDD: 30/40; healthy controls: 26/41), thus limiting the ability to statistically

infer the effects of strategy on LPP amplitudes. We also did not find evidence that anxiety

levels modulate LPP amplitudes. The prevailing view is that anxiety increases attentional

resources towards emotionally salient, aversive, stimuli thereby resulting in enhancement

of LPP amplitude (MacNamara & Hajcak, 2009). Previous research has shown enhanced LPP

to aversive pictures in healthy individuals with high self-reported anxiety (MacNamara &

Hajcak, 2009), and patients suffering from generalized anxiety disorder (MacNamara &

Hajcak, 2010; MacNamara et al., 2016). Although our results differ considerably from these

studies, they are corroborated by Weinberg et al. (2016), who found that a diagnosis of

anxiety did not modulate the LPP. The reason for these disagreements in the field is unclear

but may be related to idiosyncratic features of the study samples, including criteria used to

estimate anxiety symptoms.

Conclusion

In conclusion, exploratory analyses found significantly reduced EEG amplitudes to

unpleasant pictures in MDD, in support of the *emotional context insensitivity* hypothesis.

However, we did not find a difference in emotional reactivity between MDD participants

and healthy controls using our preregistered analysis plan, which was based on the existing

literature. Compared to preregistered LPP features, the parietal EEG cluster identified in

exploratory analyses was marginally more posterior and subtended a narrower time interval

(660-755ms). These results suggest that methodological differences between studies may

play an important role in variability in the cortical response to emotional stimuli, limiting the

generalisability of the findings. Neither emotion regulation strategy use, nor comorbid

anxiety, were found to modulate LPP amplitude. Although the pathophysiology of

depression remains poorly understood, arising possibly due to its inherent heterogeneity

(Fried & Nesse, 2015), our findings indicate that a core feature and promising biomarker of

depression may be reduced emotional reactivity to unpleasant stimuli. The presence of

heterogeneity in EEG correlates of affective processing indicates the need for standardised

EEG tests and analyses of emotional reactivity in depression.

REFERENCES

American Psychiatric Association. (2013). *Diagnostic and statistical manual of mental disorders (DSM-5®)*: American Psychiatric Pub.

Britton, J. C., Taylor, S. F., Sudheimer, K. D., & Liberzon, I. (2006). Facial expressions and complex IAPS pictures: common and differential networks. *Neuroimage, 31*(2), 906-919.

Brown, T. A., Chorpita, B. F., & Barlow, D. H. (1998). Structural relationships among dimensions of the DSM-IV anxiety and mood disorders and dimensions of negative affect, positive affect, and autonomic arousal. *Journal of abnormal psychology, 107*(2), 179.

Browning, M., Holmes, E. A., & Harmer, C. J. (2010). The modification of attentional bias to emotional information: A review of the techniques, mechanisms, and relevance to emotional disorders. *Cognitive, Affective, & Behavioral Neuroscience, 10*(1), 8-20.

Clark, L. A., & Watson, D. (1991). Tripartite model of anxiety and depression: psychometric evidence and taxonomic implications. *Journal of abnormal psychology, 100*(3), 316.

Codispoti, M., Ferrari, V., & Bradley, M. M. (2006). Repetitive picture processing: autonomic and cortical correlates. *Brain research, 1068*(1), 213-220.

Codispoti, M., Ferrari, V., & Bradley, M. M. (2007). Repetition and event-related potentials: distinguishing early and late processes in affective picture perception. *J Cogn Neurosci, 19*(4), 577-586.

Crawford, J. R., & Henry, J. D. (2003). The Depression Anxiety Stress Scales (DASS): Normative data and latent structure in a large non-clinical sample. *British journal of clinical psychology, 42*(2), 111-131.

Cuthbert, B. N., Schupp, H. T., Bradley, M. M., Birbaumer, N., & Lang, P. J. (2000). Brain potentials in affective picture processing: covariation with autonomic arousal and affective report. *Biological psychology, 52*(2), 95-111.

Delorme, A., Sejnowski, T., & Makeig, S. (2007). Enhanced detection of artifacts in EEG data using higher-order statistics and independent component analysis. *Neuroimage, 34*(4), 1443-1449.

Dunning, J. P., & Hajcak, G. (2009). See no evil: Directing visual attention within unpleasant images modulates the electrocortical response. *Psychophysiology, 46*(1), 28-33.

Foti, D., & Hajcak, G. (2008). Deconstructing reappraisal: descriptions preceding arousing pictures modulate the subsequent neural response. *J Cogn Neurosci, 20*(6), 977-988.

Foti, D., Olvet, D. M., Klein, D. N., & Hajcak, G. (2010). Reduced electrocortical response to threatening faces in major depressive disorder. *Depression and anxiety, 27*(9), 813-820.

Fried, E. I., & Nesse, R. M. (2015). Depression is not a consistent syndrome: an investigation of unique symptom patterns in the STAR* D study. *Journal of affective disorders, 172*, 96-102.

Gross, J. J., & John, O. P. (2003). Individual differences in two emotion regulation processes:
implications for affect, relationships, and well-being. *Journal of personality and social*
psychology, 85(2), 348.

Hajcak, G., Dunning, J. P., & Foti, D. (2007). Neural response to emotional pictures is
unaffected by concurrent task difficulty: an event-related potential study. *Behavioral*
neuroscience, 121(6), 1156.

Hajcak, G., Moser, J. S., & Simons, R. F. (2006). Attending to affect: appraisal strategies
modulate the electrocortical response to arousing pictures. *Emotion, 6*(3), 517.

Hajcak, G., & Nieuwenhuis, S. (2006). Reappraisal modulates the electrocortical response to
unpleasant pictures. *Cognitive, Affective, & Behavioral Neuroscience, 6*(4), 291-297.

Hajcak, G., Weinberg, A., MacNamara, A., & Foti, D. (2012). ERPs and the study of emotion.
The Oxford handbook of event-related potential components, 441, 474.

Hariri, A. R., Tessitore, A., Mattay, V. S., Fera, F., & Weinberger, D. R. (2002). The amygdala
response to emotional stimuli: a comparison of faces and scenes. *Neuroimage, 17*(1),
317-323.

Hyvärinen, A., Hoyer, P. O., & Inki, M. (2001). Topographic independent component
analysis. *Neural computation, 13*(7), 1527-1558.

Kayser, J., Bruder, G. E., Tenke, C. E., Stewart, J. E., & Quitkin, F. M. (2000). Event-related
potentials (ERPs) to hemifield presentations of emotional stimuli: differences
between depressed patients and healthy adults in P3 amplitude and asymmetry.
International Journal of Psychophysiology, 36(3), 211-236.

Kessel, E. M., Kujawa, A., Goldstein, B., Hajcak, G., Bufferd, S. J., Dyson, M., & Klein, D. N.
(2017). Behavioral observations of positive and negative valence systems in early
childhood predict physiological measures of emotional processing three years later.
Journal of affective disorders, 216, 70-77.

Lang, P., Bradley, M., & Cuthbert, B. (2008). *International affective picture system (IAPS):*
affective ratings of pictures and instruction manual. University of Florida, Gainesville.
Retrieved from

Ledgerwood, A. (2018). The preregistration revolution needs to distinguish between
predictions and analyses. *Proceedings of the National Academy of Sciences, 115*(45),
E10516-E10517.

Liu, Y., Huang, H., McGinnis-Deweese, M., Keil, A., & Ding, M. (2012). Neural substrate of the
late positive potential in emotional processing. *Journal of Neuroscience, 32*(42),
14563-14572.

Lovibond, P. F., & Lovibond, S. H. (1995). The structure of negative emotional states:
Comparison of the Depression Anxiety Stress Scales (DASS) with the Beck Depression
and Anxiety Inventories. *Behaviour research and therapy, 33*(3), 335-343.

MacNamara, A., Foti, D., & Hajcak, G. (2009). Tell me about it: neural activity elicited by
emotional pictures and preceding descriptions. *Emotion, 9*(4), 531.

MacNamara, A., & Hajcak, G. (2009). Anxiety and spatial attention moderate the
electrocortical response to aversive pictures. *Neuropsychologia, 47*(13), 2975-2980.

MacNamara, A., & Hajcak, G. (2010). Distinct electrocortical and behavioral evidence for
increased attention to threat in generalized anxiety disorder. *Depression and*
anxiety, 27(3), 234-243.

MacNamara, A., Kotov, R., & Hajcak, G. (2016). Diagnostic and symptom-based predictors of
emotional processing in generalized anxiety disorder and major depressive disorder:
An event-related potential study. *Cognitive Therapy and Research, 40*(3), 275-289.

Makeig, S., Bell, A. J., Jung, T.-P., & Sejnowski, T. J. (1996). *Independent component analysis*
of electroencephalographic data. Paper presented at the Advances in neural
information processing systems.

Maris, E., & Oostenveld, R. (2007). Nonparametric statistical testing of EEG-and MEG-data.
Journal of neuroscience methods, 164(1), 177-190.

Mavratzakis, A., Herbert, C., & Walla, P. (2016). Emotional facial expressions evoke faster
orienting responses, but weaker emotional responses at neural and behavioural
levels compared to scenes: A simultaneous EEG and facial EMG study. *Neuroimage,*
124, 931-946.

Mehmood, R. M., & Lee, H. J. (2016). A novel feature extraction method based on late
positive potential for emotion recognition in human brain signal patterns. *Computers*
& Electrical Engineering, 53, 444-457.

Montgomery, S. A., & Asberg, M. (1979). A new depression scale designed to be sensitive to
change. *Br J Psychiatry, 134*, 382-389.

Moser, J. S., Hajcak, G., Bukay, E., & Simons, R. F. (2006). Intentional modulation of
emotional responding to unpleasant pictures: an ERP study. *Psychophysiology, 43*(3),
292-296.

Moser, J. S., Krompinger, J. W., Dietz, J., & Simons, R. F. (2009). Electrophysiological
correlates of decreasing and increasing emotional responses to unpleasant pictures.
Psychophysiology, 46(1), 17-27.

Nikolin, S., Chand, N., Martin, D., Rushby, J., Loo, C. K., & Boonstra, T. (2020). EEG correlates
of affective processing in major depressive disorder. *OSF*.
doi:https://doi.org/10.17605/OSF.IO/R6XEM

Nosek, B. A., Ebersole, C. R., DeHaven, A. C., & Mellor, D. T. (2018). The preregistration
revolution. *Proceedings of the National Academy of Sciences, 115*(11), 2600-2606.

Oliver-Rodriguez, J. C., Guan, Z., & Johnston, V. S. (1999). Gender differences in late positive
components evoked by human faces. *Psychophysiology, 36*(2), 176-185.

Oostenveld, R., Fries, P., Maris, E., & Schoffelen, J.-M. (2011). FieldTrip: open source
software for advanced analysis of MEG, EEG, and invasive electrophysiological data.
Computational intelligence and neuroscience, 2011, 1.

Pernet, C., Latinus, M., Nichols, T., & Rousselet, G. (2015). Cluster-based computational
methods for mass univariate analyses of event-related brain potentials/fields: A
simulation study. *Journal of neuroscience methods, 250*, 85-93.

Renfroe, J. B., Bradley, M. M., Sege, C. T., & Bowers, D. (2016). Emotional modulation of the
late positive potential during picture free viewing in older and young adults. *PloS*
one, 11(9).

Rottenberg, J. (2005). Mood and emotion in major depression. *Current Directions in*
Psychological Science, 14(3), 167-170.

Rottenberg, J., Gross, J. J., & Gotlib, I. H. (2005). Emotion context insensitivity in major
depressive disorder. *Journal of abnormal psychology, 114*(4), 627.

Sassenhagen, J., & Draschkow, D. (2019). Cluster-based permutation tests of MEG/EEG data
do not establish significance of effect latency or location. *Psychophysiology, 56*(6),
e13335.

Schupp, H. T., Cuthbert, B. N., Bradley, M. M., Cacioppo, J. T., Ito, T., & Lang, P. J. (2000).
Affective picture processing: the late positive potential is modulated by motivational
relevance. *Psychophysiology, 37*(2), 257-261.

Schupp, H. T., Markus, J., Weike, A. I., & Hamm, A. O. (2003). Emotional facilitation of
sensory processing in the visual cortex. *Psychological science, 14*(1), 7-13.

Schupp, H. T., Öhman, A., Junghöfer, M., Weike, A. I., Stockburger, J., & Hamm, A. O. (2004).
The facilitated processing of threatening faces: an ERP analysis. *Emotion, 4*(2), 189.

Thiruchselvam, R., Blechert, J., Sheppes, G., Rydstrom, A., & Gross, J. J. (2011). The temporal
dynamics of emotion regulation: An EEG study of distraction and reappraisal.
Biological psychology, 87(1), 84-92.

Uusberg, A., Thiruchselvam, R., & Gross, J. J. (2014). Using distraction to regulate emotion:
Insights from EEG theta dynamics. *International Journal of Psychophysiology, 91*(3),
254-260.

Watson, D. (2000). *Mood and temperament*: Guilford Press.

Weinberg, A., Perlman, G., Kotov, R., & Hajcak, G. (2016). Depression and reduced neural
response to emotional images: Distinction from anxiety, and importance of symptom
dimensions and age of onset. *Journal of abnormal psychology, 125*(1), 26.

Yarkoni, T. (2019). The generalizability crisis.

Zhang, D., He, Z., Chen, Y., & Wei, Z. (2016). Deficits of unconscious emotional processing in
patients with major depression: an ERP study. *Journal of affective disorders, 199*, 13-
20.

Zhang, W., Li, H., Chen, J., Chen, N., Liu, X., Wang, D., & Shen, J. (2014). Posterior P1 and
early frontal negativity reflect developmental changes in attentional distraction
during adolescence. *Brain and cognition, 87*, 30-38.

SUPPLEMENTARY MATERIALS

624 Some participants repeated the experimental protocol over three separate session. Only

625 results from the first of these sessions is reported in the present study. To prevent carryover

626 effects between sessions, the set of pictures participants viewed at each session was

627 randomised and counterbalanced from the available parallel lists described below:

628 IAPS Picture Sets

IAPS List A	Description	Valence Mean	SD	Arousal Mean	SD	Class
6230	AimedGun	2.37	1.57	7.35	2.01	Unpleasant
2053	Baby	2.47	1.87	5.25	2.46	Unpleasant
9620	Shipwreck	2.70	1.64	6.11	2.10	Unpleasant
2745.1	Shopping	5.31	1.08	3.26	1.96	Neutral
7035	Mug	4.98	0.96	2.66	1.82	Neutral
2691	Riot	3.04	1.73	5.85	2.03	Unpleasant
7950	Tissue	4.94	1.21	2.28	1.81	Neutral
6821	Gang	2.38	1.72	6.29	2.02	Unpleasant
9520	Kids	2.46	1.61	5.41	2.27	Unpleasant
7150	Umbrella	4.72	1.00	2.61	1.76	Neutral
3300	DisabledChild	2.74	1.56	4.55	2.06	Unpleasant
2570	Man	4.78	1.24	2.76	1.92	Neutral
2688	Hunters	2.73	2.07	5.98	2.22	Unpleasant
2703	SadChildren	1.91	1.26	5.78	2.25	Unpleasant
2516	ElderlyWoman	4.90	1.43	3.50	1.88	Neutral
5532	Mushrooms	5.19	1.69	3.79	2.20	Neutral
1302	Dog	4.21	1.78	6.00	1.87	Unpleasant
2811	Gun	2.17	1.38	6.90	2.22	Unpleasant
2393	FactoryWorker	4.87	1.06	2.93	1.88	Neutral
2683	War	2.62	1.78	6.21	2.15	Unpleasant
7185	AbstractArt	4.97	0.87	2.64	2.04	Neutral
7034	Hammer	4.95	0.87	3.06	1.95	Neutral
2710	DrugAddict	2.52	1.69	5.46	2.29	Unpleasant
7100	FireHydrant	5.24	1.20	2.89	1.70	Neutral
5731	Flowers	5.39	1.58	2.74	1.95	Neutral
7037	Trains	4.81	1.12	3.71	2.08	Neutral
6260	AimedGun	2.44	1.54	6.93	1.93	Unpleasant

		Valence		Arousal		
		Mean	SD	Mean	SD	Class
7050	HairDryer	4.93	0.81	2.75	1.80	Neutral
3220	Hospital	2.49	1.29	5.52	1.86	Unpleasant
7009	Mug	4.93	1.00	3.01	1.97	Neutral
2810	Boy	4.31	1.65	4.47	1.92	Unpleasant
2840	Chess	4.91	1.52	2.43	1.82	Neutral
7590	Traffic	4.75	1.55	3.80	2.13	Neutral
5535	StillLife	4.81	1.52	4.11	2.31	Neutral
3500	Attack	2.21	1.34	6.99	2.19	Unpleasant
2455	SadGirls	2.96	1.79	4.46	2.12	Unpleasant
7705	Cabinet	4.77	1.02	2.65	1.88	Neutral
7080	Fork	5.27	1.09	2.32	1.84	Neutral
2981	DeerHead	2.76	1.94	5.97	2.12	Unpleasant
2750	Bum	2.56	1.32	4.31	1.81	Unpleasant
6020	ElectricChair	3.41	1.98	5.58	2.01	Unpleasant
2200	NeutFace	4.79	1.38	3.18	2.17	Neutral
2141	GrievingFem	2.44	1.64	5.00	2.03	Unpleasant
1930	Shark	3.79	1.92	6.42	2.07	Unpleasant
6212	Soldier	2.19	1.49	6.01	2.44	Unpleasant
2191	Farmer	5.30	1.62	3.61	2.14	Neutral
2305	Woman	5.41	1.12	3.63	2.04	Neutral
7000	RollingPin	5.00	0.84	2.42	1.79	Neutral
6200	AimedGun	2.71	1.58	6.21	2.28	Unpleasant
7500	Building	5.33	1.44	3.26	2.18	Neutral
9920	CarAccident	2.50	1.52	5.76	1.96	Unpleasant
7004	Spoon	5.04	0.60	2.00	1.66	Neutral
7010	Basket	4.94	1.07	1.76	1.48	Neutral
7235	Chair	4.96	1.18	2.83	2.00	Neutral
2700	Woman	3.19	1.56	4.77	1.97	Unpleasant
6834	Police	2.91	1.73	6.28	1.90	Unpleasant

		Valence		Arousal		
IAPS List B	Description	Mean	SD	Mean	SD	Class
6210	AimedGun	2.95	1.83	6.34	2.14	Unpleasant
2661	Baby	3.90	2.49	5.76	2.13	Unpleasant
5740	Plant	5.21	1.38	2.59	1.99	Neutral
2717	DrugAddict	2.58	1.32	5.70	2.16	Unpleasant
7493	Man	5.35	1.34	3.39	2.08	Neutral
2095	Toddler	1.79	1.18	5.25	2.34	Unpleasant
2120	AngryFace	3.34	1.91	5.18	2.52	Unpleasant

2595	Women	4.88	1.24	3.71	1.88	Neutral
9902	CarAccident	2.33	1.38	6.00	2.15	Unpleasant
9421	Soldier	2.21	1.45	5.04	2.15	Unpleasant
7043	Drill	5.17	1.26	3.68	2.09	Neutral
7003	Disk	5.00	1.22	3.07	1.98	Neutral
2900	CryingBoy	2.45	1.42	5.09	2.15	Unpleasant
7036	Shipyard	4.88	1.08	3.32	2.04	Neutral
1111	Snakes	3.25	1.64	5.20	2.25	Unpleasant
7547	Bridge	5.21	0.96	3.18	2.01	Neutral
7175	Lamp	4.87	1.00	1.72	1.26	Neutral
3230	DyingMan	2.02	1.30	5.41	2.21	Unpleasant
9050	PlaneCrash	2.43	1.61	6.36	1.97	Unpleasant
3350	Infant	1.88	1.67	5.72	2.23	Unpleasant
7233	Plate	5.09	1.46	2.77	1.92	Neutral
6244	AimedGun	3.09	1.78	5.68	2.51	Unpleasant
9430	Burial	2.63	1.59	5.26	2.55	Unpleasant
2190	Man	4.83	1.28	2.41	1.80	Neutral
7491	Building	4.82	1.03	2.39	1.90	Neutral
6831	Police	2.59	1.50	5.55	2.16	Unpleasant
9600	Ship	2.48	1.62	6.46	2.31	Unpleasant
2514	Woman	5.19	1.09	3.50	1.81	Neutral
7056	Tool	5.07	1.02	3.07	1.92	Neutral
6370	Attack	2.70	1.52	6.44	2.19	Unpleasant
7006	Bowl	4.88	0.99	2.33	1.67	Neutral
7180	NeonBuilding	4.73	1.31	3.43	1.95	Neutral
1301	Dog	3.70	1.66	5.77	2.18	Unpleasant
2102	NeuMan	5.16	0.96	3.03	1.87	Neutral
7038	Shoes	4.82	1.20	3.01	1.96	Neutral
1201	Spider	3.55	1.88	6.36	2.11	Unpleasant
5510	Mushroom	5.15	1.43	2.82	2.18	Neutral
2880	Shadow	5.18	1.44	2.96	1.94	Neutral
9941	Fire	2.91	1.54	5.83	2.14	Unpleasant
7090	Book	5.19	1.46	2.61	2.03	Neutral
7211	Clock	4.81	1.78	4.20	2.40	Neutral
2512	Man	4.86	0.84	3.46	1.75	Neutral
2130	Woman	4.08	1.33	5.02	2.00	Unpleasant
1300	PitBull	3.55	1.78	6.79	1.84	Unpleasant
2480	ElderlyMan	4.77	1.64	2.66	1.78	Neutral
7002	Towel	4.97	0.97	3.16	2.00	Neutral
2385	Girl	5.20	1.32	3.64	1.81	Neutral
3301	InjuredChild	1.80	1.28	5.21	2.26	Unpleasant

IAPS	Description	Valence Mean	SD	Arousal Mean	SD	Class
2205	Hospital	1.95	1.58	4.53	2.23	Unpleasant
3160	EyeDisease	2.63	1.23	5.36	1.79	Unpleasant
2383	Secretary	4.72	1.36	3.41	1.83	Neutral
9911	CarAccident	2.30	1.37	5.76	2.10	Unpleasant
7217	ClothesRack	4.82	0.99	2.43	1.64	Neutral
7001	Buttons	5.32	1.19	3.20	2.15	Neutral
6825	Military	2.81	1.19	5.36	2.19	Unpleasant
2800	SadChild	1.78	1.14	5.49	2.11	Unpleasant

IAPS List C	Description	Valence Mean	SD	Arousal Mean	SD	Class
8485	Fire	2.73	1.62	6.46	2.10	Unpleasant
7059	Keyring	4.93	0.81	2.73	1.88	Neutral
2495	Man	5.22	1.10	3.19	1.76	Neutral
9427	Assault	2.89	1.47	5.50	2.09	Unpleasant
6150	Outlet	5.08	1.17	3.22	2.02	Neutral
9590	Injecting	3.08	1.63	5.41	2.23	Unpleasant
8230	Boxer	2.95	1.88	5.91	2.15	Unpleasant
7546	Bridge	5.40	1.13	3.72	2.16	Neutral
9491	DeadBody	2.78	1.71	5.69	2.22	Unpleasant
7032	Shoes	4.82	1.46	3.18	1.88	Neutral
7062	Sewing	5.27	1.06	3.40	1.94	Neutral
6555	Knife	3.33	1.59	5.69	2.21	Unpleasant
6220	BoysW/Guns	3.10	1.91	5.89	2.43	Unpleasant
9120	OilFires	3.20	1.75	5.77	1.94	Unpleasant
7025	Stool	4.63	1.17	2.71	2.20	Neutral
9250	WarVictim	2.57	1.39	6.60	1.87	Unpleasant
7365	Meat	5.20	1.57	4.13	1.93	Neutral
9424	Bomb	2.87	1.62	5.78	2.12	Unpleasant
9611	PlaneCrash	2.71	1.95	5.75	2.44	Unpleasant
9621	Ship	3.22	1.76	5.76	2.05	Unpleasant
7020	Fan	4.97	1.04	2.17	1.71	Neutral
9480	Skull	3.51	2.08	5.57	1.87	Unpleasant
7030	Iron	4.69	1.04	2.99	2.09	Neutral
6250	AimedGun	2.83	1.79	6.54	2.61	Unpleasant
9909	BurningCar	2.78	1.45	5.98	2.04	Unpleasant
7255	Cracker	5.07	1.18	3.36	1.99	Neutral
2214	NeuMan	5.01	1.12	3.46	1.97	Neutral
2397	Men	4.98	1.11	2.77	1.74	Neutral

6838	Police	2.45	1.44	5.80	2.09	Unpleasant
7045	Zipper	4.97	0.76	3.32	1.96	Neutral
6190	AimedGun	3.57	1.84	5.64	2.03	Unpleasant
2396	Couple	4.91	1.05	3.34	1.83	Neutral
7053	CandleStick	5.22	0.75	2.95	1.91	Neutral
9622	Jet	3.10	1.90	6.26	1.98	Unpleasant
7130	Truck	4.77	1.03	3.35	1.90	Neutral
6830	Guns	2.82	1.81	6.21	2.23	Unpleasant
7041	Baskets	4.99	1.12	2.60	1.78	Neutral
6213	Terrorist	2.91	1.52	5.86	2.06	Unpleasant
7055	Lightbulb	4.90	0.64	3.02	1.83	Neutral
7380	RoachOnPizza	2.46	1.42	5.88	2.44	Unpleasant
9630	Bomb	2.96	1.72	6.06	2.22	Unpleasant
7224	FileCabinet	4.45	1.36	2.81	1.94	Neutral
1525	Attack Dog	3.09	1.72	6.51	2.25	Unpleasant
2038	NeuWoman	5.09	1.35	2.94	1.93	Neutral
7014	Scissors	5.15	0.97	3.25	2.03	Neutral
9800	Skinhead	2.04	1.57	6.05	2.71	Unpleasant
7017	Video	5.18	1.07	3.12	1.97	Neutral
2411	Girl	5.07	0.85	2.86	1.84	Neutral
6571	CarTheft	2.85	2.05	5.59	2.50	Unpleasant
9904	CarAccident	2.39	1.36	6.08	2.06	Unpleasant
6510	Attack	2.46	1.58	6.96	2.09	Unpleasant
7187	AbstractArt	5.07	1.02	2.30	1.75	Neutral
9321	Vomit	2.81	2.14	6.24	2.23	Unpleasant
7283	Fruit	5.50	1.84	3.81	2.01	Neutral
2890	Twins	4.95	1.09	2.95	1.87	Neutral
5531	Mushroom	5.15	1.45	3.69	2.11	Neutral

633   Emotion Regulation Questionnaire

634   An emotion regulation questionnaire adapted was used to identify which strategy, if any,

635   participants used to moderate their emotional response while passively viewing pictures.

Choose only one option:

While viewing unpleasant pictures:

1. I did not use any strategy

2. I looked on the bright side

3. I thought about something else

4. I distanced myself from the picture

5. I did not do any of the above, I did…

Exploratory Cluster Channels

Channels from the significant cluster identified by the exploratory cluster-based

permutation testing are displayed below. These are sorted as a proportion of the length of

time they were identified relative to the total duration of the cluster time interval. Channels

that were present for >75% of the cluster time interval were used in follow-up analyses.

Secondary Analyses of Emotion Regulation Strategy and Anxiety

Supplementary Figure 1. Effects of emotion regulation strategy and anxiety on LPP amplitude. A) The LPP for the preregistered centroparietal cluster and time interval in both MDD participants and healthy controls who used an emotion regulation strategy or not. Black lines show the mean, red lines show the median, light grey shaded boxes indicate the standard deviation, and dark grey regions indicate the 95% confidence interval. **B)** Association between the anxiety subscore from the 21-item DASS and LPP amplitude for MDD participants while viewing unpleasant images. Dotted line shows the regression slope ($p > 0.05$; ns).