1 **Early transmission dynamics of COVID-19 in a southern hemisphere setting: Lima-Peru:**

2 **February 29th–March 30th, 2020.**

4 César V. Munayco[1#], Amna Tariq[2*#], Richard Rothenberg[2], Gabriela G Soto-Cabezas[1], Mary F.
5 Reyes[1], Andree Valle[1], Leonardo Rojas-Mezarina[3], César Cabezas[3], Manuel Loayza[1], Peru
6 COVID-19 working group, Gerardo Chowell[2]
7
8 [#] joint first authors
9
[1] Centro Nacional de Epidemiología, Prevención y Control de Enfermedades, Peruvian Ministry

of Health, Lima, Peru.

[2] Department of Population Health Sciences, School of Public Health, Georgia State University,

Atlanta, GA, USA

[3] Instituto Nacional de Salud, Peruvian Ministry of Health, Lima, Peru.

Peru COVID-19 working group

Nombre	Area
Dante Castro Garro	Investigación Epidemiológica
Kevin Martel Vasquez	Investigación Epidemiológica
Edith Solis Castro	PREEC
Isabel Sandoval Ordinola	PREEC
Javier Masias Mimbela	PREEC
Kely Meza Cornejo	PREEC
Fabiola Caruajulca Quijano	PREEC
Lenin La Torre Rosillo	PREEC
Luis Ordoñez Ibarguen	PREEC
Mario Vasquez Dominguez	PREEC
Rommell Veintimilla Gonzalez Seminario	PREEC
Martha Calderón Silva	PREEC
Mayra Saavedra Dreyfus	PREEC
Miguel Luna Pineda	PREEC
Milagros Durand	PREEC
Noemí Janampa	PREEC
Jesus Chuquihuaccha	PREEC
Susan Mateo Lizarbe	PREEC
David Enriquez Cusi	PREEC
Idania Mamani Pilco	PREEC
Alex Jaramillo	PREEC
Karina Vargas	PREEC
Oswaldo Cabanillas	Alerta-Respuesta
Juan Arrasco	Alerta-Respuesta
María Vargas	Vigilancia en Salud Pública

| Willy Ramos | Inteligencia sanitaria |

* Corresponding author

Amna Tariq

Department of Population Health Sciences

Georgia State University School of Public Health

Atlanta GA, 30303

atariq1@student.gsu.edu

Contact number: 470-985-6352

45
46
47

Abstract

The COVID-19 pandemic that emerged in Wuhan China has generated substantial morbidity and

mortality impact around the world during the last four months. The daily trend in reported cases

has been rapidly rising in Latin America since March 2020 with the great majority of the cases

reported in Brazil followed by Peru as of April 15th, 2020. Although Peru implemented a range of

social distancing measures soon after the confirmation of its first case on March 6th, 2020, the daily

number of new COVID-19 cases continues to accumulate in this country. We assessed the early

COVID-19 transmission dynamics and the effect of social distancing interventions in Lima, Peru.

We estimated the reproduction number, R, during the early transmission phase in Lima from the

daily series of imported and autochthonous cases by the date of symptoms onset as of March 30th,

2020. We also assessed the effect of social distancing interventions in Lima by generating short-

term forecasts grounded on the early transmission dynamics before interventions were put in place.

Prior to the implementation of the social distancing measures in Lima, the local incidence curve

by the date of symptoms onset displays near exponential growth dynamics with the mean scaling

of growth parameter, p, estimated at 0.9 (95%CI: 0.9,1.0) and the reproduction number at 2.3 (95%

CI: 2.0, 2.5). Our analysis indicates that school closures and other social distancing interventions

have helped slow down the spread of the novel coronavirus, with the nearly exponential growth

trend shifting to an approximately linear growth trend soon after the broad scale social distancing

interventions were put in place by the government.

While the interventions appear to have slowed the transmission rate in Lima, the number of new

COVID-19 cases continue to accumulate, highlighting the need to strengthen social distancing and

active case finding efforts to mitigate disease transmission in the region.

Key words

COVID-19; SARS-CoV-2; Transmission potential; Short-term forecast; Reproduction number;

Generalized growth model

1. Introduction

The Coronavirus disease 2019 (COVID-19) pandemic that emerged in the city of Wuhan in China in December 2019 has invaded nearly every nation of the world, becoming the most important public health emergency of the last century after the 1918–1920 influenza pandemic (WHO, 2020). In particular, the novel Severe Acute Respiratory Syndrome Coronavirus 2 (SARS-CoV-2) has an ability to exert substantial severe disease and mortality burden especially affecting individuals older than 60 years and those with prior health conditions including hypertension, cardiovascular disease, obesity and diabetes (Adler, 2020; Team, 2020). As of April 15th, 2020, the trajectory of the pandemic varies significantly around the world ranging from relatively well contained outbreaks in Thailand, Taiwan and Hong Kong to explosive epidemics characterized by initial exponential growth periods in a few hotspots located in various countries around the world including the United States, Italy, Spain, UK, France, and Iran (Ebbs, 2020; Griffiths, 2020; Minder, 2020).

By April 15th, 2020, SARS-CoV-2 is generating local transmission in over 200 countries and over 2.2 million cases and 150 thousand deaths have been reported globally (WHO, 2020). The COVID-19 pandemic was confirmed to have reached Latin America in February 2020 with a gradual expansion in the region until March 2020 when the COVID-19 incidence curve started to grow more rapidly. The US, the country with the highest number of reported COVID-19 cases in the world, has recorded 637,196 COVID-19 cases by April 15th, 2020. In Latin America, Brazil has reported 28320 cases, the highest number of cases in the region followed by Peru with a total of 11475 cases (Worldometer, 2020).

Peru, a country located in western South America, reported its first imported case of COVID-19 in Lima, on March 6th, 2020, a Peruvian with recent travel history to France, Spain and Czech Republic (Aquino & Garrison, 2020). By April 15th, 2020, a total of 11475 cases including 254 deaths have been reported by the Peruvian government. Lima, the capital of Peru has recorded 8412 cases, the highest number of cases within Peru (MOH, 2020). To respond to the growing number of COVID-19 cases in the country, the government shuttered schools on March 11th, 2020. The next day, the government banned gatherings of more than 300 people and suspended all international flights from Europe and Asia. On March 16th, 2020, the government declared a

national emergency and closed country borders (Explorer, 2020). Subsequently, on March 17th,

2020 the president of Peru announced the beginning of community transmission of SARS-CoV-2

in the country, and ordered a curfew in the region on March 18th, 2020 to avoid night time

socializing to prevent disease transmission (Explorer, 2020; Writing, 2020).

In order to combat the spread of the COVID-19 epidemic in Lima, the capital and largest city of

Peru, estimates of the transmission potential of COVID-19 can guide the intensity of interventions

including the reproduction number, R, during the early transmission phase (Nishiura & Chowell,

2009, 2014). Moreover, using the epidemiological data and mathematical modeling, it is possible

to gauge the impact of control interventions including school closures and a national emergency

declaration in Lima by assessing short-term forecasts grounded on the trajectory of the epidemic

prior to the implementation of control interventions (Funk, Camacho, Kucharski, Eggo, &

Edmunds, 2018; Shanafelt, Jones, Lima, Perrings, & Chowell, 2018).

2. Methods

2.1. Data

We analyzed the daily number of COVID-19 confirmed cases by date of symptoms onset in Lima,

Peru by March 30th, 2020. Individual-level case details including whether the case was locally

acquired or imported were also made available from the Centro Nacional de Epidemiología

Prevención y control de Enfermedades and the National Institute of Health of the Ministry of

Health, Peru (Group, 2020). We also examined the daily testing rate and the positivity rate from

the daily number of positive and negative PCR test results by the date of reporting until March

30th, 2020.

2.2. Early growth model

We generate short-term forecasts in real time using the generalized growth model (GGM) that

relies on two parameters and characterizes the early ascending phase of the epidemic allowing to

capture a range of epidemic growth profiles including sub-exponential (polynomial) and

exponential growth. GGM characterizes epidemic growth by estimating two parameters (i) the

intrinsic growth rate, r and (ii) a dimensionless "deacceleration of growth" or "scaling of growth"

parameter, p. The latter parameter modulates the epidemic growth patterns including the sub-

exponential growth (p<1) and exponential growth dynamics (p=1). The GGM model is given by

the following differential equation:

143
$$\frac{dC(t)}{dt} = C'(t) = rC(t)^p$$

Where $C'(t)$ describes case incidence over time t. The cumulative number of cases at time t is

given by $C(t)$ while r is a positive parameter denoting the growth rate (1/time) and $p \in [0,1]$ is a

"deceleration of growth" parameter (Chowell, 2017; Viboud, Simonsen, & Chowell, 2016).

2.3. Short term forecast to assess interventions

We calibrate the GGM model to the daily case incidence by the date of symptoms onset for Lima.

We analyzed the time series data of confirmed cases by onset dates for Lima from February 29th,

2020 to March 30th, 2020. Our model was calibrated using case series from February 29th–March

15th, 2020, prior to the implementation of national emergency in Lima.

The best fit model solution is estimated by using a non-linear least square fitting approach (full

details provided in (Chowell, 2017)). This process searches for the set of model parameters that

minimizes the sum of squared differences between the observed data $y_{ti} = y_{t1}, y_{t2}, \ldots . y_{tn}$ and the

corresponding model solution given by $f(t_i, \Theta)$: where $\Theta = (r, p)$ correspond to estimated set of

parameters of the GGM model. Thus, the objective function for the best fit solution of $f(t_i, \Theta)$ is

given by

161
$$\hat{\Theta} = \arg \min \sum_{i=1}^{n} (f(t_i, \Theta) - y_{ti})^2$$

The initial condition is fixed to the first observation in the data. Next, we derive uncertainty around

the best fit model solution as well as the confidence intervals of the parameters utilizing a

parametric bootstrapping approach assuming a Poisson error structure as described in ref (Chowell,

2017).

2.4. Reproduction number from case incidence using the generalized-growth model

Generally, the reproduction number, R, quantifies the average number of secondary cases per case

during the early ascending phase of an outbreak before the implementation of interventions or

behavior changes (Anderson & May, 1991; Chowell et al., 2015; Yan & Chowell, 2019). Estimates

of the effective R indicate if the disease transmission sustains (R>1) or the disease trend is

declining (R<1). Therefore, it is necessary to maintain R<1 to contain an outbreak. Here, we

estimate the reproduction number by characterizing the early growth phase (16 day) of local cases

using the generalized-growth model (Viboud et al., 2016) and modeling the generation interval of

SARS-CoV-2 assuming a gamma distribution with a mean of 4.41 days and a standard deviation

of 3.17 days (Nishiura, Linton, & Akhmetzhanov, 2020; You et al., 2020). We simulate the

progression of local incident cases by onset dates using the calibrated GGM model and account

for the daily series of imported cases into a renewal equation given as (Nishiura & Chowell, 2009,

2014; Paine et al., 2010):

181
$$R_{t_i} = \frac{I_i}{\sum_{j=0}^{i}(I_{i-j} + \alpha\,J_{i-j})\rho_i}$$

We denote the local incidence at calendar time t_i by I_i, which is characterized using the

generalized-growth model, the imported cases at calendar time t_i by J_i, and the discretized

probability distribution of the generation interval by ρ_i. In this equation the numerator represents

the total new cases I_i, and the denominator represents the total number of cases that contribute to

the new cases I_i at time t_i. The relative contribution of imported cases to the secondary disease

transmission is represented by the parameter $0 \le \alpha \le 1$. We perform a sensitivity analyses by

setting $\alpha = 0.15$ and $\alpha = 1.0$ to assess the relative contribution of imported cases to the secondary

disease transmission (Nishiura & Roberts, 2010). This is followed by the derivation of the

uncertainty bounds around the curve of R directly from the uncertainty associated with the

parameter estimates (r, p). We estimate R for 300 simulated curves assuming a Poisson error

structure (Chowell, 2017). This method to derive early estimates of the reproduction number, R,

has been employed in several prior studies as in refs (Chowell, 2017; Tariq et al., 2020).

3. Results

3.1. COVID-19 testing and positivity rates

Figure 1 shows the daily number of positive and negative laboratory test results and the positivity

rate during the reporting period, March 4th–March 30th, 2020. The total number of PCR tests

performed for this time period were 11518 (1127 positive results and 10307 negative results). The

average daily number of PCR tests performed in Lima was estimated at ~188 between March 4th–

March 15th, 2020 whereas the number of tests performed between March 16th–March 30th, 2020

increased to ~617 tests per day, an increase of 228 % in the testing rates, perhaps reflecting an

increase number of suspected cases with respiratory symptoms. The positivity rate (percentage of

positive cases among the positive and negative cases) has ranged from 0.6-23.9 % between March

4th–March 30th, 2020.

3.2. Local and imported incidence trends

The COVID-19 epidemic curve by the date of symptoms onset, stratified by the local and imported

incidence case counts is shown in Figure 2. On average ~6 imported cases and ~162 local cases

have been reported daily between March 16th–March 30th, 2020 in Lima. A total of 2783

autochthonous cases and 151 imported cases have been reported in Lima as of March 30th, 2020.

3.3. Reproduction number, R

We estimated the reproduction number from the epidemic's early growth phase comprising the

first 16 epidemic days prior to the implementation of social distancing interventions which

includes the national emergency declaration on March 16th 2020. The local incidence curve by the

date of symptoms onset displays near exponential growth dynamics with the scaling of growth

parameter, p, estimated at 0.9 (95% CI: 0.9, 1.0) and the intrinsic growth rate, r, estimated at 0.3

(95% CI: 0.3, 0.5). The estimate of the reproduction number was estimated at 2.3 (95% CI: 2.0,

2.5) when $\alpha = 0.15$ (Figure 3). When $\alpha = 1.0$, the reproduction number slightly decreases to 2.0

(95% CI: 1.7, 2.3) (Table 1).

3.4. Assessing the impact of social distancing interventions

In order to assess the impact of social distancing interventions in Lima, including school closures

on March 11th, 2020 and the declaration of national emergency on March 16th, 2020, we generate

a 20-day ahead forecast for Lima based on the daily incidence curve up until the declaration of the

national emergency in Lima. The 16-day calibration period of the model yields an estimated

growth rate, r, at 0.8 (95% CI: 0.6, 1.1) and a scaling of growth rate parameter, p, at 0.8 (95%CI:

0.7,0.9). The 20-day ahead forecast suggests that the effect of the school closure and the national

emergency declaration slowed down the spread of the virus as shown in Figure 4. Indeed, the

scaling of growth parameter declined to 0.53 (95% CI: 0.48, 0.58), consistent with an

approximately linear incidence growth trend during the period affected by the intervention.

4. Discussion

Our estimate of the transmission potential in Lima for the first 16 days of the epidemic indicates

sustained local transmission in the region after accounting for multiple case importations with the

estimate of reproduction number, R, at ~2.3 (95% CI: 2.0, 2.5) which is comparable to estimates

of the reproduction number for China, Korea, and Iran that lie in the range of 1.5-7.1 (Hwang,

Park, Kim, Jung, & Kim, 2020; Mizumoto, Kagaya, & Chowell, 2020; Muniz-Rodriguez et al.,

2020; Read, Bridgen, Cummings, Ho, & Jewell, 2020; Shim, Tariq, Choi, Lee, & Chowell, 2020;

Wu, Leung, & Leung, 2020). In contrast, a recent study on Singapore's COVID-19 transmission

reported a lower estimate of R at ~0.7, which has been explained as a result of the early

implementation of sweeping social distancing interventions (Tariq et al., 2020).

The initial scaling of growth parameter in Lima indicates a nearly exponential growth pattern,

consistent with the early spread of the COVID-19 epidemic in Iran and the exponential growth

pattern of COVID-19 displayed by the Chinese province of Hubei (Muniz-Rodriguez et al., 2020;

Roosa et al., 2020). In comparison sub-exponential growth patterns of COVID-19 have been

observed in Singapore (p~0.7), Korea (p~0.76) and other Chinese provinces excluding Hubei

(p~0.67) as described in recent studies (Roosa et al., 2020; Shim et al., 2020; Tariq et al., 2020).

Although Lima has been quick to take aggressive measures against COVID-19, Peru remains one

of the hardest hit countries in Latin America (Tegel, 2020). Despite the closure of country borders

on March 16[th], 2020, the number of imported cases in Lima has increased with an average of ~6

imported cases reported between March 16[th]–March 30[th], 2020 compared to an average of ~4

imported cases per day before March 16[th], 2020. However, the 20-day ahead forecast of our GGM

model calibrated to first 16 epidemic days suggest that the social distancing measures, including

closure of schools and the declaration of national emergency are slowing down the virus spread in

Lima. The scaling of growth parameter, p, was estimated at ~0.5 (95%CI: 0.5,0.6) after the

implementation of social distancing measures, consistent with a linear incidence growth trend.

However, the COVID-19 case incidence continues to accumulate despite the quarantine and

lockdowns in the region highlighting the need to enhance social distancing measures to further

contain the outbreak.

The average positivity rate of COVID-19 in Lima was ~8.6% between March 4th–March 30th,

2020. This positivity rate for Lima, Peru, corresponds to the positivity rates derived from Denmark,

Germany and Canada (6-8%) (Meyer & C.Madrigal, 2020). In comparison countries like New

Zealand, South Korea and Australia have tested widely and exhibit lower positivity rates (2%)

whereas Italy and the US have shown much higher positivity rates (15-20%) for COVID-19

indicating suboptimal testing capacity in these countries (Meyer & C.Madrigal, 2020; Project,

2020). A recent study has shown that changes in testing rates over the course of the epidemic can

mask the epidemic growth rate resulting in biased epidemic trends (Omori, Mizumoto, & Chowell,

2020). Moreover, there is a substantial fraction of asymptomatic COVID-19 cases, which could

have underestimated the reproduction number derived from the daily incidence's growth trend of

symptomatic cases (Mizumoto, Kagaya, Zarebski, & Chowell, 2020; Wei et al., 2020). Our study

underscores the need for active contact tracing efforts that targets symptomatic and asymptomatic

cases, rapid isolation of infectious individuals, quarantine contacts and strict social distancing

measures to curb the spread of the virus.

5. Conclusion

In this study we estimate the early transmission potential of SARS-CoV-2 in Lima, Peru. Our

current findings point to sustained transmission of SARS-CoV-2 in the early phase of the outbreak,

with our estimate of the mean reproduction number ~2.3. The COVID-19 epidemic in Lima

followed an early exponential growth trend, which slowed down and turned into an almost linear

growth trend (p~0.5), which appears to be tied to broad scale social distancing interventions put in

place by the government. While the interventions appear to have slowed the transmission rate, the

number of new COVID-19 cases continue to accumulate, highlighting the need to continue social

distancing and active case finding efforts to mitigate disease transmission in the region.

List of abbreviations

COVID-19

SARS-CoV-2

PCR

Conflict of Interest

The authors declare no conflicts of interest.

Funding

Author Contributions

A.T., C.M. and G.C. analyzed the data. A.T. and C.M. retrieved and managed data. A.T and G.C.

wrote the first draft of the manuscript. A.T. and G.C worked on subsequent versions of the

manuscript. All authors contributed to writing and interpretation of results. All authors read and

approved the final manuscript.

Ethical approval

Data has been made available and approved for analysis by the Centro Nacional de Epidemiología

Prevención y control de Enfermedades (CDC Perú) and the National Institute of Health of the

Ministry of Health, Peru.

References

325 Adler, S. E. (2020, February 5). Why Coronaviruses hit older adults hardest. *AARP*. Retrieved
326   from
327   https://cdn.ymaws.com/www.servicecoordinator.org/resource/resmgr/images/misc_/coro
328   navirus_2020/aarp_why_coronaviruses_hit_o.pdf
329 Anderson, R. M., & May, R. M. (1991). *Infectious Diseases of Humans*. Oxford Univeristy
330   Press.
331 Aquino, M., & Garrison, C. (2020, March 6). Peru records first confirmed case of coronavirus,
332   President Vizcarra says. *Rueters*. Retrieved from https://www.reuters.com/article/us-
333   health-coronavirus-peru/peru-records-first-confirmed-case-of-coronavirus-president-
334   vizcarra-says-idUSKBN20T1S9
335 Chowell, G. (2017). Fitting dynamic models to epidemic outbreaks with quantified uncertainty:
336   A primer for parameter uncertainty, identifiability, and forecasts. *Infectious Disease
337   Modelling, 2*(3), 379-398. doi:https://doi.org/10.1016/j.idm.2017.08.001
338 Chowell, G., Abdirizak, F., Lee, S., Lee, J., Jung, E., Nishiura, H., & Viboud, C. (2015).
339   Transmission characteristics of MERS and SARS in the healthcare setting: a comparative
340   study. *BMC Medicine, 13*(1), 210. doi:10.1186/s12916-015-0450-0
341 Ebbs, W. (2020, March 20). U.S. Coronavirus Cases Explode. How Long Before the Hospital
342   System Collapses? *CCN*. Retrieved from https://www.ccn.com/us-coronavirus-cases-
343   soar-past-11000/
344 Explorer, A. (2020, April 6). Coronavirus in Peru- the latest updates. Retrieved from
345   https://amazonas-explorer.com/is-there-coronavirus-in-
346   peru/#Timeline_of_coronavirus_cases_in_Peru
347 Funk, S., Camacho, A., Kucharski, A. J., Eggo, R. M., & Edmunds, W. J. (2018). Real-time
348   forecasting of infectious disease dynamics with a stochastic semi-mechanistic model.
349   *Epidemics, 22*, 56-61. doi:10.1016/j.epidem.2016.11.003
350 Griffiths, J. (2020, April 5). Taiwan's coronavirus response is among the best globally. *CNN*.
351   Retrieved from https://www.cnn.com/2020/04/04/asia/taiwan-coronavirus-response-who-
352   intl-hnk/index.html
353 Group, C.-W. (2020). *Data set for Lima*.
354 Hwang, J., Park, H., Kim, S.-H., Jung, J., & Kim, N. (2020). Basic and effective reproduction
355   numbers of COVID-19 cases in South Korea excluding Sincheonji cases. *medRxiv*,
356   2020.2003.2019.20039347. doi:10.1101/2020.03.19.20039347
357 Meyer, R., & C.Madrigal, A. (2020, April 16). A New Statistic Reveals Why America's COVID-
358   19 Numbers Are Flat. *The Atlantic*. Retrieved from
359   https://www.theatlantic.com/technology/archive/2020/04/us-coronavirus-outbreak-out-
360   control-test-positivity-rate/610132/
361 Minder, R. (2020, April 7). Spain's Coronavirus Crisis Accelerated as Warnings Went
362   Unheeded. *The New York Times*. Retrieved from
363   https://www.nytimes.com/2020/04/07/world/europe/spain-coronavirus.html
364 Mizumoto, K., Kagaya, K., & Chowell, G. (2020). Early epidemiological assessment of the
365   transmission potential and virulence of coronavirus disease 2019 (COVID-19) in Wuhan
366   City: China, January-February, 2020. *medRxiv*, 2020.2002.2012.20022434.
367   doi:10.1101/2020.02.12.20022434
368 Mizumoto, K., Kagaya, K., Zarebski, A., & Chowell, G. (2020). Estimating the asymptomatic
369   proportion of coronavirus disease 2019 (COVID-19) cases on board the Diamond

Princess cruise ship, Yokohama, Japan, 2020. *Euro Surveill, 25*(10)*, 2000180. doi:10.2807/1560-7917.ES.2020.25.10.2000180

MOH. (2020). Sala Situacional COVID-19 Peru. Retrieved from https://covid19.minsa.gob.pe/sala_situacional.asp

Muniz-Rodriguez, K., Fung, I. C.-H., Ferdosi, S. R., Ofori, S. K., Lee, Y., Tariq, A., & Chowell, G. (2020). Transmission potential of COVID-19 in Iran. *medRxiv*, 2020.2003.2008.20030643. doi:10.1101/2020.03.08.20030643

Nishiura, H., & Chowell, G. (2009). *The Effective Reproduction Number as a Prelude to Statistical Estimation of Time-Dependent Epidemic Trends*.

Nishiura, H., & Chowell, G. (2014). Early transmission dynamics of Ebola virus disease (EVD), West Africa, March to August 2014. *Euro Surveill, 19*(36).

Nishiura, H., Linton, N. M., & Akhmetzhanov, A. R. (2020). Serial interval of novel coronavirus (2019-nCoV) infections. *medRxiv*, 2020.2002.2003.20019497. doi:10.1101/2020.02.03.20019497

Nishiura, H., & Roberts, M. G. (2010). Estimation of the reproduction number for 2009 pandemic influenza A(H1N1) in the presence of imported cases. *Eurosurveillance, 15*(29), 19622. doi:doi:https://doi.org/10.2807/ese.15.29.19622-en

Omori, R., Mizumoto, K., & Chowell, G. (2020). Changes in testing rates could mask the novel coronavirus disease (COVID-19) growth rate. *International Journal of Infectious Diseases, 94*, 116-118. doi:10.1016/j.ijid.2020.04.021

Paine, S., Mercer, G., Kelly, P., Bandaranayake, D., Baker, M., Huang, Q., . . . Hope, V. (2010). Transmissibility of 2009 pandemic influenza A(H1N1) in New Zealand: effective reproduction number and influence of age, ethnicity and importations. *Euro Surveill, 15*(24).

Project, C. T. (2020). The COVID-19 tracking project. Retrieved from https://covidtracking.com/about-project

Read, J. M., Bridgen, J. R., Cummings, D. A., Ho, A., & Jewell, C. P. (2020). Novel coronavirus 2019-nCoV: early estimation of epidemiological parameters and epidemic predictions. *medRxiv*, 2020.2001.2023.20018549. doi:10.1101/2020.01.23.20018549

Roosa, K., Lee, Y., Luo, R., Kirpich, A., Rothenberg, R., Hyman, J. M., . . . Chowell, G. (2020). Real-time forecasts of the COVID-19 epidemic in China from February 5th to February 24th, 2020. *Infectious Disease Modelling, 5*, 256-263. doi:https://doi.org/10.1016/j.idm.2020.02.002

Shanafelt, D. W., Jones, G., Lima, M., Perrings, C., & Chowell, G. (2018). Forecasting the 2001 Foot-and-Mouth Disease Epidemic in the UK. *EcoHealth, 15*(2), 338-347. doi:10.1007/s10393-017-1293-2

Shim, E., Tariq, A., Choi, W., Lee, Y., & Chowell, G. (2020). Transmission potential and severity of COVID-19 in South Korea. *International Journal of Infectious Diseases, 93*, 339-344. doi:10.1016/j.ijid.2020.03.031

Tariq, A., Lee, Y., Roosa, K., Blumberg, S., Yan, P., Ma, S., & Chowell, G. (2020). Real-time monitoring the transmission potential of COVID-19 in Singapore, March 2020. *medRxiv*, 2020.2002.2021.20026435. doi:10.1101/2020.02.21.20026435

Team, C. C.-R. (2020). *Preliminary Estimates of the Prevalence of Selected Underlying Health Conditions Among Patients with Coronavirus Disease 2019 — United States, February 12–March 28, 2020*. Retrieved from MMWR Morb Mortal Wkly Rep 2020:

https://www.cdc.gov/mmwr/volumes/69/wr/mm6913e2.htm?s_cid=mm6913e2_w#sugge
stedcitation

Tegel, S. (2020, April 17). Peru took early, aggressive measures against the coronavirus. It's still
suffering one of Latin America's largest outbreaks. *The Washington Post*. Retrieved from
https://www.washingtonpost.com/world/the_americas/peru-took-early-aggressive-
measures-against-the-coronavirus-its-still-suffering-one-of-latin-americas-largest-
outbreaks/2020/04/16/1cd783bc-8005-11ea-84c2-0792d8591911_story.html

Viboud, C., Simonsen, L., & Chowell, G. (2016). A generalized-growth model to characterize
the early ascending phase of infectious disease outbreaks. *Epidemics, 15*, 27-37.
doi:https://doi.org/10.1016/j.epidem.2016.01.002

Wei, W. E., Li, Z., Chiew, C. J., Yong, S. E., Toh, M. P., & Lee, V. J. (2020). *Presymptomatic*
Transmission of SARS-CoV-2 — Singapore, January 23–March 16, 2020. Retrieved from
https://www.cdc.gov/mmwr/volumes/69/wr/mm6914e1.htm?s_cid=mm6914e1_w

WHO. (2020). Situation Reports Coronavirus. Retrieved from
https://www.who.int/emergencies/diseases/novel-coronavirus-2019/situation-reports

Worldometer. (2020). COVID-19 coronavirus / cases. Retrieved from
https://www.worldometers.info/coronavirus/coronavirus-cases/

Writing, M. (2020, March 15). Coronavirus in Peru: Government announces mandatory
quarantine for 15 days for coronavirus. *Gestion*. Retrieved from
https://gestion.pe/peru/politica/coronavirus-en-peru-gobierno-anuncia-cuarentena-
obligatorio-por-15-dias-por-coronavirus-noticia/

Wu, J. T., Leung, K., & Leung, G. M. (2020). Nowcasting and forecasting the potential domestic
and international spread of the 2019-nCoV outbreak originating in Wuhan, China: a
modelling study. *The Lancet*. doi:10.1016/S0140-6736(20)30260-9

Yan, P., & Chowell, G. (2019). *Quantitative methods for infectious disease outbreak*
investigations. Springer.

You, C., Deng, Y., Hu, W., Sun, J., Lin, Q., Zhou, F., . . . Zhou, X.-H. (2020). Estimation of the
Time-Varying Reproduction Number of COVID-19 Outbreak in China. *medRxiv,*
2020.2002.2008.20021253. doi:10.1101/2020.02.08.20021253

Parameter	Estimated values at $\alpha = 1.0$	Estimated values at $\alpha = 0.15$
Reproduction number	2.0 (95% CI: 1.7,2.3)	2.3 (95%CI: 2.0,2.5)
Growth rate, r	0.3 (95%CI: 0.3,0.5)	
Scaling of growth parameter, p	0.9 (95%CI: 0.9,1.0)	

Table 1: Mean estimates and the corresponding 95% confidence intervals for the reproduction number in Lima, growth rate and the scaling of growth parameter during the early growth phase as of March 15th, 2020

480　Figure 1: Laboratory results of COVID-19 tests in Lima as of March 30th, 2020. Blue color

481　represents the negative test results and the yellow color represents the positive test results. The

482　orange solid line denotes the COVID-19 positivity rate in Lima.

496   Figure 2: Daily numbers of new local and imported confirmed COVID-19 cases in Lima by date

497   of symptoms onset as of March 30th, 2020.

512 Figure 3 : The reproduction number derived from the early growth phase in the number of COVID-

513 19 cases in Lima after adjusting for imported cases with $\alpha = 0.15$ using the GGM model as

514 described in the text. The reproduction number based on the incidence curve by March 15[th], 2020

515 was estimated at 2.3 (95% CI: 2.0, 2.5).

Figure 4: 20-day ahead forecast of the COVID-19 epidemic in Lima by calibrating the GGM model

until March 15[th], 2020 (vertical dashed line). Blue circles correspond to the data points, the red

solid line indicates the model's mean fit and the red dashed lines represent the 95% prediction

interval. The vertical black dashed line represents the time of the start of the forecast period. The

forecast (March 16[th]- March 30[th]) suggests that social distancing interventions have slowed down

the transmission rate.