1   **Early transmission dynamics and control of COVID-19 in a southern hemisphere setting:**

2   **Lima-Peru, February 29th-March 30th, 2020 .**

4   César V. Munayco[1#], Amna Tariq[2*#], Gabriela G Soto-Cabezas[1], Mary F. Reyes[1], Andree Valle[1],
5   Leonardo Rojas-Mezarina[3], César Cabezas[3], Manuel Loayza[1], Peru COVID-19 working group,
6   Gerardo Chowell[2]
7
8   [#] joint first authors
9
[1] Centro Nacional de Epidemiología, Prevención y Control de Enfermedades, Peruvian Ministry
of Health, Lima, Peru.
[2] Department of Population Health Sciences, School of Public Health, Georgia State University,
Atlanta, GA, USA
[3] Instituto Nacional de Salud, Peruvian Ministry of Health, Lima, Peru.

Peru COVID-19 working group

Nombre	E-mail	Area
Dante Castro Garro	dante.tcg@gmail.com	Investigación Epidemiológica
Kevin Martel Vasquez	kevin.martel@upch.pe	Investigación Epidemiológica
Edith Solis Castro	esolis@dge.gob.pe	PREEC
Isabel Sandoval Ordinola	isandoval@dge.gob.pe	PREEC
Javier Masias Mimbela	jmasias@dge.gob.pe	PREEC
Kely Meza Cornejo	kmeza@dge.gob.pe	PREEC
Fabiola Caruajulca Quijano	fcaruajulca@dge.gob.pe	PREEC
Lenin La Torre Rosillo	llatorre@dge.gob.pe	PREEC
Luis Ordoñez Ibarguen	lordonez@dge.gob.pe	PREEC
Mario Vasquez Dominguez	mvasquez@dge.gob.pe	PREEC
Rommell Veintimilla Gonzalez Seminario	rommellvgs@gmail.com	PREEC
Martha Calderón Silva	mcalderon@dge.gob.pe	PREEC
Mayra Saavedra Dreyfus	msaavedra@dge.gob.pe	PREEC
Miguel Luna Pineda	mluna@dge.gob.pe	PREEC
Milagros Durand	mdurand@dge.gob.pe	PREEC
Noemí Janampa	njanampa@dge.gob.pe	PREEC
Jesus Chuquihuaccha	jchuquihuaccha@dge.gob.pe	PREEC
Susan Mateo Lizarbe	smateo@dge.gob.pe	PREEC
David Enriquez Cusi	denriquez@dge.gob.pe	PREEC
Idania Mamani Pilco	imamani@dge.gob.pe	PREEC
Alex Jaramillo	ajaramillo@dge.gob.pe	PREEC
Karina Vargas	kvargas@dge.gob.pe	PREEC
Oswaldo Cabanillas	ocabanillas@dge.gob.pe	Alerta-Respuesta
Juan Arrasco	jarrasco@dge.gob.pe	Alerta-Respuesta
María Vargas	mvargas@dge.gob.pe	Vigilancia en Salud Pública

| Willy Ramos | wramos@dge.gob.pe | Inteligencia sanitaria |

* Corresponding author

Amna Tariq

Department of Population Health Sciences

Georgia State University School of Public Health

Atlanta GA, 30303

atariq1@student.gsu.edu

Contact number: 470-985-6352

Abstract

48
The COVID-19 pandemic that emerged in Wuhan China rapidly spread around the world. The
daily incidence trend has been rapidly rising in Latin America since March 2020 with the great
majority of the cases reported in Brazil (28320) followed by Peru (11475) as of April 15th, 2020.
Although Peru implemented social distancing measures soon after the confirmation of its first case
on March 6th, 2020, the daily number of new COVID-19 cases continues to increase. We assessed
the early COVID-19 transmission dynamics and the effect of social distancing interventions in
Lima, Peru.

We estimate the transmission potential of COVID-19, R, during the early phase of the outbreak,
from the daily series of imported and autochthonous cases by the date of symptoms onset as of
March 30th, 2020. We also assessed the effect of social distancing interventions in Lima by
generating short-term forecasts grounded on the early transmission dynamics before interventions
were put in place.

Prior to the implementation of the social distancing measures in Lima, we estimated the
reproduction number at 2.3 (95% CI: 2.0, 2.5). Our analysis indicates that school closures and
other social distancing interventions have helped stem the spread of the virus, with the nearly
exponential growth trend shifting to an approximately linear growth trend after the national
emergency declaration.

The COVID-19 epidemic in Lima followed an early exponential growth trend, which slowed down
and turned into an almost linear growth trend after broad scale social distancing interventions were
put in place by the government.

Key words
COVID-19; SARS-CoV-2; Transmission potential; Short-term forecast; Reproduction number;
Generalized growth model;

1. Introduction

The COVID-19 pandemic that emerged in the city of Wuhan in China in December 2019 (Nishiura, Jung, et al., 2020) has invaded nearly every nation of the world (Mizumoto, Kagaya, & Chowell, 2020; Worldometer, 2020), becoming the most important public health emergency of the last century after the 1918-1920 influenza pandemic. In particular, the novel coronavirus (SARS-CoV-2) has an ability to exert substantial severe disease and mortality burden especially among individuals older than 60 years and those with prior health conditions including hypertension, cardiovascular disease, obesity and diabetes (Adler, 2020; Team, 2020). As of April 15th, 2020, the trajectory of the pandemic varies significantly around the world ranging from relatively contained outbreaks in Thailand, Taiwan and Hong Kong to explosive epidemics characterized by initial exponential growth periods in some hotspots in the United States and European countries including Italy, Spain, UK, and France (Ebbs, 2020; Griffiths, 2020; Minder, 2020).

By April 15th, 2020, SARS-CoV-2 is generating local transmission in over 200 countries with over 2.2 million cases and 150 thousand deaths reported globally (WHO, 2020). The COVID-19 pandemic was confirmed to have reached Latin America in February 2020 with gradual expansion in the region until March 2020 when the COVID-19 incidence curve started to grow more rapidly. By April 15th, 2020, the US still has the highest number of COVID-19 reported cases (637196) globally whereas in Latin America, Brazil has reported the highest number of cases (28320) followed by Peru (11475) (Worldometer, 2020).

Peru, a country located in western South America, reported its confirmed imported index case in Lima on March 6th, 2020, a Peruvian with a travel history to France, Spain and Chez Republic (Aquino & Garrison, 2020). By April 15th, 2020, a total of 11475 cases including 254 deaths have been reported by the Peruvian government. Lima, the capital of Peru has the highest number of COVID-19 reported cases (8412) within Peru (MOH, 2020). To respond to the growing number of COVID-19 cases in Peru, the government shuttered schools on March 11th, 2020. The next day, the government banned gatherings of more than 300 people and suspended all international flights from Europe and Asia. On March 16th, 2020, the government declared a national emergency and closed country borders (Explorer, 2020). Subsequently, on March 17th, 2020 the president of Peru

announced the beginning of community transmission of SARS-CoV-2 in the country, and ordered
a curfew in the region on March 18[th], 2020 to avoid night time socializing to prevent disease
transmission (Explorer, 2020; Writing, 2020).

In order to combat the spread of the COVID-19 epidemic in Lima, the capital and largest city of
Peru, estimates of the transmission potential of COVID-19 can guide the intensity of interventions
including the reproduction number, R, during the early transmission phase (Nishiura & Chowell,
2009, 2014). Moreover, using the epidemiological data and mathematical modeling, it is possible
to gauge the impact of control interventions including school closures and a national emergency
declaration in Lima by assessing short-term forecasts grounded on the trajectory of the epidemic
prior to the implementation of control interventions (Funk, Camacho, Kucharski, Eggo, &
Edmunds, 2018; Shanafelt, Jones, Lima, Perrings, & Chowell, 2018).

2. Methods
2.1. Data
We analyzed the daily number of COVID-19 confirmed cases by date of symptoms onset in Lima,
Peru by March 30[th], 2020. Individual-level case details including whether the case was locally
acquired or imported were also made available from the Centro Nacional de Epidemiología
Prevención y control de Enfermedades (CDC Perú) and the National Institute of Health of the
Ministry of Health, Peru. We also examined the daily testing rate and the positivity rate from the
daily number of positive and negative PCR test results until March 30[th], 2020.

2.2. Model
We generate short term forecasts in real time using the generalized growth model (GGM) that
relies on two parameters and characterizes the early ascending phase of the epidemic allowing to
capture a range of epidemic growth profiles including sub-exponential (polynomial) and
exponential growth. GGM characterizes epidemic growth by estimating two parameters (i) the
intrinsic growth rate, r and (ii) a dimensionless "deacceleration of growth" parameter, p. The latter
parameter modulates the epidemic growth patterns including the sub-exponential growth (p<1)
and exponential growth dynamics (p=1). The GGM model is given by the following differential
equation:

$$141 \qquad \frac{dC(t)}{dt} = C'(t) = rC(t)^p$$

Where $C'(t)$ describes the incidence over time t. The cumulative number of cases at time t is given

by $C(t)$ while r is a positive parameter denoting the growth rate (1/time) and $p \in [0,1]$ is a

"deceleration of growth" parameter (Chowell, 2017; Viboud, Simonsen, & Chowell, 2016).

2.3.Short term forecast to assess interventions

We calibrate the GGM model to the daily case incidence by the date of symptoms onset for Lima.

The data by onset dates for Lima is available from February 29[th], 2020. Therefore, our model was

calibrated using case series from February 29[th]-March 15[th], 2020, prior to the implementation of

national emergency in Lima.

The best fit model solution is estimated by using a non-linear least square fitting approach (full

details provided in (Chowell, 2017)). This process searches for the set of model parameters that

minimizes the sum of squared differences between the observed data $y_{ti} = y_{t1}, y_{t2}, \ldots y_{tn}$ and the

corresponding model solution given by $f(t_i, \Theta)$: where $\Theta_{GGM} = (r, p)$ correspond to estimated set

of parameters of the GGM model. Thus, the objective function for the best fit solution of $f(t_i, \Theta)$

is given by

$$\widehat{\Theta} = \text{arg min} \sum_{i=1}^{n} (f(t_i, \Theta) - y_{ti})^2$$

The initial condition is fixed to the first observation in the data. Next, we derive uncertainty around

the best fit model solution as well as the confidence intervals of the parameters utilizing a

parametric bootstrapping approach assuming a Poisson error structure as described in ref (Chowell,

2017).

2.4. Reproduction number from case incidence using the generalized-growth model

The reproduction number, R, quantifies the variation in the average number of secondary cases

generated by a primary case during the early ascending phase of an outbreak before the

implementation of interventions or behavior changes in a completely susceptible population

(Anderson & May, 1991; Chowell et al., 2015; Nishiura, Chowell, Heesterbeek, & Wallinga, 2010;

Yan & Chowell, 2019). Estimates of R indicate if the disease transmission sustains (R>1) or the
disease trend is declining (R<1). Therefore, it is imperative to maintain the R<1 to contain an
outbreak. The reproduction number is estimated by calibrating the GGM to the early growth phase
of the epidemic (16 days) (Viboud et al., 2016), using the information on the gamma distribution
of the generation interval of SARS-CoV-2 with a mean of 4.41 days and a standard deviation of
3.17 days (Nishiura, Linton, & Akhmetzhanov, 2020; You et al., 2020). Then we simulate the
progression of local incident cases by onset dates from the calibrated GGM model incorporating
the imported cases and apply the discretized probability distribution of generation interval using
renewal equation given as (Nishiura & Chowell, 2009, 2014; Paine et al., 2010):

180
$$R_{0_i} = \frac{I_i}{\sum_{j=0}^{i}(I_{i-j} + \alpha J_{i-j})\rho_i}$$

We denote the local incidence at calendar time t_i by I_i, the incidence of imported cases at calendar
time t_i by J_i , and the discretized probability distribution of the generation interval by ρ_i. In this
equation the numerator represents the total new cases I_i, and the denominator represents the total
number of cases that contribute to the new cases I_i at time t_i. The relative contribution of imported
cases to the secondary disease transmission is represented by the parameter $0 \leq \alpha \leq 1$. We perform
a sensitivity analyses by setting $\alpha = 0.15$ and $\alpha = 1.0$ to assess the contribution of imported cases
to the secondary disease transmission (Nishiura & Roberts, 2010). This is followed by the
derivation of the uncertainty bounds around the curve of R directly from the uncertainty associated
with the parameter estimates (r, p). We estimate R for 300 simulated curves assuming a Poisson
error structure (Chowell, 2017).

3. Results

3.1. COVID-19 testing rates and positivity rate

Figure 1 shows the daily number of positive and negative laboratory test results and the positivity
rate during March 4[th]-March 30[th], 2020. The total number of PCR tests performed for this time
period were 11518 (1127 positive results and 10307 negative results). The average daily number
of PCR tests performed in Lima was estimated at ~188 between March 4[th]-March 15[th], 2020
whereas the number of tests performed between March 16[th]-March 30[th], 2020 increased to ~617
tests per day. Subsequently, the positivity rate (percentage of positive cases among the positive
and negative cases) has ranged from 0.6-23.9 % between March 4[th]-March 30[th], 2020.

3.2. Local and imported incidence trends

The COVID-19 epidemic curve by the date of symptoms onset, stratified by the local and imported incidence case counts is shown in Figure 2. On average ~6 imported cases and ~162 local cases have been reported daily between March 16th-March 30th, 2020 in Lima. A total of 2783 autochthonous cases and 151 imported cases have been reported in Lima as of March 30th, 2020.

3.3. Reproduction number, R

We estimated the reproduction number using the first 16 epidemic days prior to the implementation of social distancing interventions which includes the national emergency declaration on March 16th 2020. The local incidence curve by the date of onset displays near exponential growth dynamics with the scaling of growth parameter, p, estimated at 0.9 (95% CI: 0.8, 1.0) and the intrinsic growth rate, r, estimated at 0.3 (95% CI: 0.3, 0.5). The estimate of the reproduction number is 2.3 (95% CI: 2.0, 2.5) when $\alpha = 0.15$ (Figure 3). When $\alpha = 1.0$, the reproduction number slightly decreases to 2.0 (95% CI: 1.7, 2.3) (Table 1).

3.4. Assessing the impact of social distancing interventions

In order to assess the impact of social distancing interventions in Lima, including school closures on March 11th, 2020 and the declaration of national emergency on March 16th, 2020, we generate a 20-day ahead forecast for Lima based on the daily incidence curve until the declaration of the national emergency in Lima. The 16-day calibration period of the model yields an estimated growth rate, r, at 0.8 (95%CI: 0.6, 1.1) and a scaling of growth rate parameter, p, at 0.8 (95%CI: 0.7,0.9). The 20-day ahead forecast suggests that the effect of the school closure and the national emergency declaration slowed down the spread of the virus as shown in Figure 4. Indeed, the scaling of growth parameter declined to 0.53 (95% CI: 0.48, 0.58), consistent with an approximately linear incidence growth trend during the intervention period.

4. Discussion

The current estimates of the early transmission potential in Lima for the first 16 days of the epidemic indicates sustained local transmission in the region after accounting for multiple case importations with the estimate of reproduction number R at ~2.3 (95% CI: 2.0, 2.5) which is

comparable to estimates of the reproduction number for China, Korea, and Iran that lie in the range
of 1.5-7.1 (Hwang, Park, Kim, Jung, & Kim, 2020; Mizumoto, Kagaya, & Chowell, 2020; Muniz-
Rodriguez et al., 2020; Read, Bridgen, Cummings, Ho, & Jewell, 2020; Shim, Tariq, Choi, Lee,
& Chowell, 2020; Wu, Leung, & Leung, 2020). In contrast, a recent study on Singapore's COVID-
19 transmission reported a lower estimate of R at ~0.7, which has been explained as a result of the
implementation of a sweeping social distancing interventions (Tariq et al., 2020).
238
The initial scaling of growth parameter in Lima indicates a nearly exponential growth pattern,
consistent with the early spread of the COVID-19 epidemic in Iran and the exponential growth
pattern of COVID-19 displayed by the Chinese province of Hubei (Muniz-Rodriguez et al., 2020;
Roosa et al., 2020). In comparison sub-exponential growth patterns of COVID-19 have been
observed in Singapore (p~0.7), Korea (p~0.76) and other Chinese provinces excluding Hubei
(p~0.67) as described in recent studies (Roosa et al., 2020; Shim et al., 2020; Tariq et al., 2020).
245
Although Lima has been quick to take aggressive measures against COVID-19, Peru remains one
of the hardest hit countries in Latin America (Tegel, 2020). Despite the closure of country borders
on March 16[th], 2020 imported cases in Lima have increased with an average of ~6 imported cases
reported between March 16[th]-March 30[th], 2020 in comparison to an average of ~4 imported cases
reported per day before March 16[th], 2020. As of April 2[nd], 2020, Peru has adopted the gender-
based quarantine measures for a week, where men and women were assigned particular days of
the week to leave the house to buy groceries and medicine, in an attempt to further slow the spread
of COVID-19 in the region (Westcott et al., 2020). The 20-day ahead forecast of our GGM model
calibrated to 16 days suggest that the social distancing measures, including closure of schools and
the declaration of national emergency are helping slow down the virus spread in Lima. Moreover,
the scaling of growth parameter, p was estimated at ~0.5 (95%CI: 0.5,0.6) after the implementation
of social distancing measures, consistent with a linear incidence growth trend. However, the
COVID-19 case incidence continues to accumulate despite the quarantine and lockdowns in the
region highlighting the need to enhance social distancing measures to further contain the outbreak.
260
The average positivity rate of COVID-19 in Lima was ~8.6% between March 4[th]-March 30[th], 2020.
This positivity rate for Lima, Peru, corresponds to the positivity rates derived from Denmark,

Germany and Canada (6-8%) (Meyer & C.Madrigal, 2020). In comparison countries like New

Zealand, South Korea and Australia have exhibited lower positivity rates (2%) whereas Italy and

the US have shown much higher positivity rates (15-20%) for COVID-19 (Meyer & C.Madrigal,

2020). Moreover, there is a substantial fraction of asymptomatic COVID-19 cases, which could

have underestimated our estimates that relied on the daily incidence's growth trend from

symptomatic cases (Mizumoto, Kagaya, Zarebski, & Chowell, 2020; Wei et al., 2020). On the

other hand, it is not known up to what extent asymptomatic cases transmit the virus relative to

symptomatic cases. Our study underscores the need for active contact tracing efforts, isolation and

strict social distancing measures to contain the outbreak and mitigate active disease transmission

in Peru.

5. Conclusion

In this study we estimate the early transmission potential of SARS-CoV-2 in Lima, Peru. Our

current findings point to sustained transmission of SARS-CoV-2 in the early phase of the outbreak,

with our estimate of the mean reproduction number ~2.3. The COVID-19 epidemic in Lima

followed an early exponential growth trend, which slowed down and turned into an almost linear

growth trend (p~0.5) after broad scale social distancing interventions were put in place by the

government. While the interventions have slowed the virus spread, the number of new COVID-19

cases continue to accumulate, highlighting the need to continue social distancing and active case

finding efforts to mitigate disease transmission in the region.

List of abbreviations

COVID-19

SARS-CoV-2

PCR

Conflict of Interest

The authors declare no conflicts of interest.

Funding

G.C. is partially supported from NSF grants 1610429 and 1633381 and R01 GM 130900.

Author Contributions

A.T., C.V. and G.C. analyzed the data. A.T. and C.M. retrieved and managed data. A.T and G.C.
296
wrote the first draft of the manuscript. A.T. and G.C worked on subsequent versions of the
297
manuscript. All authors contributed to writing and interpretation of results. All authors read and
298
approved the final manuscript.
299

Ethical approval

Data has been made available and approved for analysis by the Centro Nacional de Epidemiología
303
Prevención y control de Enfermedades (CDC Perú) and the National Institute of Health of the
304
Ministry of Health, Peru.
305

References

Adler, S. E. (2020, February 5). Why Coronaviruses hit older adults hardest. *AARP*. Retrieved
from
https://cdn.ymaws.com/www.servicecoordinator.org/resource/resmgr/images/misc_/coro
navirus_2020/aarp_why_coronaviruses_hit_o.pdf
Anderson, R. M., & May, R. M. (1991). *Infectious Diseases of Humans*. Oxford Univeristy
Press.
Aquino, M., & Garrison, C. (2020, March 6). Peru records first confirmed case of coronavirus,
President Vizcarra says. *Rueters*. Retrieved from https://www.reuters.com/article/us-
health-coronavirus-peru/peru-records-first-confirmed-case-of-coronavirus-president-
vizcarra-says-idUSKBN20T1S9
Chowell, G. (2017). Fitting dynamic models to epidemic outbreaks with quantified uncertainty:
A primer for parameter uncertainty, identifiability, and forecasts. *Infectious Disease*
Modelling, 2(3), 379-398. doi:https://doi.org/10.1016/j.idm.2017.08.001
Chowell, G., Abdirizak, F., Lee, S., Lee, J., Jung, E., Nishiura, H., & Viboud, C. (2015).
Transmission characteristics of MERS and SARS in the healthcare setting: a comparative
study. *BMC Medicine, 13*(1), 210. doi:10.1186/s12916-015-0450-0
Ebbs, W. (2020, March 20). U.S. Coronavirus Cases Explode. How Long Before the Hospital
System Collapses? *CCN*. Retrieved from https://www.ccn.com/us-coronavirus-cases-
soar-past-11000/
Explorer, A. (2020, April 6). Coronavirus in Peru- the latest updates. Retrieved from
https://amazonas-explorer.com/is-there-coronavirus-in-
peru/#Timeline_of_coronavirus_cases_in_Peru
Funk, S., Camacho, A., Kucharski, A. J., Eggo, R. M., & Edmunds, W. J. (2018). Real-time
forecasting of infectious disease dynamics with a stochastic semi-mechanistic model.
Epidemics, 22, 56-61. doi:10.1016/j.epidem.2016.11.003
Griffiths, J. (2020, April 5). Taiwan's coronavirus response is among the best globally. *CNN*.
Retrieved from https://www.cnn.com/2020/04/04/asia/taiwan-coronavirus-response-who-
intl-hnk/index.html
Hwang, J., Park, H., Kim, S.-H., Jung, J., & Kim, N. (2020). Basic and effective reproduction
numbers of COVID-19 cases in South Korea excluding Sincheonji cases. *medRxiv*,
2020.2003.2019.20039347. doi:10.1101/2020.03.19.20039347
Meyer, R., & C.Madrigal, A. (2020, April 16). A New Statistic Reveals Why America's COVID-
19 Numbers Are Flat. *The Atlantic*. Retrieved from
https://www.theatlantic.com/technology/archive/2020/04/us-coronavirus-outbreak-out-
control-test-positivity-rate/610132/
Minder, R. (2020, April 7). Spain's Coronavirus Crisis Accelerated as Warnings Went
Unheeded. *The New York Times*. Retrieved from
https://www.nytimes.com/2020/04/07/world/europe/spain-coronavirus.html
Mizumoto, K., Kagaya, K., & Chowell, G. (2020). Early epidemiological assessment of the
transmission potential and virulence of coronavirus disease 2019 (COVID-19) in Wuhan
City: China, January-February, 2020. *medRxiv*, 2020.2002.2012.20022434.
doi:10.1101/2020.02.12.20022434
Mizumoto, K., Kagaya, K., Zarebski, A., & Chowell, G. (2020). Estimating the asymptomatic
proportion of coronavirus disease 2019 (COVID-19) cases on board the Diamond

370   Princess cruise ship, Yokohama, Japan, 2020. *Euro Surveill, 25*(10), 2000180.
371   doi:10.2807/1560-7917.ES.2020.25.10.2000180
372 MOH. (2020). Sala Situacional COVID-19 Peru. Retrieved from
373   https://covid19.minsa.gob.pe/sala_situacional.asp
374 Muniz-Rodriguez, K., Fung, I. C.-H., Ferdosi, S. R., Ofori, S. K., Lee, Y., Tariq, A., & Chowell,
375   G. (2020). Transmission potential of COVID-19 in Iran. *medRxiv*,
376   2020.2003.2008.20030643. doi:10.1101/2020.03.08.20030643
377 Nishiura, H., & Chowell, G. (2009). *The Effective Reproduction Number as a Prelude to*
378   *Statistical Estimation of Time-Dependent Epidemic Trends*.
379 Nishiura, H., & Chowell, G. (2014). Early transmission dynamics of Ebola virus disease (EVD),
380   West Africa, March to August 2014. *Euro Surveill, 19*(36).
381 Nishiura, H., Chowell, G., Heesterbeek, H., & Wallinga, J. (2010). The ideal reporting interval
382   for an epidemic to objectively interpret the epidemiological time course. *Journal of the*
383   *Royal Society, Interface, 7*(43), 297-307. doi:10.1098/rsif.2009.0153
384 Nishiura, H., Jung, S., Linton, N., Kinoshita, R., Yang, Y., Hayashi, K., . . . Akhmetzhanov, R.
385   (2020). The Extent of Transmission of Novel Coronavirus in Wuhan, China, 2020.
386   *Journal of Clinical Medicine, 9*(2), 330. Retrieved from https://www.mdpi.com/2077-
387   0383/9/2/330
388 Nishiura, H., Linton, N. M., & Akhmetzhanov, A. R. (2020). Serial interval of novel coronavirus
389   (2019-nCoV) infections. *medRxiv*, 2020.2002.2003.20019497.
390   doi:10.1101/2020.02.03.20019497
391 Nishiura, H., & Roberts, M. G. (2010). Estimation of the reproduction number for 2009
392   pandemic influenza A(H1N1) in the presence of imported cases. *Eurosurveillance,*
393   *15*(29), 19622. doi:doi:https://doi.org/10.2807/ese.15.29.19622-en
394 Paine, S., Mercer, G., Kelly, P., Bandaranayake, D., Baker, M., Huang, Q., . . . Hope, V. (2010).
395   Transmissibility of 2009 pandemic influenza A(H1N1) in New Zealand: effective
396   reproduction number and influence of age, ethnicity and importations. *Euro Surveill,*
397   *15*(24).
398 Read, J. M., Bridgen, J. R., Cummings, D. A., Ho, A., & Jewell, C. P. (2020). Novel coronavirus
399   2019-nCoV: early estimation of epidemiological parameters and epidemic predictions.
400   *medRxiv*, 2020.2001.2023.20018549. doi:10.1101/2020.01.23.20018549
401 Roosa, K., Lee, Y., Luo, R., Kirpich, A., Rothenberg, R., Hyman, J. M., . . . Chowell, G. (2020).
402   Real-time forecasts of the COVID-19 epidemic in China from February 5th to February
403   24th, 2020. *Infectious Disease Modelling, 5*, 256-263.
404   doi:https://doi.org/10.1016/j.idm.2020.02.002
405 Shanafelt, D. W., Jones, G., Lima, M., Perrings, C., & Chowell, G. (2018). Forecasting the 2001
406   Foot-and-Mouth Disease Epidemic in the UK. *EcoHealth, 15*(2), 338-347.
407   doi:10.1007/s10393-017-1293-2
408 Shim, E., Tariq, A., Choi, W., Lee, Y., & Chowell, G. (2020). Transmission potential and
409   severity of COVID-19 in South Korea. *International Journal of Infectious Diseases, 93*,
410   339-344. doi:10.1016/j.ijid.2020.03.031
411 Tariq, A., Lee, Y., Roosa, K., Blumberg, S., Yan, P., Ma, S., & Chowell, G. (2020). Real-time
412   monitoring the transmission potential of COVID-19 in Singapore, March 2020. *medRxiv*,
413   2020.2002.2021.20026435. doi:10.1101/2020.02.21.20026435
414 Team, C. C.-R. (2020). *Preliminary Estimates of the Prevalence of Selected Underlying Health*
415   *Conditions Among Patients with Coronavirus Disease 2019 — United States, February*

12–March 28, 2020. Retrieved from MMWR Morb Mortal Wkly Rep 2020:
https://www.cdc.gov/mmwr/volumes/69/wr/mm6913e2.htm?s_cid=mm6913e2_w#sugge
stedcitation
Tegel, S. (2020, April 17). Peru took early, aggressive measures against the coronavirus. It's still
suffering one of Latin America's largest outbreaks. *The Washington Post.* Retrieved from
https://www.washingtonpost.com/world/the_americas/peru-took-early-aggressive-
measures-against-the-coronavirus-its-still-suffering-one-of-latin-americas-largest-
outbreaks/2020/04/16/1cd783bc-8005-11ea-84c2-0792d8591911_story.html
Viboud, C., Simonsen, L., & Chowell, G. (2016). A generalized-growth model to characterize
the early ascending phase of infectious disease outbreaks. *Epidemics, 15*, 27-37.
doi:https://doi.org/10.1016/j.epidem.2016.01.002
Wei, W. E., Li, Z., Chiew, C. J., Yong, S. E., Toh, M. P., & Lee, V. J. (2020). *Presymptomatic*
Transmission of SARS-CoV-2 — Singapore, January 23–March 16, 2020. Retrieved from
https://www.cdc.gov/mmwr/volumes/69/wr/mm6914e1.htm?s_cid=mm6914e1_w
Westcott, B., Regan, H., Renton, A., Picheta, R., Wagner, M., & Hayes, M. (2020, April 2).
April 2 coronavirus news. *CNN.* Retrieved from https://www.cnn.com/world/live-
news/coronavirus-pandemic-04-02-20-intl/index.html
WHO. (2020). Situation Reports Coronavirus. Retrieved from
https://www.who.int/emergencies/diseases/novel-coronavirus-2019/situation-reports
Worldometer. (2020). COVID-19 coronavirus / cases. Retrieved from
https://www.worldometers.info/coronavirus/coronavirus-cases/
Writing, M. (2020, March 15). Coronavirus in Peru: Government announces mandatory
quarantine for 15 days for coronavirus. *Gestion.* Retrieved from
https://gestion.pe/peru/politica/coronavirus-en-peru-gobierno-anuncia-cuarentena-
obligatorio-por-15-dias-por-coronavirus-noticia/
Wu, J. T., Leung, K., & Leung, G. M. (2020). Nowcasting and forecasting the potential domestic
and international spread of the 2019-nCoV outbreak originating in Wuhan, China: a
modelling study. *The Lancet.* doi:10.1016/S0140-6736(20)30260-9
Yan, P., & Chowell, G. (2019). *Quantitative methods for infectious disease outbreak*
investigations. Springer.
You, C., Deng, Y., Hu, W., Sun, J., Lin, Q., Zhou, F., . . . Zhou, X.-H. (2020). Estimation of the
Time-Varying Reproduction Number of COVID-19 Outbreak in China. *medRxiv,*
2020.2002.2008.20021253. doi:10.1101/2020.02.08.20021253

Parameter	Estimated values at $\alpha = 1.0$	Estimated values at $\alpha = 0.15$
Reproduction number	2.0 (95% CI: 1.7,2.3)	2.3 (95%CI: 2.0,2.5)
Growth rate, r	0.3 (95%CI: 0.3,0.5)	
Scaling of growth parameter, p	0.9 (95%CI: 0.8,1.0)	

458   Table 1: Mean estimates and the corresponding 95% confidence intervals for the reproduction

459   number, growth rate and the scaling of growth parameter during the early growth phase as of

460   March 15th, 2020

Figure 1: Laboratory results for the COVID-19 cases in Lima as of March 30th, 2020. Blue color

represents the negative cases and the yellow color represents the positive cases. The orange solid

line represents the positivity rate of the cases in Lima.

Figure 2: Local and imported incidence for confirmed COVID-19 cases in Lima by the date of symptoms onset as of March 30th, 2020.

Figure 3 : The reproduction number with 95% CI estimated by adjusting for the imported cases $\alpha = 0.15$ using the GGM model. The reproduction number based on the incidence curve by March 15th, 2020 is at 2.3 (95% CI: 2.0, 2.5).

516 Figure 4: 20-day ahead forecast of the COVID-19 epidemic in Lima by calibrating the model until

517 March 15th, 2020. Blue circles correspond to the data points, the red solid line indicates the best

518 model fit and the red dashed lines represent the 95% prediction interval. The vertical black dashed

519 line represents the time of the start of the forecast period.