

Spatial Inequities in COVID-19 outcomes in 3 US Cities

Usama Bilal^{1,2}, Sharrelle Barber^{1,2}, Ana V. Diez-Roux^{1,2}

1.- Urban Health Collaborative, Drexel Dornsife School of Public Health, Philadelphia, PA

2.- Department of Epidemiology and Biostatistics, Drexel Dornsife School of Public Health, Philadelphia, PA

Corresponding Author: Usama Bilal, MD MPH PhD

Address: 3600 Market St. Suite 730, Philadelphia, PA, 19104

Email: ubilal@drexel.edu

Phone: +1 267-359-6378

Abstract

Background: Preliminary evidence has shown wide inequities in COVID-19 related deaths in the US. We explored the emergence of spatial inequities in COVID-19 testing, positivity, and incidence in New York City, Philadelphia, and Chicago.

Methods: We used zip code-level data on cumulative tests and confirmed cases by date for each city and computed testing, positivity, and incidence indicators. We linked these to 2014-2018 American Community Survey data on income, education, race/ethnicity, occupation, health insurance, and overcrowding, and computed a summary index. We computed associations between using Poisson models. We also examined clusters of high and low incidence using the G^* statistic.

Results: Through May 18th, there were wide inequities in positivity and incidence, with less advantaged neighborhoods having a higher incidence (RR=1.36 [95% CrI 1.18;1.57], 1.17 [1.11;1.23], and 1.10 [0.98;1.23], per 1 SD increase in the summary index in Chicago, NYC and Philadelphia, respectively). In all three cities inequities in incidence increased as the pandemic advanced, while inequities in positivity remained stable. In contrast the social patterning of testing changed over time: testing was inversely associated with disadvantage early in the pandemic but was either not associated or positively associated with disadvantage later in the pandemic. We also found clusters of high and low incidence, co-located with areas of high and low disadvantage.

Conclusions: We found wide spatial inequities in COVID-19 positivity and incidence in three large metropolitan areas of the US. In health crises health inequities become magnified and reflect a longstanding history of racial and economic injustice.

Keywords: COVID-19, SARS-COV-2, urban health, health disparities, spatial inequities, neighborhoods, health equity

Introduction

As of May 26th, 2020, the COVID-19 pandemic had taken the lives of more than 340,000 people worldwide, while in the US deaths are approaching 100,000¹. Cities across the globe have emerged as especially vulnerable to COVID-19 and as locations where social and spatial inequities in COVID-19 incidence and mortality are quickly emerging. For example, there is a much higher incidence rate of COVID-19 in the poor compared to the wealthier neighborhoods of Barcelona, one of the most affected cities in Spain². Reports from lower- and middle- income countries are also revealing higher COVID rates in informal settlements within cities³.

Preliminary research has shown very wide inequities in COVID-19 incidence and mortality in the US. Around 20% of US counties have a majority Black population, and they account for more than half of all COVID-19 cases⁴. In states reporting data by race/ethnicity, Black Americans are only 13% of the population but represent 27% of the COVID-19 related deaths.⁵ These inequities are especially worrisome in some US cities with very high incidence. In New York City, both Blacks and Latinxs have double the mortality rate as compared to whites after age-adjustment⁶, in Chicago 50% of deaths have occurred in Blacks, who make only 30% of the population⁷, while in Milwaukee Blacks account for 69% of deaths while accounting for only 27% of the population⁸.

Characterizing social and spatial inequities in cities is critical to understanding the racial inequities that have emerged and developing appropriate interventions and policies to prevent COVID-19 deaths in the future. Yet it is rendered complex during an evolving pandemic because of the interrelated nature of access to testing and diagnosis and because the social patterning of the pandemic is likely to change as it advances through the population. We used data from

three large US cities to (1) characterize spatial and social inequities in testing, positivity, and incidence and (2) examine how the social patterning has evolved in different cities as the pandemic progressed through them.

Methods

Setting

We used data on cumulative total numbers of tests and confirmed cases by zip code of residence from Chicago, New York City (NYC), and Philadelphia. For Chicago, we downloaded data compiled by the Illinois Department of Public Health⁹ and made available by the Chicago Observer¹⁰, and obtained daily cumulative data from April 18th through May 18th for the entire state of Illinois, and restricted the data to zip codes that intersected with the city of Chicago.

We followed a similar approach for New York City, where we downloaded daily cumulative data compiled by the NYC Department of Health and Mental Hygiene in their GitHub repository¹¹ from April 1st through May 18th. For Philadelphia, we downloaded data from the Philadelphia Department of Public Health¹² on April 24th, including all tests and confirmed cases prior to that date, by zip code and result date. We then calculated cumulative number of confirmed cases and total tests by zip code and date from March 21st through April 23rd. From thereon we followed an approach similar to the other three cities, and downloaded daily cumulative number of total tests and confirmed cases by zip code¹².

Predictors

We linked zip code data to the 2014-2018 American Community Survey. We obtained data on variables that we hypothesized could be linked to differential exposure to SARS-COV-2 or to differential testing practices. Specifically, we calculated six variables reflecting income (median household income [log]), race/ethnicity (% non-Hispanic White), education (% with college education), occupation (% working in service jobs [food preparation and serving, personal care, and other service occupations]), health insurance status (% uninsured), and housing (%

overcrowded households [>1 person per room]). To provide a summary of all six variables, we calculated extracted the first component from a principal component analysis (PCA) with all six variables (Appendix Table 1 shows the loadings for each variable and city). Median household income, % non-Hispanic White, and % college education loaded negatively on this index, while % working in service jobs, % uninsured, and % overcrowded households loaded positively. Therefore, a higher value in this summary index represents a higher concentration of relative disadvantage, while a lower value represents a higher concentration of relative advantage.

Outcomes

As the main outcomes, we computed three indicators that reflect different testing practices and infection burden: (1) testing per capita (total tests/population); (2) positivity rate¹³ (confirmed cases/total tests); and (3) incidence (confirmed cases/population).

Analysis

We conducted our analysis in three steps. First, we conducted a graphical assessment of correlations between each of the seven variables (six predictors and the summary index) and the three outcomes (testing, positivity, and incidence).

Second, similar to other work¹⁴, we estimated the strength of the associations between each predictor and outcome using a Poisson model. To make coefficients comparable, we standardized all predictors by subtracting the mean and dividing over the standard deviation (SD) for each city. To account for spatial autocorrelation of the outcomes, we fitted a conditional autoregressive generalized linear mixed model. The resulting coefficients and standard errors can be used to estimate rate ratios of testing, positivity and incidence (and 95% credible intervals) per 1 SD increase in each predictor. Third, to explore temporal trends in

inequities, we repeated the second step for every date from April 1st through May 18th, 2020 and plotted rate ratios for each outcome cumulative through each date associated with a one SD higher summary index, separately for each city.

Fourth, to show spatial patterns in the outcomes we mapped zip code-level outcomes and the summary index for each city, for the last available date (May 18th, 2020). To show clusters of high and low incidence, testing, positivity, and disadvantage, we computed the G* star statistic, and show clusters of high or low levels with a Bonferroni-adjusted p-value of $p < 0.05$.

All analyses were conducted using R v4.0. Code for replication is available at:

https://github.com/usamabila/COVID_Disparities

Results

We included a total of 177, 84, and 46 zip codes in NYC, Chicago and Philadelphia, respectively. For NYC, we had cumulative testing data for 44 days from April 1st through May 18th; for Chicago we had cumulative testing data for 30 days from April 18th through May 18th; for Philadelphia we had cumulative testing data for 54 days, from March 21st through May 18th.

Cumulative data through May 18th, 2020, showed that COVID-19 testing outcomes were not correlated with zip code-level socioeconomic variables (**Figure 1, Appendix Figures 1-4**). In Chicago, NYC and Philadelphia, a 1 SD higher zip code-level summary index was not associated with higher testing rates (RR=0.99, 95% CrI 0.91 to 1.08, RR=1.01, 95% CrI 0.98 to 1.06, RR=0.95, 95% CrI 0.87 to 1.03, for Chicago, NYC, and Philadelphia respectively) (**Table 1**).

We found strong inequities in positivity rates and incidence for the three cities by May 18th. A higher zip code-level median household income, % college educated, and % non-Hispanic white, and a lower % uninsured, % service workers, and % overcrowding, were all associated with a higher positivity rate for all three cities. Specifically, a 1 SD higher zip code-level summary index was associated with a 33%, 13%, and 14% higher positivity rate and a 36%, 17%, and 10% higher incidence in Chicago, NYC and Philadelphia, respectively (**Table 1**).

These inequities were dynamic and evolved over time (**Figure 2**). Cumulatively through April 1, New York had a weak inequity in incidence and testing, and a strong inequity in positivity rates, whereby higher value of the summary index was associated with higher incidence and positivity but lower testing. As the pandemic evolved and testing became more widespread, the inequity in testing was reversed, so that by late April cumulative testing per capita was higher in less advantaged neighborhoods. Given that the inequity in positivity rate

stayed constant over time, the inequity in incidence became stronger. Chicago followed a similar pattern, with a reversal of the inequity in testing, a stable inequity in the positivity rate, and a strengthening of the inequity in incidence.

While Philadelphia followed similar trends to the other cities, data availability in Philadelphia allowed us to explore the inequities earlier in the epidemic. Specifically, during all of April in Philadelphia there was a strong inequity in testing, with less advantaged zip codes receiving a much lower amount of testing (RR=0.79, 95% CrI 0.72 to 0.87 by April 1st) (**Figure 2**). However, this association changed over time, and by May 18th the inequity in testing was very weak or non-existent (RR=0.94, 95% CrI 0.87 to 1.02). Although the inequity in each city differed over time, we found that inequities in testing and incidence were similar in NYC and Philadelphia when dates were matched based on the levels of cumulative incidence at the city level, while they were stronger overall for Chicago at any given time (**Appendix Figure 5**).

There was a clear spatial pattern of incidence on May 18th, and this pattern strongly follows the spatial patterning of advantage, especially in New York City and Chicago (**Figure 3**). There are clusters of high and low incidence in the three cities (**Figure 4**). In Chicago, there is a large cluster of high incidence in the West and Southwest side of the city, and a cluster of low incidence in the North side and Central parts of the city (**Figure 4, Panel A**). In New York, there is a cluster of low incidence in most of Manhattan and parts of Brooklyn, while there are several clusters of high incidence in most of the Bronx and Staten Island, and parts of Brooklyn and Queens (**Figure 4, Panel B**). In Philadelphia, there was a cluster of low incidence in most of Center City and other areas along the Delaware river, while there were clusters of high

incidence in West, North and Northeast Philadelphia (**Figure 4, Panel C**). **Appendix Figures 6-8** show the spatial clustering of positivity rates, testing, and disadvantage.

Discussion

We documented large spatial inequities in COVID-19 outcomes through May 18 in three large US cities with more deprived neighborhoods having a higher incidence and positivity rates. These inequities were strongest in Chicago and similar in New York City and Philadelphia. We also found clusters of high and low incidence, co-located with areas of low and high advantage, respectively. In all three cities, inequities in incidence increased as the pandemic advanced, while inequities in positivity rates remained stable. In contrast the social patterning of testing changed over time: testing was more common in more advantage neighborhoods early in the pandemic, while it became more common in less advantaged neighborhoods later in the pandemic.

Potential explanations for inequities in incidence include a differential exposure to the virus and a differential susceptibility to infection due to a longstanding history of structural racism and residential segregation¹⁵⁻¹⁷. Differential exposures could emerge from occupation, such as healthcare¹⁸, personal care or service occupations¹⁹, lack of telecommuting options²⁰, mass transit use²¹, and overcrowding within households²². Utility disruptions leading to a lack of access to running water can also introduce barriers to hand washing in cities like Detroit²³. Whether there are factors associated with differential susceptibility to infection is still unclear, but prior research on respiratory viruses has documented that stress linked to disadvantage may increase the likelihood of developing disease after exposure^{24,25}.

We found inequities in testing at the beginning stages of the epidemic by which deprived neighborhoods had less per capita testing, although these inequities were reduced as the epidemic progressed. This change in the social patterning of testing was most pronounced

in Philadelphia, for which data earlier in the pandemic was available. Barriers to testing when resources are constrained can include unequal location of testing sites²⁶, lack of vehicle ownership²⁷, lack of health insurance²⁸, a usual source of care for referrals²⁹, and potential mistrust of the medical system³⁰.

We also found clusters of high and low incidence that were mostly co-located with clusters of high and low disadvantage, respectively. These include areas of concentrated poverty and with a history of extreme racial segregation and discrimination, including West and North Philadelphia, the West Side of Chicago, and The Bronx in NYC.

The consequences of these inequities in infection rates are reflected in preliminary reports of wide inequities in deaths due to COVID-19. Vulnerability to severe disease and death by COVID-19 are related to the presence of previous comorbidities, such as cardiovascular disease, diabetes and hypertension³¹. Since these comorbidities are more prevalent in people of lower socioeconomic status and racial/ethnic minorities^{32,33}, it is expected that, at equal levels of exposure, these groups will suffer more severe consequences from COVID-19. However other factors may also affect the severity of disease and the case-fatality rates including access to and quality of health care and the role of other factors including co-occurring social factors (e.g. stressors) and environmental factors (e.g. air pollution). In fact, a study with 17 million records in the UK has shown that, even after adjusting for a number of comorbidities, racial/ethnic minorities and people living in socioeconomically deprived areas had a higher risk of death after infection³¹.

We also found that inequities in incidence became stronger over time. It is possible that the social patterning of infection has been changing over time as the pandemic progressed,

beginning in wealthier areas (possibly linked to business travel³⁴) and subsequently shifting to more deprived areas. The greater testing in less deprived areas in all three cities early in the epidemic could in part reflect this, given that testing was initially strongly linked to symptoms. However, the strong association of positivity with disadvantage even early in the pandemic suggest that access to testing was, at least initially, lower in more deprived areas. As a result, incidence rates in more deprived neighborhoods early in the pandemic could be underestimated.

An important limitation of our study is the likely underestimation of inequities in incidence due to the lack of systematic widespread testing. In addition, zip codes are very imperfect proxies for neighborhoods. Heterogeneity in the sociodemographic composition within zip codes likely led to underestimation of inequities in our analyses, leading to a lower power to detect associations and spatial clusters. However, zip codes they represent easy-to-collect data in the middle of a public health emergency when more detailed geocoding is less available.

Conclusion

We found wide spatial inequities in COVID-19 outcomes in three large metropolitan areas of the US. Communities of color and of low socioeconomic status have long borne the burden of adverse health linked to discrimination, structural racism, xenophobia and systematic disinvestment leading to a lack of material resources. In health crises, such as this one, all these social problems become magnified, reflecting a longstanding history of racial and economic injustice.

References

1. Dong E, Du H, Gardner L. An interactive web-based dashboard to track COVID-19 in real time. *The Lancet Infectious Diseases* 2020.
2. Burgen S, Jones S. Poor and vulnerable hardest hit by pandemic in Spain. *The Guardian* 2020 April 1st.
3. Corburn J, Vlahov D, Mberu B, et al. Slum health: arresting COVID-19 and improving well-being in urban informal settlements. *Journal of Urban Health* 2020:1-10.
4. Millett GA, Jones AT, Benkeser D, et al. Assessing Differential Impacts of COVID-19 on Black Communities. *Annals of Epidemiology* 2020.
5. THE COLOR OF CORONAVIRUS: COVID-19 DEATHS BY RACE AND ETHNICITY IN THE U.S. 2020. (Accessed May 15th, 2020, at <https://www.apmresearchlab.org/covid/deaths-by-race>.)
6. COVID-19: Data. 2020. (Accessed May 15th, 2020, at <https://www1.nyc.gov/site/doh/covid/covid-19-data.page>.)
7. THE FIRST 100: COVID-19 Took Black Lives First. It Didn't Have To. 2020. (Accessed May 18th, 2020, at <https://features.propublica.org/chicago-first-deaths/covid-coronavirus-took-black-lives-first/>.)
8. Rast J. Milwaukee's Coronavirus Racial Divide: A Report on the Early Stages of COVID-19 Spread in Milwaukee County: University of Wisconsin-Milwaukee; 2020.
9. COVID-19 Statistics. 2020. (Accessed May 15th, 2020, at <http://www.dph.illinois.gov/covid19/covid19-statistics>.)
10. COVID-19 In Illinois - ZIP code lookup. 2020. (Accessed May 19th, 2020, at <https://observablehq.com/@chicagoreporter/covid-19-in-illinois-zip-code-lookup>.)

11. NYC Coronavirus (COVID-19) data. (Accessed April 8th, 2020, at <https://github.com/nychealth/coronavirus-data>.)
12. COVID-19 Testing and data. 2020. (Accessed April 8th, 2020, at <https://www.phila.gov/programs/coronavirus-disease-2019-covid-19/testing-and-data/>.)
13. Randhawa AK, Fisher LH, Greninger AL, et al. Changes in SARS-CoV-2 Positivity Rate in Outpatients in Seattle and Washington State, March 1-April 16, 2020. JAMA : the journal of the American Medical Association 2020.
14. Chen JT, Krieger N. Revealing the unequal burden of COVID-19 by income, race/ethnicity, and household crowding: US county vs. ZIP code analyses 2020.
15. Bailey ZD, Krieger N, Agénor M, Graves J, Linos N, Bassett MT. Structural racism and health inequities in the USA: evidence and interventions. The Lancet 2017;389:1453-63.
16. Rothstein R. The color of law: A forgotten history of how our government segregated America: Liveright Publishing; 2017.
17. Williams DR, Collins C. Racial residential segregation: a fundamental cause of racial disparities in health. Public health reports 2016.
18. Chou R, Dana T, Buckley DI, Selph S, Fu R, Totten AM. Epidemiology of and Risk Factors for Coronavirus Infection in Health Care Workers: A Living Rapid Review. Annals of Internal Medicine 2020.
19. Tomer A, Kane JW. How to protect essential workers during COVID-19. Brookings Report, Brookings Institute March 2020.
20. Valentino-DeVries J, Lu D, Dance GJX. Location Data Says It All: Staying at Home During Coronavirus Is a Luxury. The New York Times 2020 April 3rd.

21. Anderson M. Who relies on public transit in the US. Pew Research Center 2016.
22. Burr JA, Mutchler JE, Gerst K. Patterns of residential crowding among Hispanics in later life: Immigration, assimilation, and housing market factors. *Journals of Gerontology Series B: Psychological Sciences and Social Sciences* 2010;65:772-82.
23. Wahowiak L. Access to water surfaces as human rights issue as poor in Detroit lose services: Sanitation at risk. American Public Health Association; 2014.
24. Cohen S, Tyrrell DAJ, Smith AP. Psychological Stress and Susceptibility to the Common Cold. *New England Journal of Medicine* 1991;325:606-12.
25. Kiecolt-Glaser JK, Glaser R, Gravenstein S, Malarkey WB, Sheridan J. Chronic stress alters the immune response to influenza virus vaccine in older adults. *Proceedings of the National Academy of Sciences of the United States of America* 1996;93:3043-7.
26. Rader B, Astley CM, Sy KTL, et al. Geographic access to United States SARS-CoV-2 testing sites highlights healthcare disparities and may bias transmission estimates. *Journal of Travel Medicine* 2020.
27. Percent of households without a vehicle by race/ethnicity: United States, 2015. 2020. at [https://nationalequityatlas.org/indicators/Car_access.\)](https://nationalequityatlas.org/indicators/Car_access.)
28. Sohn H. Racial and ethnic disparities in health insurance coverage: Dynamics of gaining and losing coverage over the life-course. *Population research and policy review* 2017;36:181-201.
29. Ganguli I, Shi Z, Orav EJ, Rao A, Ray KN, Mehrotra A. Declining use of primary care among commercially insured adults in the United States, 2008–2016. *Annals of Internal Medicine* 2020;172:240-7.

30. King WD. Examining African Americans' mistrust of the health care system: expanding the research question. Commentary on " Race and trust in the health care system". Public Health Reports 2003;118:366.
31. Williamson E, Walker AJ, Bhaskaran KJ, et al. OpenSAFELY: factors associated with COVID-19-related hospital death in the linked electronic health records of 17 million adult NHS patients. medRxiv 2020:2020.05.06.20092999.
32. Kershaw KN, Albrecht SS. Racial/ethnic residential segregation and cardiovascular disease risk. Current cardiovascular risk reports 2015;9:10.
33. Raifman M, Raifman J. Disparities in the population at risk of severe illness from covid-19 by race/ethnicity and income. American journal of preventive medicine 2020.
34. Forster P, Forster L, Renfrew C, Forster M. Phylogenetic network analysis of SARS-CoV-2 genomes. Proceedings of the National Academy of Sciences 2020;117:9241-3.

Acknowledgements

The authors want to acknowledge help by Alyssa Furukawa on data collection, Ran Li on designing some of the visualizations, and Dr. Rene Najera for useful code to calculate spatial clustering.

Funding

UB was supported by the Office of the Director of the National Institutes of Health under award number DP5OD26429. The funding sources had no role in the analysis, writing or decision to submit the manuscript.

Conflicts of interest

The authors declare no conflict of interest.

Table 1: Relative rates of cumulative testing, positivity and incidence associated with zip code socioeconomic characteristics in Chicago, NYC and Philadelphia, using cumulative data up to May 18th, 2020.

City	Variable	Tests per 1,000	Positivity Rate	Incidence
Chicago	Median Household Income	1.02(0.92;1.14)	0.84(0.73;0.95)	0.83(0.71;1.02)
	% Non-Hispanic White	0.96(0.86;1.04)	0.79(0.70;0.91)	0.77(0.66;0.87)
	% College-educated	0.96(0.87;1.07)	0.77(0.68;0.86)	0.75(0.63;0.88)
	% Uninsured	1.00(0.91;1.08)	1.34(1.22;1.44)	1.33(1.15;1.50)
	% Service Worker	0.95(0.86;1.02)	1.25(1.11;1.41)	1.14(0.90;1.34)
	% Overcrowding	1.03(0.96;1.17)	1.22(1.13;1.32)	1.29(1.13;1.45)
	Summary disadvantage Index	0.99(0.91;1.08)	1.33(1.20;1.45)	1.36(1.18;1.57)
New York City	Median Household Income	0.94(0.88;1.00)	0.91(0.88;0.93)	0.85(0.80;0.90)
	% Non-Hispanic White	0.99(0.94;1.03)	0.89(0.87;0.91)	0.89(0.84;0.94)
	% College-educated	0.89(0.84;0.95)	0.85(0.82;0.88)	0.76(0.72;0.80)
	% Uninsured	0.97(0.93;1.02)	1.07(1.05;1.10)	1.04(1.00;1.08)
	% Service Worker	1.01(0.97;1.06)	1.09(1.07;1.12)	1.10(1.04;1.14)
	% Overcrowding	1.03(1.00;1.07)	1.05(1.03;1.07)	1.09(1.04;1.13)
	Summary disadvantage Index	1.02(0.98;1.06)	1.13(1.10;1.16)	1.17(1.11;1.23)
Philadelphia	Median Household Income	1.01(0.95;1.08)	0.89(0.84;0.94)	0.89(0.79;0.99)
	% Non-Hispanic White	0.95(0.87;1.08)	0.84(0.79;0.90)	0.81(0.72;0.90)
	% College-educated	1.03(0.93;1.12)	0.86(0.80;0.93)	0.89(0.78;1.01)
	% Uninsured	0.93(0.86;1.00)	1.11(1.04;1.19)	1.05(0.91;1.18)
	% Service Worker	0.94(0.88;1.00)	1.02(0.95;1.09)	0.97(0.88;1.08)
	% Overcrowding	0.93(0.87;0.99)	1.07(1.01;1.14)	0.99(0.91;1.10)
	Summary disadvantage Index	0.95(0.87;1.03)	1.14(1.07;1.22)	1.10(0.98;1.23)

Footnote: Rate ratios (95% CrI) for each 1 SD increase in the predictor. Median Household Income was log-transformed before standardization.

Figure legends

Figure 1: Scatterplots showing the relation between six socioeconomic variables and three COVID-19 testing outcomes at the zip code level in Chicago, NYC, and Philadelphia. Data is cumulative up to May 18th, 2020.

Footnote: solid lines are linear fits of outcome on predictor, for each city separately

Figure 2: Trends in association between zip-code disadvantage and COVID-19 testing outcomes by city


Footnote: rate ratios (and 95% credible intervals) for each outcome associated with a 1 SD increase in the summary index


Figure 3: Spatial distribution of disadvantage and COVID-19 incidence in Chicago, New York City and Philadelphia

Footnote: testing, positivity rate, and incidence are proportional to circle area


Figure 4: spatial clusters of low and high incidence

Footnote: clusters calculated using the G^* statistic, with a Bonferroni-adjusted threshold of $p < 0.05$







Tests per capita



Chicago
Positivity rate






Cumulative Incidence






medRxiv preprint doi: <https://doi.org/10.1101/2020.05.01.20087833>; this version posted May 27, 2020. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity. It is made available under a [CC-BY 4.0 International license](#).


New York City




Philadelphia




Chicago



New York City



Philadelphia



Cluster ■ High incidence ■ Low incidence