Testing lags and emerging COVID-19 outbreaks in federal penitentiaries: A view from Canada

Alexandra Blair¹, Abtin Parnia¹, Arjumand Siddiqi^{1,2}

- 1. Dalla Lana School of Public Health, University of Toronto, Toronto Canada.
- 2. Gillings School of Global Public Health, University of North Carolina-Chapel Hill, Chapel Hill, USA

ABSTRACT


Objectives: To provide the first known comprehensive analysis of COVID-19 outcomes in a federal penitentiary system. We examined the following COVID-19 outcomes within federal penitentiaries in Canada and contrasted them with estimates for the overall population in the penitentiaries' respective provincial jurisdictions: testing, prevalence, the proportion recovered, and fatality.

Methods: Data for prisons were obtained from the Correctional Service of Canada and, for the general population, from the Esri COVID-19 Canadian Outbreak Tracking Hub. Data were retrieved between March 30 and April 21, 2020, and are accurate to this date. Penitentiary-, province- and sex-specific frequency statistics for each outcome were calculated.

Results: Data on 50 of 51 penitentiaries (98%) were available. Of these, 72% of penitentiaries reported fewer tests per 1000 population than the Canadian general population average (16 tests/1000 population), and 24% of penitentiaries reported zero tests. Penitentiaries with high levels of testing were those that already had elevated COVID-19 prevalence. Five penitentiaries reported an outbreak (at least one case). Hardest hit penitentiaries were those in Quebec, Ontario, and British Columbia, with some prisons reporting COVID-19 prevalence of 30% to 40%. Of these, two were women's prisons. Female prisoners were over-represented among cases (31% of cases overall, despite representing 5% of the total prison population).

Conclusion: Increased sentinel or universal testing may be appropriate given the confined nature of prison populations. This, along with rigorous infection prevention control practices and the potential release of prisoners, will be needed to curb current outbreaks and those likely to come.

GRAPHICAL SUMMARY



- Between 20% and 57% fewer tests per 1000 population have been conducted in federal prisons in Saskatchewan, New Brunswick, Nova Scotia and Alberta than in the general population of those provinces.
- Though Alberta, Manitoba, Saskatchewan, New Brunswick and Nova Scotia are reporting lower counts of COVID-19 cases, these are also the regions reporting the lowest levels of testing.
- Case incidence has been highest in federal prisons in Quebec, Ontario, and British
 Columbia, where a total of five prisons are experiencing outbreaks (1 or more cases).
 These regions are those reporting the highest levels of testing higher than the testing
 levels in the general population.

43

Keywords: Prison; Canada; COVID-19; Health equity

- 46 Number of Figures: 5
- Number of Supplementary data tables: 7

1. BACKGROUND

| 49 | |
|----|---|
| 50 | As we are (re)learning from the evidence of COVID-19 outbreaks in long-term care facilities, |
| 51 | housing large groups of individuals in confined, institutionalized environments is a recipe for |
| 52 | large-scale, human-to-human transmission of infectious disease. Prisons have similar dynamics |
| 53 | of confinement, of residents who are dependent on the institutions' conditions and policies for |
| 54 | basic living and hygiene and, of clinical susceptibility due to underlying health conditions [1-3]. |
| 55 | Epidemiologic findings from past outbreaks of respiratory diseases such as influenza, |
| 56 | adenoviruses, and tuberculosis in high-income countries highlight the higher likelihood of |
| 57 | infection transmission and disease incidence within prisons compared to the general population. |
| 58 | [1-3]. Early reports suggest that several penitentiaries in the United States are experiencing |
| 59 | COVID-19 outbreaks [4], though, as far as we know, no comprehensive analysis of theses |
| 60 | penitentiaries has yet been conducted, and very little is known about COVID-19 in prison |
| 61 | contexts in other high-income countries. In this study, we use Canadian data to provide the first |
| 62 | known examination of COVID-19 statistics for a federal penitentiary system. In doing so, we |
| 63 | also bring front-and-center a major health equity issue, because Indigenous and racialized |
| 64 | communities are over-represented within the Canadian prison system [5]. |
| 65 | |
| 66 | In the context of the COVID-19 pandemic, there are specific factors that place prisoner |
| 67 | populations at particularly high risk of COVID-19 infection, and of a higher risk for severe |
| 68 | (compared to mild) COVID-19 outcomes. These include housing multiple prisoners to one cell |
| 69 | ("double-bunking")[6], the ageing population within prisons—particularly in federal prisons[7], |
| 70 | a high prevalence of chronic disease comorbidities and immuno-compromised health status |
| 71 | associated with substance use and blood-borne infections[8], and the regular entrance of |
| 72 | custodial and health care staff from communities with likely community-based transmission. |
| 73 | |
| 74 | At the time of writing (late April 2020), Canada has been documenting the COVID-19 pandemic |
| 75 | on its soil for almost two months. In late March 2020, the Correctional Service of Canada (CSC) |
| 76 | Canada's federal body responsible for incarcerated individuals serving sentences of at least two |
| 77 | years, had indicated that they had taken steps to prevent and were prepared to respond to |
| 78 | COVID-19 outbreaks inside its federal penitentiaries [9]. With this study, we aim to offer a |

baseline situational assessment of COVID-19 testing and burden in Canadian federal penitentiaries, from which future analyses can be based. This study aims to summarize and compare the prevalence of testing, test positivity, COVID-19 prevalence, case fatality, and the proportion of cases recovered within federal penitentiaries in Canada, by province and for Canada overall, and to contrast these with rates in penitentiaries' respective provincial jurisdictions.

2. METHOD

Data and study population

In an unprecedented act of transparency, CSC made COVID-19 testing and case numbers publicly available for each federal penitentiary on their website. Data up until April 21, 2020, were used [10]. Equivalent values for the Canadian population were extracted from the COVID-19 Canada Outbreak Tracker Resource Hub [11]. For each indicator, described below, we calculated prison-, sex-, and province-specific estimates. Prison population denominators were approximated by the maximum prison capacity for each penitentiary [12]. For the general population, denominators (i.e., total general population) were obtained from Statistics Canada population estimates for the first quarter of 2020 [13].

Additionally, to provide a timeline for the number of tests and cases in prisons with possible outbreaks (5 or more cases at the time of analysis), we used the Wayback Machine (archive.org) to obtain all previous copies of CSC's COVID-19 reporting webpage [10]. The earliest version of the CSC webpage was created on April 1, 2020, which provided cumulative data up until March 30, 2020. There were eight additional updates, which reported on data from April 7, 8,10, 12, 13, 15, 17, and 18, 2020. Cumulative data at each of these dates were merged with the data from April 21, 2020, to create a time-series ranging from March 30 to April 21, 2020. For calendar dates at which no CSC updates were made available, we filled in the information with the data from the previous observation.

| 109 | Given that several federal penitentiaries are in fact multiplex sites—either with distinct buildings |
|---|--|
| 110 | or units operating under different security levels (e.g. minimum, medium or maximum) or |
| 111 | offering distinct services (e.g. treatment facilities)—and that sub-population capacities were not |
| 112 | always available for each separate unit, we chose to group several multi-complex facilities in our |
| 113 | analyses. This was the case of Quebec's Federal Training Center Multi-Level Unit and Minimum |
| 114 | security facility; British Columbia's Pacific Institution, Pacific Institution Regional Treatment |
| 115 | Center and Pacific Institution Reception Center; Ontario's Millhaven Institution, Millhaven |
| 116 | Regional Hospital and Millhaven Regional Treatment Center, as well as Collins Bay Minimum |
| 117 | Institution and Collins Bay Regional Treatment Center, and Joyceville Institution and Joyceville |
| 118 | Minimum Institution. With these groupings, we recorded 51 facilities. We were able to analyze |
| 119 | data from 50 of these facilities (98%) (all data in Supplementary File 1). |
| 120 | |
| 121 | Measures |
| 122 | |
| 123 | Total tests and cases |
| 124 | The total number of tests performed was obtained directly from the CSC and the Canada |
| 125 | Outbreak Tracker Resource Hub websites. CSC reports on the number of "positive tests". We |
| 126 | considered all "positive tests" as confirmed cases. |
| | |
| 127 | |
| 127 128 | Tests per 1000 population |
| | Tests per 1000 population Tests per 1000 population were estimated by dividing the total number of tests performed by the |
| 128 | |
| 128 129 | Tests per 1000 population were estimated by dividing the total number of tests performed by the |
| 128 129 130 | Tests per 1000 population were estimated by dividing the total number of tests performed by the total population in each facility, in the prisoner population of each province, and the general |
| 128 129 130 131 | Tests per 1000 population were estimated by dividing the total number of tests performed by the total population in each facility, in the prisoner population of each province, and the general |
| 128 129 130 131 132 | Tests per 1000 population were estimated by dividing the total number of tests performed by the total population in each facility, in the prisoner population of each province, and the general population of each province, respectively, and multiplying the fraction by 1000. |
| 128 129 130 131 132 133 | Tests per 1000 population were estimated by dividing the total number of tests performed by the total population in each facility, in the prisoner population of each province, and the general population of each province, respectively, and multiplying the fraction by 1000. *Test-positive rate* |
| 128 129 130 131 132 133 134 | Tests per 1000 population were estimated by dividing the total number of tests performed by the total population in each facility, in the prisoner population of each province, and the general population of each province, respectively, and multiplying the fraction by 1000. *Test-positive rate* We estimated the proportion of tests that were positive by dividing the total number of positive |

Tested COVID-19 prevalence

We estimated the prevalence of COVID-19 as identified through tests in the penitentiary and provincial populations by dividing the total number of positive tests (confirmed cases) by the population of each facility, the total prison population by province, and the total provincial population, respectively. This indicator can also be considered the cumulative incidence proportion.

Proportion recovered

To offer a crude estimate of the evolution of the epidemics inside and outside of prisons, we estimated the proportion of cases that had recovered. This was done by dividing the number of cases who had recovered by the total number of positive tests (total confirmed cases). Operational definitions of recovered cases vary across jurisdictions in Canada, and CSC does not define their measure in their reporting. Recovered cases are those who either received confirmatory negative test results or for whom 10 to 14 days have elapsed since the start of their symptoms and they are symptom-free for at least 2 to 3 days [14, 15]. Therefore, we interpret recovered case numbers with caution.

Population categories in prisons with outbreaks: Susceptible, Infected, Recovered and Died An outbreak of COVID-19 within confined populations, such as long-term care homes, can be declared after a single resident or staff member is confirmed positive [16]. Thus, we consider prisons with one or more reported cases of COVID-19 among its prisoner population as those experiencing outbreaks. To describe the evolution of testing and incidence in prisons with outbreaks across our study period, we classified each prison's population into four categories, for each calendar day of the study period. We estimated the number of prisoners who were "susceptible" to infection by subtracting the total number of confirmed active, recovered, and deceased cases from the maximum population capacity of each prison. Prisoners considered "infected" were those with positive tests who had yet to recover or die. Prisoners declared to be "recovered" were assumed to no longer present active infections or be susceptible to infection. Lastly, the total number of prisoners who had "died" were those whose death was declared to

3. RESULTS

result from COVID-19 complications.

170 171 3.1 Testing inside versus outside federal prisons 172 173 An analysis of testing within all federal penitentiaries illustrates both the heterogeneity of 174 COVID-19 testing within prisons in Canada and a low if not a complete absence of testing in 175 many facilities (Figure 1). Twelve of the 50 facilities studied—nearly one in five (24%)—had 176 recorded a complete absence of testing (indicated by hollow bars in Figure 1). Approximately 177 72% of all facilities (36/50) recorded fewer tests than the Canadian general population average of 178 16 tests per 1000 population. Facilities with higher levels of testing per 1000 population, tended 179 to be those that had already reported a higher COVID-19 prevalence (Figure 1). 180 181 On average across provinces, New Brunswick, Nova Scotia, Alberta, and Saskatchewan have 182 been testing less inside prisons than in the general population (Figure 2). The opposite is true for 183 Quebec, Ontario, and British Columbia, which were likely responsible for driving the average of 184 34 tests per 1000 population observed for the prison population, overall. 185 186 3.2 Prevalence of COVID-19 inside versus outside federal prisons 187 188 The high positive test rates and estimated prevalence of COVID-19 in prisons in Quebec and 189 British Columbia point to emerging outbreaks in these provinces (Figure 2). Prevalence was 10 190 times higher in Quebec, 6 times higher in British Columbia, and 2 times higher in Ontario within 191 federal prisons compared to the general population of each province (Figure 2). 192 193 Many of the hardest-hit penitentiaries are near major city centers. In Quebec, this was the 194 Federal Training Center complex in Laval, just north of Montreal (51 cases, 7% COVID-19 195 prevalence) and, Joliette Institution for Women in Joliette, a 1-hour drive away from Montreal 196 (51 cases, 39% prevalence). However, Port Cartier Institution in Côte Nord (a relatively remote 197 region of the province) also reported 15 cases (Figure 1). In British Columbia, most affected was 198 the Mission Medium Security Institution, a 1-hour drive away from Vancouver (64 cases, 30% 199 prevalence) (Figure 1). Ontario's Grand Valley Institution for Women, located in Kitchener (1-200 hour drive away from Toronto) also experienced a smaller outbreak of 8 cases (6% prevalence).




Figure 1: Test-positive proportion, case prevalence, and testing per 1000 population for all federal penitentiaries.




Figure 2: Testing per 1000 population, test-positive rate, and prevalence of positive tests, for federal prison and general populations, by province and in Canada overall

In the five institutions with one or more cases, we found that testing was largely reactive, with the onset of concentrated testing efforts occurring after COVID-19 outbreaks had already established. An exception to this observation was Quebec's Federal Training Center, which was the only facility recording negative tests before the observations of positive tests (Figure 3). Further, though institutions all recorded a gradual increase in the number of tests performed over time, the high proportion of positive tests throughout the study period suggests that these prisons likely did not implement more intensive, wide-spread testing throughout the prison populations once a case was observed. An exception to this, however, was Ontario's Grand Valley Women's Institution, which recorded a significant jump in testing once the first cases were confirmed, with a majority of tests returning negative since April 7, 2020.

3.3 Proportion of cases recovered and fatality inside versus outside prisons

The proportion of cases that had recovered inside federal penitentiaries ranged from 0% to 33% (Figure 1). The Susceptible, Infected, and Recovered plots (Figure 4) illustrate what proportion of prisoners each prison with one or more cases have become infected and have recovered or died throughout the study period. Of the five prisons with outbreaks, Quebec's Joliette Women's Institution and British Columbia's Mission Medium Institution have the seen largest proportion of their population become infected and subsequently recover. Smaller proportions of cases have recovered in Quebec's Federal Training Centre and Port-Cartier and Ontario's Grand Valley Women's Institution, and plot results suggest that a large proportion of prisoners remain susceptible in these institutions (Figure 4).




Figure 3: Testing patterns and outcomes between March 30 and April 21, 2020, in five penitentiaries with one or more recorded COVID-19 cases. * Indicated date at which data was reported by Correctional Services Canada.




Figure 4: Number of susceptible, infected, recovered cases and deaths between March 30 and April 21, 2020, in penitentiaries with one or more recorded COVID-19 cases. * Indicates date at which data was reported by CSC.

In contrast to cases in the general population, a smaller proportion of cases in federal penitentiaries in Quebec, Ontario, and British Columbia had recovered (Figure 2). In both Ontario and British Columbia, the proportion of individuals who had recovered within prisons was half that of the proportion of cases recovered in the community. With the available evidence, it is difficult to know precisely what may be driving these differences in percent recovered inside versus outside of prisons. One explanation is that outbreaks inside prisons may have started more recently than those within the general public. Another possibility is that health and sanitary conditions within prisons may lead to lags in recovery. Further, this cross-sectional summary of differences must be interpreted with caution, since the proportion of cases that are recovered will vary according to the number of new incident cases observed in both settings.

At the time of data collection, one of the 189 laboratory-confirmed cases across all federal penitentiaries had died (0.5%), a 67% higher proportion than that observed in the general population, (0.3%, 1,966 of 614,340 cases). Case fatality estimates should be interpreted with caution, however, given the likely underestimation of the true number of cases (denominator value) in both settings.

3.4 Testing, prevalence and proportion recovered by sex

Among the 50 facilities included in these analyses, six are penitentiaries for women (Nova Institution for Women in Nova Scotia, Joliette Institution in Quebec, Grand Valley Institution for Women in Ontario, Edmonton Institution for Women in Alberta, Fraser Valley Institution in British Columbia, and Okimaw Ohci Healing Lodge in Saskatchewan). Together they hold up to 785 women (5% of the total capacity of 16,401 federal prisoners in Canada).




Figure 5: Distribution of women among federal prisons and prison COVID-19 cases (Canada, April 21, 2020).

At the time of data collection, there were 59 cases of COVID-19 in women's penitentiaries. These represented 31% of all cases in federal penitentiaries (N=189) (Figure 5), suggesting that women, and women's penitentiaries, are over-represented among COVID-19 cases inside federal prisons. COVID-19 prevalence was 8 times higher among women's prisons (8% prevalence) than prisons for men (1% prevalence) and 80 times higher than in the general Canadian population (0.1%).

4. DISCUSSION

Key findings

The data suggest that at least five federal penitentiaries, one in British Columbia, three in Quebec and one in Ontario, are experiencing COVID-19 outbreaks. Each of these penitentiaries is reporting high levels of testing, compared to provincial and national rates, and this testing appears to be a reaction to the emergence of outbreaks that have already begun, rather than a proactive prevention effort. Though at least thirteen (26%) of facilities appear to be experiencing very low if any COVID-19 cases, these are also facilities that have reported low levels of testing per 1000 population, so it is unclear whether the low case counts reflect true case prevalence. We observed lower levels of testing than the Canadian general population average in 26 penitentiaries (72%) and a total absence of testing in 12 penitentiaries (24%). Our study also suggests a smaller proportion of cases inside prisons have recovered compared to cases in the general population. This may indicate that the spread of COVID-19 is lagging in penitentiaries, and the worst is yet to come. It may also indicate that prisoners face suboptimal conditions for recovery. We also found that at least three of the current penitentiary outbreaks are in proximity to city centers (e.g. Montreal and Vancouver), and higher COVID-19 prevalence has been observed in women's (8% prevalence) compared to men's prisons (1% prevalence), overall.

Strengths and limitations

The primary strength of this study is its use of the best available data on COVID-19 testing and case incidence in Canadian Federal penitentiaries and the general Canadian population to give an

early situational report on testing and infection-related outcomes within these facilities. This study is a baseline assessment from which future analyses using updated data can be based. A limitation of this study is our use of the maximum potential capacity of each penitentiary as the population denominator for rate calculations, rather than the 'true' population at the time of data collection. We expect that these denominators are likely too large, given that prisons may not be at their full capacity and pressures to release non-violent offenders to reduce the number of susceptible individuals within prisons [17]. This may have led to the under-estimation of testing and infection prevalence estimates. Further, this study was not able to also count the number of tests and cases identified within custodial and health staff within these same penitentiaries, which is an important contributing factor to disease transmission. While other deaths in federal penitentiaries were recorded by CSC during the study period (e.g. [18]), only one case was reported as COVID-19 related. It is unclear whether other deaths occurring within CSC facilities have or will be tested for COVID-19 post-mortem. Lastly, these findings may not be generalizable to provincial, remand, or immigration detention facilities, which may see far more movement in and out given the shorter duration of sentences.

Implications

These early findings have several implications for public health and emergency outbreak response inside federal penitentiaries. The observation that at least five Canadian penitentiaries are experiencing outbreaks highlights the importance that all penitentiary populations and custodial and health care staff uphold rigorous prevention control practices (and be given the resources to do so), including regular hand hygiene and cleaning practices, the use of personal protection equipment, and physical distancing measures [19]. Scholars and legal experts have also emphasized the need to consider the release of prisoners to reduce the proportion of susceptible individuals within prisons [20].

Our findings of sparse testing across several penitentiaries in Canada suggests that a more proactive testing approach may be needed to help curb potential future outbreaks. This is especially the case given studies suggesting that up to 60% of COVID-19 cases may be asymptomatic [21-23]. Testing based on symptom-presence alone may thus need to be

325 reconsidered within the confined spaces of prisons, as many cases may go undetected as long as 326 testing is not expanded. 327 328 Relevant testing strategies could include universal testing or, active, sentinel-based testing. 329 While universal testing [19] [24] is time and resource-intensive but is much more feasible within 330 small confined populations than in the general public. In the context of COVID-19, this strategy 331 has been used navy ships [21] and homeless shelters [25, 26] to identify all cases and adapt 332 outbreak control interventions accordingly. In Canadian penitentiaries, universal testing may be 333 most pertinent in prisons that have already reported confirmed cases. At the time of writing, 334 British Columbia's Mission Institution penitentiary — where one of the five outbreaks is 335 unfolding — had reported planning universal testing of all inmates and staff [27]. However, none 336 of the other institutions had indicated plans to use such an approach. 337 338 As an alternative to universal testing, active sentinel surveillance involves identifying a subset of 339 prisons in which regular testing for COVID-19 among prisoners and staff, regardless of symptom 340 presence, could be conducted. Test results would be used for monitoring in the months to come 341 until a vaccine or effective treatment is identified. This approach may be most relevant in 342 provinces with low numbers of new daily COVID-19 cases (e.g. currently New Brunswick, Nova 343 Scotia, Saskatchewan and Manitoba currently) or in later in the pandemic, when community 344 transmission subsides and testing all prisoners may be too costly and inefficient. 345 These types of proactive surveillance activities may be particularly important in the context of 346 347 alternative options to reduce the spread of COVID-19, such as long-term cell-based confinement 348 for the entire prison population. Confinement has been associated with severe mental health risk 349 [28]—particularly for Indigenous and racialized populations [6]—and has come under 350 considerable scrutiny in Canada in recent years [29]. 351 352 Lastly, studies on the effectiveness of universal and sentinel-based testing strategies are 353 warranted, as are mathematical models that explore the epidemiologic consequence of 354 interventions (or lack thereof) within federal prisons on COVID-19 transmission, morbidity and 355 death, as has been done in other country settings [30].

| 356 | |
|-----------------------------------|--|
| 357 | 5. CONCLUSION |
| 358 | |
| 359 | Gaps in COVID-19 testing in several jurisdictions and recorded outbreaks in several |
| 360 | penitentiaries in Canada suggest that federal prisons represent an emerging battleground agains |
| 361 | the COVID-19 pandemic. With an ageing population that is particularly at high risk due to the |
| 362 | prevalence of underlying chronic and infectious health conditions, prisoners represent a |
| 363 | potentially neglected segment of society that is at high risk of bearing a high burden of COVID |
| | |
| 364 | 19 related morbidity and mortality. As indigenous and racialized communities are over- |
| 365 | represented within the Canadian prison system, COVID-19 prevention inside penitentiaries |
| 366 | represents a pressing public health and health equity issue. |
| 367 | |
| 368 | |
| 369 | AUTHOR CONTRIBUTIONS |
| 370 | |
| 371 | AB and AP conceptualized the study, with input from AS. AB and AP analyzed the data. AB |
| 372373 | and AP drafted the manuscript. AS responded to drafts of the manuscript. |
| 374 | FUNDING |
| 375 | |
| 376 | AB receives postdoctoral funding from the Fonds de Recherche du Québec-Santé. AS is |
| 377 | supported by the Canada Research Chair in Population Health Equity. |
| 378 | |
| 379 | CONFLICT OF INTEREST |
| 380 | N |
| 381 382 | None |
| 383 | ETHICS COMMITTEE APPROVAL |
| 384 | |
| 385 | None required |
| 386 | |
| 387 | |
| 388 389 | REFERENCES |
| 390 | 1. Young, L.C., et al., Summer outbreak of respiratory disease in an Australian prison due |
| 391 392 | to an influenza A/Fujian/411/2002(H3N2)-like virus. Epidemiol Infect, 2005. 133 (1): p. 107-12. |

- Parcell, B.J., et al., *Prison and community outbreak of severe respiratory infection due to adenovirus type 14p1 in Tayside, UK.* J Public Health (Oxf), 2015. **37**(1): p. 64-9.
- 395 3. Dolan, K., et al., *Global burden of HIV, viral hepatitis, and tuberculosis in prisoners and detainees.* The Lancet, 2016. **388**(10049): p. 1089-1102.
- Williams, T., B. Weiser, and W.K. Rashbaum, 'Jails Are Petri Dishes': Inmates Freed as the Virus Spreads Behind Bars, in The New York Times. 2020: New York, USA.
- 399 5. Owusu-Bempah, A. and S. Wortley, *Race, crime, and criminal justice in Canada*, in *The Oxford handbook of ethnicity, crime, immigration*. 2014, Oxford UP: Oxford. p. 321-359.
- Zinger, I., Annual Report of the Office of the Correctional Investigator 2014-2015. 2015,
 Office of the Correctional Investigator: Ottawa, Canada.
- Zinger, I. and M. Landry, *Aging and dying in prison: an investigation into the* experiences of older individuals in federal custody. 2019, Office of the Correctional Investigator: Ottawa, Canada.
- 406 8. Zinger, I., *Annual Report of the Office of the Correctional Investigator 2018-2019*. 2018, 407 Office of the Correctional Investigator: Ottawa, Canada.
- 408 9. Davis, D., Correctional Service of Canada takes steps to reduce risk of coronavirus entering prisons, in Global News. 2020.
- 410 10. CSC. Correctional Service of Canada: Inmate COVID-19 testing in federal correctional
 411 institutions, April 21, 2020. 2020 [cited 2020 April 21, 2020]; Available from:
 412 https://www.csc-scc.gc.ca/001/006/001006-1014-en.shtml.
- 413 11. EsriCanada. *COVID-19 Canadian Outbreak Tracker*. 2020 [cited 2020 April 22]; 414 Available from: https://resources-covid19canada.hub.arcgis.com/app/eb0ec6ffdb654e71ab3c758726c55b68.
- 416 12. CSC. Facilities and Security: Institutional profiles. 2013 [cited 2020 April 21]; 417 Available from: https://www.csc-scc.gc.ca/institutions/index-eng.shtml.
- 418 13. StatisticsCanada. *Population estimates, quarterly: Table: 17-10-0009-01 (formerly*419 *CANSIM 051-0005)*. 2020 [cited 2020 April 21]; Available from:
 420 https://www150.statcan.gc.ca/t1/tbl1/en/tv.action?pid=1710000901.
- 421 14. HealthLinkBC. *Coronavirus disease (COVID-19)*. 2020 [cited 2020 April 24]; Available from: https://www.healthlinkbc.ca/health-feature/coronavirus-disease-covid-19.
- INSPQ, COVID-19: Recommandations pour la levée des mesures d'isolement des
 travailleurs de la santé. 2020, Institut National de Santé Publique du Québec, Comité sur les infections nosocomiales du Québec: Quebec, Canada.
- 426 16. PHAC. Infection Prevention and Control for COVID-19: Interim Guidance for Long
 427 Term Care Homes. 2020 [cited 2020 April 29]; Available from:
- https://www.canada.ca/en/public-health/services/diseases/2019-novel-coronavirusinfection/prevent-control-covid-19-long-term-care-homes.html.
- 430 17. Harris, K., *Bill Blair asks prison, parole heads to consider releasing some inmates to stop spread of COVID-19*, in *CBC News*. 2020: Ottawa.
- 432 18. CSC. Death of an inmate from Dorchester Penitentiary. 2020 [cited 2020 April 29]; 433 Available from: https://www.canada.ca/en/correctional-service/news/2020/04/death-of-
- 434 <u>an-inmate-from-dorchester-penitentiary.html.</u>
- World Health, O., *Preparedness, prevention and control of COVID-19 in prisons and other places of detention (2020), Interim guidance 15 March 2020.* 2020.
- 437 20. Kinner, S.A., et al., *Prisons and custodial settings are part of a comprehensive response to COVID-19*. Lancet Public Health, 2020. **5**(4): p. e188-e189.

- 439 21. Stewart, P. and I. Ali, *Coronavirus clue? Most cases aboard U.S. aircraft carrier are symptom-free*, in *Reuters*. 2020: Washington, USA.
- 441 22. Nishiura, H., et al., *Estimation of the asymptomatic ratio of novel coronavirus infections* (*COVID-19*). medRxiv, 2020: p. 2020.02.03.20020248.
- 443 23. Kimball, A., et al., Asymptomatic and Presymptomatic SARS-CoV-2 Infections in
 444 Residents of a Long-Term Care Skilled Nursing Facility King County, Washington,
 445 March 2020. MMWR Morb Mortal Wkly Rep, 2020. 69(13): p. 377-381.
- ECDC, Strategies for the surveillance of COVID-19, in Technical Report: 9 April 2020.
 ECDC, Strategies for the surveillance of COVID-19, in Technical Report: 9 April 2020.
 European Centre for Disease Prevention and Control.
- Mosites, E., Assessment of SARS-CoV-2 Infection Prevalence in Homeless Shelters—
 Four US Cities, March 27–April 15, 2020. MMWR. Morbidity and Mortality Weekly
 Report, 2020. 69.
- 451 26. Karedes, D., *CDC reviewing 'stunning' universal testing results from Boston homeless shelter*, in *Boston 25 News*. 2020, Boston 25 News: Boston, USA.
- 453 27. Lovgreen, T., *All inmates and staff at Mission Institution being tested for COVID-19*, in *CBC News*. 2020: British Columbia, Canada.
- 455 28. Arrigo, B.A. and J.L. Bullock, *The Psychological Effects of Solitary Confinement on Prisoners in Supermax Units: Reviewing What We Know and Recommending What Should Change*. International Journal of Offender Therapy and Comparative Criminology, 2007. **52**(6): p. 622-640.
- Senate, Proceedings of the Standing Senate Committee on Social Affairs, Science and
 Technology Issue No. 60 Evidence May 15, 2019, in Issue No. 60 Evidence May 15,
 2019. 2019, Canadian Senate: Ottawa, Canada.
- 462 30. Lofgren, E., et al., The Epidemiological Implications of Incarceration Dynamics in Jails
 463 for Community, Corrections Officer, and Incarcerated Population Risks from COVID-19.
 464 medRxiv, 2020: p. 2020.04.08.20058842.
 465

7. SUPPLEMENTARY FILE

Federal Prison Population Data – as of April 21, 2020

| Province | Federal penitentiary | Population | Cases | Total tests | Deaths | Recovered |
|----------|--|------------|-------|-------------|--------|-----------|
| NB | Atlantic Institution | 331 | 0 | 1 | 0 | 0 |
| | Dorchester Medium Institution | 397 | 0 | 4 | 0 | 0 |
| | Dorchester Minimum Institution | 302 | 0 | 1 | 0 | 0 |
| NS | Nova Institution for Women | 99 | 0 | 3 | 0 | 0 |
| | Springhill Institution | 636 | 0 | 4 | 0 | 0 |
| QC | Federal Training Centre Multi-Level | 764 | 51 | 76 | 0 | 0 |
| | Unit-Minimum | | | | | |
| | Regional Reception Centre | 599 | 0 | 5 | 0 | 0 |
| | Archambault Institution Medium | 284 | 0 | 1 | 0 | 0 |
| | Archambault Institution Minimum | 215 | 0 | 1 | 0 | 0 |
| | Cowansville Institution | 599 | 0 | 1 | 0 | 0 |
| | Donnacona Institution | 451 | 0 | 5 | 0 | 0 |
| | Drummond Institution | 369 | 0 | 4 | 0 | 0 |
| | Joliette Institution | 132 | 51 | 76 | 0 | 16 |
| | La Macaza Institution | 240 | 0 | 1 | 0 | 0 |
| | Port-Cartier Institution | 237 | 15 | 21 | 0 | 3 |
| ONT | Bath Institution & Regional Treatment | 516 | 0 | 11 | 0 | 0 |
| | Center Beaver Creek Medium Institution | 516 | 0 | 2 | 0 | 0 |
| | Beaver Creek Minimum Institution | 201 | 0 | 1 | 0 | 0 |
| | Collins Bay Maximum Institution | 96 | 0 | 0 | 0 | 0 |
| | | 482 | 0 | | 0 | - |
| | Collins Bay Medium Institution | 182 | 0 | 0 | 0 | 0 |
| | Collins Bay Minimum Inst. & Regional Treatment Ctr | 182 | 0 | 1 | 0 | U |
| | Grand Valley Institution for Women | 215 | 8 | 108 | 0 | 2 |
| | Joyceville Institution & Minimum | 752 | 0 | 6 | 0 | 0 |
| | Institution | | | | | |
| | Millhaven Institution-Regional Hospital- | 496 | 0 | 5 | 0 | 0 |
| | Treatment Ctr | | | | | |
| | Warkworth Institution | 537 | 0 | 26 | 0 | 0 |
| AB | Bowden Institution | 600 | 0 | 17 | 0 | 0 |
| | Drumheller Institution | 704 | 0 | 2 | 0 | 0 |
| | Edmonton Institution | 324 | 0 | 5 | 0 | 0 |
| | Edmonton Institution for Women | 167 | 0 | 0 | 0 | 0 |
| | Grande Cache Institution | 243 | 0 | 2 | 0 | 0 |
| | Grierson Centre | 30 | 0 | 0 | 0 | 0 |
| | Pê Sâkâstêw Centre | 60 | 0 | 1 | 0 | 0 |
| SASK | Okimaw Ohci Healing Lodge | 60 | 0 | 2 | 0 | 0 |
| | Regional Psychiatric Centre | 204 | 0 | 11 | 0 | 0 |
| | Saskatchewan Maximum Penitentiary | 176 | 0 | 0 | 0 | 0 |
| | Saskatchewan Medium Penitentiary | 457 | 0 | 7 | 0 | 0 |
| | Saskatchewan Minimum Penitentiary | 176 | 0 | 0 | 0 | 0 |
| | Willow Cree Healing Lodge | 80 | 0 | 0 | 0 | 0 |
| MAN | Stony Mountain Maximum Institution | 96 | 0 | 0 | 0 | 0 |
| | Stony Mountain Medium Institution | 484 | 0 | 12 | 0 | 0 |
| | Stony Mountain Minimum Institution | 217 | 0 | 0 | 0 | 0 |
| BC | Fraser Valley Institution | 112 | 0 | 3 | 0 | 0 |

| | Kent Institution | 378 | 0 | 0 | 0 | 0 | | | | |
|------------|---|-------|-----|-----|---|----|--|--|--|--|
| | Kwìkwèxwelhp Healing Lodge | 50 | 0 | 0 | 0 | 0 | | | | |
| | Matsqui Institution | 446 | 0 | 0 | 0 | 0 | | | | |
| | Mission Medium Institution | 216 | 64 | 105 | 1 | 21 | | | | |
| | Mission Minimum Institution | 324 | 0 | 2 | 0 | 0 | | | | |
| | Mountain Institution | 440 | 0 | 4 | 0 | 0 | | | | |
| | Pacific Institution-Regional Treatment- | 509 | 0 | 21 | 0 | 0 | | | | |
| | Reception Ctr | | | | | | | | | |
| | William Head Institution | 200 | 0 | 3 | 0 | 0 | | | | |
| Canada | Overall | 16401 | 189 | 564 | 1 | 42 | | | | |
| Accessed A | Accessed April 21, 2020: https://www.csc-scc.gc.ca/001/006/001006-1014-en.shtml | | | | | | | | | |

General Population Data – as of April 21, 2020

| Province | Population | Positive tests | Total tests | Deaths | Recovered | | | | | | |
|------------------|--|----------------|-------------|--------|-----------|--|--|--|--|--|--|
| New Brunswick | 779,993 | 118 | 11281 | 0 | 102 | | | | | | |
| Nova Scotia | 977,457 | 772 | 22927 | 12 | 330 | | | | | | |
| Quebec | 8,537,674 | 20965 | 174490 | 1134 | 4291 | | | | | | |
| Ontario | 14,711,827 | 12245 | 174170 | 659 | 6221 | | | | | | |
| Alberta | 4,413,146 | 3095 | 109015 | 61 | 1273 | | | | | | |
| Saskatchewan | 1,181,666 | 320 | 24811 | 4 | 252 | | | | | | |
| Manitoba | 1,377,517 | 255 | 20319 | 6 | 150 | | | | | | |
| British Columbia | 5,110,917 | 1724 | 65545 | 87 | 1041 | | | | | | |
| Canada (Overall) | 37,894,799 | 39805 | 614340 | 1966 | 13908 | | | | | | |
| | Accessed April 21, 2020: http://ppenhub-esrica-apps.opendata.arcgis.com/datasets/covid-19-canada-1; Population data, | | | | | | | | | | |

Data on facilities with one or more cases as of April 21, 2020

Data were available for March 30 and April 7-8, 10, 12, 13, 15, 17-18, 21 of 2020.

Port Cartier Institution, Quebec (Maximum Population Estimate: 237 prisoners)

| Date | Cumulative tes | st totals (N) | Population categories | | | | | | |
|--------|----------------|---------------|-----------------------|--------------|-------|-------------|----------------|-----------|--------|
| | Positive | | | | | | Infected | | |
| (2020) | (Cases) | Negative | Pending | Inconclusive | Total | Susceptible | (Active Cases) | Recovered | Deaths |
| 30-03 | 2 | 0 | 3 | 0 | 5 | 235 | 2 | 0 | 0 |
| 31-03 | 2 | 0 | 3 | 0 | 5 | 235 | 2 | 0 | 0 |
| 01-04 | 2 | 0 | 3 | 0 | 5 | 235 | 2 | 0 | 0 |
| 02-04 | 2 | 0 | 3 | 0 | 5 | 235 | 2 | 0 | 0 |
| 03-04 | 2 | 0 | 3 | 0 | 5 | 235 | 2 | 0 | 0 |
| 04-04 | 2 | 0 | 3 | 0 | 5 | 235 | 2 | 0 | 0 |
| 05-04 | 2 | 0 | 3 | 0 | 5 | 235 | 2 | 0 | 0 |
| 06-04 | 2 | 0 | 3 | 0 | 5 | 235 | 2 | 0 | 0 |
| 07-04 | 7 | 3 | 4 | 0 | 14 | 230 | 7 | 0 | 0 |
| 08-04 | 9 | 4 | 1 | 0 | 14 | 228 | 9 | 0 | 0 |
| 09-04 | 9 | 4 | 1 | 0 | 14 | 228 | 9 | 0 | 0 |
| 10-04 | 10 | 4 | 0 | 0 | 14 | 227 | 10 | 0 | 0 |
| 11-04 | 10 | 4 | 0 | 0 | 14 | 227 | 10 | 0 | 0 |
| 12-04 | 10 | 4 | 1 | 0 | 15 | 227 | 10 | 0 | 0 |
| 13-04 | 10 | 4 | 1 | 0 | 15 | 227 | 10 | 0 | 0 |
| 14-04 | 10 | 4 | 1 | 0 | 15 | 227 | 10 | 0 | 0 |
| 15-04 | 10 | 5 | 4 | 0 | 19 | 227 | 10 | 0 | 0 |
| 16-04 | 10 | 5 | 4 | 0 | 19 | 227 | 10 | 0 | 0 |
| 17-04 | 14 | 5 | 0 | 0 | 19 | 223 | 12 | 2 | 0 |
| 18-04 | 14 | 5 | 0 | 0 | 19 | 223 | 12 | 2 | 0 |
| 19-04 | 14 | 5 | 0 | 0 | 19 | 223 | 12 | 2 | 0 |
| 20-04 | 14 | 5 | 0 | 0 | 19 | 223 | 12 | 2 | 0 |
| 21-04 | 15 | 5 | 1 | 0 | 21 | 222 | 12 | 3 | 0 |

Federal Training Center, Quebec (Maximum Population Estimate: 764)

| Date | Cumulative tes | t totals (N) | | | | Population categories | | | | |
|--------|----------------|--------------|---------|--------------|-------|-----------------------|-------------------------|-----------|--------|--|
| | Positive | | | | Total | | | | | |
| (2020) | (Cases) | Negative | Pending | Inconclusive | tests | Susceptible | Infected (Active Cases) | Recovered | Deaths | |
| 30-03 | 0 | 1 | 0 | 0 | 1 | 764 | 0 | 0 | 0 | |
| 31-03 | 0 | 1 | 0 | 0 | 1 | 764 | 0 | 0 | 0 | |
| 01-04 | 0 | 1 | 0 | 0 | 1 | 764 | 0 | 0 | 0 | |
| 02-04 | 0 | 1 | 0 | 0 | 1 | 764 | 0 | 0 | 0 | |
| 03-04 | 0 | 1 | 0 | 0 | 1 | 764 | 0 | 0 | 0 | |
| 04-04 | 0 | 1 | 0 | 0 | 1 | 764 | 0 | 0 | 0 | |
| 05-04 | 0 | 1 | 0 | 0 | 1 | 764 | 0 | 0 | 0 | |
| 06-04 | 0 | 1 | 0 | 0 | 1 | 764 | 0 | 0 | 0 | |
| 07-04 | 0 | 4 | 0 | 0 | 4 | 764 | 0 | 0 | 0 | |
| 08-04 | 0 | 4 | 0 | 0 | 4 | 764 | 0 | 0 | 0 | |
| 09-04 | 0 | 4 | 0 | 0 | 4 | 764 | 0 | 0 | 0 | |
| 10-04 | 1 | 4 | 1 | 0 | 6 | 763 | 1 | 0 | 0 | |
| 11-04 | 1 | 4 | 1 | 0 | 6 | 763 | 1 | 0 | 0 | |
| 12-04 | 5 | 4 | 3 | 0 | 12 | 759 | 5 | 0 | 0 | |
| 13-04 | 5 | 5 | 2 | 0 | 12 | 759 | 5 | 0 | 0 | |
| 14-04 | 5 | 5 | 2 | 0 | 12 | 759 | 5 | 0 | 0 | |
| 15-04 | 25 | 9 | 15 | 0 | 49 | 739 | 25 | 0 | 0 | |
| 16-04 | 25 | 9 | 15 | 0 | 49 | 739 | 25 | 0 | 0 | |
| 17-04 | 37 | 12 | 10 | 0 | 59 | 727 | 37 | 0 | 0 | |
| 18-04 | 37 | 12 | 10 | 0 | 59 | 727 | 37 | 0 | 0 | |
| 19-04 | 37 | 12 | 10 | 0 | 59 | 727 | 37 | 0 | 0 | |
| 20-04 | 37 | 12 | 10 | 0 | 59 | 727 | 37 | 0 | 0 | |
| 21-04 | 51 | 14 | 11 | 0 | 76 | 713 | 51 | 0 | 0 | |

Joliette Women's Institution, Quebec (Maximum Population Estimate: 132 prisoners)

| Date | Cumulative test to | otals (N) | | | | Population c | ategories | | |
|--------|--------------------|-----------|---------|--------------|-------------|--------------|-------------------------|-----------|--------|
| (2020) | Positive (Cases) | Negative | Pending | Inconclusive | Total tests | Susceptible | Infected (Active Cases) | Recovered | Deaths |
| 30-03 | 0 | 0 | 7 | 0 | 7 | 132 | 0 | 0 | 0 |
| 31-03 | 0 | 0 | 7 | 0 | 7 | 132 | 0 | 0 | 0 |
| 01-04 | 0 | 0 | 7 | 0 | 7 | 132 | 0 | 0 | 0 |
| 02-04 | 0 | 0 | 7 | 0 | 7 | 132 | 0 | 0 | 0 |
| 03-04 | 0 | 0 | 7 | 0 | 7 | 132 | 0 | 0 | 0 |
| 04-04 | 0 | 0 | 7 | 0 | 7 | 132 | 0 | 0 | 0 |
| 05-04 | 0 | 0 | 7 | 0 | 7 | 132 | 0 | 0 | 0 |
| 06-04 | 0 | 0 | 7 | 0 | 7 | 132 | 0 | 0 | 0 |
| 07-04 | 10 | 0 | 6 | 0 | 16 | 122 | 10 | 0 | 0 |
| 08-04 | 10 | 0 | 6 | 0 | 16 | 122 | 10 | 0 | 0 |
| 09-04 | 10 | 0 | 6 | 0 | 16 | 122 | 10 | 0 | 0 |
| 10-04 | 16 | 0 | 0 | 0 | 16 | 116 | 16 | 0 | 0 |
| 11-04 | 16 | 0 | 0 | 0 | 16 | 116 | 16 | 0 | 0 |
| 12-04 | 17 | 0 | 0 | 0 | 17 | 115 | 17 | 0 | 0 |
| 13-04 | 17 | 0 | 0 | 0 | 17 | 115 | 17 | 0 | 0 |
| 14-04 | 17 | 0 | 0 | 0 | 17 | 115 | 17 | 0 | 0 |
| 15-04 | 48 | 17 | 10 | 0 | 75 | 84 | 48 | 0 | 0 |
| 16-04 | 48 | 17 | 10 | 0 | 75 | 84 | 48 | 0 | 0 |
| 17-04 | 51 | 25 | 0 | 0 | 76 | 81 | 41 | 10 | 0 |
| 18-04 | 51 | 25 | 0 | 0 | 76 | 81 | 41 | 10 | 0 |
| 19-04 | 51 | 25 | 0 | 0 | 76 | 81 | 41 | 10 | 0 |
| 20-04 | 51 | 25 | 0 | 0 | 76 | 81 | 41 | 10 | 0 |
| 21-04 | 51 | 25 | 0 | 0 | 76 | 81 | 35 | 16 | 0 |

Mission Medium Security Institution (Maximum Population Estimate: 216 prisoners)

| Date | Cumulative tes | st totals (N) | | | | Population c | ategories | | |
|--------|----------------|---------------|---------|--------------|-------|--------------|-----------|-----------|--------|
| | Positive | | | | Total | | | | |
| (2020) | (Cases) | Negative | Pending | Inconclusive | tests | Susceptible | | Recovered | Deaths |
| 30-03 | 0 | 0 | 0 | 0 | 0 | 216 | 0 | 0 | 0 |
| 31-03 | 0 | 0 | 0 | 0 | 0 | 216 | 0 | 0 | 0 |
| 01-04 | 0 | 0 | 0 | 0 | 0 | 216 | 0 | 0 | 0 |
| 02-04 | 0 | 0 | 0 | 0 | 0 | 216 | 0 | 0 | 0 |
| 03-04 | 0 | 0 | 0 | 0 | 0 | 216 | 0 | 0 | 0 |
| 04-04 | 0 | 0 | 0 | 0 | 0 | 216 | 0 | 0 | 0 |
| 05-04 | 0 | 0 | 0 | 0 | 0 | 216 | 0 | 0 | 0 |
| 06-04 | 0 | 0 | 0 | 0 | 0 | 216 | 0 | 0 | 0 |
| 07-04 | 11 | 1 | 17 | 0 | 29 | 205 | 11 | 0 | 0 |
| 08-04 | 14 | 1 | 21 | 0 | 36 | 202 | 14 | 0 | 0 |
| 09-04 | 14 | 1 | 21 | 0 | 36 | 202 | 14 | 0 | 0 |
| 10-04 | 25 | 6 | 15 | 0 | 46 | 191 | 25 | 0 | 0 |
| 11-04 | 25 | 6 | 15 | 0 | 46 | 191 | 25 | 0 | 0 |
| 12-04 | 35 | 6 | 14 | 0 | 55 | 181 | 35 | 0 | 0 |
| 13-04 | 41 | 6 | 16 | 0 | 63 | 175 | 41 | 0 | 0 |
| 14-04 | 41 | 6 | 16 | 0 | 63 | 175 | 41 | 0 | 0 |
| 15-04 | 54 | 10 | 6 | 0 | 70 | 162 | 54 | 0 | 0 |
| 16-04 | 54 | 10 | 6 | 0 | 70 | 162 | 54 | 0 | 0 |
| 17-04 | 60 | 12 | 19 | 2 | 93 | 156 | 59 | 0 | 1 |
| 18-04 | 60 | 23 | 8 | 2 | 93 | 156 | 44 | 15 | 1 |
| 19-04 | 60 | 23 | 8 | 2 | 93 | 156 | 44 | 15 | 1 |
| 20-04 | 60 | 23 | 8 | 2 | 93 | 156 | 44 | 15 | 1 |
| 21-04 | 64 | 27 | 12 | 2 | 105 | 152 | 42 | 21 | 1 |

Grand Valley Institution for Women (Maximum Population Estimate: 112 prisoners)

| Date | Cumulative tes | st totals (N) | | | | Population categories | | | | |
|--------|----------------|---------------|---------|--------------|-------|-----------------------|----------------|-----------|--------|--|
| | Positive | | | | Total | | Infected | | | |
| (2020) | (Cases) | Negative | Pending | Inconclusive | tests | Susceptible | (Active Cases) | Recovered | Deaths | |
| 30-03 | 1 | 1 | 0 | 0 | 2 | 214 | 1 | 0 | 0 | |
| 31-03 | 1 | 1 | 0 | 0 | 2 | 214 | 1 | 0 | 0 | |
| 01-04 | 1 | 1 | 0 | 0 | 2 | 214 | 1 | 0 | 0 | |
| 02-04 | 1 | 1 | 0 | 0 | 2 | 214 | 1 | 0 | 0 | |
| 03-04 | 1 | 1 | 0 | 0 | 2 | 214 | 1 | 0 | 0 | |
| 04-04 | 1 | 1 | 0 | 0 | 2 | 214 | 1 | 0 | 0 | |
| 05-04 | 1 | 1 | 0 | 0 | 2 | 214 | 1 | 0 | 0 | |
| 06-04 | 1 | 1 | 0 | 0 | 2 | 214 | 1 | 0 | 0 | |
| 07-04 | 7 | 24 | 5 | 0 | 36 | 208 | 7 | 0 | 0 | |
| 08-04 | 8 | 27 | 1 | 0 | 36 | 207 | 8 | 0 | 0 | |
| 09-04 | 8 | 27 | 1 | 0 | 36 | 207 | 8 | 0 | 0 | |
| 10-04 | 8 | 40 | 35 | 0 | 83 | 207 | 8 | 0 | 0 | |
| 11-04 | 8 | 40 | 35 | 0 | 83 | 207 | 8 | 0 | 0 | |
| 12-04 | 8 | 80 | 0 | 0 | 88 | 207 | 8 | 0 | 0 | |
| 13-04 | 8 | 80 | 1 | 0 | 89 | 207 | 8 | 0 | 0 | |
| 14-04 | 8 | 80 | 1 | 0 | 89 | 207 | 8 | 0 | 0 | |
| 15-04 | 8 | 81 | 17 | 0 | 106 | 207 | 8 | 0 | 0 | |
| 16-04 | 8 | 81 | 17 | 0 | 106 | 207 | 8 | 0 | 0 | |
| 17-04 | 8 | 99 | 1 | 0 | 108 | 207 | 6 | 2 | 0 | |
| 18-04 | 8 | 99 | 1 | 0 | 108 | 207 | 6 | 2 | 0 | |
| 19-04 | 8 | 99 | 1 | 0 | 108 | 207 | 6 | 2 | 0 | |
| 20-04 | 8 | 99 | 1 | 0 | 108 | 207 | 6 | 2 | 0 | |
| 21-04 | 8 | 100 | 0 | 0 | 108 | 207 | 6 | 2 | 0 | |