

1 Updated estimates of comorbidities associated with risk for COVID-19 complications based on  
2 US data

3 Mary L. Adams, MS, MPH, Joseph Grandpre, PhD, MPH, David L. Katz, MD, MPH, FACPM,  
4 FACP, FACLM.

5 Author affiliations: On Target Health Data LLC, Suffield, CT, USA (M.L. Adams); Wyoming  
6 Department of Health, Cheyenne, WY, USA (J. Grandpre); True Health Initiative, Derby, CT,  
7 USA (D. L. Katz).

8 Corresponding author: Mary Adams, 247 N Stone St, West Suffield, CT 06093, (860-370-9035).

9 Abstract (149 words)

10 We updated previous estimates ([wwwnc.cdc.gov/eid/article/26/8/20-0679\\_article](http://wwwnc.cdc.gov/eid/article/26/8/20-0679_article)) of adults with  
11 any underlying condition increasing risk of complications from COVID-19 using recent US  
12 hospitalization data instead of mortality data from China. This substitutes obesity for cancer in  
13 the definition and increased the percentage of adults reporting  $\geq 1$  condition to 56.0% (95% CI  
14 55.7-56.4). When controlled for all measures listed, factors increasing odds of reporting any of  
15 the underlying conditions include being male, older, African American, American Indian,  
16 household income  $< \$25,000$ ,  $<$  high school education, underinsurance, living in the South or  
17 Midwest (vs. West), plus the risk factors of ever smoking, sedentary lifestyle, and inadequate  
18 fruit and vegetable consumption. Population-attributable risk for the listed risk factors was  
19 13.0%, 12.6%, and 15.0% respectively. Results have potential implications for policies based on  
20 risk-stratification of the population and for improvement of risk status through lifestyle change.  
21 National support for a “health promotion” campaign would be timely.

22 Word count: Abstract: 149; text: 2,510

23 Keywords: COVID-19; risk factors; complications; hospitalizations; attributable risk

24

25

26 One of the novel aspects of the current coronavirus pandemic is that information is  
27 constantly being updated as more studies are published. We had earlier reported estimates of  
28 comorbidities affecting risk for complications from COVID-19 that were based on mortality data  
29 from China (1). Newly released US data on hospitalizations for COVID-19 from 14 states  
30 indicate a slightly different list of conditions affecting risk, with obesity added and cancer  
31 subtracted (2). Those preliminary results indicate that 89% of those hospitalized for COVID-19  
32 had an underlying condition, the most common being hypertension, obesity, chronic lung disease  
33 (including asthma and chronic obstructive pulmonary disease), diabetes, and cardiovascular  
34 disease (2). Similar results were found for ICU admissions on a smaller sample and without  
35 obesity (3). The Guidelines for Opening up America Again released by the White House on  
36 April 16, 2020 (4) include a similar list to define vulnerable individuals, except that list includes  
37 elderly (undefined) and those with a compromised immune system and omits cardiovascular  
38 disease.

39 Our main objective for this study was to use population based US data to estimate the  
40 fraction of adults in the community who might be a risk of hospitalization for COVID-19 (2) due  
41 to reporting any of the most common underlying conditions listed above. Secondly, we sought  
42 to determine the demographic and risk factors that increased the likelihood of having one or  
43 more of these underlying conditions. And thirdly, using the information about potentially  
44 modifiable risk factors obtained through our second objective, we estimated population  
45 attributable-risk for conditions increasing risk for COVID-19 hospitalization.

46

47 Methods

48 We used publicly available 2017 Behavioral Risk Factor Surveillance System (BRFSS)  
49 data (5) from telephone surveys of 444,649 randomly selected adults ages 18 and older in the 50  
50 states and the District of Columbia (DC). The BRFSS includes only non-institutionalized adults  
51 so residents of prisons, nursing homes and assisted living facilities are among those not  
52 surveyed. We chose to use 2017 data in order to include hypertension which was not addressed  
53 in 2018. Data were adjusted for the probability of selection and weighted to be representative of  
54 the adult population in each state by age, gender, race/ethnicity, marital status, education, home  
55 ownership, and telephone type and included weights and stratum variables needed for analysis.  
56 Results were not age adjusted in order to reflect the age distribution of each state rather than a  
57 standard population. The median response rate for cell phone and land line surveys combined  
58 was 47.2%, ranging from 33.9% in California to 61.1% in Utah (6). Reliability and validity of  
59 the BRFSS have been found to be moderate to high for many survey measures, in particular  
60 those used here which can be checked versus medical records (7).

61 The key variable was a composite measure including adults reporting they were ever told  
62 they had diabetes (excluding if only when pregnant), asthma (and still had it), chronic obstructive  
63 pulmonary disease (COPD), hypertension, cardiovascular disease (CVD; ever told they had a  
64 heart attack, angina, coronary heart disease, or a stroke) or they were obese with body mass  
65 index  $\geq 30.0$  based on self-reported height and weight. The number of conditions was counted for  
66 each respondent and adults who reported at least one were considered to be at risk of  
67 hospitalization due to an underlying condition. Risk factors included ever smoking 100  
68 cigarettes, sedentary lifestyle, defined as no leisure time physical activity in the past month, and  
69 inadequate fruit and vegetable consumption, defined as consuming the combination  $<5$  times per  
70 day based on responses to five separate questions, with responses to the question about fried

71 potatoes or French fries excluded. A composite measure was also created for risk factors by  
72 counting the number for each respondent and determining those with each number of risks from  
73 0-3.

74 Demographic measures included age group (18-29, 30-39, 40-49, 50-59, 60-69, 70-79, and  
75 80+ years, which was created by combining 5 year age groups provided in the data set), self-  
76 reported race/ethnicity (non-Hispanic white, Black or African American, Hispanic of any race,  
77 American Indian/Alaska native, Asian/Pacific Islander, and other), less than a high school  
78 education (vs. high school graduate or higher), household income < \$25,000 (vs. all other  
79 incomes including unknown), being underinsured (having no health insurance coverage or  
80 reporting a time in the past year when the needed health care but could not get it due to cost),  
81 state of residence which included DC, whether or not they lived within the center city of a  
82 Metropolitan Statistical Area (MSA), and census region (Northeast, Midwest, South, or  
83 West)(8).

84  
85 Stata version 14.1 (Stata Corp LP, College Station, TX) was used for analysis to account for the  
86 complex sample design of the BRFSS. Point estimates and 95% confidence intervals are  
87 reported using the weights, stratum, and PSU variables supplied in the data set (5). Missing  
88 values were excluded from analysis. Population-attributable risk (PAR) was used to compare the  
89 contributions of individual risk factors to the composite measure of any of the 6 underlying  
90 conditions and to each of the 6 component conditions (9). PAR estimates take into account not  
91 only the relative risk of people with that risk factor developing the outcome, but also the  
92 prevalence of the risk factor in the population (10). PARs were estimated in Excel using Levin's

93 formula (11) and adjusted odds ratios (AOR) obtained from logistic regression instead of  
94 unadjusted relative risk.

## 95 Results

96 Sample was described earlier (1). Overall, 56.0% (95% CI 55.7-56.4) of respondents had  
97 one or more of the underlying conditions, ranging from 33.7% among 18-29 year olds to 79.1%  
98 for those 80 years and older. Prevalence rates of the separate conditions were 8.5% for CVD,  
99 6.6% for COPD, 9.1% for asthma, 10.8% for diabetes, 32.4% for hypertension, and 30.1% for  
100 obesity. While the percentage of adults with any of the conditions increased with age (Table 1),  
101 60.7% of the total was younger than age 60. Rates also varied by all the measures in Table 1  
102 except underinsurance. State rates ranged from 45.6% in DC to 68.8% in West Virginia. Using  
103 the definition for vulnerable individuals according to the Guidelines for Opening up America  
104 (4) which adds age (we used age 65 as elderly) and removes CVD, that prevalence rate is 60.4%  
105 (95% CI 60.0-60.7).

106 Prevalence rates for the risk factors used in PAR estimates for all ages were 40.4% for ever  
107 smoking, 26.6% for sedentary lifestyle, and 84.1% for inadequate fruit and vegetable  
108 consumption. Among ever smokers of all ages, 40.7% currently smoke, including 50.6% of those  
109 ages 18-59 and 22.8% of adults ages 60 and older. Those estimates result in current smoking  
110 rates of 16.5%, 18.6%, and 11.2% respectively.

111 Results of multiple logistic regression controlled for age, gender, education, income,  
112 race/ethnicity, underinsurance, region, and the 3 behavioral risk factors (Table 2) indicate that  
113 male gender, increasing age, less education and income, being underinsured, having each of the  
114 risk factors, being African American, or American Indian/Alaska Native and living in either the

115 Midwest or South significantly increased the odds of having an underlying condition shown to be  
116 associated with COVID-19 hospitalizations. Hispanic ethnicity had no effect while being  
117 Asian/Pacific Islander decreased the odds of having any of the conditions. Adding a variable for  
118 living in the center city of an MSA had no effect when controlled for the other measures.  
119 Including the composite risk factor measure in the logistic regression model in place of the  
120 separate risks confirmed the step-wise increase with an AOR of 2.56 (2.38-2.76) for adults with  
121 all 3 risk factors compared with those reporting none.

122 PAR estimates for the 3 potentially modifiable risk factors, estimated using the AORs in  
123 Table 2 are shown in Table 3. Among all adults, these 3 risk factors were estimated to account  
124 for 40.6% of the total risk of any underlying condition, 34.2% of total risk for adults 18-59 and  
125 59.7% of total risk for adults age 60 and older. The total attributable risk for all adults for the  
126 separate conditions ranged from 18.7% for asthma to 69.2% for COPD. (Table 3).

## 127 Discussion

128 We estimate that 56.0% of US adults, with a wide range across age groups and states, have  
129 one or more underlying conditions that increase risk of hospitalization due to COVID-19. Our  
130 previous definition of adults at risk of complications from COVID-19 (1) that excluded obesity  
131 and added cancer based on data for China where obesity is much less prevalent (13) resulted in  
132 an estimate of 45.4%. Using the definition of vulnerable individuals in the White House  
133 Guidelines (4) resulted in an estimate for those at risk of 60.4%. Thus, except for measures that  
134 exclude obesity, it appears that over 50% of all adults are at risk for hospitalization or can be  
135 considered vulnerable individuals. In addition, while rates increase with increasing age group  
136 and age is the key predictor of reporting an underlying condition, over half of adults with

137 underlying conditions using *any* of these definitions are < 60 years. Although our definition of  
138 underlying conditions does not include every condition reported among the 89% of persons  
139 hospitalized for COVID-19 (2) who reported any, our population based prevalence of 56% for  
140 any of the conditions is consistent with any of these 6 conditions increasing risk for  
141 hospitalization. The underlying chronic conditions increasing risk for COVID-19 hospitalizations  
142 are also very similar to those increasing risk for seasonal influenza complications (14).

143 All the underlying conditions used in our measures are conditions for which behavioral risk  
144 factors have been well established (9, 12). Our list of underlying conditions includes obesity,  
145 diabetes, and hypertension, which are also considered risk factors in some studies (9, 12). Causal  
146 relationships between risk factors and outcomes can be complicated; for example, previous  
147 results (9) have shown that obesity contributes 37.8% to attributable-risk for diabetes, 30.9% to  
148 hypertension and 16.4% to asthma, while hypertension and diabetes contribute 29.3% and 6.7%  
149 respectively to CVD. For simplification, we limited our list of risk factors to lifetime smoking,  
150 sedentary lifestyle, and inadequate fruit and vegetable consumption. Each selected risk factor  
151 was associated with increased likelihood of reporting any of the 6 underlying conditions with  
152 AORs of 1.21 to 1.54. In addition, the unadjusted results for increasing number of risk factors  
153 indicate a step-wise increase in the percentage of adults reporting any of the underlying  
154 conditions with each additional risk factor as shown in Table 1. This result was confirmed by  
155 logistic regression and provides new and informative data on factors that predict increased  
156 likelihood of reporting any of the underlying conditions.

157 The results on the risk factors suggest the impact is consequential, with total PARs for the 3  
158 risk factors of 40.6% overall, 34.2% for adults 18-59 and 59.7% for adults ages 60 and older. Of  
159 course the smoking measure included lifetime smoking in order to capture effects among former

160 smokers, while current smoking rates are much lower at 16.5% overall. The results suggest the  
161 potential for considerable improvement by reducing just these 3 risk factors, or even 2 of the 3.  
162 For example, increasing fruit and vegetable consumption and initiating an exercise program such  
163 as walking could potentially lower the future risk of developing any of these underlying  
164 conditions, with total elimination lowering risk by as much as 46.0% for adults ages 60 and older  
165 and 21.6% for those 18-59 years.

166 In addition to the risk factors mentioned above, a long list of variables increased the  
167 likelihood of reporting an underlying condition when controlled for other factors. Foremost  
168 among them are being older, male, and African American, all groups which preliminary  
169 hospitalization data (2) suggested were disproportionately affected by COVID-19. Results from  
170 this study indicate that even when controlled for age, sex, underinsurance, income, region, and  
171 the 3 risk factors, men, African-Americans, and adults age 60 and older are all more likely to  
172 report any of the underlying conditions increasing risk for hospitalizations, with AORs of 1.15,  
173 1.67 and  $\geq 4.79$  respectively. Thus it appears part of the reason these groups were  
174 disproportionately hospitalized for COVID-19 (2) is because they are more likely to report any  
175 of the underlying conditions. More study will be needed to determine if other factors increasing  
176 likelihood of underlying conditions such as less education and income, and being underinsured  
177 or American Indian, are confirmed as increasing hospitalization rates among adults with  
178 COVID-19.

179 In this study we grouped states into regions based on the 4 census regions (8) which clearly  
180 showed that living in either the Midwest or South increased risk of reporting an underlying  
181 condition compared with the West (or Northeast). This result is consistent with studies showing  
182 that obesity rates are also highest in these regions (15). And while living in a city (center city of

183 an MSA) was associated with reporting an underlying condition in unadjusted analysis, once  
184 controlled for demographics and risk factors, urban residence was no longer statistically  
185 significant. On the other hand, underinsurance was not associated with reporting an underlying  
186 condition in unadjusted analysis but was a statistically significant predictor of reporting an  
187 underlying condition once controlled for all measures in the model. Underinsurance has the  
188 potential to present a barrier to hospitalization for any reason.

### 189 Limitations

190 Our study does not address possible differences in contracting the disease, only the risk of  
191 hospitalization among those with COVID-19, based on preliminary US results for underlying  
192 conditions (2,3). Only non-institutionalized adults are surveyed so 1.3 million adults in nursing  
193 homes (16) were excluded which almost certainly underestimates those with underlying  
194 conditions who are included in hospitalization data. Data are self-reported and reliability and  
195 validity can vary for different measures tested (7). But as long as a respondent was told they had  
196 a chronic condition, validity was high. The variable for living in the center city of an MSA was  
197 missing for a large number of respondents. Reverse causality may be a factor in some of the PAR  
198 estimates, especially for sedentary lifestyle. Age groups used for analysis did not match those  
199 used for weighting data but that should have a minimal effect on results. Low response rates  
200 could introduce bias but, as noted, validity appears high for the measures used in this study. A  
201 dearth of denominator data addressing the actual prevalence of coronavirus infection and  
202 immunity in the United States makes it difficult to draw reliable conclusions about overall rates  
203 of hospitalization or death among those exposed.

204 Conclusion: We estimate 56.0% of US adults are at risk of needing hospitalization for COVID-  
205 19 due to underlying conditions. Such estimates will vary depending on exact criteria used to  
206 define underlying conditions but represent a substantial fraction of all US adults. These  
207 underlying conditions are, in turn, associated with modifiable risk factors including ever  
208 smoking, being sedentary and inadequate fruit and vegetable consumption. The three risk factors  
209 were estimated to contribute 40.6% of attributable-risk for all adults reporting any of the  
210 underlying conditions in this study including 34.2% of attributable-risk among adults ages 18-59  
211 and 59.7% for adults ages 60 and older. These results suggest the potential for policies based on  
212 risk-stratification of the population and for possible improvement of risk status through lifestyle  
213 change. A national focus on, and support for, a “health promotion” campaign would be timely.

214

215 Funding: Data collection, analysis, and interpretation of data for this study were supported  
216 by the Centers for Disease Control and Prevention (CDC) Grant/Cooperative Agreement number  
217 1U58DP006069-01.

218

219

220

221

222 References

- 223 1. Adams ML, Katz DL, Grandpre J. Population-based estimates of concurrent conditions  
224 affecting risk for complications from coronavirus disease, United States. *Emerg Infect*  
225 *Dis.* 2020 Aug [date cited]. <https://doi.org/10.3201/eid2608.200679>
- 226 2. Garg S, Kim L, Whitaker M, et al. Hospitalization Rates and Characteristics of Patients  
227 Hospitalized with Laboratory-Confirmed Coronavirus Disease 2019 — COVID-NET, 14  
228 States, March 1–30, 2020. *MMWR Morb Mortal Wkly Rep* 2020;69:458–464. DOI:  
229 <http://dx.doi.org/10.15585/mmwr.mm6915e3>
- 230 3. Preliminary Estimates of the Prevalence of Selected Underlying Health Conditions  
231 Among Patients with Coronavirus Disease 2019 — United States, February 12–March  
232 28, 2020. *MMWR Morb Mortal Wkly Rep* 2020;69:382–386. DOI:  
233 <http://dx.doi.org/10.15585/mmwr.mm6913e2>
- 234 4. The White House (Washington, DC). Guidelines Opening Up America Again. April 16,  
235 2020. Available at:  
236 [https://www.whitehouse.gov/openingamerica/?fbclid=IwAR2m39LPMgNUAcDLvjRR3](https://www.whitehouse.gov/openingamerica/?fbclid=IwAR2m39LPMgNUAcDLvjRR3DHLh1Lhwxlc-o3BN3Eous0FSeagdsuj2Jwuf3U)  
237 [DHLh1Lhwxlc-o3BN3Eous0FSeagdsuj2Jwuf3U](https://www.whitehouse.gov/openingamerica/?fbclid=IwAR2m39LPMgNUAcDLvjRR3DHLh1Lhwxlc-o3BN3Eous0FSeagdsuj2Jwuf3U) Accessed April 18, 2020.
- 238 5. Behavioral Risk Factor Surveillance System (BRFSS) Survey Data and Documentation.  
239 [https://www.cdc.gov/brfss/data\\_documentation/index.htm](https://www.cdc.gov/brfss/data_documentation/index.htm). Accessed March 3, 2020.
- 240 6. Behavioral Risk Factor Surveillance System 2017 Summary Data Quality Report,  
241 Centers for Disease Control and Prevention, Atlanta, GA. June 13, 2018  
242 [https://www.cdc.gov/brfss/annual\\_data/2015/pdf/2015-](https://www.cdc.gov/brfss/annual_data/2015/pdf/2015-sdqr.pdf)  
243 [sdqr.pdf](https://www.cdc.gov/brfss/annual_data/2017/pdf/2017-sdqr-508.pdf)[https://www.cdc.gov/brfss/annual\\_data/2017/pdf/2017-sdqr-508.pdf](https://www.cdc.gov/brfss/annual_data/2017/pdf/2017-sdqr-508.pdf) , accessed  
244 November 16, 2018.

- 245 7. Pierannunzi C, Hu SS, Balluz L. A systematic review of publications assessing reliability  
246 and validity of the Behavioral Risk Factor Surveillance System (BRFSS), 2004-2011.  
247 BMC Med Res Methodol. 2013 Mar 24;13:49. Doi: 10.1186/1471-2288-13-49.
- 248 8. US Census Bureau. Geographic Terms and Concepts – Census Divisions and Census  
249 Regions. [https://www.census.gov/geo/reference/gtc/gtc\\_census\\_divreg.html](https://www.census.gov/geo/reference/gtc/gtc_census_divreg.html) accessed  
250 January, 27, 2018.
- 251 9. Adams M, Grandpre J, Katz D, Shenson D. The impact of key modifiable risk factors on  
252 leading chronic conditions. *Prev Med.* 2019 Mar;120:113-118.  
253 <https://doi.org/10.1016/j.ypmed.2019.01.006>.
- 254 10. Brownson RC, Remington PL, Wegner, MV. *Chronic Disease Epidemiology and*  
255 *Control.* 4th ed. Washington, DC: American Public Health Association; 2016.
- 256 11. Rückinger S, von Kries R and Toschke AM. An illustration of and programs estimating  
257 attributable fractions in large scale surveys considering multiple risk factors. *BMC Med*  
258 *Res Methodol.* 2009; 9: 7. doi: 10.1186/1471-2288-9-7.
- 259 12. Adams ML, Grandpre J, Katz DL, Shenson D. Linear association between number of  
260 modifiable risk factors and multiple chronic conditions: Results from the Behavioral Risk  
261 Factor Surveillance System. *Prev Med.* 2017 Dec;105:169-175. doi:  
262 10.1016/j.ypmed.2017.09.013. Epub 2017 Sep 14.
- 263 13. Wu, Y. Overweight and obesity in China. *BMJ.* 2006 Aug 19; 333(7564): 362–363.  
264 doi: 10.1136/bmj.333.7564.362.
- 265 14. Influenza (flu): People at High Risk For Flu Complications. Centers for Disease Control  
266 and Prevention. (Atlanta, Georgia).

267 [https://www.cdc.gov/flu/highrisk/index.htm?CDC\\_AA\\_refVal=https%3A%2F%2Fwww.cdc.gov%2Fflu%2Fabout%2Fdisease%2Fhigh\\_risk.htm](https://www.cdc.gov/flu/highrisk/index.htm?CDC_AA_refVal=https%3A%2F%2Fwww.cdc.gov%2Fflu%2Fabout%2Fdisease%2Fhigh_risk.htm).

268 [cdc.gov%2Fflu%2Fabout%2Fdisease%2Fhigh\\_risk.htm](https://www.cdc.gov/flu/highrisk/index.htm?CDC_AA_refVal=https%3A%2F%2Fwww.cdc.gov%2Fflu%2Fabout%2Fdisease%2Fhigh_risk.htm).

269 15. Centers for Disease Control and Prevention (Atlanta, GA). Overweight and obesity:

270 obesity prevalence maps. 2018. <https://www.cdc.gov/obesity/data/prevalence-maps.html>.

271 Accessed April 29, 2020.

272 16. Centers for Disease Control and Prevention (Atlanta, GA). National Center for Health

273 Statistics; Nursing Home Care. <https://www.cdc.gov/nchs/fastats/nursing-home-care.htm>.

274

Table 1. Demographics of adults with any of 6 underlying conditions<sup>a</sup>;

2017 Behavioral Risk Factor Surveillance System.

<b>Population Group</b>	<b>Percent</b>	<b>95% CI</b>	<b>Sample Size</b>
Total	56.0	55.7-56.3	398,865
<b>Gender</b>			
Males	56.7	56.2-57.2	183,097
Females	55.3	54.8-55.8	215,768
<b>Age (years)</b>			
18-29	33.7	32.9-34.5	42,434
30-39	45.0	44.2-45.9	44,676
40-49	53.6	52.7-54.5	49,449
50-59	63.4	62.6-64.1	73,839
60-69	71.4	70.8-72.1	90,088
70-79	78.8	78.0-79.6	63,627
80+	79.1	77.9-80.2	31,915
<b>Race/ethnicity</b>			
White (non-Hispanic)	56.7	56.3-57.0	307,282
African-American	65.8	64.7-66.8	31,939
Hispanic	52.3	51.2-53.5	26,837
American Indian/Alaska Native	67.7	65.3-70.1	7,468
Asian/Pacific Islander	34.1	32.1-36.3	9,108
Other	57.0	55.0-59.0	10,089
<b>Education</b>			
<High school	66.8	65.6-67.9	26,450
High school	59.6	58.9-60.2	107,876
Some college	57.2	56.5-57.8	111,267
College graduate	46.4	45.9-46.9	152,651
<b>Household income</b>			
< \$15,000	67.5	66.2-68.7	31,888
\$15K-\$24,999	63.6	62.7-64.6	54,891
\$25K-\$49,999	58.9	58.1-59.6	85,409
\$50K-\$74,999	55.6	54.7-56.6	55,794
\$75K+	47.9	47.3-48.5	116,138
Unknown	54.8	53.9-55.8	54,903

**Census Region**

Northeast	54.0	53.3-54.8	67,564
Midwest	57.4	56.9-58.0	119,303
South	58.9	58.3-59.5	122,132
West	51.6	50.8-52.4	90,056

**Relation to center city of MSA**

Center City	65.2	64.2-66.1	53,318
Not center city	55.1	54.7-55.5	345,737

**Underinsured**

Yes	56.4	55.6-57.3	58,967
No	56.0	55.6-56.4	337,892

**Risk Factors**

**Ever smoked**

Yes	64.3	63.8-64.8	170,149
No	50.5	50.0-51.0	219,779

**Sedentary lifestyle**

Yes	67.8	67.2-68.5	100,438
No	52.3	51.9-52.7	276,037

**Eat <5 a day**

Yes	57.0	56.6-57.4	300,938
No	50.0	49.0-50.9	58,286

**Number of risk factors**

0	42.8	41.6-44.0	31,529
1	48.9	48.4-49.5	146,808
2	62.1	61.5-62.7	133,833
3	73.2	72.3-74.1	43,147

**State**

AL	65.6	63.9-67.3	6,083
AK	56.5	53.4-59.5	2,961
AZ	55.5	54.4-56.6	13,824
AR	65.1	62.5-67.7	4,713
CA	50.2	48.7-51.7	8,304
CO	47.6	46.3-48.9	8,812

CT	54.0	52.5-55.4	9,302
DE	59.2	56.9-61.5	3,554
DC	45.6	43.5-47.6	4,012
FL	57.4	55.7-59.0	19,402
GA	57.3	55.5-59.1	5,247
HI	52.5	50.9-54.1	7,226
ID	53.3	51.2-55.3	4,535
IL	56.8	55.1-58.6	5,208
IN	59.7	58.5-60.9	12,676
IA	58.6	57.2-59.9	6,991
KS	58.0	57.1-58.9	19,304
KY	62.7	60.9-64.5	7,826
LA	61.9	60.0-63.9	4,328
ME	58.4	56.8-60.0	9,141
MD	56.4	55.0-57.9	12,056
MA	52.1	50.1-54.2	6,086
MI	59.4	58.1-60.6	9,863
MN	49.9	48.9-50.9	15,155
MS	63.3	61.1-65.5	4,627
MO	57.9	56.2-59.5	6,828
MT	51.4	49.5-53.3	5,409
NE	56.6	55.3-57.9	14,059
NV	55.9	53.3-58.4	3,427
NH	54.5	52.4-56.6	5,077
NJ	55.2	53.5-56.8	10,160
NM	56.1	54.3-58.0	5,908
NY	51.5	50.2-52.8	10,937
NC	59.2	57.2-61.1	4,386
ND	55.4	53.7-57.1	6,430
OH	59.4	58.0-60.8	11,138
OK	62.7	61.0-64.4	5,891
OR	54.7	53.0-56.4	4,757
PA	57.3	55.6-59.0	6,022
RI	56.9	54.8-58.9	4,987
SC	61.6	60.2-63.0	10,191
SD	55.2	52.9-57.5	6,382
TN	61.1	59.2-63.0	5,215
TX	56.5	54.5-58.4	10,972
UT	47.4	46.1-48.7	9,115
VT	54.2	52.4-56.0	5,852
VA	55.8	54.3-57.3	8,630

WA	53.6	52.4-54.8	11,694
WV	68.8	67.1-70.5	4,999
WI	55.4	53.5-57.3	5,269
WY	53.6	51.7-55.6	4,084

275 <sup>a</sup> Cardiovascular disease, diabetes, chronic obstructive pulmonary disease,

276 asthma, hypertension, and/or obesity.

277

Table 2. Multiple logistic regression results for outcome of reporting any of 6 underlying conditions <sup>a</sup>; 2017 Behavioral Risk Factor Surveillance System; N=346,198.

<b>Group</b>	<b>Adjusted Odds Ratio</b>	<b>95% CI</b>
Females (referent)		
Males	1.15	1.12-1.19
Age 18-29 (referent)		
30-39	1.57	1.49-1.67
40-49	2.21	2.09-2.34
50-59	3.27	3.10-3.46
60-69	4.79	4.53-5.06
70-79	7.19	6.73-7.68
80+	7.01	6.34-7.76
Never smoked (referent)		
Ever smoked	1.37	1.33-1.42
Not sedentary (referent)		
Sedentary	1.54	1.48-1.60
Fruit & vegetables 5 X/day (referent)		
< 5 time/day	1.21	1.16-1.26
High school grad/GED (referent)		
Not high school graduate	1.23	1.14-1.31
Income \$25,000+ (referent)		
Income <\$25,000	1.46	1.40-1.53
Not underinsured (referent)		
Underinsured	1.11	1.06-1.16
White non-Hispanic (referent)		
African American	1.67	1.57-1.76
Hispanic any race	1.03	0.97-1.10
American Indian/AK Native	1.67	1.46-1.92
Asian/Pacific Islander	0.57	0.51-0.63
Other	1.27	1.15-1.40
West region (referent)		
Northeast	0.98	0.94-1.03
Midwest	1.14	1.09-1.19
South	1.13	1.08-1.19

278 <sup>a</sup> Cardiovascular disease, diabetes, chronic obstructive pulmonary disease,

279 asthma, hypertension, and/or obesity.

Table 3. Population attributable-risk (PAR) for selected outcomes.

2017 Behavioral Risk Factor Surveillance System; Maximum N=372,041

Outcome	PAR	PAR	PAR	Sum of PARs
	Ever smoked	Sedentary lifestyle	Fruits/vegetables < 5X/day	
Any of 6 conditions <sup>a</sup> all adults	13.0%	12.6%	15.0%	40.6%
Any of 6 conditions <sup>a</sup> age <60	12.6%	9.8%	11.8%	34.2%
Any of 6 conditions <sup>a</sup> age 60+	13.6%	21.6%	24.4%	59.7%
Diabetes	5.0%	11.5%	15.6%	32.1%
Hypertension	8.5%	8.5%	13.1%	30.2%
COPD	52.5%	16.6%	0.0%	69.2%
Asthma	11.8%	6.9%	0.0%	18.7%
CVD	22.3%	8.7%	0.0%	31.0%
Obesity	0.0%	14.5%	13.8%	28.3%

280

281 Abbreviations: PAR: population attributable risk; COPD: chronic obstructive pulmonary disease; CVD: cardiovascular disease.

282 <sup>a</sup> Cardiovascular disease, diabetes, chronic obstructive pulmonary disease, asthma, hypertension, and/or obesity, conditions

283 found to be associated with US hospitalizations for COVID-19.