Temperature and relative humidity are not major contributing factor on the occurrence
of COVID-19 pandemic: An observational study in 57 countries

Md. Abdus Sobur,[1] Md. Saiful Islam,[1]Md. Emdadul Haque,[2]Md. Taohidul Islam,[3]Antonio
Toniolo,[4] Md. Tanvir Rahman[1]*

[1]Department of Microbiology and Hygiene, Faculty of Veterinary Science, Bangladesh
Agricultural University, Mymensingh 2202, Bangladesh.

[2]Faculty of Fisheries, Bangladesh Agricultural University, Mymensingh 2202, Bangladesh.

[3]Department of Medicine, Faculty of Veterinary Science, Bangladesh Agricultural
University, Mymensingh 2202, Bangladesh.

[4]Global Virus Network, University of Insubria, 21100 Varese, Italy.
*Corresponding Author: Dr. Md. Tanvir Rahman

Professor, Department of Microbiology and Hygiene, Faculty of Veterinary Science,
Bangladesh Agricultural University, Mymensingh-2202, Bangladesh.

Phone: +8809167401-6 Ext. 6385, Mobile. +8801913323307, Fax: +8809161510.

Email: tanvirahman@bau.edu.bd

Abstract

The world searching for hope has already experienced a huge loss of lives due to COVID-19
caused by SARS-CoV-2 started in Wuhan, China. There are speculations that climatic
conditions can slowdown the transmission of COVID-19.Findings from the early outbreak
indicated the possible association of air temperature and relative humidity in COVID-19
occurrence in China. Current study focused on whether climatic conditions(temperature and
relative humidity)are having any influence in the occurrence of COVID-19 when the
outbreak has been classified as pandemic. To determine the effect of daily average
temperature and average relative humidity on log-transformed total daily cases of COVID-19,
polynomial regression as a quadratic term and linear regression were done. Linear regression
analysis was also carried out to explore the same effect on selected countries. Present study
observed no correlation between the climatic conditions (the daily average temperature and
relative humidity) and the number of cases of COVID-19. Similar result was found in relation
between daily average temperature and average number of cases per day in country-wise
analysis. However, about 93.5% cases of COVID-19 occurred between 1^0C to 16^0C and the
average number of cases per day was lower in high temperature country than low temperature
country with exceptions. The minimum effect of summer temperature may not be effective to
control the pandemic rather need to apply the control measures of COVID-19.

Keywords: COVID-19; temperature; relative humidity; contact; transmission

1. Introduction

COVID-19 caused by SARS-CoV-2 is an enveloped, positive sense single stranded RNA virus of the family *Coronaviridae*, a major pathogen of respiratory illness (Pal et al., 2020). Since the outbreak of COVID-19in Wuhan, China on the late December, 2019 (Lu et al., 2020),it has rapidly spread throughout China and all over the world (Gorbalenya, 2020; Arab-Mazar et al., 2020; Chen et al., 2020; Huang et al., 2020; Wang et al., 2020a; Holshue et al., 2020; Wang et al., 2020b).The major contributing factor for this new epidemic was the importation of COVID-19 cases from epidemic countries. So far, more than 200 countries and territories have been affected with 1,395,136 confirmed cases and 81,580 deaths, togetherwith cessation of normal social operations and massive economic loss (WHO, 2020a).Earlier, WHO, observing the current situation across the world, declared COVID-19 as "a pandemic" (WHO, 2020b).

The experience of SARS and MERS outbreak differs with COVID-19 in extensiveness of spread, and death rate but finds similarity in sign-symptoms and transmission pattern (Wang et al., 2020c). Seasonality of respiratory viral disease is an old theme that recognized winter as an important factor of influenza like diseases transmission (Sobsey and Meschke, 2003; Wolkoff, 2018). The transmission of virus is under the influence of climate conditions e.g., temperature and relative humidity (Hemmes, et al., 1960,Dalziel, et al. 2018).Several studies found droplet, aerosol, and direct and indirect contact can act as the major factors in the transmission of SARS like viruses (Peng et al., 2020; Rodríguez-Morales et al., 2020; To et al., 2020).Survivability of SARS virus in environment and surfaces is associated with temperature and relative humidity(Chan et al., 2011). High temperature reduces aerosol but cannot inhibit contact transmission immediately (Lowen et al., 2008). SARS-CoV-1 and SARS-CoV-2 are able to survive in air and surface for sizable time periods, thus enabling the virus more capable to affect human (VanDoremalen et al., 2020). Survival of SARS-CoV-2 in air and surfacesassociated with climate with is still unclear.

Although some studies found relationships of climatic conditions with the occurrence of COVID-19 cases (Wang et al., 2020d, Wang et al., 2020e, Zhu and Xie, 2020), they had limited option to include high temperature country as the outbreak was manly circulating in temperate zone during the early stage of outbreak.Therefore, this study aimed to understand the possible role of temperature and relative humidityin the occurrence of COVID-19 in 57 countries from tropical and temperate zone.

2. Materials and methods

2.1. Data

Countries having at least 500 cumulative cases by March 28, 2020 were selected forstudy. Once a country reported at least 10 cases on particular day was considered for data recordingfor that day. Date wise metrological parameter, like temperature and relative humidity were obtained from 57 countries to find out a possible link between COVID-19 cases and metrological data. The highest and the lowest temperature were recorded and the average of these two data was considered as the daily temperature. The daily relative

humidity was recorded as AM (Before mid-day) and PM (After mid-day). The average was
considered as the daily relative humidity.

2.2. Source of Data

We extracted information from national and international reports of the outbreak and
metrological data from relevant websites. The daily case reports were documented from
"WHO", "Worldometer", and "Kaiser Family Foundation"websites (WHO, 2020c;
Worldometer, 2020; KFF, 2020). The daily temperature and relative humidity data were
recorded from "The Weather Channel", "Timeanddate" and "Accuweather" websites (The
Weather Channel, 2020; Timeanddate, 2020, Accuweather 2020). All the data taken for this
study were from January 9, 2020 to March 25, 2020.

2.3. Statistical Analysis

Natural Log-transformation was performed on total number of daily COVID-19
confirmed cases of 57 countries. Polynomial regression as a quadratic term was done to find
out the relationship between daily average temperature and log transformed daily cases of
COVID-19 as we observed a non-linear relationship during initial data exploration (Bradley
and Srivastava, 1979).However, the linear regression was used to find out the effect of daily
average relative humidity on COVID-19 cases (Matthews, 2014).For a certain country, the
natural log of the average number of cases per day was used, and the linear regression was
done to find out the effect of temperature and relative humidity on the occurrence of COVID-
19 cases.

3. Results

The data on average temperature, average relative humidity, and log transformed
confirmed cases per day of COVID-19 in 57 countries are listed in Supplementary Table.
Here, we included 444,405 cases from 57 countries (January 9, 2020 to March 25, 2020).The
daily average lowest and highest temperature was respectively -5^{0}C in Finland and 29.9^{0}C in
Thailand. Percent relative humidity varied from 30 in Saudi Arabia to 78.8 in Indonesia. Data
suggest relative humidity was fluctuating with large deviation in every country. Except China
(January, February, March), Italy (February, March), Iran (February, March), South Korea
(February, March), France (February, March) and Japan (February, March) all other countries
had COVID-19 cases in March. In China, the early outbreak was in January when average
cases per day were lower than in February and March. Similarly, Italy, Iran, South Korea,
France and Japan had lower average cases per day in February compared to March. Among
the countries, Italy had the lowest and highest average cases per day respectively in February
(average monthly temperature 7.5^{0}C) and March (average monthly temperature 9.4^{0}C).

In polynomial regression as a quadratic term,no correlation (R² = 0.031) between the
daily average temperature and the number of cases of COVID-19 was found. The analysis
showed that lgN of cases increased as the average temperature rose and started to decline
moderately when temperature reached the peak (Fig. 1).The analysis showed that 98.8%
proportionate cases occurred between -0.5^{0}C to 30^{0}Cand, more restrictively, 93.5% cases
occurred between 1^{0}C to 16^{0}C (Fig. 1). The average relative humidity was also not correlated

($R^2 = 0.0272$) with the occurrence of COVID-19 in linear regression (Fig. 2). However, most
of the low temperature countries had relatively higher average number of cases per day than
high temperature countries (Fig. 3). There were also exceptions. For country-wise analysis,
average monthly observations were used for each country. In linear regression country-wise
analysis revealed no correlation ($R^2 = 0.0602$) between temperature and occurrence of
COVID-19 cases per day (Fig. 3).

4. Discussion

This is the first study that consider the pandemic situation (57 countries including
high and low temperature countries)to draw conclusions on possible association of
temperature and relative humidity with COVID-19 cases. Therefore, the findings of this study
demand importance. No correlation between the daily average temperature and the
occurrence of cases of COVID-19 was observed. Earlier studies found correlation between
temperature and the occurrence of COVID-19 (Wang et al., 2020d, Wang et al., 2020e, Zhu
et al 2020).Wang et al., (2020d) showed that most COVID-19 cases in China occurred
between 0^0C to 10^0C. Another study analyzed data (January 20, to February 4, 2020) of
COVID-19 cases and revealed that the majority of cases occurred with temperature around
10^0C (Wang et al., 2020e).But all these studies obtained data from the early stage of the
outbreak, at best till February when most of the affected countries were in winter. In our
study, temperature varied from -24^0C to 30.5^0C in 57 countries. The huge variation of
temperature in analysis found no correlation with COVID-19 cases. We found 93.5%
COVID-19 cases were between 1^0C to 16^0C as most of cases were from China, Italy, USA,
Iran, Germany, France and Spain where the duration of outbreak is prolong (Supplementary
Table).

In country-wise analysis, no correlation was observed between average daily
temperature and average per day cases of COVID-19. However, in general, countries having
high temperature experienced relatively lower occurrence of COVID-19 (Fig. 3) with
exceptions. This is may be due to high temperature countries are at early stage of outbreak.
With the progress of outbreak the number of cases may increase exponentially if proper
control measures are not taken. The transmission was exponential in some developed low-
temperature countries like USA, Italy, Spain, Germany, and France where large scale
community transmission was reported (WHO, 2020a). On the other hand, similar countries
like South Korea, Japan took steps during early stage of the outbreak being quite successful
in limiting virus transmission (Financial Times, 2020). The average daily temperature in
Hong Kong (21.7°C) and Brazil (23.1°C) was quite similar, but Brazil had relatively higher
number of daily average cases (Supplementary Table). These findings suggest all courtiers
for applying proper control measures. However, these exceptions can be interpreted taking
into consideration that virus transmission in an outbreak largely depends on the interventions
taken by each particular government. In addition, duration of outbreak, population density,
other metrological variables, level of awareness in people, and medical facilities are not same
for all countries. Among so many factors, transmission of COVID-19 virus from
asymptomatic is carrier very important. A study has been carried out on the passengers of
Diamond Princess cruise ship, where 50.47% COVID-19 patients were asymptomatic carrier

(Mizumoto et al., 2020).Additional study is required to consider many more variables. A variable relative humidity was considered in our study. Though Wang et al., (2020d) found a correlation between relative humidity and the occurrence of COVID-19 cases, we could not confirm this association.A recent study in China Jakarta, Indonesiaalso showed no effect of relative humidity on COVID-19 occurrence (Tosepu et al., 2020). This observed variation in results may be due to a variation in the investigated time periods and data size in the different studies.In addition we observed inconsistent relative humidity in most of the countries in our study.

The stability SARS virus was studied in laboratory, at 22-25°C SARS-CoV is stable up to five days on plastic surface while rapidly losing activity at 38°C or more (Chan et al., 2011). SARS-CoV-2 survive in air (3 hours), on plastic and stainless steel surfaces (up to 72 hours) thus contact transmission is facilitated (Van Doremalen et al., 2020), but investigations of this type associated temperature and humidity are scarce for SARS-CoV-2. Preliminary evidences indicate that high temperature affect virus transmissibility through droplet and aerosol (Lowen et al., 2008), but contact transmission cannot be ignored (Van Doremalen et al, 2020). Therefore, "high temperature" itself is not enough to combat the pandemic. The primary epicenter in Wuhan, Chinaalready has proved that the virus is controllable in a low-temperature region (WHO, 2020d). We suggest that all countries should follow control measures similar to those adopted in China, taking into account that temperature increments will not be enough.

5. Conclusions

To the best of our knowledge, this is the first study describing temperature and relative humidity has no major effect to describe on the current COVID-19 pandemic. More studies needed to which variables are contributing most. Current study suggests all countries not to depend on summer rather to implement possible all control measures of COVID-19.

Acknowledgement

The author would like to acknowledge Dr. AMM Taufiqur Rahman, Adunik Sadar Hospital, Naogaon, Bangladesh for his critical reading and comments on this manuscript.

Conflict of interest

All the authors report no conflicts of interest in this paper.

Funding

No finding.

Author's contribution

MTR and MAS designed the study. MAS, MSI and MEH collected data. MAS, MEH and MTI analyzed and interpreted the data. MAS, MSI, AT, MTI and MTR drafted the manuscript. MAS, MTI, AT and MTR critically reviewed and updated the manuscript to its final version. The final version of the manuscript was approved by all authors.

Figure and Table legends

Figure 1. The relationship between average temperature and COVID-19 cases (natural log transformed) in 57 countries

Figure 2.The relationship between average relative humidity and COVID-19 cases (natural log transformed) in 57 countries

Figure 3. Country-wise log transformed average cases of COVID-19 per day v.s. average temperature in 57 countries

Supplementary Table legend:

Supplementary Table: Average temperature, relative humidity and COVID-19 cases in 57 countries

Figure and Supplementary Table legend:

Figure 1. The relationship between average temperature and COVID-19 cases (natural log transformed) in 57 countries

Figure 2.The relationship between average relative humidity and COVID-19 cases (natural log transformed) in 57 countries

Figure 3. Country-wise log transformed average cases of COVID-19 per day v.s. average temperature in 57 countries.

Supplementary Table legend:

Supplementary Table: Average temperature, relative humidity and COVID-19 cases in 57 countries

Figure 1. The relationship between average temperature and COVID-19 cases (natural log

transformed) in 57 countries.

240   Figure 2.The relationship between average relative humidity and COVID-19 cases (natural

241   log transformed) in 57 countries.

Figure 3. Country-wise log transformed average cases of COVID-19 per day v.s. average

temperature in 57 countries.

251   Supplementary Table 1. Average temperature, relative humidity and COVID-19 cases in 57 countries

Country	Month	Number of Reports	Temperature				Relative Humidity				COVID-19 Cases			
			Minimum	Maximum	Average	Standard Deviation	Minimum	Maximum	Average	Standard Deviation	Minimum	Maximum	Average	Standard Deviation
Argentina	March	12	19.5	24.5	22.1	1.9	60.5	84.0	72.5	6.9	2.4	4.8	3.3	0.9
Australia	March	19	12.5	21.5	16.4	2.3	58.5	89.5	69.0	8.2	2.5	6.3	4.2	1.3
Austria	March	21	.5	15.5	7.7	4.4	46.0	78.5	60.0	9.0	2.5	6.8	4.9	1.4
Belgium	March	22	3.5	13.0	7.3	2.3	38.0	93.5	71.7	16.3	2.3	6.5	4.9	1.2
Brazil	March	14	20.5	25.5	23.1	2.0	64.0	80.5	72.9	4.9	2.9	5.9	4.8	1.1
Canada	March	18	-9.5	2.5	-3.4	3.5	48.5	88.0	65.8	11.7	2.3	6.6	4.4	1.4
Chile	March	13	15.5	22.0	19.5	1.7	40.0	70.5	52.3	8.2	2.4	5.4	4.1	1.0
China	Feb	373	-24.0	18.0	6.3	6.4	15.0	96.5	71.2	14.9	2.3	9.5	3.7	1.3
China	Jan	143	-12.0	14.5	5.0	3.4	37.5	96.0	70.0	14.2	2.3	6.8	3.3	0.9

China	March	25	7.5	20.0	13.9	3.7	44.5	85.5	67.3	9.9	2.5	6.4	4.2	0.9
Colombia	March	10	15.0	17.5	16.6	0.8	72.5	83.5	76.9	4.3	2.4	4.6	3.5	0.8
Croatia	March	10	.5	12.5	7.1	5.1	42.5	86.5	63.3	14.8	2.4	4.4	3.5	0.7
Czechia	March	15	-1.0	11.0	5.3	4.1	43.5	90.5	64.2	14.2	3.1	5.6	4.5	0.8
Denmark	March	19	1.5	7.0	4.7	1.6	54.5	93.0	73.6	12.5	2.5	5.5	4.2	0.8
D Republic	March	8	24.0	25.0	24.6	0.3	61.5	76.0	70.9	4.6	2.3	4.5	3.6	0.8
Ecuador	March	10	10.5	16.5	13.4	1.9	62.5	77.5	70.5	5.0	3.0	5.5	4.6	0.7
Egypt	March	10	16.5	27.5	20.9	3.7	34.0	55.5	43.1	7.6	2.6	4.1	3.6	0.4
Estonia	March	11	-2.5	5.5	0.9	2.5	53.0	89.5	65.7	11.9	2.8	4.1	3.3	0.5
Finland	March	14	-12.0	.5	-5.0	3.5	55.0	89.5	70.8	10.6	2.7	4.6	3.9	0.7
France	Feb	1	10.5	10.5	10.5		88.5	88.5	88.5		2.9	2.9	2.9	
France	March	25	5.5	14.5	9.1	2.8	36.0	89.5	74.2	14.8	3.0	8.3	6.2	1.4
Germany	March	25	3.0	12.5	6.5	2.8	34.0	85.0	63.1	13.8	3.3	8.4	6.4	1.7
Greece	March	19	7.5	16.5	11.0	2.5	36.0	73.0	57.0	11.8	2.3	4.6	3.5	0.7

Hong Kong	March	10	19.5	24.5	21.7	1.8	68.0	87.0	78.1	5.7	2.3	3.9	3.2	0.5
Iceland	March	16	-3.5	2.5	-0.2	1.9	47.0	86.5	72.6	10.0	2.4	4.6	3.5	0.7
India	March	15	21.5	30.5	27.0	2.7	63.0	73.0	70.1	3.0	2.5	4.8	3.4	0.8
Indonesia	March	14	26.0	29.0	28.0	0.9	71.0	85.5	78.8	3.5	2.6	4.7	3.8	0.7
Iran	Feb	16	4.5	15.0	9.3	2.9	37.5	91.0	63.4	14.3	2.3	4.2	3.0	0.5
Iran	March	32	7.0	17.0	11.9	2.7	19.5	85.5	41.2	17.0	2.4	7.7	6.1	1.6
Iraq	March	13	12.5	21.5	18.0	2.5	33.5	76.5	56.1	12.8	2.3	3.9	2.9	0.5
Ireland	March	16	3.5	10.5	6.4	2.4	68.0	86.0	77.8	5.7	2.5	5.5	4.2	1.0
Israel	March	18	5.5	18.5	11.3	3.5	50.0	83.5	71.6	8.5	2.5	6.2	4.2	1.2
Italy	Feb	4	6.5	9.5	7.5	1.4	63.0	85.0	74.1	9.0	2.3	3.4	2.8	0.6
Italy	March	25	4.5	15.0	9.4	2.9	40.0	80.5	62.6	11.1	5.8	8.8	7.7	0.9
Japan	Feb	14	3.0	12.5	8.7	2.7	37.0	91.5	59.6	16.7	2.3	4.6	3.6	0.8
Japan	March	24	6.5	16.5	10.6	2.7	32.5	87.5	60.9	15.2	2.7	4.7	3.7	0.5
Luxembourg	March	13	3.5	12.0	6.8	3.3	32.0	81.5	63.1	18.2	2.5	5.5	4.2	1.1

Malaysia	March	19	27.5	30.5	29.4	1.0	62.0	87.0	78.1	6.9	2.3	5.4	4.1	1.0
Mexico	March	12	17.5	21.5	19.6	1.2	21.5	55.0	40.6	10.4	2.4	4.2	3.3	0.6
Netherlands	March	22	5.0	10.5	7.5	1.7	42.5	91.5	73.2	14.7	2.6	6.7	5.2	1.1
Norway	March	22	-2.0	7.5	3.0	2.5	38.5	93.5	71.5	17.1	3.0	5.5	4.7	0.9
Pakistan	March	11	16.5	20.5	18.6	1.4	62.5	77.5	68.3	5.3	3.1	5.4	4.4	0.6
Panama	March	11	26.5	28.5	28.1	0.6	64.5	75.0	67.6	2.9	2.5	4.6	3.4	0.7
Peru	March	11	13.0	16.0	14.4	1.0	48.0	92.5	77.4	11.8	2.3	4.5	3.5	0.7
Philippines	March	15	27.5	30.0	28.9	0.7	52.5	71.0	63.1	5.5	2.3	4.7	3.6	0.9
Poland	March	15	-2.0	10.5	4.2	4.6	41.0	77.5	60.5	13.6	2.8	5.0	4.0	0.8
Portugal	March	17	11.5	16.5	14.2	1.4	59.0	86.0	74.0	7.3	1.9	6.5	4.4	1.5
Qatar	March	11	20.0	25.5	23.2	1.8	42.0	72.5	60.4	9.0	2.3	5.5	3.2	1.0
Romania	March	14	1.5	14.5	8.2	4.3	37.5	87.5	59.9	15.9	2.3	5.4	3.8	0.9
Russia	March	12	-3.0	6.5	1.3	3.3	36.0	75.0	54.3	11.8	2.6	5.1	3.7	0.7
Saudi Arabia	March	12	23.0	29.5	25.5	2.1	20.0	51.5	30.0	10.1	2.7	5.3	4.0	0.8

Serbia	March	11	-.5	14.0	7.5	5.3	45.0	91.0	60.1	13.1	2.3	4.4	3.3	0.7
Singapore	March	17	26.5	30.0	28.7	0.9	69.0	84.0	76.0	4.8	2.3	4.3	3.2	0.6
Slovenia	March	14	.5	13.5	6.9	4.4	41.0	87.0	60.6	14.4	2.4	4.0	3.5	0.4
South Africa	March	11	18.5	22.5	20.9	1.2	51.0	81.0	65.3	8.3	2.4	5.0	3.7	1.0
South Korea	Feb	6	5.0	8.0	6.6	1.4	40.0	52.5	45.8	5.8	2.9	5.3	4.3	1.0
South Korea	March	25	4.5	14.5	8.5	2.8	29.5	80.5	49.9	13.2	3.6	6.7	5.1	0.8
Spain	March	25	6.5	16.5	11.2	2.7	44.5	82.0	63.3	11.5	3.3	8.9	6.5	1.9
Swedan	March	21	-1.0	6.5	2.7	2.2	33.5	94.0	72.3	16.4	2.6	5.5	4.6	0.7
Switzerland	March	24	1.0	11.0	6.0	2.8	51.0	88.5	71.3	9.9	2.7	7.2	5.4	1.4
Thailand	March	12	26.5	31.0	29.9	1.4	70.0	87.5	75.0	4.9	2.4	5.2	4.0	0.9
Turkey	March	11	-.5	11.5	6.0	4.7	45.5	80.5	59.4	9.4	2.5	6.3	4.9	1.2
UK	March	23	5.0	12.5	8.0	1.9	44.5	92.0	73.0	14.1	2.4	7.3	5.2	1.4
USA	March	24	3.0	16.0	9.0	3.4	31.5	83.0	52.2	15.9	3.0	9.5	6.5	2.1

References:

Accuweather, 2020. https://www.accuweather.com/en (accessed 26 March, 2020).

Arab-Mazar, Z., Sah, R., Rabaan, A.A., Dhama, K., Rodríguez-Morales, A.J., 2020. Mapping the incidence of the COVID-19 hotspot in Iran–Implications for Travellers. Travel. Med. Infect. Dis. 101630. https://doi.org/10.1016/j.tmaid.2020.101630.

Bradley, R.A., Srivastava, S.S., 1979. Correlation in polynomial regression. Am. Stat. 33, 11-14.https://doi.org/10.1080/00031305.1979.10482644.

Chan, K.H., Peiris, J.S., Lam, S.Y., Poon, L.L., Yuen, K.Y., Seto, W.H., 2011. The effects of temperature and relative humidity on the viability of the SARS coronavirus. Adv. Virol. 2011, 734690. https://doi.org/10.1155/2011/734690.

Chen, N., Zhou, M., Dong, X., Qu J., Gong, F., Han, Y., Qiu Y., Wang, J., Liu, Y., Wei, Y., Yu, T., 2020. Epidemiological and clinical characteristics of 99 cases of 2019 novel coronavirus pneumonia in Wuhan, China: a descriptive study. The Lancet. 395, 507-513. https://doi.org/10.1016/S0140-6736(20)30211-7.

Dalziel, B.D., Kissler, S., Gog, J.R., Viboud, C., Bjørnstad, O.N., Metcalf, C.J., Grenfell, B.T., 2018. Urbanization and humidity shape the intensity of influenza epidemics in US cities. Science. 362, 75-9. https://doi.org/10.1126/science.aat6030.

Financial Times, 2020. Containing coronavirus: lessons from Asia. https://www.ft.com/content/e015e096-6532-11ea-a6cd-df28cc3c6a68 (Accessed7April, 2020).

Gorbalenya, A.E., 2020. Severe acute respiratory syndrome-related coronavirus–The species and its viruses, a statement of the Coronavirus Study Group. BioRxiv. https://doi.org/10.1101/2020.02.07.937862.

Hemmes, J.H., Winkler, K., Kool, S.M., 1960. Virus survival as a seasonal factor in influenza and poliomyelitis. Nature. 188, 430-431. https://doi.org/10.1038/188430a0.

Holshue, M.L., DeBolt, C., Lindquist, S., Lofy, K.H., Wiesman, J., Bruce, H., Spitters, C., Ericson, K., Wilkerson, S., Tural, A., Diaz, G., 2020. First case of 2019 novel coronavirus in the United States. N. Engl. J. Med. 382, 929-936. https://doi.org/10.1056/NEJMoa2001191.

Huang, C., Wang, Y., Li, X., Ren, L., Zhao, J., Hu, Y., Zhang, L., Fan, G., Xu, J., Gu, X., Cheng, Z., 2020. Clinical features of patients infected with 2019 novel coronavirus in Wuhan, China. The Lancet. 395, 497-506. https://doi.org/10.1016/S0140-6736(20)30183-5.

KFF, 2020. Global Health Policy. COVID-19 Coronavirus Tracker. https://www.kff.org/global-health-policy/fact-sheet/coronavirus-tracker/ (accessed 26 March, 2020).

Lowen, A.C., Steel, J., Mubareka, S., Palese, P., 2008. High temperature (30 C) blocks aerosol but not contact transmission of influenza virus. J.Virol. 82, 5650-5652. https://doi.org/10.1128/JVI.00325-08.

Lu, H., Stratton, C.W., Tang, Y.W., 2020. Outbreak of Pneumonia of Unknown Etiology in
Wuhan China: the Mystery and the Miracle. J.Med.Virol. 92, 401–402.
https://doi.org/10.1002/jmv.25678.

Matthews, D.E., 2014. Multiple linear regression. Wiley StatsRef: Statistics Reference
Online. https://doi.org/10.1002/9781118445112.stat05766.

Mizumoto, K., Kagaya, K., Zarebski, A., Chowell, G., 2020. Estimating the asymptomatic
proportion of coronavirus disease 2019 (COVID-19) cases on board the Diamond Princess
Cruise ship, Yokohama, Japan, 2020. Eurosurveillance. 25, 2000180.
https://doi.org/10.2807/1560-7917.ES.2020.25.10.2000180.

Pal, M., Berhanu, G., Desalegn, C., Kandi, V., 2020. Severe Acute Respiratory Syndrome
Coronavirus-2 (SARS-CoV-2): An Update. Cureus. 26, e7423.
https://doi.org/10.7759/cureus.7423.

Peng, X., Xu, X., Li, Y., Cheng, L., Zhou, X., Ren, B., 2020. Transmission routes of 2019-
nCoV and controls in dental practice. Int. J. Oral. Sci. 12, 1-6.
https://doi.org/10.1038/s41368-020-0075-9.

Rodríguez-Morales, A.J., MacGregor, K., Kanagarajah, S., Patel, D., Schlagenhauf, P., 2020.
Going global-Travel and the 2019 novel coronavirus. Travel. Med. Infect. Dis. 33,101578.
https://doi.org/10.1016/j.tmaid.2020.101578.

Sobsey, M.D., Meschke, J.S., 2003. Virus survival in the environment with special attention
to survival in sewage droplets and other environmental media of fecal or respiratory origin.
Report for the World Health Organization, Geneva, Switzerland.

The Weather Channel, 2020. Monthly Weather, https://weather.com/ (accessed 26 March,
2020).

Timeanddate, 2020. World Temperatures — Weather around the World,
https://www.timeanddate.com/weather/ (accessed 26 March, 2020).

To, K.K., Tsang, O.T., Chik-Yan Yip, C., Chan, K.H., Wu, T.C., Chan, J., Leung, W.S.,
Chik, T.S., Choi, C.Y., Kandamby, D.H., Lung, D.C., 2020. Consistent detection of 2019
novel coronavirus in saliva. Clin. Infect. Dis. https://doi.org/10.1093/cid/ciaa149.

Tosepu, R., Gunawan J., Effendy, D.S., Ahmad L.O.A.I., Lestari, H., Bahar, H., Asfian, P.,
2020. Correlation between weather and Covid-19 pandemic in Jakarta, Indonesia. Sci. Total.
Environ. https://doi.org/10.1016/j.scitotenv.2020.138436.

Van Doremalen, N., Bushmaker, T., Morris, D., Holbrook, M., Gamble, A., Williamson, B.,
Tamin, A., Harcourt, J., Thornburg, N., Gerber, S., Lloyd-Smith, J., 2020. Aerosol and
surface stability of HCoV-19 (SARS-CoV-2) compared to SARS-CoV-1.N. Engl. J. Med.
https://doi.org/10.1056/NEJMc2004973.

Wang, C., Horby, P.W., Hayden, F.G., Gao, G.F., 2020a. A novel coronavirus outbreak of
global health concern. The Lancet. 395, 470-473. https://doi.org/10.1016/S0140-
6736(20)30185-9.

Wang, D., Hu, B., Hu, C., Zhu, F., Liu, X., Zhang, J., Wang, B., Xiang, H., Cheng, Z., Xiong,
Y., Zhao, Y., 2020b. Clinical characteristics of 138 hospitalized patients with 2019 novel
coronavirus–infected pneumonia in Wuhan, China. JAMA. 323, 1061-1069.
https://doi.org/10.1001/jama.2020.1585.

Wang, J., Tang, K., Feng, K., Lv, W., 2020d. High Temperature and High Humidity Reduce
the Transmission of COVID-19. http://dx.doi.org/10.2139/ssrn.3551767.

Wang, M., Jiang, A., Gong, L., Luo, L., Guo, W., Li, C., Zheng, J., Li, C., Yang, B., Zeng, J.,
Chen, Y., 2020e. Temperature significantly change COVID-19 Transmission in 429 cities.
MedRxiv. https://doi.org/10.1101/2020.02.22.20025791.

Wang, Y., Wang, Y., Chen, Y., Qin, Q., 2020c. Unique epidemiological and clinical features
of the emerging 2019 novel coronavirus pneumonia (COVID‐19) implicate special control
measures. J.Med.Virol. 2020, 1-9. https://doi.org/10.1002/jmv.25748.

Wolkoff, P., 2018. Indoor air humidity, air quality, and health–An overview. Int.J.Hyg.
Environ. Health. 221, 376-390. https://doi.org/10.1016/j.ijheh.2018.01.015.

World Health Organization (WHO), 2020a. Novel Coronavirus (2019‐nCoV) situation
report-79. https://www.who.int/docs/default-source/coronaviruse/situation-reports/20200408-
sitrep-79-covid-19.pdf?sfvrsn=4796b143_6 (accessed9April, 2020).

World Health Organization (WHO), 2020b. Coronavirus disease (COVID-2019) press
briefings: WHO Emergencies Press Conference on coronavirus disease outbreak – 11 March
2020. https://www.who.int/emergencies/diseases/novel-coronavirus-2019/media-
resources/press-briefings (accessed 7 April, 2020).

World Health Organization (WHO), 2020c. Coronavirus disease (COVID-19) Pandemic.
https://www.who.int/emergencies/diseases/novel-coronavirus-2019 (accessed 8 April, 2020).

World Health Organization (WHO), 2020d. Report of the WHO-China Joint Mission on
Coronavirus Disease 2019 (COVID-19). https://www.who.int/docs/default-
source/coronaviruse/who-china-joint-mission-on-covid-19-final-report.pdf (accessed 7 April,
2020).

Worldometer, 2020. COVID-19 coronavirus outbreak.
https://www.worldometers.info/coronavirus/ (accessed7April, 2020).

Zhu, Y., and Xie, J., 2020. Association between ambient temperature and COVID-19
infection in 122 cities from China. Sci. Total. Environ. 138201.
https://doi.org/10.1016/j.scitotenv.2020.138201.