

Social determinants of COVID-19 mortality at the county level

Rebecca K. Fielding-Miller, PhD^{1*}

Maria E. Sundaram, PhD²

Kimberly Brouwer, PhD¹

¹ University of California, San Diego, School of Medicine, San Diego, CA

² Emory University, Rollins School of Public Health, Atlanta, GA

*Corresponding author:

email: rfieldingmiller@health.ucsd.edu

Abstract

The United States is currently the global epicenter of the COVID-19 pandemic. Emerging data suggests that social determinants of health may be key drivers of the epidemic, and that minorities, migrants, and essential workers may bear a disproportionate degree of risk. We used publicly accessible datasets to build a series of spatial autoregressive models assessing county level associations between COVID-19 mortality and (1) Percentage of Non-English speaking households, (2) percentage of individuals engaged in hired farm work, (3) percentage of uninsured individuals under the age of 65, and (3) percentage of individuals living at or below the poverty line. Across all counties (n=2940), counties with more farmworkers, more residents living in poverty, higher density, and more residents over the age of 65 had significantly higher levels of mortality. In urban counties (n=114), only county density was significantly associated with mortality. In non-urban counties (n=2826), counties with more non-English speaking households and more farm workers had significantly higher levels of mortality, as did counties with higher levels of poverty and more residents over the age of 65. More uninsured residents was significantly associated with decreased reported COVID-19 mortality. Individuals who do not speak English, individuals engaged in farm work, and individuals living in poverty may be at heightened risk for COVID-19 mortality in non-urban counties. Mortality among the uninsured may be being systematically undercounted in county and national level surveillance.

1 **Introduction**

2 A novel coronavirus responsible for COVID-19 respiratory disease is causing a global
3 pandemic which has already resulted in nearly 5 million cases and over 300,000 deaths
4 since early January(1). The United States currently has more cases than any other
5 nation in the world, with over 2 million cases and 113,000 deaths as of June 11,
6 2020(1). Preliminary data indicates that existing health inequities in the United States
7 are likely linked to COVID-19 morbidity and mortality(2).

8
9 Both infectious and non-communicable disease tends to impact marginalized
10 populations at disproportionate rates. While demographically disaggregated data is not
11 currently available at the national level, data from county and state level entities suggest
12 that COVID-19 may follow similar patterns. In the State of California, Latinos make up
13 approximately 39% of the total population but represent just over 53% of total cases(3).
14 Similarly, in New York City, Black/African American and Hispanic residents have
15 significantly higher rates of COVID-19 illness and mortality than white residents, with a
16 nearly doubled risk of mortality for Black/African American residents compared to white
17 residents(4). While more granular data are not yet available to assess which risk factors
18 may be leading to these disparities in morbidity and mortality, journalistic reporting early
19 analyses suggest that language barriers, poor working conditions among essential
20 workers – who are more likely to be immigrants and/or racial/ethnic minorities(5) - and
21 concerns about immigration status may be creating particular risk among racial and
22 ethnic immigrants across the United States(6, 7).

23

24 We sought to assess the associations between COVID-19 mortality and immigrant and
25 farm worker population at the county level. We hypothesized that counties with more
26 immigrants and farm workers would report higher COVID-19 mortality, adjusting for
27 poverty, insurance rates, population age, and density at the county level.

28

29 **Methods**

30 We built a series of spatial autoregressive models to assess county-level associations
31 between COVID-19 mortality and: (1) Percentage of Non-English speaking households
32 (defined as households in which no one 14 years or older reports speaking English at
33 least “very well”) and (2) percentage of individuals engaged in hired farm work(8) in the
34 county as of 2018. To account for potential confounders, we adjusted our analyses for
35 the percentage of uninsured individuals under the age of 65, percentage of individuals
36 living at or below the poverty line, percentage of residents age 65 or older, and county
37 density, measured as number of residents per square mile.

38

39 COVID-19 mortality data was sourced from county public health agencies, aggregated
40 and made publicly available by the New York Times(9). The proportion of households
41 with limited English speaking ability was drawn from the American Community Survey’s
42 (ACS) 2014 5-year estimate, percentages of individuals living below poverty, and
43 percentage of residents over the age of 65 were from 2017 ACS data. The percentage
44 of farmworkers was taken from the US Bureau of Economic Analysis. Percent uninsured
45 was based on the US Census Small Area Health Insurance Estimates (SAHIE)

46 program's 2018 estimates. Density was measured as number of individuals per square
47 mile, based on US census data.

48
49 In addition to hypothesized predictors and potential confounders, we adjusted our
50 models to account for the stage of the local epidemic by including a variable for the
51 number of "high risk" days a county had experienced. We calculated this number by
52 subtracting the total number of days in which a county had been under a shelter in place
53 order from the total number of days since the first reported case.

54
55 Counties with 1000 residents or more per square mile were coded as urban, counties
56 with less than 1000 residents per square mile were coded as non-urban. While there are
57 many ways to classify counties, we chose to use 1000 people per square mile for two
58 reasons. First, the US census uses this cutpoint to designate census blocks as urban
59 vs. non-urban. Second we felt that doing so allowed us to more clearly delineate major
60 metropolitan areas and their associated resources and public health infrastructures from
61 neighboring suburban or exurban counties.

62
63 We first built a series of simple linear regression models to assess the bivariate
64 association between number of deaths within a county and our hypothesized predictors,
65 adjusting for days since 1st case and SIP order. We then constructed a spatial
66 contiguity matrix, and checked the assumption that residuals were distributed spatially
67 using a Moran's I test.

68

69 We next built three separate spatial autoregressive models to assess the association
 70 between number of deaths and our hypothesized social determinants, adjusting for
 71 potential confounders, and fitted the model with a spatial lag of the dependent variable
 72 based on our contiguity matrix. Our first model assessed relationships across all
 73 counties. We then stratified our analyses to measure the association between mortality
 74 and our hypothesized predictors in urban and non-urban counties.

75

76 Results

77 This analysis encompassed 2,941 counties across all 50 states. As of June 11, 2020,
 78 the number of deaths reported in the NY Times aggregated dataset ranged from 0 to
 79 21,436 per county, with a median of 1 an interquartile range (IQR) of 0-8¹. We classified
 80 115 counties as urban and 2826 counties as non-urban. Deaths in urban counties
 81 ranged from 0 – 21436, with a median of 210 and an IQR of 66-62. Deaths in rural
 82 counties ranged from 0-844 with a median of 1 and an IQR of 0-6 (table 1).

Table 1: Primary predictor and covariates of interest across all counties and stratified by urban and non-urban

	All counties (n=2941)		non-urban counties (n=2826)		urban counties (n=115)	
	<i>median</i>	<i>IQR</i>	<i>median</i>	<i>IQR</i>	<i>median</i>	<i>IQR</i>
<i>deaths</i>	1	0-8	1	0-6	210	66.0-662.0
<i>% Farm workers</i>	2.2	0.9 – 4.6	2.4	1.0 – 4.8	0.1	0.0 – 0.1
<i>% Non-English speakers</i>	5.0	2.9 – 10.2	4.8	2.8 – 9.4	19.0	11.4 – 29.8
<i>% Residents uninsured</i>	10.5	7.4 – 14.5	10.6	7.4 – 14.6	8.3	5.9 – 12.6
<i>% Residents in poverty</i>	15.1	11.4 – 9.5	15.2	11.5 – 19.7	13.0	8.9 – 16.7
<i>Residents per square mile</i>	46.9	19.9 – 115.0	44.7	18.8-101.5	1754.9	1313.4-2715.3
<i>% Residents Over 65</i>	16.3	11.0 – 14.5	16.5	14.1-19.0	12.5	11.0-14.5

¹ Within this dataset, the 5 boroughs/counties of New York are treated as a single entity. We have done the same in these analyses, assigning all 5 counties the values associated with New York County.

83 The Moran's I test was statistically significant at $p < 0.01$ in each simple (non-spatial)
84 regression, with the exception of associations between density and mortality (Moran's I
85 $p = 0.22$) and non-English speakers and mortality (Moran's I p -value = 0.31) in urban
86 counties, indicating a significant spatial pattern to associations between our
87 hypothesized predictors and mortality.

88

89 In our fully adjusted model of all counties, the percentage of farm workers in a county,
90 poverty, population density, and the percentage of residents over the age of 65 were all
91 significantly associated with higher levels of COVID-19 mortality. Each additional
92 percentage of people living in poverty was associated with 3.49 additional deaths within
93 that county ($p < 0.01$), and an additional 0.63 deaths in neighboring counties via a
94 'spillover' effect ($p < 0.05$). Overall, each additional percentage increase of individuals
95 living in poverty within a county was associated with 6.24 additional COVID-19 deaths
96 ($p = 0.001$). Each additional person per square mile was associated with 0.33 additional
97 deaths within a county and 0.06 indirect deaths, for a total of 0.39 additional deaths (p
98 < 0.001). Across all counties, each percentage increase in residents over the age of 65
99 was associated with 4.20 additional deaths within that county ($p = 0.01$) and an
100 additional 0.75 deaths in neighboring counties ($p = 0.023$). Across all counties in the
101 United States, each additional percentage increase in farmworkers was associated with
102 5.29 additional deaths within a county ($p = 0.003$) and 0.95 additional deaths in
103 neighboring counties ($p = 0.02$)

104

105 In urban counties (n=114), only population density was significantly associated with
106 higher mortality. In these counties, each additional person per square mile was
107 associated with 0.35 additional deaths within that county ($p < 0.001$).

108
109 In non-urban counties, all of our hypothesized social determinants were significantly
110 associated with higher levels of mortality. Each increase in the percentage of
111 farmworkers residing in a county was associated with 0.74 additional deaths ($p = 0.02$).
112 Each additional percentage of non-English speaking households was associated with
113 0.62 additional deaths ($p < 0.001$), 0.49 deaths within a given county, and 0.14
114 attributable to 'spillover' effects to neighboring counties ($p = 0.002$). As in the all county
115 and urban models, density, poverty, and the percentage of residents over the age of 65
116 were all significantly associated with higher mortality. Contrary to our initial hypotheses,
117 the percentage of uninsured individuals was associated with lower reported COVID19
118 mortality. In rural areas, each increase in the percentage of uninsured individuals was
119 associated with a direct effect of 0.69 fewer deaths within the county ($p < 0.001$) and
120 0.19 fewer deaths in neighboring counties ($p = 0.007$).

121

122

Table 2: Full spatial regression models for all counties and stratified by urban/rural

	All counties (n=2940)				non-urban counties (n=2826)				urban counties (n=114)			
	<i>b direct</i>	<i>b indirect</i>	<i>b total</i>	<i>p-value</i>	<i>b direct</i>	<i>b indirect</i>	<i>b total</i>	<i>p-value</i>	<i>b direct</i>	<i>b indirect</i>	<i>b total</i>	<i>p-value</i>
% <i>Farm workers</i>	5.29	0.95	6.24	0.003	0.57	0.16	0.74	0.02	1956.83	130.54	2087.37	0.20
% <i>Non-English speakers</i>	0.13	0.03	0.15	0.83	0.49	0.14	0.62	<0.001	16.04	1.07	17.11	0.21
% <i>Residents uninsured</i>	-1.64	-0.30	-1.93	0.25	-0.69	-0.19	-0.89	<0.001	-50.56	-3.37	-53.93	0.27
% <i>Residents in poverty</i>	3.50	0.63	4.12	0.001	0.32	0.09	0.40	0.01	59.86	3.99	63.85	0.10
<i>Residents per square mile</i>	0.33	0.06	0.39	<0.001	0.17	0.05	0.21	<0.001	0.35	0.02	0.38	<0.001
% <i>Residents Over 65</i>	4.20	0.75	4.95	0.01	0.61	0.17	0.78	0.002	45.82	3.06	48.87	0.50

123 **Discussion**

124 Although we cannot draw conclusions about individual risk profiles, our findings do
125 suggest that that farm work may create unique risk factors and that farmworkers may
126 require additional protections, such as personal protective equipment and/or targeted
127 outreach. Immigrants provide approximately 75% of all farm labor in the United
128 States(8). Among those engaged in crop work specifically, nearly three quarters are
129 migrants and approximately half are undocumented(8). Undocumented status may
130 impede an individual's willingness or ability to seek healthcare, or their ability to request
131 additional protections from an employer if they worry doing so could result in their own
132 deportation or that of a family member(10). Farm labor is considered essential work, but
133 there are reports of inadequate personal protective equipment and inadequate social
134 distancing guidelines or enforcement(6). There have also been several high profile
135 outbreaks at meat processing facilities across the country. While the farmworker
136 classification that we used in this analysis does not include individuals engaged in
137 meatpacking, the category does include individuals engaged in livestock care. It is
138 reasonable to assume that some of the risk we identify here may be attributable to the
139 fact that counties with high numbers of farmworkers are also likely to have high
140 numbers of workers in animal processing facilities.

141
142 The negative association we found between insured status and mortality is a point of
143 concern. The CDC has noted higher than expected numbers of death across the United
144 States throughout April in recent months, suggesting that COVID-19 mortality is
145 potentially higher than what has thus far been captured by state and county level

146 surveillance(11). It is possible that this association represents a gap in testing and
147 linkage to care among the uninsured, and/or a gap in ascertaining deaths due to
148 COVID-19 among uninsured individuals.

149

150 **Conclusion**

151 COVID-19 mortality appears to be statistically significantly associated with social
152 determinants of health at the county level, and these relationships may be more
153 pronounced in non-urban counties. Individuals who do not speak English, individuals
154 engaged in farm work, and individuals living in poverty may be at heightened risk for
155 COVID-19 mortality in non-urban counties.

156

157

158 **Acknowledgments:** This work was supported by the National Institute of Mental
159 Health, grant K01MH112436 and a National Institute on Minority Health and Health
160 Disparity Loan Repayment Contract, L60-MD011114

REFERENCES

1. Dong E, Du H, Gardner L. An interactive web-based dashboard to track COVID-19 in real time. *The Lancet Infectious Diseases*. 2020;20(5):533-4.
2. Dorn AV, Cooney RE, Sabin ML. COVID-19 exacerbating inequalities in the US. *The Lancet*. 2020;395(10232):1243-4.
3. Who is getting infected with COVID-10: State of California; 2020 [updated May 17, 2020. Available from: <https://update.covid19.ca.gov/>.
4. Rates of cases, hospitalizations, and deaths by race/ethnicity group. New York City Health Department; 2020 May 14, 2020.
5. Bier D. Immigrants are about 1/3 of California's "essential workers". *The Cato Institute*; 2020 March 30, 2020
6. Borunda A. Farmworkers risk coronavirus infection to keep the U.S. fed. *National Geographic*. 2020 April 10, 2020.
7. Data show COVID-19 is hitting essential workers and people of color hardest [press release]. *ACLU of Massachusetts*, April 8, 2020 2020.
8. Farm Labor: *USDA*; 2020 [updated April 22, 2020. Available from: <https://www.ers.usda.gov/topics/farm-economy/farm-labor/-size>.
9. Data from *The New York Times*, based on reports from state and local health agencies. *The New York Times*. New York, NY2020.
10. Philbin MM, Flake M, Hatzenbuehler ML, Hirsch JS. State-level immigration and immigrant-focused policies as drivers of Latino health disparities in the United States. *Social Science & Medicine*. 2018;199:29-38.
11. Excess Deaths Associated with COVID-19: *Centers for Disease Control*; 2020 [updated April 29, 2020. Available from: https://www.cdc.gov/nchs/nvss/vsrr/covid19/excess_deaths.htm.