

1 **Title Page**

2

3 **Manuscript Title**

4 Analysis of SARS-CoV-2 transmission in different settings, among cases and close contacts
5 from the Tablighi cluster in Brunei Darussalam

6

7 **Manuscript type:** Original Research Article

8

9 **Abstract word count:** 150 words

10 **Manuscript word count:** 3510 words

11

12 **Author Affiliations:**

13 1. Liling CHAW, PhD

14 PAPRSB Institute of Health Sciences, Universiti Brunei Darussalam, Brunei

15

16 2. Wee Chian KOH, PhD

17 Centre for Strategic and Policy Studies, Brunei Darussalam

18

19 3. Sirajul Adli JAMALUDIN, MSc

20 Environmental Health Division, Ministry of Health, Brunei Darussalam

21

22 4. Lin NAING, MD

23 PAPRSB Institute of Health Sciences, Universiti Brunei Darussalam, Brunei

24

25 5. Mohammad Fathi ALIKHAN, MSc

26 Disease Control Division, Ministry of Health, Brunei Darussalam

27

28 6. Justin WONG, MPhil

29 Disease Control Division, Ministry of Health, Brunei Darussalam

30

31 **Corresponding author**

32 Liling Chaw, PhD

33 PAPRSB Institute of Health Sciences, Universiti Brunei Darussalam

34 Jalan Tungku Link Gadong BE1410, Brunei Darussalam

35 Email: liling.chaw@ubd.edu.bn

36

37 **Author contributions**

38 Study conceptualisation and design: LC, JW

39 Data acquisition: JW, SJ, FA

40 Data analysis: LC, WCK, LN

41 Data interpretation: LC, WCK, LN, JW

42 Critical revision of the manuscript for important intellectual content: LC, JW, WCK

43 All authors contributed to the writing of the first and final draft.

44

45 **Source of funding**

46 This study does not receive any form of financial support.

47 **Conflict of interest**

48 None to be declared.

49

50 **Article summary line**

51 We highlight the variability of SARS-CoV-2 transmission across different settings, identify
52 settings at highest risk, and characterize the role of environmental, behavioural, and host factors
53 in driving SARS-CoV-2 transmission.

54

55 **Running title**

56 SARS-CoV2 transmission across settings in Brunei

57

58 **Biographical sketch**

59 Dr Liling Chaw is a lecturer at the PAPRSB Institute of Health Sciences, Universiti Brunei
60 Darussalam, Bandar Seri Begawan. Her primary research interests include the burden and
61 transmission dynamics of infections diseases, particularly on acute respiratory diseases.

62

63

64

65

66 **Abstract**

67

68 We report the transmission dynamics of SARS-CoV-2 across different settings from the initial
69 COVID-19 cluster in Brunei, arisen from 19 attendees at the Malaysian Tablighi Jama'at
70 gathering and resulted in 52 locally transmitted cases. Highest non-primary attack rates(ARs)
71 were observed at a subsequent local religious gathering (14.8% [95%CI: 7.1,27.7]) and in the
72 household (10.6% [95%CI: 7.3,15.1]). Household ARs of symptomatic cases were higher (14.4%
73 [95%CI: 8.8,19.9]) than asymptomatic (4.4% [95%CI: 0.0,10.5]) or presymptomatic cases (6.1%
74 [95%CI: 0.3,11.8]). Low ARs (<1%) were observed for workplace and social settings. Our
75 analysis highlights that SARS-CoV-2 transmission varies depending on environmental,
76 behavioural and host factors. We identify 'red flags' of potential super-spreading events, namely
77 densely populated gatherings for prolonged periods in enclosed settings, presence of individuals
78 with recent travel history, and group behaviours such as communal eating, sleeping and sharing
79 of personal hygiene facilities. We propose differentiated testing strategies that account for
80 transmission risk.

81

82

83

84 INTRODUCTION

85

86 Initially reported on December 31, 2019, cases of coronavirus disease (COVID-19) have since
87 escalated significantly, prompting the World Health Organisation's declaration of a pandemic on
88 March 11. A rapid response by the global scientific community has described many aspects of
89 the SARS-CoV-2 virus. Its primary mode of transmission is through respiratory droplets, and
90 fomite and aerosol transmission may also play a role (1). Asymptomatic transmission has been
91 observed (2) and the peak infective period appears to be within the first few days of symptom
92 onset (3). Estimates suggest a basic reproduction number (R_0) of 2–3 in the early stages of an
93 outbreak (4). While the R_0 is valuable in assessing the spread of the outbreak, it can obscure
94 individual heterogeneity in the level of infectivity, among persons and in different settings (5, 6).
95 Modelling of the 2003 SARS outbreak indicates that >70% of transmission occurred due to
96 super-spreading events (SSE) (7). Early reports suggest that similar dynamics may be in play in
97 the explosive propagation of SARS-CoV-2 (8).

98

99 As part of their mitigation strategies, many countries have adopted stringent yet blunt
100 'lockdowns' of whole cities and provinces (9) and are implementing exit strategies. In moving
101 towards a more targeted approach, countries should be able to answer two major questions
102 regarding the transmission dynamics of SARS-CoV-2: (i) what are the environmental,
103 behavioural, and host factors that drive transmission? and, (ii) what are the most effective
104 interventions to control this? To address these, we report on analysis of a transmission chain in
105 Brunei Darussalam that resulted from an international SSE.

106

107 Brunei (population 459,500) (10) detected its first COVID-19 case on March 9, arising from an
108 Islamic religious gathering (Tablighi Jama'at) in Kuala Lumpur, Malaysia, a known SSE lasting
109 four days and attended by >16,000 people, including international participants (11). Of the 135
110 confirmed cases in Brunei reported as of first week of April, 71 cases (52.6%) have an
111 epidemiological link to the Tablighi event (Figure 1).

112
113 Figure 1 – Epidemic curve for the first 135 COVID-19 cases in Brunei Darussalam, by Tablighi
114 (green bars) and other imported (orange bars) clusters.

115
116 As SARS-CoV-2 is a novel infection in a naive population, an outbreak investigation of this
117 event can provide insights into the transmission dynamics of the disease and the effectiveness of
118 outbreak control measures. The thorough nature of Brunei's contact tracing provides a rare
119 opportunity to study the epidemiological and transmission characteristics of SARS-CoV-2 in the
120 community setting.

121
122 **METHODS**

123
124 **Surveillance and case identification**

125
126 Brunei's Ministry of Health has responsibility for communicable disease surveillance. Since
127 January 23, testing criteria have been implemented for suspected COVID-19 cases. Initially,
128 individuals with acute respiratory symptoms, and a travel history to a high-risk area were tested
129 for SARS-CoV-2. Over the next weeks, the program expanded to include: (i) contacts of a

130 confirmed case (regardless of symptoms); (ii) individuals admitted to an inpatient facility with
131 pneumonia; and (iii) individuals who present to a health facility with acute respiratory illness for
132 the second time within 14 days. On March 21, Brunei started testing and isolating all travellers
133 and returning residents. On March 25, SARS-CoV-2 sampling at selected sentinel health centers
134 was introduced, followed by mandatory random screening for selected groups of foreign workers
135 on April 7.

136
137 A confirmed case is defined as a person who tested positive for SARS-CoV-2 through real-time
138 reverse transcriptase polymerase chain reaction (RT-PCR) test on nasopharyngeal (NP) swab
139 (12). The first positive case in Brunei was detected on March 9, having met the testing criteria as
140 a person with fever and cough with a recent travel history to Kuala Lumpur.

141
142 **Epidemiological investigation**

143
144 Under the Infectious Disease Act, the Ministry of Health conducted epidemiological
145 investigation and data collection for each case and close contact using WHO's First Few Cases
146 protocol (13). The first case was interviewed for demographic characteristics, clinical
147 symptoms, travel history, activity mapping, and contact history. Upon identification of his
148 participation at the Tablighi event in Malaysia, and we found that a number of other Bruneians
149 had also participated at the same event. We subsequently obtained the details of all Bruneian
150 participants.

151

152 NP swabs were collected from all identified participants and tested with RT-PCR. Those who
153 tested positive were admitted to the National Isolation Centre (NIC), while those tested negative
154 were quarantined for 14 days from their return to the country at a designated community
155 quarantine facility where symptom and temperature screening was conducted daily. Those who
156 developed symptoms were re-tested. Activity mapping of confirmed cases was conducted, and
157 contact tracing initiated.

158
159 A close contact is defined as any person living in the same household, or someone within one
160 meter of a confirmed case in an enclosed space for more than 15 minutes. All close contacts of
161 confirmed cases were tested with RT-PCR. Those who tested positive were admitted to NIC,
162 while those tested negative, were placed under home quarantine for 14 days from last exposure
163 to the confirmed case. For individuals under home quarantine, their compliance and health status
164 were monitored daily, through video calls or face-to-face assessments. Those who developed
165 symptoms during home quarantine were re-tested.

166
167 **Clinical management**

168
169 All confirmed cases were treated and isolated at NIC and followed-up until recovery. We
170 obtained clinical information on their history (including any prior presentation to health
171 services), examination, laboratory and radiological results from digital inpatient records on the
172 national health information system database. Additionally, oral history was taken to ascertain
173 whether they had symptoms up to 14 days prior to diagnosis. Cases were discharged following
174 two consecutive negative specimens collected at ≥ 24 -hour intervals.

175

176 **Cases**

177

178 We categorised cases into two groups: primary (those presumably infected at the Tablighi event
179 in Malaysia), and non-primary (those who did not attend the Tablighi event but had an
180 epidemiological link to the primary cases).

181

182 For each case, the symptom status was recorded and classified as: (i) symptomatic, if symptoms
183 were reported during or prior to NP swab collection; (ii) presymptomatic, if symptoms were
184 reported after NP sampling but during admission; or (iii) asymptomatic, if no symptoms were
185 ever reported up to discharge.

186

187 **Close contacts**

188

189 We classified the type of close contact into five settings: household, relatives, workplace, social,
190 and local religious gathering.

191

192 ‘Household’ is defined as those living in the same household. Among the household members,
193 their relationship with a case was also recorded and classified as spouse, child, and others
194 (including other familial relationship or housekeepers living in the same household). ‘Relatives’
195 is defined as those who live outside the household but are related to the case. ‘Workplace’ is
196 defined as contacts encountered in the workplace or school. ‘Social’ is defined as those
197 encountered during travel or in social events. Lastly, ‘local religious gathering’ is defined as

198 those who attended a local religious event in Brunei on March 5, which ran throughout the night
199 with participants staying overnight.

200

201 **Data analysis**

202

203 Group comparison was done between primary and non-primary cases using Chi-square, Fisher's
204 Exact, or Mann-Whitney's tests as appropriate. Incubation period was calculated from cases
205 where dates of exposure and symptom onset were clear. Serial interval (SI) was calculated by
206 subtracting the date of symptom onset of the infectee from the infector; only symptomatic and
207 pre-symptomatic infector-infectee pairs with clear epidemiological links were included.

208

209 For each setting, the attack rate (AR) was calculated by dividing the number of positive contacts
210 by the total number of close contacts (that is, the proportion of contacts that tested positive). To
211 identify risk factors of infection, log-binomial regression analysis was applied to estimate the
212 risk ratio for gender, age, and setting. Further stratification was done to assess differences in
213 infector symptom status across settings. The 95% confidence interval (95% CI) was estimated
214 using the normal-approximation method, or binomial method if the count was less than five.

215

216 The mean observed reproductive number, R , and the distribution of personal reproductive
217 numbers in each setting were calculated from the number of infected close contacts caused by
218 each primary case. The 95% CI was estimated based on the Poisson distribution (14).

219

220 All analyses were conducted using Microsoft Excel and R (ver. 3.6.3) (15). A *p*-value <0.05 was
221 considered as statistically significant. Ethical approval was obtained from the University
222 Research Ethics Committee, Universiti Brunei Darussalam (Ref:
223 UBD/OAVCR/UREC/Apr2020-05).

224

225

226 **RESULTS**

227

228 **Epidemiological characteristics**

229

230 Seventy-five individuals in Brunei attended the Tablighi event in Malaysia. Of these, 19 were
231 positive for SARS-CoV-2 resulting in 52 additional cases transmitted locally, bringing the total
232 cluster size to 71. Figure 2 illustrates the epidemiological links in the cluster by generation in the
233 transmission chain, and by their symptomatic status. There were 32 (45.1%), 15 (21.1%), and 5
234 (7.0%) cases in generations one, two, and three, respectively.

235

236 Figure 2 – The Tablighi cluster in Brunei Darussalam, with epidemiological links illustrated by
237 generation and symptomatic status

238

239 Table 1 shows the demographic and clinical characteristics of the cases in the cluster. The
240 median age was 33.0 years [Interquartile range (IQR)= 29.0 years], majority male (64.8%,
241 n=46), and 5 cases (7.1%) had pre-existing chronic conditions. Compared to non-primary cases,
242 primary cases were significantly older and predominantly male. More than three-quarter of the

243 cases (77.4%, n=55) were diagnosed and immediately admitted to the NIC within 5 days of
244 symptom onset or NP swab taken (data not shown).

245
246 A substantial proportion of cases were presymptomatic (31.0%, n=22) or asymptomatic (12.7%,
247 n=9). 40 (56.3%) patients reported symptoms during contact tracing investigation. The most
248 commonly reported symptoms were fever, cough, and sore throat. Only one (1.4%) and two
249 (2.8%) cases were critical and severe, respectively.

250
251 Table 1 - Demographic and clinical characteristics of COVID-19 cases in Tablighi cluster,
252 Brunei

253
254 The incubation period was calculated from eight cases that had confirmed epidemiological links
255 and had attended the March 5 religious gathering in Brunei. Using their attendance date as the
256 exposure date, the median incubation period was 4.5 days (IQR= 2.75 days, range= 1 to 11
257 days). Based on 35 symptomatic infector-infectee pairs, the mean (SD) SI was 4.26 (4.27) days,
258 ranging from -4 to 17 days. Four pairs (11.4%) had negative SI values. The SI distribution
259 resembles a normal distribution (Supplementary Figure S1).

260
261 **Transmission characteristics**

262
263 Of the 1755 close contacts in the Tablighi cluster, 51 local transmissions were detected, giving
264 an overall non-primary AR of 2.9% (95% CI: 2.2, 3.8). Case 121 (see Figure 2) was excluded
265 from this analysis because he was not detected during contact tracing. Highest AR was observed

266 among spouse (41.9% [95% CI: 24.1, 60.7]), followed by the local religious gathering (14.8%
267 [95% CI: 7.1, 27.7]) and children (14.1% [95% CI: 7.8, 23.8]). The overall household AR is
268 10.6% (95% CI: 7.3, 15.1).

269

270 Multiple log-binomial regression analyses revealed that type of close contact was the only
271 statistically significant variable ($p < 0.001$; Table 2). When compared to social contacts, spouses
272 of positive cases had the highest adjusted risk ratio of getting the infection (45.20 [95% CI: 16.8,
273 156.1]), followed by the local religious gathering attendees (15.60 [95% CI: 4.8, 59.9]), and their
274 children (14.09 [95% CI: 4.8, 51.5]).

275

276 Table 2 - Risk factors of SARS-COV2 infection among close contacts

277

278 ARs also differed by symptom status of the infector (Table 3). ARs in households where the
279 infectors were symptomatic (14.4%) were higher than those who were asymptomatic (4.4%) or
280 presymptomatic (6.1%). AR for the local religious gathering could not be calculated as three
281 primary cases at the event had different symptom status; hence we could not ascertain how
282 transmission occurred. In the household setting, symptomatic cases have 2.66 times higher risk
283 of transmitting to their close contacts, when compared to asymptomatic and presymptomatic
284 (crude risk ratio: 2.66 [95% CI: 1.12, 6.34], Table S1).

285

286 Table 3 - Attack rates in different settings, stratified by symptom status of the infector

287

288 The mean observed R was highest in the local religious gathering (2.67), followed by the
289 household setting (0.67 [95% CI: 0.44, 0.96]) (Table 4). The distribution of the observed R in the
290 household setting was skewed towards zero (Figure S2). 71.4% of household infections (20 of 28
291 positive contacts) were from 16.7% of cases (7 of 42 possible links).

292

293 Table 4 – Characteristics and mean observed R for each setting

294

295

296 **DISCUSSION**

297

298 We characterise the Tablighi Jama'at cluster that started the COVID-19 epidemic in Brunei. Our
299 analysis reveals several key findings. First, SSE plays an important role in SARS-CoV-2
300 transmission. Second, there is high transmission variability across different settings. Third,
301 transmission varies between symptomatic vs. asymptomatic and presymptomatic cases, and we
302 highlight the potential for silent chains of transmission.

303

304 **SSE**

305

306 Within this cluster, 38% of all cases were participants at an SSE: 19 (26.7%) from the Tablighi
307 event in Malaysia, and eight (11.3%) from the local religious gathering in Brunei. Notably, 19 of
308 the 75 Brunei attendees at the Tablighi event tested positive. Assuming a representative sample,
309 this would suggest an AR of 25%, implying that approximately 4,000 cases (of an estimated
310 16,000 participants) may have been infected at that Malaysian event. Moreover, we find that the

311 highest overall non-primary AR and mean observed R was at the local religious gathering (AR=
312 14.8%, R= 2.67), which was higher than that observed in the household setting (AR= 10.6%, R=
313 0.67). These observations suggest a role for mass gatherings in facilitating SARS-CoV2
314 transmission.

315
316 The Tablighi is an apolitical Islamic movement with a presence in nearly 200 countries. Tablighi
317 adherents usually travel to gather at annual international events each lasting several days.
318 Communal prayers, meals and speeches form part of these events. In Malaysia, the participants
319 stayed and slept at the mosque, and several participants were deputised to cook meals and clean.
320 Smaller gatherings, usually every week or so take place in their home countries, with other
321 adherents. Over the course of this investigation, we identified several common characteristics at
322 both the local religious gathering and the Tablighi event in Malaysia (11). First, significant
323 numbers of people gathered in an enclosed area for a prolonged period. Second, attendees had a
324 recent travel history – the Tablighi event in Malaysia drew participants from across the world,
325 while the local religious gathering had at least three individuals who had recently returned from
326 Malaysia. Third, communal sharing of sleeping areas and toilets, and shared dining were
327 observed. We propose that these three characteristics are hallmarks for the development of SSE
328 for SARS-CoV-2 transmission and can be used as ‘red flags’ by health authorities in their risk
329 assessment and mitigation strategies for preventing and detecting high risk activities including
330 mass gatherings, and other institutional settings such as care homes, prisons and dormitories.

331

332 **Variability of non-primary AR across different settings**

333

334 To a lesser degree, our observations on the within-household transmission are similar to that
335 observed for the two religious gatherings. Out of 16 household contacts who subsequently
336 became first generation cases, 10 (62.5%) of these were from just three primary cases. As such,
337 even within similar settings, we can expect wide variability in transmission patterns. This
338 observation supports our finding of a moderately high household AR but an observed R of less
339 than one, suggesting that transmission is driven by a relatively small number of cases (5). High
340 ARs in spouses and children reflect intimate relationships with high degree of interaction, close
341 proximity, and in the case of the spouse, sleeping in the same room. Concordant with our SSE
342 findings, we suggest that encounters among groups of people that involve close proximity in
343 enclosed settings for prolonged time periods (at least for one night) is a main driver of SARS-
344 CoV-2 transmission.

345
346 Our overall non-primary AR result of 10.6% in the household setting is comparable to other
347 studies that used contact-tracing datasets (16-19). A study near Wuhan, China (20) reported a
348 higher AR of 16.3%; however, they detected 56.2% of their cases more than five days after
349 symptom onset. By contrast, 77.4% of the cases in our study were detected and isolated within
350 five days of symptom onset, suggesting that early case isolation can reduce AR. The aggressive
351 testing of contacts strategy employed may have contributed to this.

352
353 We note the low non-primary AR (<1%) and mean observed R (<0.3) for workplace and social
354 settings. While moderate physical distancing was implemented in Brunei following the
355 identification of this cluster, there was no community quarantine or lockdown, public services
356 and businesses remained open, and no internal movement restrictions were imposed.

357

358 Combined with our observations on the role of SSE in driving SARS-CoV-2 transmission, we
359 suggest that in areas with limited community transmission full lockdown measures that adopt a
360 blunt approach by restricting all movement can be avoided, in favour of a more targeted
361 approach that includes a combination of case isolation, contact tracing, and moderate levels of
362 physical distancing that take into account the ‘red flags’ for mass gatherings identified earlier.
363 However, this approach is resource intensive and only feasible where there is sufficient public
364 health capacity. The high asymptomatic proportion suggest that even with best efforts at contact
365 tracing, potential for widespread community transmission is clear. Once established, suppression
366 necessitates the implementation of broader physical distancing measures (9, 21). Nonetheless, if
367 done effectively, contact tracing and case isolation approaches are shown to control the COVID-
368 19 outbreak during its early stage (22). Modelling studies using South Korean data showed that
369 less extreme physical distancing measures can help to suppress the outbreak (23).

370

371 **Symptomatic vs. asymptomatic and presymptomatic transmission**

372

373 We identified several environmental (settings) and behavioural factors that potentially account
374 for higher ARs observed in mass gatherings and in the household. In order to assess the impact
375 of host factors in driving transmission, we compared non-primary AR in symptomatic vs.
376 asymptomatic and presymptomatic individuals, considering the high proportion of asymptomatic
377 (12.7%) and presymptomatic (31.0%) of cases.

378

379 While there are case reports of presumptive asymptomatic and presymptomatic transmission (24,
380 25), observational studies quantifying such transmission are few. A study from Ningbo, China
381 analysed the overall ARs in symptomatic vs. asymptomatic cases and did not find significant
382 difference between the two groups (26). Another study, re-interpreted the same data and
383 theorized that under certain conditions, symptomatic cases could be more transmissible than
384 asymptomatic ones (27). In fact, our overall crude risk ratio for symptomatic cases showed no
385 significant difference when compared with asymptomatic and/or presymptomatic cases (Table 3,
386 Table S1). However, we suggest that this masks the true picture in transmissibility when
387 different settings are taken into account.

388
389 In our study, we do not find a significant difference in AR in non-household settings. These
390 settings usually practice some form of non-pharmaceutical interventions (NPI)—individuals with
391 moderate and severe symptoms may be on medical leave, and it is reasonable to expect some
392 physical distancing would be practiced by contacts of persons who display visible symptoms.
393 This is less feasible within the household setting; hence we suggest that transmission occurs
394 more frequently at the household level where control measures are less practical. We observed
395 that the household AR for symptomatic cases (14.4%) is higher than that of asymptomatic
396 (4.1%) or presymptomatic cases (6.1%), suggesting that the presence of symptoms is a host
397 factor in driving transmission.

398
399 The higher household AR observed among symptomatic cases suggest that testing for such cases
400 should be prioritised, especially in low resource areas with limited testing capacity. Nonetheless,
401 an AR of 4.4% and 6.1% in asymptomatic and presymptomatic cases, respectively, is not

402 negligible. As such, our findings have several implications for high resource areas with greater
403 testing capacity. First, it strengthens the argument for testing household contacts in the absence
404 of symptoms. Second, there is a need to allow for ‘slack’ in the surveillance system as the high
405 proportion of asymptomatic cases pose challenges for rapid detection and isolation. We
406 recommend that even in countries with highly developed testing and tracing capacities, moderate
407 levels of social distancing should be implemented to account for this. Third, proactive testing of
408 travellers, attendees at ‘red flag’ events, and institutional settings may be necessary to contain
409 COVID-19 spread.

410
411 This study has several limitations. First, as a retrospective study based on a contact tracing
412 dataset, the index case determination or the direction of transmission may be uncertain,
413 particularly as a substantial proportion of cases are asymptomatic. Moreover, without accounting
414 for outside sources of infection, setting-specific SARs could have been overestimated (although
415 no community transmission has been detected in Brunei). Viral sequencing can
416 confirm homology between the strains infecting the index and secondary cases across the various
417 settings, however this was not conducted for all cases. Second, we have not accounted for other
418 potential environmental factors such as the relative size of the household, time spent at home
419 with others, air ventilation, and indirect transmission through fomites. Third, we do not have
420 information on NPIs practiced by the close contacts; presumably, individuals would take
421 precautions during an outbreak. Fourth, symptom status of the cases was reported during their
422 swab collection date. We assume this to be reflective of their actual condition when their close
423 contacts were exposed, however, this may not be necessarily true for all cases. Finally, the

424 generalizability of our results are limited due to no community transmission, a lack of cases in
425 settings such as residential care facilities and dormitories, and small number of cases.

426
427 The main strength of our study is the availability of a complete contact tracing dataset at the
428 national level. Since all contacts were tested, it is reasonable to assume that this study more
429 accurately detects SARS-CoV-2 transmission than those that only test symptomatic contacts. In
430 conclusion, our analysis highlights the variability of SARS-CoV-2 transmission across different
431 settings and in particular, the role of SSEs. We identify ‘red flags’ for the development of
432 potential SSEs, and describe environmental, behavioural, and host factors that drive transmission.
433 Overall, we provide evidence that a combination of case isolation, contact tracing, and moderate
434 physical distancing measures is an effective approach to containment.

435

436 **Acknowledgement**

437 The authors would like to thank Mr Haji Mohamad Ruzaimi Haji Rosli from Corporate
438 Communications, Ministry of Health Brunei, for his assistance in designing the figure for this
439 manuscript.

440

441

442

443

444 Figure and table captions:

445 Figure 1 – Epidemic curve for the first 135 COVID-19 cases in Brunei Darussalam, by Tablighi
446 (green bars) and other imported (orange bars) clusters.

447 Figure 2 – The Tablighi cluster in Brunei Darussalam, with epidemiological links illustrated by
448 generation and symptomatic status

449 Table 1 - Demographic and clinical characteristics of COVID-19 cases in Tablighi cluster,
450 Brunei

451 Table 2 - Risk factors of SARS-COV2 infection among close contacts

452 Table 3 - Attack rates in different settings, stratified by symptom status of the infector

453 Table 4 – Characteristics and mean observed R for each setting

454

455

456

457 Supplementary information:

458 **Figure S1.** Distribution of the serial interval, fitted with a normal distribution

459 **Figure S2.** Distribution of the household observed R

460 Table S1. Attack rates in different settings, stratified by symptom status of the infector
461 (symptomatic versus presymptomatic and asymptomatic)

462

463

464

465

466

467 **References**


468


469

470

- 471 1. Ong SWX, Tan YK, Chia PY, Lee TH, Ng OT, Wong MSY, et al. Air, Surface
472 Environmental, and Personal Protective Equipment Contamination by Severe Acute Respiratory
473 Syndrome Coronavirus 2 (SARS-CoV-2) From a Symptomatic Patient. *JAMA*. 2020.
474 2. Furukawa N, Brooks J, Sobel J. Evidence Supporting Transmission of Severe Acute
475 Respiratory Syndrome Coronavirus 2 While Presymptomatic or Asymptomatic. *Emerging*
476 *Infectious Disease journal*. 2020;26(7).
477 3. He X, Lau EHY, Wu P, Deng X, Wang J, Hao X, et al. Temporal dynamics in viral
478 shedding and transmissibility of COVID-19. *Nature Medicine*. 2020 2020/04/15.
479 4. Kucharski AJ, Russell TW, Diamond C, Liu Y, Edmunds J, Funk S, et al. Early dynamics
480 of transmission and control of COVID-19: a mathematical modelling study. *The Lancet*
481 *Infectious Diseases*. 2020.
482 5. Liu Y, Eggo RM, Kucharski AJ. Secondary attack rate and superspreading events for
483 SARS-CoV-2. *The Lancet*. 2020;395(10227):e47.
484 6. Stein ML, van der Heijden PGM, Buskens V, van Steenberghe JE, Bengtsson L,
485 Koppeschaar CE, et al. Tracking social contact networks with online respondent-driven
486 detection: who recruits whom? *BMC Infectious Diseases*. 2015 2015//;15(1):1-12.
487 7. Li Y, Yu ITS, Xu P, Lee JHW, Wong TW, Ooi PL, et al. Predicting Super Spreading
488 Events during the 2003 Severe Acute Respiratory Syndrome Epidemics in Hong Kong and
489 Singapore. *Am J Epidemiol*. 2004;160(8):719-28.
490 8. !!! INVALID CITATION !!! [Frieden, 2020 #4083].
491 9. Lau H, Khosrawipour V, Kocbach P, Mikolajczyk A, Schubert J, Bania J, et al. The
492 positive impact of lockdown in Wuhan on containing the COVID-19 outbreak in China. *Journal of*
493 *Travel Medicine*. 2020.
494 10. Department of Economic Planning and Development. Mid-year population estimates for
495 Brunei Darussalam, 2019. 2020 [cited 2020 22 April]; Available from:
496 <http://www.deps.gov.bn/SitePages/Population.aspx>
497 11. Mat NFC, Edinur HA, Razab MKAA, Safuan S. A Single Mass Gathering Resulted in
498 Massive Transmission of COVID-19 Infections in Malaysia with Further International Spread.
499 *Journal of Travel Medicine*. 2020.
500 12. Wong J, Koh WC, Momin RN, Alikhan MF, Fadillah N, Naing L. Probable causes and
501 risk factors for positive SARS-CoV-2 test in recovered patients: Evidence from Brunei
502 Darussalam. *Journal of Medical Virology*. 2020 2020/06/19;n/a(n/a).
503 13. World Health Organization. The First Few X (FFX) Cases and contact investigation
504 protocol for 2019-novel coronavirus (2019-nCoV) infection, version 2. 2020.
505 14. Liu Y, Gayle AA, Wilder-Smith A, Rocklöv J. The reproductive number of COVID-19 is
506 higher compared to SARS coronavirus. *Journal of Travel Medicine*. 2020;27(2).
507 15. R Core Team. R: A language and environment for statistical computing. R Foundation
508 for Statistical Computing, Vienna, Austria; 2020.
509 16. Cheng H-Y, Jian S-W, Liu D-P, Ng T-C, Huang W-T, Lin H-H. High transmissibility of
510 COVID-19 near symptom onset. *medRxiv*. 2020:2020.03.18.20034561.
511 17. Luo L, Liu D, Liao X-I, Wu X-b, Jing Q-I, Zheng J-z, et al. Modes of contact and risk of
512 transmission in COVID-19 among close contacts. *medRxiv*. 2020:2020.03.24.20042606.

- 513 18. Bi Q, Wu Y, Mei S, Ye C, Zou X, Zhang Z, et al. Epidemiology and Transmission of
514 COVID-19 in Shenzhen China: Analysis of 391 cases and 1,286 of their close contacts.
515 medRxiv. 2020:2020.03.03.20028423.
- 516 19. Jing Q-L, Liu M-J, Yuan J, Zhang Z-B, Zhang A-R, Dean NE, et al. Household
517 Secondary Attack Rate of COVID-19 and Associated Determinants. medRxiv.
518 2020:2020.04.11.20056010.
- 519 20. Li W, Zhang B, Lu J, Liu S, Chang Z, Cao P, et al. The characteristics of household
520 transmission of COVID-19. *Clinical Infectious Diseases*. 2020.
- 521 21. Sjödin H, Wilder-Smith A, Osman S, Farooq Z, Rocklöv J. Only strict quarantine
522 measures can curb the coronavirus disease (COVID-19) outbreak in Italy, 2020.
523 *Eurosurveillance*. 2020;25(13):2000280.
- 524 22. Hellewell J, Abbott S, Gimma A, Bosse NI, Jarvis CI, Russell TW, et al. Feasibility of
525 controlling COVID-19 outbreaks by isolation of cases and contacts. *The Lancet Global Health*.
526 2020.
- 527 23. Park SW, Sun K, Viboud C, Grenfell BT, Dushoff J. Potential roles of social distancing in
528 mitigating the spread of coronavirus disease 2019 (COVID-19) in South Korea. medRxiv.
529 2020:2020.03.27.20045815.
- 530 24. Wei WE, Li Z, Chiew CJ, Yong SE, Toh MP, Lee VJ. Presymptomatic Transmission of
531 SARS-CoV-2 — Singapore, January 23–March 16, 2020. *MMWR Morb Mortal Wkly Rep*. 2020.
- 532 25. Qian G, Yang N, Ma AHY, Wang L, Li G, Chen X, et al. COVID-19 Transmission Within
533 a Family Cluster by Presymptomatic Carriers in China. *Clinical Infectious Diseases*. 2020.
- 534 26. Chen Y, Wang A, Yi B, Ding K, Wang H, Wang J, et al. The epidemiological
535 characteristics of infection in close contacts of COVID-19 in Ningbo city
536 . *Chinese Journal of Epidemiology*. 2020;41.
- 537 27. He D, Zhao S, Lin Q, Zhuang Z, Cao P, Wang MH, et al. The relative transmissibility of
538 asymptomatic cases among close contacts. *International Journal of Infectious Diseases*.
539 2020;94:145-7.
- 540







Tablighi Jamaat Religious Gathering
Kuala Lumpur, Malaysia
28 February - 1 March 2020

Legend:

-  Symptomatic Case
-  Presymptomatic Case
-  Asymptomatic Case



Local Religious Gathering
Brunei Darussalam
5 March

