

1 **Associations between ambient air pollutants exposure and case**  
2 **fatality rate of COVID-19: a multi-city ecological study in China.**

3  
4 Tiantian Zhang<sup>1#</sup> Guoli Zhao<sup>2#</sup> Li Luo<sup>1</sup> Yongzhen Li<sup>1</sup> Wenming Shi<sup>3\*</sup>

5  
6 1. School of Public Health, Key Lab of Public Health Safety of the Ministry of  
7 Education and NHC Key Lab of Health Technology Assessment, Fudan University,  
8 Shanghai 200032, China.

9 2. Institutes of Brain Science, Zhongshan Hospital, Fudan University, Shanghai  
10 200032, China.

11 3. Clinical and Translational Research Center of Shanghai First Maternity & Infant  
12 Hospital, School of Medicine, Tongji University, Shanghai 201204, China

13  
14

15 **# These two author contributed equally to this article.**

16

17 **\* Correspondence:** Wenming Shi, MSc. E-mail: [Shiwmfd@126.com](mailto:Shiwmfd@126.com)

18

19

20 **Type:** Research Letter

21

22 **Abstract**

23 **Background:** Environmental factors, including air pollution, can strongly impact on  
24 spatio-temporal patterns of infectious diseases outbreak. In this study, we aimed to  
25 investigate the association and correlation between ambient air pollutants and case  
26 fatality rate (CFR) of the novel coronavirus disease (COVID-19) in China.

27 **Methods:** Publicly accessible data on COVID-19 average CFR were utilized in the  
28 data analysis. The ambient daily air pollutants including fine particulate matter  
29 (PM<sub>2.5</sub>), inhalable particles (PM<sub>10</sub>) and nitrogen dioxide (NO<sub>2</sub>) during the period from  
30 December 25, 2019 to March 5, 2020 were obtained from National Air Quality  
31 Real-time Publishing System of China. Ecological analysis was performed to explore

32 the association and correlation between the cumulative average exposure of ambient  
33 air pollutants at different lag days (14 and 28 days) and average CFR in China outside  
34 Hubei and cities in Hubei province via model fitting.

35 **Results:** The average case fatality rate was highest in Wuhan city (4.53%) and the  
36 cumulative average exposure of ambient PM<sub>2.5</sub>, PM<sub>10</sub> and NO<sub>2</sub> at lag 28 days was  
37 55.8±12.1µg/m<sup>3</sup>, 66.8±9.2µg/m<sup>3</sup>, 20.7±4.4µg/m<sup>3</sup>, respectively in Hubei province  
38 during the study period. Ecological analysis showed that ambient PM<sub>2.5</sub>, PM<sub>10</sub> and  
39 NO<sub>2</sub> exposure at both lag 14 and 28 days was positively correlated with average CFR  
40 in China outside Hubei (province-level). For city-level analysis in Hubei, significant  
41 associations were only found between cumulative ambient NO<sub>2</sub> exposure and average  
42 CFR(r=0.693 for Lag0-14, r=0.697 for Lag0-28, respectively) during the same period.

43 **Conclusion:** Our findings suggested ambient PM<sub>2.5</sub>, PM<sub>10</sub> and NO<sub>2</sub> exposure,  
44 especially at 28 lag days, positively associated with the case fatality rate of  
45 COVID-19 in China. These results could help provide guidance for identifying  
46 potential exposure window and preventing and controlling the epidemic.

47

48 **Keywords:** COVID-19; Air pollution; Case fatality rate; Ecological study; China

49

50

51

52

53

54

55

56

57

## 58 **Introduction**

59       The novel coronavirus disease (COVID-19) is an ongoing global epidemic event,  
60 which caused by severe acute respiratory syndrome coronavirus 2 (SARS- CoV-2)[1].  
61 Respiratory droplets and human-to-human contact are the major routes of  
62 transmission of COVID-19. Some studies showed that environmental factors might  
63 influence the transmissions and risk of COVID-19[2, 3]. Since COVID-19 is a  
64 respiratory infectious disease and SARS-CoV-2 could remain viable in aerosols for  
65 hours [4], it's interesting to investigate the associations between air pollution and  
66 COVID-19 infection ,and identify possible indicator of air pollutants for blocking the  
67 epidemic. However, the study on the effects of air pollutants on the death risk of  
68 COVID-19 infection was still limited and inconsistent in China. In this study, we  
69 aimed to investigate the associations between ambient air pollutants exposure and  
70 case fatality rate (CFR) of COVID-19 in Mainland China, to provide scientific  
71 guidance for identifying potential exposure time window and preventing and  
72 controlling the epidemic.

73

## 74 **Methods**

75       For this multi-city ecological study , we collected the daily cumulative confirmed  
76 and death numbers of COVID-19 cases information in China during the study period  
77 from January 22 to March 5, 2020 from the National Health Commissions  
78 ([http://www.nhc.gov.cn/xcs/xxgzbd/gzbd\\_index.shtml](http://www.nhc.gov.cn/xcs/xxgzbd/gzbd_index.shtml)) and provincial Health  
79 Commissions of China (<http://wjw.hubei.gov.cn/bmdt/ztzl/fkxxgzbdgrfyyq/index.shtml>). Due on no death case was reported in 5 provinces (Xizang, Qinghai, Shanxi,  
80 Jiangsu and Ningxia) by March 5, 2020 and island-province factor (Hainan), a total of  
81 25 provinces covering 256 cities (243 cities outside Hubei, 13 main cities in Hubei)  
82 were included in our study. The average case fatality rate of COVID-19 was  
83 calculated in China outside Hubei at province-level, and in each city of Hubei  
84 province during the same period. Due on the data were all from public websites, the  
85 Ethical approval was not needed in the study.

86

87 The data on daily average concentrations of air pollutants including fine  
88 particulate matter (PM<sub>2.5</sub>), inhalable particles (PM<sub>10</sub>) and nitrogen dioxide (NO<sub>2</sub>)  
89 during the period from December 25, 2019 to March 5, 2020 was obtained from  
90 National Air Quality Real-time Publishing System of China. This platform provided  
91 the real-time criteria air pollutants in all state-controlled monitoring stations. To  
92 control for the potential confounders, the ambient daily average temperature, relative  
93 humidity for each city was obtained from China Meteorological Data Service Center  
94 ([http:// data. cma. cn/en](http://data.cma.cn/en)).

95 The temporal cumulative average exposure of ambient PM<sub>2.5</sub>, PM<sub>10</sub> and NO<sub>2</sub> at  
96 different lag days (14 days and 28 days) before the daily case reports was evaluated  
97 among the cities in Hubei province (city-level) and in China outside Hubei  
98 (province-level).

99 Ecological analysis was conducted to explore the temporal associations between  
100 the cumulative average exposure of ambient air pollutants and COVID-19 case  
101 fatality rates in China outside Hubei (province-level) and in Hubei province  
102 (city-level). Multiple regression models were fitted to the data among 24 provinces of  
103 China and city-level analysis in Hubei province alone during the study period.

## 104 **Results**

105 Among the cities in Hubei province, the cumulative average exposure level of  
106 the ambient PM<sub>2.5</sub>, PM<sub>10</sub> and NO<sub>2</sub> at lag 28 days was 55.8±12.1 μg/m<sup>3</sup>, 66.8±9.2 μg/m<sup>3</sup>,  
107 20.7±4.4 μg/m<sup>3</sup>, respectively during the study period. The average case fatality rate  
108 was highest in Wuhan city (4.53%), followed by Jingmen city and Ezhou city in  
109 Hubei province during the study period. For provinces outside Hubei, the average  
110 CFR was highest in Heilongjiang province and lowest in Zhejiang province of China.

111 The linear regression analysis showed a positive association between air  
112 pollutants exposure and average case fatality rate in China. As indicted in Figure 2,  
113 the associations are all significant between average CFR and cumulative average  
114 exposure to ambient PM<sub>2.5</sub>, PM<sub>10</sub> and NO<sub>2</sub> in both lag 14 and 28 days at  
115 province-level in China outside Hubei. The correlation coefficient of NO<sub>2</sub> was highest

116 (r=0.739, P<0.001) among the three air pollutants at lag 28 days (Figure 2). For Hubei  
117 province, city-level analysis showed a positive association between cumulative  
118 average exposure of NO<sub>2</sub> and average CFR (r=0.693 for Lag0-14, r=0.697 for  
119 Lag0-28, respectively), while no significant associations were observed between  
120 ambient PM<sub>2.5</sub> and PM<sub>10</sub> exposure and average CFR in cities of Hubei province.

## 121 **Discussion**

122 Our study suggested that ambient particulate matter and NO<sub>2</sub> exposure was  
123 associated with the increased case fatality of COVID-19 in Mainland China. These  
124 findings could provide evidence that air pollution is an important factor in COVID-19  
125 infections.

126 Previous studies has indicated that air pollution associated with a variety of  
127 respiratory disease including SARS, influenza, asthma and COPD[5-8]. A recent  
128 time-series study showed that ambient particulate matter, NO<sub>2</sub> exposure was  
129 associated with the increased confirmed case number of COVID-19[9]. A study  
130 conducted in multiply countries indicated that long-term exposure to ambient NO<sub>2</sub>  
131 could be a contributor factor to COVID-19 fatality[10]. Consistent with prior studies,  
132 our results indicate ambient NO<sub>2</sub> exposure, especially at 28 lag days, might increase  
133 the CFR of COVID-19. The biological explanation might be that air pollutants could  
134 compromise lung function and increase COVID-19 fatality[8].

135 To our knowledge , this was one of the few studies performed in Chinese cities  
136 to explore the temporal associations between ambient air pollutants exposure and case  
137 fatality rate of COVID-19. The high correlation between ambient NO<sub>2</sub> and CFR of  
138 COVID-19, especially at 28 lag days exposure, might help identifying which  
139 pollutants in certain exposure time window play the dominate the role in increasing  
140 the death risk of COVID-19 in China.

141 However, our study has some limitations. First, ecological study designs were  
142 adopted in the study, other city-level factors including implementation ability of  
143 control policy of COVID-19, the availability of medical resources might confound our  
144 findings. Second, the study period may not represent a whole air pollutants pattern

145 associated with the average case fatality rate of COVID-19. Furthermore ,the  
146 exposure misclassification might bias the study result due to the different activity  
147 patterns of people. For all these limitations, the findings should be interpreted with  
148 caution before extension to other population and research.

149

## 150 **Conclusions**

151 Our study suggest positive associations between ambient PM<sub>2.5</sub>, PM<sub>10</sub> and NO<sub>2</sub>  
152 exposure and case fatality rate of COVID-19 in China. The 28-day lag effects of air  
153 pollutants can help to identify the potential exposure window that have the high  
154 adverse impact on the COVID-19 and improve the public practice for preventing the  
155 epidemic.

156

## 157 **Author Contributions**

158 TZ, GZ, and WS designed the project, processed and analyzed the data, and  
159 wrote the manuscript. LL and YL collected the data and edited the manuscript. All  
160 authors critically reviewed and approved the final version of the manuscript

161

## 162 **Conflict of interests**

163 The authors declare no competing financial interest.

164

## 165 **Acknowledgment**

166 This study was supported by the National Natural Science Foundation of China  
167 (Grant nu :71874033) .

168

169

170

171


172

173


174 **References:**

- 175 1. Bi Q, Wu Y, Mei S, Ye C, Zou X, Zhang Z, Liu X, Wei L, Truelove SA, Zhang T,  
176 Gao W, Cheng C, Tang X, Wu X, Wu Y, Sun B, Huang S, Sun Y, Zhang J, Ma T, Lessler  
177 J, Feng T. Epidemiology and transmission of COVID-19 in 391 cases and 1286 of their  
178 close contacts in Shenzhen, China: a retrospective cohort study. *LANCET INFECT DIS*  
179 2020.
- 180 2. Sobral M, Duarte GB, Da PSA, Marinho M, de Souza MA. Association between  
181 climate variables and global transmission of SARS-CoV-2. *SCI TOTAL ENVIRON*  
182 2020: 729: 138997.
- 183 3. Shi P, Dong Y, Yan H, Zhao C, Li X, Liu W, He M, Tang S, Xi S. Impact of  
184 temperature on the dynamics of the COVID-19 outbreak in China. *SCI TOTAL*  
185 *ENVIRON* 2020: 728: 138890.
- 186 4. van Doremalen N, Bushmaker T, Morris DH, Holbrook MG, Gamble A,  
187 Williamson BN, Tamin A, Harcourt JL, Thornburg NJ, Gerber SI, Lloyd-Smith JO, de  
188 Wit E, Munster VJ. Aerosol and Surface Stability of SARS-CoV-2 as Compared with  
189 SARS-CoV-1. *N Engl J Med* 2020: 382(16): 1564-1567.
- 190 5. Belgrave D, Granell R, Turner SW, Curtin JA, Buchan IE, Le Souef PN, Simpson  
191 A, Henderson AJ, Custovic A. Lung function trajectories from pre-school age to  
192 adulthood and their associations with early life factors: a retrospective analysis of three  
193 population-based birth cohort studies. *Lancet Respir Med* 2018; 6(7): 526-534.
- 194 6. Lei H, Li Y, Xiao S, Lin CH, Norris SL, Wei D, Hu Z, Ji S. Routes of transmission  
195 of influenza A H1N1, SARS CoV, and norovirus in air cabin: Comparative analyses.  
196 *INDOOR AIR* 2018; 28(3): 394-403.
- 197 7. Kan HD, Chen BH, Fu CW, Yu SZ, Mu LN. Relationship between ambient air  
198 pollution and daily mortality of SARS in Beijing. *BIOMED ENVIRON SCI* 2005; 18(1):  
199 1-4.
- 200 8. Cui Y, Zhang ZF, Froines J, Zhao J, Wang H, Yu SZ, Detels R. Air pollution and  
201 case fatality of SARS in the People's Republic of China: an ecologic study. *Environ*  
202 *Health* 2003; 2(1): 15.
- 203 9. Zhu Y, Xie J, Huang F, Cao L. Association between short-term exposure to air

204 pollution and COVID-19 infection: Evidence from China. *SCI TOTAL ENVIRON*  
205 2020: 727: 138704.  
206 10. Ogen Y. Assessing nitrogen dioxide (NO<sub>2</sub>) levels as a contributing factor to  
207 coronavirus (COVID-19) fatality. *SCI TOTAL ENVIRON* 2020: 726: 138605.



**Figure 1.** Map of the distribution of the average case fatality rate of COVID-19 in Hubei Province and in Mainland China.



**Figure 2.** The correlation and associations between air pollutants exposure at different lag days and average case fatality rate of COVID-19 in Hubei province (city-level) and in Mainland China (province-level).