Systematic Review and Meta-analysis of the Effectiveness and Safety of Hydroxychloroquine in COVID-19.

Tzu-Han Yang[1,a], Chian-Yin Chou[1,2,a], Yi-Fan Yang[3], Yi-Ping Yang[4,5], Chian-Shiu Chien[4,5], Aliaksandr A. Yarmishyn[4,5], Tzu-Ying Yang[6], Shih-Hwa Chiou[4,5], Yuh-Lih Chang[1,2,7]

[1] Department of Pharmacy, Taipei Veterans General Hospital, Taipei, Taiwan.

[2] Faculty of Pharmacy, School of Pharmaceutical Sciences, National Yang-Ming University, Taipei, Taiwan.

[3] Department of Internal Medicine, National Taiwan University Hospital, Taipei, Taiwan

[4] Department of Ophthalmology, Taipei Veterans General Hospital, Taipei, Taiwan.

[5] Department of Medical Research, Taipei Veterans General Hospital, Taipei, Taiwan

[6] Department of dietetics and nutrition, Taipei Veterans General Hospital, Taipei, Taiwan.

[7] Department and Institute of Pharmacology, National Yang-Ming University, Taipei, Taiwan.

[a] Both authors contributed equally

All authors take responsibility for all aspects of the reliability and freedom from bias of

the data presented and their discussed interpretation.

Correspondence: Dr. Yuh-Lih Chang, Department of Pharmacy, Taipei Veterans

General Hospital, 201, Section 2, Shi-Pai Road, Taipei 112, Taiwan

(ylchang@vghtpe.gov.tw).

Declaration of interest: The authors report no conflict of interest.

Keywords: COVID-19; SARS-CoV-2; hydroxychloroquine; azithromycin; systematic

review.

Paper length: 3779 words   Title word count: 13   Abstract word count: 248

Take home: This study demonstrates that the use of hydroxychloroquine with or without

azithromycin might have benefits in positive-to-negative conversion of SARS-CoV-2 and

reduction of progression rate, but was associated with increased mortality in COVID-19.

Running title: Hydroxychloroquine for treating COVID-19

Abstract.

Backgrounds.

Since COVID-19 outbreak, various agents have been tested but no proven effective therapies have been identified. This has led to a lot of controversies among associated researches. Hence, in order to address the issue of using hydroxychloroquine in treating COVID-19 patients, we conducted a systematic review and meta-analysis.

Methods.

A thorough search was carried out to find relevant studies in MEDLINE, medRxiv, PubMed, Cochrane Database, China Academic Journals Full-text Database and Web of Science. Two investigators independently reviewed 274 abstracts and 23 articles. The trials which evaluated hydroxychloroquine for treatment of COVID-19 were included for this systematic review. Two investigators assessed quality of the studies and data extraction was done by one reviewer and cross checked by the other.

Results.

Five trials involving 677 patients were included while conducting the meta-analysis. Compared with the control group, hydroxychloroquine with or without azithromycin showed benefits in positive-to-negative conversion of SARS-CoV-2 (odds ratio [OR], 1.95 [95% CI,0.19 to 19.73] and a reduction in progression rate (OR, 0.89 [95% CI, 0.58 to 1.37]), but without demonstrating any statistical significance. This systematic review has also suggested a possible synergistic effect of the combination therapy which included hydroxychloroquine and azithromycin. However, the use of hydroxychloroquine alone

was associated with increased mortality in COVID-19 patients.

Conclusion.

The use of hydroxychloroquine with or without azithromycin for treatment of COVID-19 patients, seems to be effective. The combination of hydroxychloroquine and azithromycin has shown synergic effects. However, mortality rate was increased when the treatment was conducted with hydroxychloroquine.

Keywords.

COVID-19; SARS-CoV-2; hydroxychloroquine; azithromycin; systematic review

INTRODUCTION

In December 2019, the sudden outbreak of newly detected coronavirus (SARS-CoV-2) resulted in spread of coronavirus disease (COVID-19) to other countries, leading to a global pandemic. This virus was initially reported in the Hubei province, Wuhan, China [1]. As of April 27, 2020, more than 3 million people have been rapidly affected by COVID-19 worldwide and over two hundred thousand people have died in more than 200 countries due to this deadly disease [2]. As COVID-19 is spreading at a fast pace therefore without wasting time researchers are working tirelessly to find an effective treatment for this disease. However, despite all the efforts there is no proven, effective, pharmacological treatment available currently except following infection control and preventive strategies such as personal protective measures and social distancing,.

Various active clinical trials are underway to detect effectiveness of ribavirin, remdesivir, lopinavir/ritonavir and hydroxychloroquine for COVID-19 treatment. The pathogenic mechanisms of SARS-CoV-2 might be related to its impact on immune system since SARS-CoV-2 can infect T cells [3]. Therefore, recently conducted studies have drawn a lot of attention to the possible benefit of hydroxychloroquine, an antimalarial agent with an immunomodulatory activity which has been used at a large-scale for the treatment of rheumatic disease mainly because of its relatively low price and considerable safety.

Colson *et al.* has speculated that hydroxychloroquine might be the choice in treating COVID-19 patients [4]. Some *in vitro* studies have demonstrated that hydroxychloroquine has a potent antiviral activity against SARS-CoV-2 [5, 6]. Based on the outcome seen in some *in vitro* studies, numerous clinical studies were launched but only few of these which were published, presented relevant results and highlighted the efficacy of different treatment modalities [7-15]. Additionally, a lot of controversial findings are noticed among the researches. A study conducted by Gautret *et al.* has supported that hydroxychloroquine is effective in reducing viral load [8], while there are a few studies which disagree with this [7]. Some research studies have even reported harmful effects of hydroxychloroquine [10, 16]. Moreover, most of the published studies had small sample size [7-9, 13] or conducted a non-randomized clinical trial (non-RCT), which are major limitations in deriving a sound outcome [8, 10-12, 14, 15]. Many review articles which have discussed the use of hydroxychloroquine in COVID-19 patients showed limitation as they included only few relevant studies, significant Chinese studies were omitted and analysis of treatment effects was not quantified while formulating the results [17-19]. Recently, two systematic review with meta-analysis were published which had small sample size [20] and there was no clear clarity with regard to the effects of using hydroxychloroquine alone, or when combined with azithromycin, for the treatment of COVID-19 [21]. These systematic review did not include any national trials

and also excluded population of non-Hispanic black while conducting their research [20, 21].

On April 24, FDA issued a warning against use of hydroxychloroquine and chloroquine for treating patients with COVID-19 due to its serious adverse effect, such as heart arrhythmias being reported in some patients. Therefore, efficacy and safety of hydroxychloroquine for patient with COVID-19 is still uncertain. In view of severe outbreak of COVID-19 and rapid increase of studies being conducted, it is crucial to systematically evaluate the relevant studies in a rigorous manner. Hence, we conducted a systematic review and meta-analysis to demonstrate the significance of present evidence regarding benefits and safety of hydroxychloroquine use, for the treatment of COVID-19 patients.

METHODS

Data Search

We systematically searched the MEDLINE, medRxiv, PubMed, China Academic Journals Full-text Database, Cochrane database and Web of Science database up till April 27, 2020, with no language restrictions taken into consideration. In addition, relevant articles were also obtained from the reference list of reports which were identified using

this search strategy. The main key words included the following terms, "COVID-19", "SARS-CoV-2" and "hydroxychloroquine".

Study Selection

Two reviewers reviewed 251 abstracts and 23 full-text articles independently. Our study included only comparative studies that examined the efficacy or safety of hydroxychloroquine with or without azithromycin in comparison with standard treatment based on factors such as virological cure, degree of progression to severe illness and all-cause mortality as identified in COVID-19 patients. Out of all the selected search material, 257 articles were excluded due to their inappropriate study design (review, editorial News and letter). Among these, we also excluded a RCT as the results showed that 8 % (n=6) of the patients assigned to hydroxychloroquine group did not receive any dose of hydroxychloroquine and this might have caused misclassification bias in the study [22].

Quality Assessment

One reviewer extracted study-level data into standardized working tables, and the other checked data accuracy. Two independent reviewers critically appraised the eligible articles by using the Modified Downs and Black risk assessment scale [23]. This scale consists of 27 items which assesses different study characteristics such as internal validity,

statistical power and external validity. Publication bias was also assessed by using a funnel plot.

Data Synthesis and Statistical Analysis

The efficacy outcome of interest in this article was virological clearance and progression of disease. The safety outcome which we planned to investigate through this review, was mortality. We analyzed the main beneficial outcome based on the consistency and availability of results which were reported across various recruited trials such as the level of virological clearance noticed after the treatment, which was defined as a negative result of SARS-COV-2 after detecting it by real-time reverse transcription-PCR. We performed the meta-analysis by using Mantel Hazel method for dichotomous data. In our present study, either fixed-effect model or random effect model was used according to the level of heterogeneity. The odds ratio (OR) was used as the common measure of association across studies. We assessed statistical heterogeneity through the I^2 index. Meta-analysis was undertaken by using Cochrane Collaboration Review Manager (RevMan) software version 5.3.

Subgroup Analysis

Subgroup analysis was carried out by including the clinical status (symptomatic or

asymptomatic) and age so as to assess the impact of these variables on outcome.

RESULTS

Description of Included Trials

After selecting the relevant articles we found five eligible comparative trials that focused

on the benefits of hydroxychloroquine alone or in combination with azithromycin, for the

treatment of COVID-19. There were a total of 677 participants, and each trial included in

this research had sample size ranging from 30 to 368 [7-11]. Supplementary table 1 shows

the main characteristics of the eligible studies included in the research. Follow-up

duration of included trials ranged from 6 to 7 days. Among the included trials, two trials

which were conducted in China studied hydroxychloroquine effect at a dose of 400 mg

when given daily for 5 days [7, 9] and other French trial studied hydroxychloroquine

outcome at a doses of 600 mg given daily with or without azithromycin (500 mg at day 1

and 250 mg at days 2 to 5) [8]. In addition, another French trial tested hydroxychloroquine

effect when given at a dose of 600 mg daily [11] and another trial investigated benefits of

hydroxychloroquine, with or without azithromycin but dosage was not mentioned in the

research [10]. Among the five trials which were included, four trials mentioned their

exclusion criteria as - the patients who had known allergy to hydroxychloroquine or any

other known contraindication in terms of usage of the study drugs, such as retinopathy or

pregnancy [7-9, 11]. However, one trial was conducted without considering any exclusion criteria [10] and where the average age of participants was reported to be 62.6 years.

Quality of Included Studies

The risk of bias assessment is reported in supplementary figure 1. The average Downs and Black score was 19, with a range between 18 and 22 (A higher score indicates less bias).Various studies have shown different degree of bias wherein the RCT conducted by Chen *et al*. has displayed the lowest risk of bias [9] and the open-label non-RCT had shown the highest risk of bias [8]. One Chinese trial was described as randomized, although the method of randomization, allocation concealment and blinding of assessors during outcome evaluation was not mentioned [7] whereas another randomized control trial properly described the method of randomization and double-blinding [9]. With regard to the study design, one French trial was designed as open-label non- randomized trial [8] and two other studies were planned as controlled retrospective studies [10, 11]. A trial reported a loss of follow-up by participants while undergoing treatment hence they were excluded during analysis planning due to their withdrawal from the study [8].

Effect on Antiviral Clearance

Antiviral clearance was calculated using results obtained from PCR which were

documented as SARS-CoV-2 carriage during the follow-up period after treatment was over. For conducting the meta-analysis, we used one-time observation of negative results seen in PCR which were carried forward during analysis. Two studies have reported treatment effects based on antiviral clearance [7, 8].

Only two trials were included for meta-analysis, assessing a total of 66 participants. We found no significant evidence to verify whether hydroxychloroquine with or without azithromycin used for the treatment was effective in reducing viral carriage (OR 1.95 [95% CI 0.19 to 19.73]). There was substantial and statistically significant heterogeneity present while conducting this analysis ($I^2 = 62\%$) (figure 2A). As for the treatment undertaken by using hydroxychloroquine alone, the meta-analysis showed an increment by 1.74 times as compared with control group (95% CI 0.51 to 5.91; $I^2 = 35\%$) while evaluating the viral clearance (figure 2B).

One trial compared the effectiveness of combination therapy of hydroxychloroquine and azithromycin with control group and the results indicated that there was a significant reduction in viral load when combination therapy was used (OR 27.18 [95% CI 1.29 to 574.32]) (figure 2C) [8]. Based on intention-to-treat method, we also included 6 patients in the research who did not follow up after the treatment was over during the French trial

[8]. The pooled analysis implied a trend that treatment by hydroxychloroquine with or without azithromycin was effective in reducing viral load in Covid-19 patients (OR 1.72 [95% CI 0.26 to 11.50]; $I^2 =$49%) (figure 2D). A statistically significant reduction of the viral carriage was noticed at D6-post inclusion while performing a French trial [8]. However, another trial conducted in Chinese patients indicated a shift towards favoring the control group in reducing viral load at D7-post inclusion [7]. A Chinese study reported that the use of hydroxychloroquine as compared to the control group where only standard treatment was followed, could shorten clinical recovery time (TTCR) and promote development of pneumonia (CT) [9].

Subgroups Analysis

Subgroup analysis was undertaken to explore the impact of different factors relating to the context of the study that may influence the effectiveness of the intervention including, clinical status and age.

Clinical status

A recent trial conducted by Gautret *et al.* which included 6 asymptomatic COVID-19 patients and 60 patients who showed symptoms of upper respiratory tract infection (URTI) or lower respiratory tract infection (LRTI) [8]. Another trial conducted in China included

30 moderately affected COVID-19 patients who displayed symptoms such as fever or respiratory tract infection [7]. Subgroup analysis performed according to baseline clinical status of patients did not demonstrate any difference related to the effect of the intervention. A meta-analysis which included patients with symptomatic clinical status (e.g. URTI or LRTI) and had used hydroxychloroquine alone or in combination with azithromycin for treatment, showed no significant evidence in the effect of the viral elimination as seen by the value of odd's ratio (OR 3.33 [95% CI 0.08 to 146.36]; I^2 =80%) (supplementary figure 2A). Among the selected studies there were two studies that included symptomatic patients who used hydroxychloroquine monotherapy [7, 8]. When meta-analysis was conducted by including these two trials, no significant difference between intervention and control groups was seen, in addition substantial statistical heterogeneity was identified ($I^2 = 69\%$) (supplementary figure 2B). A meta-analysis has reported that in patients with symptomatic clinical status, usage of hydroxychloroquine along with azithromycin for treatment of COVID-19 has showed beneficial outcome with regard to seroconversion (OR 99.67 [95% CI 3.52 to 2818.12]) (supplementary figure 2C).

Age

Out of all the included trials in our present research, one trial [8] has included only adults

while the other trial [7] has included 4 participants younger than 18 years-old. To examine the treatment effects on adults, we excluded those 4 patients and conducted subgroup analysis. Data extracted from adult-specific subgroup analysis showed limited benefit of using hydroxychloroquine alone or in combination with azithromycin for the purpose of viral elimination (OR 3.59 [95% CI 0.07 to 183.82]; I^2=82%) (supplementary figure 3A). The result also showed that there was no significant evidence that could support that hydroxychloroquine, when used without azithromycin for the treatment, was effective in reducing viral carriage in adults (OR 2.71 [95% CI 0.09 to 80.15]; I^2=75%) (supplementary figure 3B). A pooled analysis of adult patients highlighted that using hydroxychloroquine plus azithromycin for the treatment showed significant benefits in promoting viral seroconversion (OR 99.67 [95% CI 3.52 to 2818.12]) (supplementary figure 3C).

Antiviral Effect of Hydroxychloroquine Monotherapy or Azithromycin-based Combination Therapy

We conducted another meta-analysis to compare the antiviral effects of using hydroxychloroquine alone, or when combined with azithromycin and the results showed that there is an inclination of more benefits being detected with regard to reduction in viral load in the combination group (OR 9.94 [95% CI 0.47 to 210.41]) (figure 3).

Progression to Severe Illness

During the research period, progression to severe illness as seen in COVID-19 patients in association with the study drug was noticed in all the five included trials [7-11]. Although, quantitatively lower rate of progression to severe illness was reported in the hydroxychloroquine usage groups, whether with or without azithromycin combination, but there was no significant difference documented when compared with control groups (OR 0.89 [95% CI 0.58 to 1.37]; I^2 =30%) (figure 4A). We analyzed the efficacy of treatment in association with worsening of disease and the research findings showed a more favorable outcome in the study drug groups. (figure 4B and 4C).

All-cause Mortality

For analyzing all-cause mortality, we included three trials which had relevant results associated with regard to mortality [7, 8, 10]. The research findings concluded that hydroxychloroquine use, whether in combination with azithromycin or not, was associated with increased mortality in COVID-19 patients (figure 5A-C)

Death or Progression to Severe Illness

The overall pooled OR was 1.00 for death or progression to severe illness in patients

treated with hydroxychloroquine, with or without azithromycin versus control groups for four studies that had complete data regarding the same [7-9, 11] (95% CI 0.27-3.75, I^2 =33%) (figure 6A). However, the pooled OR was 1.37 when a comparison of hydroxychloroquine alone versus the control group was conducted (95% CI 0.09-21.97, I^2 =59%) (figure 6B).

Publication Bias

In total, five trials were included in the present study and were appropriately evaluated for publication bias by conducting funnel plot analysis. As shown in supplementary figure 4, there was no obvious publication bias reported.

DISCUSSION

To solve the COVID-19 pandemic, it is urgently required to find effective pharmacological agents. Hydroxychloroquine is a repositioning candidate, still with reports of unclear efficacy and safety issue when used for the treatment of COVID-19. Therefore, we conducted a systematic review and meta-analysis to address not only the efficacy and safety issue of hydroxychloroquine, but also to explore its efficacy when used in combination with azithromycin for treating COVID-19 patients. The present study yielded several important findings. Firstly, hydroxychloroquine might show benefits in virological clearance of SARS-CoV-2 and a reduction in the risk of disease progression, although no statistical significant findings were reported. Secondly, the combination therapy of hydroxychloroquine and azithromycin seems to show a synergistic effect. Thirdly, use of hydroxychloroquine was associated with an increased mortality. To the best of our knowledge, this study is the first systematic review and meta-analysis which has addressed the issue of combining hydroxychloroquine and azithromycin in COVID-19 patients, by summarizing available results extracted from clinical trials.

Our results have suggested that hydroxychloroquine use, whether in the form of monotherapy or combined with azithromycin, has shown a beneficial response in conversion of nucleic acid in COVID-19 patients as compared to the control groups.

Consistent findings were also noted in groups treated with hydroxychloroquine alone or groups where hydroxychloroquine and azithromycin-based combination therapy was used. While conducting subgroup analysis with regard to symptomatic and adult subpopulations, no statistically significant difference was found between drug treatment group and control group in terms of virological cure. These results were based on the outcome reported in only two trials which had their own limitations such as small sample size (66 patients), short observation period (6-7 days) and short duration of drug treatment (5-10 days) [7, 8]. Another small case series which included more severely infected COVID-19 patients reported an unfavorable viral elimination with the combined use of hydroxychloroquine and azithromycin [13]. However, Gautret *et al.* conducted a similar study which included 80 patients and reported that hydroxychloroquine when co-administered with azithromycin displayed a prompt reduction in viral load, with nearly 80-90% negative cases reported at Day7-8 [12]. Another cohort study which took place in France, included 1061 new COVID-19 patients who were treated with hydroxychloroquine and azithromycin. The study results showed excellent clinical outcome and virological cure in 973 patients on day 10 (91.7%) with death reported in few patients (0.75%) [15]. A study which was conducted in Iraq reported that 79 patients were treated with a protocol which included hydroxychloroquine and other antiviral drugs for dealing with diverse COVID-19 categories, showed a promising rate of complete

clinical recovery [14]. In the present study, we found that hydroxychloroquine monotherapy might decrease COVID-19 patients' progression to worse illness but without any reports of statistical significance. Although our results showed higher rate of viral elimination and lower rate of disease worsening in patients treated with combination of hydroxychloroquine and azithromycin, there was no significant difference in the outcome when compared with control groups.

The anti-inflammatory effect of hydroxychloroquine on the immune system has been well recognized. The antiviral mechanisms of hydroxychloroquine in treating COVID-19 patients has also been discovered, including increasing pH values of endosomes or lysosomes, stopping the glycosylation of angiotensin converting enzyme (ACE) 2 receptor, inhibiting antigen processing and following major histocompatibility complex (MHC)-II presentation to T cells and interrupting the interaction between cytosolic viral DNA/RNA and toll-like receptors (TLRs) [24, 25]. Azithromycin, a member of the macrolide family, has antibacterial, anti-inflammatory, and antiviral properties [26, 27]. Although azithromycin has been found to block influenza virus internalization into host cells, its mechanism of action against SARS-CoV-2 is still uncertain [28]. Regarding the antiviral effects of using combination of hydroxychloroquine and azithromycin, Andreani *et al.* demonstrated that such combination showed a synergistic effect against SARS-CoV-

2 when research was conducted *in vitro* [29].

In terms of safety issue, two RCT trials and one case series reported mild adverse effects such as rash and diarrhea among patients [7, 9, 11]. . One comparative study which has used hydroxychloroquine alone, found that 9.5% of patients experienced ECG change, leading to its discontinuation [11]. Another case series which was conducted in France, involving 11 patients reported that one patient developed QT prolongation [13]. Chorin *et al*. also found that 11% of patient receiving combination therapy (hydroxychloroquine and azithromycin) suffered from severe QT prolongation (>300 milliseconds) [16]. Notably, reports have suggested that both azithromycin and hydroxychloroquine are associated with QT prolongation, and their combined use may potentiate this adverse effect [30, 31].

In addition, we found that the incidence of mortality favored control group rather than the use of hydroxychloroquine with or without azithromycin as demonstrated with an OR of 2.54 in the pooled analysis. Most of the included trials had small sample size which might have impacted the significance of our results [8, 9]. Among the included studies, one of them investigated by Magagnoli *et al*. had few findings which were different from the rest. Here, subjects with contraindication to usage of hydroxychloroquine or had

significant comorbidities were not excluded from the patients population, comprised only men whose median age was over 65 years, and majorly who were non-Hispanic black [10]. Reports have indicated that the average age of COVID-19 patients was about 50 years, and the percentage distribution among male and female was nearly equal [30-32] Since the mortality risk for COVID-19 varies by age, the current evidence suggests that older people are at a higher risk of getting affected from COVID-19 than younger ones [33, 34] and it might be related to multiple comorbidities commonly seen in older people [35]. Higher rates of COVID-19 -related hospitalization among the elder population and non-Hispanic black have also been found in the United States [36]. The mortality rate in different countries might be also associated with socio-economic and political factors [37]. Therefore, our present results of pooled analysis might be biased as we also included the demographic composition of patients, mentioned in the study conducted by Magagnoli *et al.*

Despite the pertinent information demonstrated above, our systematic review has its own limitations. As far as the lack of compiling sufficient data with regard to virological cure as efficacy outcome is concerned, we tried to solve this problem by conducting a search for unpublished literature and non-English journals to increase the quantity of relevant integrated literature.. Most studies which are published are mainly observational studies

and case series with substantial methodology concerns such as no framework of exclusion criteria, and therefore susceptible to bias and confounding. Therefore, as there are limited related RCTs in the literature at present, only those were included in our study. In the lack of experimental random allocation to the intervention, the effects of the hydroxychloroquine might be overestimated. Furthermore, the two single-center RCTs which were conducted in China had some limitations as it included small sample size, did not conduct an international trial, and only limited data was available regarding the viral load. In our present study, bias might be present since little information was provided on demographic data, comorbidity, severity of disease, and the adverse effects of hydroxychloroquine or azithromycin such as QT prolongation. In addition, we have not performed a dose–response meta-analysis, due to the lack of original data. Finally, the limited number of available studies which were included, made it unworkable to evaluate the true effect of hydroxychloroquine. Therefore, the present results should be interpreted cautiously, and further large, prospective RCTs needs to be conducted in a methodologically rigorous manner.

CONCLUSION

The present study has suggested that hydroxychloroquine alone, or in combination with azithromycin for COVID-19 treatment showed beneficial outcome but also higher

mortality. The combination of hydroxychloroquine and azithromycin has demonstrated synergistic effects. In future, large, randomized, placebo-controlled trials with longer follow-up are needed to further confirm these findings.

ACKNOWLEDGMENTS

None.

DISCLOSURES

None.

FUNDING

REFERENCES

1. Steven S, Yen Ting L, Chonggang X, et al. High Contagiousness and Rapid Spread of Severe Acute Respiratory Syndrome Coronavirus 2. *Emerg Infect Dis* 2020: 26(7).

2. Worldometers. COVID-19 coronavirus pandemic. 2020: Available from: https://www.worldometers.info/coronavirus/.

3. Wang X, Xu W, Hu G, et al. SARS-CoV-2 infects T lymphocytes through its spike protein-mediated membrane fusion. *Cell Mol Immunol* 2020.

4. Colson P, Rolain J, Lagier J, et al. Chloroquine and hydroxychloroquine as available weapons to fight COVID-19. *Int J Antimicrob Agents* 2020: 55(4): 105932.

5. Yao X, Ye F, Zhang M, et al. In Vitro Antiviral Activity and Projection of Optimized Dosing Design of Hydroxychloroquine for the Treatment of Severe Acute Respiratory Syndrome Coronavirus 2 (SARS-CoV-2). *Clin Infect Dis* 2020.

6. Liu J, Cao R, Xu M, et al. Hydroxychloroquine, a less toxic derivative of chloroquine, is effective in inhibiting SARS-CoV-2 infection in vitro. *Cell discovery* 2020: 6: 16.

7. Chen J, Liu D, Liu L, et al. A pilot study of hydroxychloroquine in treatment of patients with common coronavirus disease-19 (COVID-19). *J Zhejiang Univ (Med Sci)* 2020: Published online March 6, 2020. doi:10.3785/j.issn.1008-9292.2020.03.03.

8. Gautret P, Lagier J, Parola P, et al. Hydroxychloroquine and azithromycin as a treatment of COVID-19: results of an open-label non-randomized clinical trial. *Int J Antimicrob Agents* 2020: Published online March 20, 2020. doi:10.1016/j. ijantimicag.2020.105949: 105949.

9. Chen Z, Hu H, Zhang Z, et al. Efficacy of hydroxychloroquine in patients with COVID-19: results of a randomized clinical trial. *medRxiv* 2020: preprint. [https://doi.org/10.1101/2020.1103.1122.20040758].

10. Magagnoli J, Narendran S, Pereira F, et al. Outcomes of hydroxychloroquine usage in United States veterans hospitalized with Covid-19. *medRxiv* 2020: preprint. [https://doi.org/10.1101/2020.1104.1116.20065920].

11. Mahevas M, Tran V, Roumier M, et al. No evidence of clinical efficacy of hydroxychloroquine in patients hospitalized for COVID-19 infection with oxygen requirement: results of a study using routinely collected data to emulate a target trial. *medRxiv* 2020: preprint. [https://doi.org/10.1101/2020.1104.1110.20060699].

12. Gautret P, Lagier J, Parola P, et al. Clinical and microbiological effect of a

combination of hydroxychloroquine and azithromycin in 80 COVID-19 patients with at least a six-day follow up: A pilot observational study. *Travel Med Infect Dis* 2020: 101663.

13. Molina J, Delaugerre C, Le Goff J, et al. No Evidence of Rapid Antiviral Clearance or Clinical Benefit with the Combination of Hydroxychloroquine and Azithromycin in Patients with Severe COVID-19 Infection. *Med Mal Infect* 2020.

14. Allawi J, Abbas H, Rasheed J, et al. The first 40-days experience and clinical outcomes in the management of coronavirus covid-19 crisis. Single center preliminary study. *Journal of the Faculty of Medicine Baghdad* 2020: 61(3,4): https://doi.org/10.32007/j fac med bagdad.%y.32613,32004%p.

15. Million M, Lagier J-C, Gautret P, et al. Full-length title: Early treatment of COVID-19 patients with hydroxychloroquine and azithromycin: A retrospective analysis of 1061 cases in Marseille, France. *Travel Med Infect Dis* 2020: 101738.

16. Chorin E, Dai M, Shulman E, et al. The QT Interval in Patients with SARS-CoV-2 Infection Treated with Hydroxychloroquine/Azithromycin. *medRxiv* 2020: preprint. [https://doi.org/10.1101/2020.1104.1102.20047050].

17. Gbinigie K, Frie K. Should chloroquine and hydroxychloroquine be used to treat COVID-19? A rapid review. *BJGP Open* 2020: bjgpopen20X101069.

18. Singh A, Singh A, Shaikh A, et al. Chloroquine and hydroxychloroquine in the treatment of COVID-19 with or without diabetes: A systematic search and a narrative review with a special reference to India and other developing countries. *Diabetes Metab Syndr* 2020: 14(3): 241-246.

19. Kapoor K, Kapoor A. Role of Chloroquine and Hydroxychloroquine in the Treatment of COVID-19 Infection- A Systematic Literature Review. *medRxiv* 2020: preprint. [https://doi.org/10.1101/2020.1103.1124.20042366].

20. Sarma P, Kaur H, Kumar H, et al. Virological and Clinical Cure in Covid-19 Patients Treated with Hydroxychloroquine: A Systematic Review and Meta-Analysis. *J Med Virol* 2020.

21. Shamshirian A, Hessami A, Heydari K, et al. Hydroxychloroquine Versus COVID-19: A Rapid Systematic Review and Meta-Analysis. *medRxiv* 2020: preprint. [https://doi.org/10.1101/2020.1104.1114.20065276].

22. Tang W, Cao Z, Han M, et al. Hydroxychloroquine in patients with COVID-19: an open-label, randomized, controlled trial. *medRxiv* 2020: preprint. [https://doi.org/10.1101/2020.1104.1110.20060558].

23. Downs S, Black N. The feasibility of creating a checklist for the assessment of the methodological quality both of randomised and non-randomised studies of health care interventions. *J Epidemiol Community Health* 1998: 52(6): 377-384.

24. Alia E, Grant-Kels JM. Does Hydroxychloroquine Combat COVID-19? A Timeline of Evidence. *J Am Acad Dermatol* 2020.

25. Sinha N, Balayla G. Hydroxychloroquine and covid-19. *Postgrad Med J* 2020.

26. Schogler A, Kopf BS, Edwards MR, et al. Novel antiviral properties of azithromycin in cystic fibrosis airway epithelial cells. *Eur Respir J* 2015: 45(2): 428-439.

27. Gielen V, Johnston SL, Edwards MR. Azithromycin induces anti-viral responses in bronchial epithelial cells. *Eur Respir J* 2010: 36(3): 646-654.

28. Ohe M, Shida H, Jodo S, et al. Macrolide treatment for COVID-19: Will this be the way forward? *Biosci Trends* 2020.

29. Andreani J, Le Bideau M, Duflot I, et al. In vitro testing of combined hydroxychloroquine and azithromycin on SARS-CoV-2 shows synergistic effect. *Microb Pathog* 2020: 145: 104228.

30. Mercuro NJ, Yen CF, Shim DJ, et al. Risk of QT Interval Prolongation Associated With Use of Hydroxychloroquine With or Without Concomitant Azithromycin Among Hospitalized Patients Testing Positive for Coronavirus Disease 2019 (COVID-19). *JAMA cardiology* 2020.

31. Bessiere F, Roccia H, Deliniere A, et al. Assessment of QT Intervals in a Case Series of Patients With Coronavirus Disease 2019 (COVID-19) Infection Treated With Hydroxychloroquine Alone or in Combination With Azithromycin in an Intensive Care Unit. *JAMA cardiology* 2020.

32. Wu J, Liu J, Zhao X, et al. Clinical Characteristics of Imported Cases of Coronavirus Disease 2019 (COVID-19) in Jiangsu Province: A Multicenter Descriptive Study. *Clin Infect Dis* 2020.

33. Zhai P, Ding Y, Wu X, et al. The epidemiology, diagnosis and treatment of COVID-19. *Int J Antimicrob Agents* 2020: 105955.

34. ChinaCDC. The epidemiological characteristics of an outbreak of 2019 novel coronavirus diseases (COVID-19) in China. *Zhonghua Liu Xing Bing Xue Za Zhi* 2020: 41(2): 145-151.

35. Zhou F, Yu T, Du R, et al. Clinical course and risk factors for mortality of adult inpatients with COVID-19 in Wuhan, China: a retrospective cohort study. *Lancet* 2020: 395(10229): 1054-1062.

36. Garg S, Kim L, Whitaker M. Hospitalization Rates and Characteristics of Patients Hospitalized with Laboratory-Confirmed Coronavirus Disease 2019 — COVID-NET, 14 States, March 1–30, 2020. *MMWR Morb Mortal Wkly Rep* 2020(69): 458–464.

37. Sorci G, Faivre B, Morand S. Why does COVID-19 case fatality rate vary among

countries? *medRxiv* 2020: preprint.
[https://doi.org/10.1101/2020.1104.1117.20069393].

Figure 1. Prisma flow chart of included studies

Figure 2. Seroconversion of SARS-CoV-2. (HCQ ± AZI: hydroxychloroquine with or without azithromycin; HCQ + AZI: combination of hydroxychloroquine and azithromycin)

Figure 3. Seroconversion of SARS-CoV-2 (Hydroxychloroquine plus azithromycin versus hydroxychloroquine alone).

(A) Progression to severe illness (HCQ ± AZI vs. control)

Footnotes
(1) We included all HCQ treated patients using intention-to-treat method (20 HCQ with or without AZI + 6 lost to follow up)
(2) Outcomes based on pre-ventilation treatment
(3) Disease progression was defined as acute respiratory distress syndrome on day 7 (using weighted proportion data)

(B) Progression to severe illness (HCQ alone vs. control)

Footnotes
(1) We included all HCQ treated patients using intention-to-treat method (14 HCQ + 6 lost to follow up)
(2) Outcomes based on pre-ventilation treatment

(C) Progression to severe illness (HCQ + AZI vs. control)

Footnotes
(1) Outcomes based on pre-ventilation treatment

Figure 4. Progression to severe illness.

(A) All-cause mortality (HCQ ± AZI vs. control)

Study or Subgroup	HCQ ± AZI Events	HCQ ± AZI Total	Control Events	Control Total	Weight	Odds Ratio M-H, Fixed, 95% CI
Chen J 2020	0	15	0	15		Not estimable
Gautret P 2020 (1)	1	26	0	16	3.6%	1.94 [0.07, 50.56]
Magagnoli J 2020	52	210	18	158	96.4%	2.56 [1.43, 4.58]
Total (95% CI)		251		189	100.0%	2.54 [1.43, 4.50]
Total events	53		18			

Heterogeneity: Chi² = 0.03, df = 1 (P = 0.87); I² = 0%
Test for overall effect: Z = 3.18 (P = 0.001)

Favours HCQ ± AZI   Favours control

Footnotes
(1) We included all HCQ treated patients using intention-to-treat method (20 HCQ with or without AZI + 6 lost to follow up)

(B) All-cause mortality (HCQ alone vs. control)

Study or Subgroup	HCQ Events	HCQ Total	Control Events	Control Total	Weight	Odds Ratio M-H, Fixed, 95% CI
Chen J 2020	0	15	0	15		Not estimable
Gautret P 2020 (1)	1	20	0	16	4.9%	2.54 [0.10, 66.59]
Magagnoli J 2020 (2)	27	97	18	158	95.1%	3.00 [1.55, 5.81]
Total (95% CI)		132		189	100.0%	2.98 [1.56, 5.70]
Total events	28		18			

Heterogeneity: Chi² = 0.01, df = 1 (P = 0.92); I² = 0%
Test for overall effect: Z = 3.29 (P = 0.0010)

Favours HCQ   Favours control

Footnotes
(1) We included all HCQ treated patients using intention-to-treat method (14 HCQ + 6 lost to follow up)
(2) Outcomes based on treatment exposure

(C) All-cause mortality (HCQ + AZI vs. control)

Study or Subgroup	HCQ + AZI Events	HCQ + AZI Total	Control Events	Control Total	Weight	Odds Ratio M-H, Random, 95% CI
Gautret P 2020	0	6	0	16		Not estimable
Magagnoli J 2020 (1)	25	113	18	158	100.0%	2.21 [1.14, 4.28]
Total (95% CI)		119		174	100.0%	2.21 [1.14, 4.28]
Total events	25		18			

Heterogeneity: Not applicable
Test for overall effect: Z = 2.35 (P = 0.02)

Favours HCQ + AZI   Favours control

Footnotes
(1) Outcomes based on treatment exposure

Figure 5. All-cause mortality.

(A) Death or clinical worsening of disease (HCQ ± AZI vs. control)

Study or Subgroup	HCQ ± AZI Events	Total	control Events	Total	Weight	Odds Ratio M-H, Random, 95% CI
Chen J 2020	1	15	0	15	13.1%	3.21 [0.12, 85.20]
Chen Z 2020	0	31	4	31	15.4%	0.10 [0.00, 1.88]
Gautret P 2020 (1)	4	26	0	16	15.2%	6.60 [0.33, 131.25]
Mahevas M 2020 (2)	16	84	19	89	56.3%	0.87 [0.41, 1.82]
Total (95% CI)		156		151	100.0%	1.00 [0.27, 3.75]
Total events	21		23			

Heterogeneity: Tau² = 0.66; Chi² = 4.46, df = 3 (P = 0.22); I² = 33%
Test for overall effect: Z = 0.00 (P = 1.00)

Favours HCQ ± AZI   Favours control

Footnotes
(1) We included all HCQ treated patients using intention-to-treat method (20 HCQ with or without AZI + 6 lost to follow up)
(2) Use weighted proportion data

(B) Death or clinical worsening of disease (HCQ alone vs. control)

Study or Subgroup	HCQ Events	Total	Control Events	Total	Weight	Odds Ratio M-H, Random, 95% CI
Chen J 2020	1	15	0	15	31.6%	3.21 [0.12, 85.20]
Chen Z 2020	0	31	4	31	34.4%	0.10 [0.00, 1.88]
Gautret P 2020 (1)	4	20	0	16	34.0%	9.00 [0.45, 180.84]
Total (95% CI)		66		62	100.0%	1.37 [0.09, 21.97]
Total events	5		4			

Heterogeneity: Tau² = 3.54; Chi² = 4.87, df = 2 (P = 0.09); I² = 59%
Test for overall effect: Z = 0.22 (P = 0.82)

Favours HCQ   Favours control

Footnotes
(1) We included all HCQ treated patients using intention-to-treat method (14 HCQ + 6 lost to follow up)

Figure 6. Death or clinical worsening of disease.