Title Page

Low testosterone levels in H7N9 infected men correlate with high cytokine

responses and lethal outcome: a retrospective cohort study

Yongkun Chen[1*], Tian Bai[2,3*], Sebastian Beck[2*], Stephanie Stanelle-Bertram[2], Tao Chen[3], Jie

Dong[3], Jing Yang[3], Lijie Wang[3], Dayan Wang[3], Yuelong Shu[1,3,§], Gülsah Gabriel[2,4,§]

1   School of Public Health (Shenzhen), Sun Yat-sen University, Guangdong 510275, P.R. China

2   Viral Zoonoses-One Health, Heinrich Pette Institute, Leibniz Institute for Experimental Virology,

Hamburg, Germany

3   National Institute for Viral Disease Control and Prevention, Collaboration Innovation Center for

Diagnosis and Treatment of Infectious Diseases, Chinese Center for Disease Control and Prevention;

Key Laboratory for Medical Virology, National Health Commission of the People's Republic of China,

Beijing 102206, P.R. China

4   Institute of Virology, University of Veterinary Medicine, Hannover, Germany

*: equally contributed first authors

§: equally contributed senior authors

Correspondence to:

Gülsah Gabriel, PhD: guelsah.gabriel@leibniz-hpi.de or Yuelong Shu, PhD: shuylong@mail.sysu.edu.cn

Keywords: avian influenza A(H7N9) virus, testosterone, estradiol, cytokine, sex difference

Running title: Testosterone levels in H7N9 infection

Key point: Avian H7N9 infection presents a higher incidence in infected men compared to

women. Low testosterone levels in H7N9 infected men correlate with high-level cytokine

expression posing a high risk factor for lethal outcome.

Abstract

Background. Human infections with avian influenza A (H7N9) virus emerged in East China in March 2013. In contrast to seasonal influenza A viruses, H7N9 infections present a strong sex bias. Over the five epidemic waves in China, ~70% of all H7N9 cases were observed in men. We and others have independently shown that influenza A virus infections may significantly lower testosterone levels in male mice. In this study, we sought to analyze the role of testosterone in disease outcome in H7N9 infected men.

Methods. We systematically analyzed a total of n=330 human samples obtained from male and female subjects of two age groups (18-49 years and ≥50 years) with laboratory-confirmed H7N9 influenza, seasonal influenza (H1N1, H3N2) or negative control cohorts including H7N9 virus-negative close contacts and influenza virus-negative poultry workers in dependency of sex. The level of testosterone, estradiol as well as cytokines and chemokines were measured and compared in all study participants. We used regression analysis to evaluate the association of sex hormones and cytokines in dependency of sex and disease outcome.

Results. We analyzed sex hormones and innate immune responses in H7N9 cases (n=98), their H7N9 virus-negative close contacts (n=71), influenza virus-negative poultry workers (n=108) and mild seasonal (H1N1, H3N2) influenza cases (n=53). Samples were collected between 2014 and 2017. All control groups showed a similar median age within H7N9 cases except for the seasonal influenza group with a younger median age. In H7N9 infected men, testosterone levels were strongly reduced compared to male H7N9 virus-negative close contacts or males with seasonal influenza. Low testosterone levels in H7N9 infected men correlated with high inflammatory cytokine levels, e.g. IL-6, and lethal outcome in 18-49 year olds. No significant differences were detected in estradiol levels in H7N9 infected men. In H7N9 infected women (≥ 50 years), estradiol levels were elevated compared to H7N9 virus-negative close contacts without significantly affecting disease outcome.

Conclusions. Here, we show that low testosterone levels pose a poor prognostic marker in H7N9 infected men. Thus, treatment of H7N9 infected patients should consider sex-specific mitigation strategies.

Introduction

In early 2013, human infections with avian influenza A (H7N9) virus were first reported in China[1]. Since then, the H7N9 avian influenza A virus caused five epidemic waves from 2013 to 2017. Exposure at live poultry markets was considered a major risk factor for human infections with H7N9 virus. A primary epidemiological study revealed that 71% of all H7N9 cases were males in the first wave[2]. The high degree of sex bias in men consistently accounted with similar proportions (68%-71%) during the entire five H7N9 epidemic waves[3]. Currently, very little knowledge exists on the complex interplay of biological sex and gender specific behavior as contributors to influenza disease outcome. The World Health Organization (WHO) Western Pacific team proposed gender-associated practices and norms, such as more frequent exposure of men to birds, may pose one of the reasons for sex-specific H7N9 incidence[4]. However, a follow-up study suggested increased risk in older men is not due to higher exposure time to live poultry market[5].

Sex hormones, particularly testosterone and estradiol are important biological factors that affect sex differences in immune responses[6–9]. We have recently shown that H1N1 infection itself reduces testosterone levels in male but not female mice. Conversely, treatment of female mice with testosterone protected the majority of H1N1 infected females from lethal outcome[10]. In line, aged male mice with low testosterone levels were reported to undergo elevated pulmonary inflammation and severe disease upon H1N1 influenza virus infection compared to young male mice[11]. Conversely, other murine models showed that low concentrations of estradiol in female mice was associated with enhanced pro-inflammatory responses and H1N1 influenza pathogenesis[12]. In contrast, high estradiol levels protected female mice from lethal influenza infection by dampening inflammatory cytokine responses[12]. Thus, there is increasing evidence from small animal models that sex hormones play an important role in influenza disease outcome.

However, very little knowledge exists on the role of sex hormones in infectious diseases in humans and their association with disease outcome. Thus, in this study, we retrospectively analyzed the impact of sex hormones on influenza disease outcome in males and females.

98 **Materials and Methods**
99

Ethics statement

Sampling from laboratory-confirmed H7N9 avian influenza cases, seasonal influenza cases,

close contacts of H7N9 cases and poultry workers was reviewed and approved by the Ethics

Committee of the National Institute for Viral Disease Control and Prevention, China CDC.

Study participants

Stored serum samples from acute phase of laboratory-confirmed H7N9 hospitalized patients

were collected from 2014 to 2017 through the influenza surveillance network in the local

Center for Disease Control and Prevention (CDC), China. H7N9 patients (n=98) aged above

18 years with complete epidemiological information (age, sex, illness onset date, sample

collection date, outcome and antiviral treatment) and required amount of serum were enrolled

in this study. Once a patient was confirmed with H7N9 influenza virus, blood sample and

throat swabs of the close contacts and epidemiological linked poultry workers were collected

through the influenza surveillance network of the local CDC, China. Swabs were first

screened by real-time RT-PCR to identify influenza A and B viruses in the local CDC or the

Chinese National Influenza Center. Positive samples for influenza A virus were further tested

for influenza A (H7N9) virus. No influenza A (H7N9) virus was detected from close contacts

and poultry workers. A total of 71 close contacts and 108 poultry workers were included in

this study as control groups by matching age or sex of H7N9 cases. Archived plasma samples

from laboratory-confirmed outpatients infected with influenza A (H1N1) pandemic 2009 or

influenza A (H3N2) viruses were collected from 2015-2016 influenza season in China.

Plasma samples and throat swabs were collected simultaneously when the patient went to

outpatient care. Blood samples were archived when seasonal influenza virus was detected by

real-time RT-PCR. Seasonal influenza cases (2009 pandemic H1N1: n=22, H3N2: n=31)

were selected as another control group in this study. All samples from study participants were

stored at -80 °C until further analysis.

Hormone quantification

The total concentration of estradiol (17β-estradiol) and testosterone was measured in the

clinical laboratory of the Sino-Japan Friendship Hospital (Beijing, China) from collected

serum or plasma samples of all study participants using the Beckman Coulter DxI 800 Access

Immunoassay System. The majority of circulating estradiol and testosterone are bound to the

sex hormone binding globulin (SHBG), but it also exists loosely bound to albumin and in the free status. We used the Access Estradiol Assay (REF 33540, BECKMAN COULTER) and Access Testosterone Assay (REF 33560, BECKMAN COULTER) which use paramagnetic particles in a chemiluminescent immunoassay for the quantitative determination of total estradiol and testosterone level (bound and unbound).

Cytokine and chemokine measurement

Serum or plasma samples of H7N9 and seasonal influenza cases were analyzed for a panel of 29 cytokines and chemokines (EGF, Eotaxin, G-CSF, GM-CSF, IFN-α2, IFN-γ, IL-10, IL12-P40, IL12-P70, IL-13, IL-15, IL-17A, IL-1RA, IL-1α, IL-1β, IL-2, IL-3, IL-4, IL-5, IL-6, IL-7, IL-8, IP10, MCP-1, MIP-1α, MIP-1β, TNF-α, TNF-β, VEGF) by Luminex bead based multiplex assay using human cytokine/chemokine magnetic bead panel 96 well plate assay (HCYTOMAG-60K, Millipore, USA) according to manufacturer's instruction. A smaller panel including 21 selected cytokines and chemokines, where significant differences were observed earlier (GM-CSF, IFN-α2, IFN-γ, IL-10, IL12-P40, IL12-P70, IL-13, IL-17A, IL-1RA, IL-1α, IL-1β, IL-5, IL-6, IL-7, IL-8, IP10, MCP-1, MIP-1α, MIP-1 β, TNF-α, TNF-β) were analyzed for close contacts of H7N9 cases and poultry workers using the same measurements. For undetectable cytokines and chemokines (below the lowest detection limit) in all groups, we assigned a value as 0·1 for analysis.

Statistical analysis

Statistical significance was determined with GraphPad Prism 5 v.5·03 (GraphPad Software, Inc.). Difference of sex hormones, cytokines/chemokines among multiple groups was analyzed using One-Way ANOVA (Kruskal-Wallis test) and selected Dunn's as a post-test to compare each group. Since the results of cytokine/chemokines exhibited deviated scale among all groups, all values in the statistical analysis and figures were used after Ln-transformation. The difference of sex hormones and cytokines/chemokines between survival and dead H7N9 cases in each age group was analyzed using the unpaired, two-tailed Student's *t*-test (non-parametric). Association between sex hormones and cytokine/chemokines were determined using linear regression and correlation analysis (Pearson). Since the results of cytokine/chemokines exhibited deviated scale than testosterone, all values in the statistical analysis and figures were used after Ln-transformation in comparison of cytokines/chemokines in all groups and linear regression with testosterone.

Statistical significance was defined as $p < 0.05$ (* $p < 0.05$, ** $p < 0.01$,
*** $p < 0.001$).**Results**

Participant characteristics

A total of $n=44$ avian H7N9 influenza positive cases of reproductive age (18-49 years) were
enrolled with a median age of 42 years (IQR 34-46 years) (Table 1). A total of $n=54$ avian
H7N9 influenza positive cases were included in those ≥50 years old with the median age of
61 years (IQR 56-67 years) (Table 1). The male H7N9 cases accounted for 75% in the
younger and 70% in the older age groups, which is consistent with previous epidemiological
studies based on larger laboratory-confirmed H7N9 cohorts. Blood samples of H7N9 patients
were collected within acute phases of 6·5 days (IQR 5-8·3 days) and 7·5 days (IQR 6-9 days)
after illness onset. H7N9 cases confirmed in 2017 accounted for over 77% of all cases
enrolled since 2014, in this study, presenting most of the participants from South China (data
not shown). For all control groups, the median age is similar in close contacts and poultry
workers compared to H7N9 cases in both younger and older age groups (Table 1). However,
the age of seasonal influenza cases in younger age groups was lower compared to H7N9
cases. Consistent with H7N9 cases, the proportion of males in all control groups is higher
than females, except for the seasonal influenza cases with a similar proportion in both sexes.

Low testosterone levels in H7N9 infected men are associated with lethal outcome
In order to assess the role of sex hormones in the observed sex bias of H7N9 cases, we first
measured testosterone concentrations in all cohorts. Testosterone levels were strongly
reduced in H7N9 infected men of both age groups assessed compared to virus negative H7N9
close contacts, poultry workers and those with seasonal influenza (Figure 1A). Low
testosterone levels strongly correlated with lethal outcome in H7N9 infected men in the age
group of 18-49 years olds (Figure 1B). No significant differences in testosterone levels were
observed in H7N9 infected women compared to their close contacts and poultry workers.
However, testosterone levels were significantly increased in those infected with seasonal
influenza compared to all other female groups (Figure 1C). However, no significant impact of
testosterone levels was observed regarding H7N9 disease outcome in women (Figure 1D).
These data show that low testosterone levels in H7N9 infected men of 18-49 years of age
correlate with an enhanced risk for lethal outcome.

High estradiol levels in H7N9 infected men is not associated with disease outcome

Conversely, we measured estradiol levels as a predominant female hormone in all cohorts. Seasonal influenza infected men of 18-49 years of age presented slightly but significantly increased estradiol levels compared to poultry workers, H7N9 cases and their H7N9 virus-negative close contacts (Figure 2A). No differences were detected among all groups of ≥50 years of age (Figure 2A). However, estradiol levels were not associated with disease outcome in H7N9 infected men (Figure 2B). In H7N9 infected women of ≥50 years of age, estradiol levels were significantly elevated compared H7N9 virus-negative close contacts (Figure 2C). Increased estradiol levels, however, were not significantly associated with disease outcome in females but a tendency towards an increased risk for lethal outcome may be speculated among both age groups (Figure 2D). These findings suggest that estradiol does not significantly affect disease outcome in H7N9 infected men.

H7N9 infection induces a cytokine storm in men and women

Next, we assessed the cytokine and chemokine responses in H7N9 infected patients compared to their close contacts, poultry workers or those infected with seasonal influenza. In general, H7N9 infection resulted in highly increased cytokine and chemokine levels compared to their H7N9-negative close contacts or poultry workers (Figure S1 and S2). Seasonal influenza virus infected patients also presented elevated cytokine and chemokine levels but in general less than those infected with H7N9 (Figure S1 and S2). Thus, H7N9 induced cytokine storm is detected in men as well as women.

Storm cytokine levels in H7N9 infected men are associated with lethal outcome

We then analyzed whether certain cytokines might be associated with differential disease outcome. In H7N9 infected men, elevated levels of G-CSF, GM-CSF, IL-8, IL-10, IL-15, MCP-1 and TNF-α were associated with death (Figure 3). In H7N9 infected women, elevated levels Eotaxin correlated with survival (Figure 3P), whereas elevated MIP-1ß (Figure 3R) and EGF (Figure 3T) levels were associated with death. However, direct comparison of cytokine and chemokine responses in H7N9 infected men and women revealed important differences (Figure S3). Lethal outcome in H7N9 infected men correlated with elevated Eotaxin and MIP-1ß levels in the 18-49 year olds and with elevated IL-7 in the ≥50 year olds unlike in age-matched women with H7N9 influenza (Figure S3A-C). Conversely, survival in H7N9 infected men correlated with elevated IL-7 and MIP-1ß levels (Figure S3B and S3C).

233 These findings suggest that differential cytokines are associated with disease outcome in
234 males and females.

Low testosterone levels correlate with cytokine storm in H7N9 infected men

We could show that (1) high cytokine levels and (2) low testosterone levels correlate with
lethal outcome in H7N9 infected men. In order to assess potential mechanistic links between
individual cytokines and testosterone, we compared their individual levels in the same patient
with disease outcome. In H7N9 infected men (18-49 years old), among all assessed cytokines,
particularly IL-6, IL-8, IL-10, IL-1RA and GM-CSF levels were negatively associated with
testosterone concentrations in H7N9 survivors with the exception of TNF-α. Thus, a
significant number of H7N9 surviving men had high testosterone and low cytokine levels
(Figure 4A-E). In contrast, all men who succumbed to H7N9 influenza presented low
testosterone and high cytokine levels. In H7N9 infected men ≥50 years, a similar correlation
was observed for the same cytokines as described for the 18-49 years olds (Figure 4 G-L).
However, death was not significantly associated with low testosterone and high cytokine
levels. In H7N9 infected women, positive linear association between cytokine and
testosterone levels was observed unlike the observations in men (Figure S4). These findings,
suggest that in men, expression levels of cytokines, such as IL-6, increases with reducing
testosterone levels suggestive of potentially interlinked mechanisms.

Discussion

Avian influenza A (H7N9) virus is of public health importance that has caused 1,567 laboratory-confirmed cases and 615 deaths since early 2013 in China[13]. Exposure to poultry was considered a predominant risk factor for H7N9 infections based on epidemiological and virological studies[14,15]. Although many aspects of H7N9 infection were studied, systematic analysis of the male bias of H7N9 avian influenza detected across five waves in China were lacking.

Several animal studies illustrated that sex hormones pose biological variables that may contribute to sex differences in influenza pathogenesis. But there are still large gaps regarding evidence from human cohorts. In this study, we retrospectively selected cohorts (18-49 years old and those ≥50 years of age) from laboratory-confirmed H7N9 (*n*=98) and seasonal influenza (*n*=53) cases, epidemiological linked poultry workers (*n*=108) and close contacts of H7N9 cases that were virus-negative (*n*=71). We systematically analyzed the association of the main estrogen (estradiol) and androgen (testosterone) levels and correlated to cytokine and chemokine levels known to be hallmarks of influenza disease outcome in both sexes.

We observed a strong suppression of testosterone levels in H7N9 infected men in both age groups (18-49 years and ≥50 years). One limitation of this study is that we have no information regarding potential comorbidities in these groups, such as diabetes type II or obesity, which are known to lower testosterone levels in men[16–19]. However, our data is in line with data reported in murine models, where infection of healthy male mice with influenza A virus was shown to reduce testosterone levels by independent laboratories[10]˙[12]. Interestingly, treatment of female and male mice with testosterone protected animals from lethal outcome[10]. However, testosterone treatment of elderly male mice was less effective[20]. These data strongly hint towards an active reduction of testosterone levels by influenza A virus infection by a yet unknown mechanism.

For chronic infections, such as HCV or HIV, it was repeatedly shown that these infections cause low testosterone levels likely due to constant inflammation in infected individuals[21,22]. In line with chronic conditions, such as type II diabetes, it is believed that chronic inflammatory diseases can cause low testosterone levels. Since testosterone plays

key roles in not only fertility but also innate and adaptive immunity, one would hypothesize

that all these "testosterone level" decreasing hits may increase vulnerability towards

infections in general.

Elevated pro-inflammatory cytokines and chemokines, such as TNF-α, IL-8, IFN-α2, IFN-γ,

IL-6, MCP-1, IP10, and anti-inflammatory cytokine IL-10 were detected in both male and

female H7N9 cases without major statistically significant differences. However, testosterone

levels were negatively associated with specific cytokines and chemokines, such as IL-6, and

survival rates in H7N9 infected men. This is particularly interesting, since high IL-6 levels

were repeatedly reported to be a poor prognostic marker for severe infections, including

coronavirus disease 2019[23]. Albeit the underlying mechanism causing testosterone

deficiency is not known, there is increasing evidence that testosterone levels may modulate

inflammatory cytokines[24–27]. It would be interesting to see whether other acute infections,

such as the current pandemic Severe Acute Respiratory Syndrome Coronavirus 2(SARS-

CoV-2), may also cause low testosterone levels.

Clearly, future investigations are required to unravel the underlying mechanism of low

testosterone levels in H7N9 infected men. Increasing knowledge on the regulation of sex

hormones in acute virus infections will provide an important basis to understand diseases

with sex disparity and to develop sex specific therapies against viral infections.

Funding

This work was supported by the National Key Research and Development Program of China

[2016YFC1200200 to Y.L.S. and 2016YFD0500208 to D.Y.W], the Guangdong Province

Science and Technology Innovation Strategy Special Fund [2018A030310337 to Y.K.C] and

the German Free and Hanseatic City of Hamburg [to HPI (G.G.)] as well as the German

Federal Ministry of Health [to HPI (G.G.)].

Author contributions

GG, YLS and SSB conceived, designed and overviewed the experiments. YLS and DYW

organized and coordinated the field investigations. TB and YKC collected and organized

specimens. SB and TB established testosterone and estradiol measurements. TB, YKC and

SB conducted testosterone and estradiol measurements. YKC conducted cytokine and

chemokine measurements and real-time RT PCR on outpatients for seasonal influenza viruses.

JD performed real-time RT PCR on close contacts of H7N9 patients and poultry workers. TC,

JY, LJW and DYW provided and overviewed the epidemiology information from study

participants. TB and SSB analyzed the data. SB, TB and SSB developed all figures. TB and

GG wrote the manuscript. All authors revised the manuscript.

Declaration of Interests

All authors declare no competing interests.

Acknowledgement

We thank Chinese National Influenza Surveillance Network and all staff at local CDCs in

China for their contribution on sample collection and diagnosis for H7N9 and seasonal

influenza A viruses. We would also like to thank staff at Sino-Japan Friendship Hospital for

the measurement of testosterone and estradiol. Finally, we thank the study participants and

their families.

Figure Legends

Figure 1: Testosterone expression levels in H7N9 IAV infected patients compared to control groups. (A, C) Total testosterone expression levels were measured in H7N9 IAV infected male (A) and female (C) patients (H7N9 cases, filled squares) and compared to control groups: i) seasonal influenza cases (triangles), ii) poultry workers (circles), and iii) H7N9 close contacts (open squares). (B, D) Total testosterone expression levels were measured in H7N9 IAV infected male (B) and female (D) patients (H7N9 cases) and shown in dependency of disease outcome (survival, open circles; death, filled circles). (A-D) Samples from both sexes were divided into two groups based on age: 18-49 and $\geq$ 50 years (yrs) old. Data represent the mean $\pm$ 95% confidence interval (CI). Numbers of samples are summarized in Table 1. Significant differences among all study groups were calculated using either one-way ANOVA (Kruskal-Wallis test, Dunn's post-hoc-test) (A, C) or the unpaired, two-tailed Student's *t*-test (B, D). Statistical significant differences were further classified into three groups: * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$.

Figure 2: Estradiol expression levels in H7N9 IAV infected patients compared to control groups. (A, C) Total estradiol expression levels were measured in H7N9 IAV infected male (A) and female (C) patients (H7N9 cases, filled squares) and compared to control groups: i) seasonal influenza cases (triangles), ii) poultry workers (circles), and iii) H7N9 close contacts (open squares). (B, D) Total estradiol expression levels were measured in H7N9 IAV infected male (B) and female (D) patients (H7N9 cases) and shown in dependency of disease outcome (survival, open circles; death, filled circles). (A-D) Samples from both sexes were divided into two groups based on age: 18-49 and $\geq$50 years (yrs) old. Data represent the mean $\pm$ 95% confidence interval (CI). Numbers of samples are summarized in Table 1. Significant differences among all study groups were calculated using either one-way ANOVA (Kruskal-Wallis test, Dunn's post-hoc-test) (A, C) or the unpaired, two-tailed Student's *t*-test (B, D). Statistical significant differences were further classified into three groups: * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$.

Figure 3: Cytokine and chemokine expression levels in H7N9 IAV infected patients in dependency of disease outcome. Shown are the significant expression levels of cytokines and chemokines in the sera of H7N9 IAV infected male and female patients who either survived or succumbed to the infection (survival and death, respectively). Samples were

divided into two groups based on age: 18-49 and ≥50 years (yrs) old. The measurement was

carried out using a multiplex immunoassay. Analytes included were: G-CSF (granulocyte-

colony stimulating factor), GM-CSF (granulocyte-macrophage colony-stimulating factor),

IL-8 (interleukin 8), IL-10 (interleukin 10), IL-15 (interleukin 15), MCP-1/CCL2 (monocyte

chemoattractant protein 1), TNF-α (tumor necrosis factor alpha), eotaxin, MIP-1β/CCL4

(macrophage inflammatory protein 1 beta), EGF (epidermal growth factor). Values are shown

as means ± SD. Statistical significant differences between groups were determined using the

unpaired, two-tailed Student's *t*-test and divided into: $* p < 0.05$, $** p < 0.01$,

$*** p < 0.001$.

Figure 4: Correlative analysis of testosterone-modulated inflammatory immune

responses in H7N9 IAV infected male patients. Testosterone expression levels measured in

the sera from H7N9 IAV infected male patients (A-F, age group 18-49 yrs old; G-L, age

group ≥ 50 yrs old) were plotted over the expression levels of a selection of six significant

individual inflammatory genes: IL-6 (interleukin 6, A and G), IL-8 (interleukin 8, B and H),

IL-10 (interleukin 10, C and I), IL-1RA (interleukin 1 receptor antagonist, D and J), GM-CSF

(granulocyte-macrophage colony-stimulating factor, E and K), and TNF-α (tumor necrosis

factor alpha, F and L). Cytokines/Chemokine expression values were used after Ln (*natural*

logarithm) transformation. Linear regression was performed for the two individual groups,

survival (green) and death (red), and shown are the respective *r* (correlation coefficient) and *p*

values of the regression lines. A *p* value < 0.05 was considered statistically significant, i.e.

strong positive or negative correlation between testosterone and cytokine/chemokine

expression.

Table

Table 1:Age and sex characteristics of study participants

Age group	Study participants	Age, years		Sex (%)	
		Median	**IQR**	**Male**	**Female**
18-49 years	Seasonal influenza cases(*n*=28)	26·5	23·3-32·8	16 (57%)	12 (43%)
	Poultry workers (*n*=43)	41	36-46·5	32 (74%)	11 (26%)
	H7N9 close contacts (*n*=40)	35	27-41·3	29 (73%)	11 (27%)
	H7N9 cases (*n*=44)	42	34-46	33 (75%)	11 (25%)
≥50 years	Seasonal influenza cases (*n*=25)	61	56-67	12 (48%)	13 (52%)
	Poultry workers (*n*=65)	55	53-62	45 (69%)	20 (31%)
	H7N9 close contacts (*n*=31)	62	54-70·5	16 (52%)	15 (48%)
	H7N9 cases (*n*=54)	61	56-67	38 (70%)	16 (30%)

Data are median (Interquartile Range, IQR) or *n*(%).

Reference

1.　Gao R, Cao B, Hu Y, et al. Human infection with a novel avian-origin. N Engl J Med
2013; 368:1888–1897.

2.　Li Q, Zhou L, Zhou M, et al. Epidemiology of human infections with avian influenza
A(H7N9) virus in China. N Engl J Med **2014**; 370:520–532.

3.　Wang X, Jiang H, Wu P, et al. Evolving epidemiology of human infections with avian
influenza A(H7N9) virus across five epidemic waves in mainland China, 2013–17: an
epidemiological study of laboratory-confirmed case series. Lancet **2018**; 17:822–832.

4.　Arima Y, Vong S. Human infections with avian influenza A(H7N9) virus in China:
preliminary assessments of the age and sex distribution. West Pacific Surveill response
J WPSAR **2013**; 4:1–3.

5.　Rivers C, Lum K, Lewis B, Eubank S. Estimating human cases of avian influenza
A(H7N9) from poultry exposure. PLoS Curr **2013**; 5:1–12.

6.　Bupp MRG, Potluri T, Fink AL, Klein SL. The confluence of sex hormones and aging
on immunity. Front Immunol **2018**; 9.

7.　Klein SL, Flanagan KL. Sex differences in immune responses. Nat. Rev. Immunol.
2016;

8.　Gabriel G, Arck PC. Sex, immunity and influenza. J Infect Dis **2014**; 209:93–99.

9.　Academy T, Academy R, Trakt SS. Sex Hormones and inmmunit of infection. 2010.

10.　Tuku B, Stanelle-bertram S, Sellau J, et al. Testosterone Protects Against Severe
Influenza by Reducing the Pro-Inflammatory Cytokine Response in the Murine Lung.
2020; 11:1–7.

11.　Vom Steeg LG, Vermillion MS, Hall OJ, et al. Age and testosterone mediate influenza
pathogenesis in male mice. Am J Physiol - Lung Cell Mol Physiol **2016**; 311:L1234–
L1244.

12.　Robinson DP, Lorenzo ME, Jian W, Klein SL. Elevated 17β-estradiol protects females
from influenza a virus pathogenesis by suppressing inflammatory responses. PLoS
Pathog **2011**; 7.

13.　World Health Organization. Human infection with avian influenza A(H7N9) virus –
China: Update. https://www.who.int/csr/don/05-september-2018-ah7n9-china/en/ **2018**;

14.　Liu D, Shi W, Shi Y, et al. Origin and diversity of novel avian influenza A H7N9
viruses causing human infection: Phylogenetic, structural, and coalescent analyses.
Lancet **2013**; 381:1926–1932.

15.　Shi JZ, Deng GH, Liu PH, et al. Isolation and characterization of H7N9 viruses from

live poultry markets - Implication of the source of current H7N9 infection in humans.

Chinese Sci Bull **2013**; 58:1857–1863.

16. Angele MK, Knöferl MW, Ayala A, Bland KI, Chaudry IH. Testosterone and estrogen

differently effect Th1 and Th2 cytokine release following trauma-haemorrhage.

Cytokine **2001**; 16:22–30.

17. Cheung KKT, Luk AOY, So WY, et al. Testosterone level in men with type 2 diabetes

mellitus and related metabolic effects: A review of current evidence. J Diabetes

Investig **2015**; 6:112–123.

18. Yao QM, Wang B, An XF, Zhang JA, Ding L. Testosterone level and risk of type 2

diabetes in men: A systematic review and meta-analysis. Endocr Connect **2018**;

7:220–231.

19. Jensen TK, Andersson AM, Jørgensen N, et al. Body mass index in relation to semen

quality and reproductive hormones among 1,558 Danish men. Fertil Steril **2004**;

82:863–870.

20. Landon G, Attreed SE, Zirkin B, Klein SL. Testosterone treatment of aged male mice

improves some but not all aspects of age-associated increases in in fl uenza severity.

Cell Immunol **2019**; :103988. Available at:

https://doi.org/10.1016/j.cellimm.2019.103988.

21. Gomes AR, Souteiro P, Silva CG, et al. Prevalence of testosterone deficiency in HIV-

infected men under antiretroviral therapy. BMC Infect Dis **2016**; 16:1–7. Available at:

http://dx.doi.org/10.1186/s12879-016-1892-5.

22. Chaudhury CS, Mee T, Chairez C, et al. Testosterone in Men With Chronic Hepatitis

C Infection and After Hepatitis C Viral Clearance. Clin Infect Dis **2019**; 69:571–576.

23. Tay MZ, Poh CM, Rénia L, Macary PA, Ng LFP. The trinity of COVID-19: immunity,

inflammation and intervention. Nat Rev Immunol **2020**;

24. Maggio M, Basaria S, Ble A, et al. Correlation between testosterone and the

inflammatory marker soluble interleukin-6 receptor in older men. J Clin Endocrinol

Metab **2006**; 91:345–347.

25. Traish A, Bolanos J, Nair S, Saad F, Morgentaler A. Do Androgens Modulate the

Pathophysiological Pathways of Inflammation? Appraising the Contemporary

Evidence. J Clin Med **2018**; 7:549.

26. Morooka N, Ueguri K, Yee KKL, Yanase T, Sato T. Androgen-androgen receptor

system improves chronic inflammatory conditions by suppressing monocyte

chemoattractant protein-1 gene expression in adipocytes via transcriptional regulation.

Biochem Biophys Res Commun **2016**; 477:895–901. Available at:

http://dx.doi.org/10.1016/j.bbrc.2016.06.155.

27.   Posma E, Moes H, Heineman MJ, Faas MM. The Effect of Testosterone on Cytokine

Production in the Specific and Non-specific Immune Response. Am J Reprod

Immunol **2004**; 52:237–243.

Figure 1

Figure 2

Figure 3

Figure 4