Article tittle: "Clinical, Radiological and Therapeutic Characteristics of Patients with COVID-

19 in Saudi Arabia"

Authors' names:

Mohammed Shabrawishi, M.D,[1,2] Manal M Al-Gethamy, M.D,[3] Abdallah Y Naser, PhD,[4] Maher A Ghazawi,

M.D,[5] Ghaidaa F Alsharif, M.D,[2] Elaf F Obaid, M.D,[2] Haitham A Melebari, M.D,[6] Dhaffer M Alamri,[3] Ahmad

S Brinji, M.D,[5] Fawaz H Al Jehani, M.D,[7] Wail Almaimani, M.D,[7] Rakan A Ekram, M.D,[8] Kasim H Alkhatib,

M.D,[9] Hassan Alwafi, M.D[10]

Authors affiliations:

[1]Department of Respiratory Medicine, Al Noor Specialist Hospital, Mecca, Saudi Arabia.

[2]Department of Internal Medicine, Al Noor Specialist Hospital, Mecca, Saudi Arabia.

[3]Department of Infection Prevention and Control Programme, Al Noor Specialist Hospital, Mecca, Saudi
Arabia.

[4]Faculty of Pharmacy, Isra University, Amman, Jordan.

[5]Department of Radiology, Al Noor Specialist Hospital, Mecca, Saudi Arabia.

[6]Department of Internal Medicine, King Faisal General Hospital, Mecca, Saudi Arabia.

[7]Al Noor Specialist Hospital, Mecca, Saudi Arabia.

[8]School of Public Health and Health Informatics, Umm Al Qura University, Mecca, Saudi Arabia.

[9]Intensive Care Department, Al Noor Specialist Hospital, Mecca, Saudi Arabia.

[10]Department of Clinical Pharmacology and Toxicology, Faculty of Medicine, Umm Al Qura University,
Mecca, Saudi Arabia.

21
Corresponding author

Dr.Hassan Alwafi

Department of Clinical Pharmacology and Toxicology

Umm Alqura University, Faculty of Medicine

Mecca, Saudi Arabia

Post Code 24381

T +966555544319

Email: hhwafi@uqu.edu.sa

Number of words in abstract: 249 words

Number of words in main body text: 2,633 words

Number of tables in main body text: 4 table

Number of figures in main body text: 3 figures

Number of references in main body text: 34

Abstract (249)

BACKGROUND

Coronavirus disease 2019 (COVID-19) is a rapidly spreading global pandemic. The clinical

characteristics of COVID-19 has been reported; however, there are limited researches that

investigated the clinical characteristics of COVID-19 in the Middle East. The aim of this study is to

investigate the clinical, radiological and therapeutic characteristics of patients diagnosed with

COVID19 in Saudi Arabia.

METHODS

This study is a retrospective single-centre case series study. We extracted data for patients who were

admitted to the Al-Noor Specialist hospital with a PCR confirmed SARS-COV-2 between 12th and

31st of March 2020. Descriptive statistics were used to describe patients' characteristics. Continuous

data were reported as mean ± SD. Chi-squared test/Fisher test were used as appropriate to compare

proportions for categorical variables.

RESULTS

A total of 150 patients were hospitalised for COVID-19 during the study period. The mean age was

46.1 years (SD: 15.3 years). Around 61.0% (n= 90) were males and six patients (3.9%) reported

working in the healthcare sector. The most common comorbidities were hypertension (28.8%, n= 42)

and diabetes mellitus (26.0%, n= 38). The majority of the patients, 64.4% (n = 96) had a recent

contact history with a COVID patient. Regarding the severity of the hospitalised patients, 105 patients

(70.0%) were mild, 29 (19.3%) were moderate, and 16 patients (10.7%) were severe or required ICU

care. From the 105 mild patients, around 31.3% (n= 47) were asymptomatic.

CONCLUSION

This case series provides clinical, radiological and therapeutic characteristics of hospitalised patients

with confirmed COVID-19 in Saudi Arabia.

KEYWORDS

Clinical Characteristics, COVID-19, Retrospective case series, Radiological SARS-Cov2

1. INTRODUCTION

In early December 2019, a cluster of acute pneumonia of unknown aetiology has been identified in Wuhan, China (1). The pathogen has been identified as a new RNA virus from the betacoronavirus family, and has been named as severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) (2). The respiratory illness caused by the 2019 novel coronavirus disease (COVID-19) is highly infectious, and therefore, the World Health Organization (WHO) has characterized the diseases a pandemic infection (3). As of April 25, 2020, more than 2,700,000 confirmed cases were reported worldwide, and it has spread from Wuhan to more than 200 countries across the world (4).

The Kingdom of Saudi Arabia (KSA) is the largest country in the Arabian Peninsula and it is located in the South West part of Asia (5). In a historical decision, KSA has suspended Umrah and all religious visits to the country in an attempt to prevent and delay the spread of COVID-19 in KSA. However, in March 2, 2020, Saudi Arabia has confirmed its first case of COVID-19 which was imported from Iran (4). Several other local clusters were identified later with the majority of the cases being linked to recent travel history.

In recent studies, the clinical features and severity of COVID-19 has been described to be similar of other respiratory viruses such as severe acute respiratory syndrome (SARS) and Middle East respiratory syndrome (MERS) (6, 7). Symptoms can range from mild flu-like symptoms to acute respiratory distress syndrome (ARDS) (8). However, the characteristics and the course of the disease in Middle Eastern populations remains unclear. Exploring the clinical characteristics of patients diagnosed with COVID-19 in Saudi Arabia is important knowing that there are many visitors who travel to Saudi Arabia for religious purposes. Beside this, there is a high air traffic for other purposes in this country which was estimated to be around 39 million people in 2018. In 2019, around 7.5 million Muslim entered the holy city of Mecca for Umrah purposes (9). This highlights how crucial is to have a deeper exploration on the characteristics of patients diagnosed with this widespread infection in this region. To address the above knowledge gaps and giving the ongoing spread of COVID-19 in the Middle East, this study aims to describe the clinical, radiological, and therapeutic characteristics of COVID-19 in a selected cohort of patients in Mecca, Saudi Arabia.

2. METHODS
2.1. Study Design and participants

This was a retrospective single-centre case series study of 150 patients diagnosed with COVID-19. We extracted data for patients who were admitted to Al-Noor Specialist hospital with a polymerase chain reaction (PCR) confirmed SARS-COV-2 between 12th and 31st of March 2020. Al-Noor Specialist hospital in Mecca, Saudi Arabia is a 500-bedder specialist and teaching hospital in the centre of the holy city of Mecca. It delivers tertiary care throughout the Mecca region of Saudi Arabia

96   and its part of the Ministry of Health services (10, 11). All patients enrolled in this study were

97   diagnosed with COVID-19 through real time (RT)-PCR obtained through nasopharyngeal swabs. All

98   data including outcomes, mortality and length of stay were monitored up to 8th April 2020.

2.2. Data Collection

Data were extracted from both paper and electronic records using a unique medical record number

(MRN) for each patient. All data were reviewed and checked by a medical team including; two

medical residents and a consultant Pulmonologist. Extracted data included patients' demographics,

comorbidities, history of recent travel and history of contact with a confirmed COVID19 patient in the

past two weeks. In addition, clinical signs, symptoms, radiological findings and pharmacological

treatment received were collected. The radiological examinations were interpreted by a certified

consultant radiologist who was blinded from the clinical presentation of the patients. The severity

assessment of the chest x-ray (CXR) were estimated subjectively. All data were collected at the time

of the admission.

2.3. Study variables

Data regarding the clinical progression and severity of the disease were reported as the worst

classification reached at any point during hospitalisation. We further classified the severity of the

disease based on the following criteria; 1) mild disease was defined as patients with upper respiratory

tract symptoms (as rhinorrhoea, sore throat, headache, myalgia, body pain, low grad fever and or dry

cough) with absent of clinical or radiological finding of pneumonia; 2) moderate disease defined as

symptomatic patients with radiological sign of pneumonia; 3) severe disease defined as confirmed

COVID-19 pneumonia with any of the following respiratory rate $\geq$30/min, blood oxygen saturation

$\leq$93% at rest, PaO2/FiO2 ratio <300, lung infiltration >50% of the lung field, and 4) critically sever

disease defined as any of the following: respiratory failure required invasive mechanical ventilation,

shock or organ failure require admission to the intensive care unit.

2.4. Ethics

121   This study was approved by the institutional ethics board at the Ministry of Health in Saudi Arabia

122   (No. H-02-K-076-0420-286). Patients were informed that their clinical data will be used for clinical or

123   research purposes with keeping all their personal information confidential.

2.5. Statistical Analysis

125   Descriptive statistics were used to describe patients' demographic characteristics, radiological

126   findings, medications use, and comorbidities. Continuous data were reported as mean $\pm$ SD, and

127   categorical data were reported as percentages (frequencies). Independent sample t test was used to

compare the mean value for continuous variables. Chi-squared test/Fisher test were used as appropriate to compare proportions for categorical variables. Logistic regression analysis was used to identify predictors of clinical characteristics. A confidence interval of 95% ($p < 0.05$) was applied to represent the statistical significance of the results and the level of significance was assigned as 5%. SPSS (Statistical Package for the Social Sciences) version 25.0 software (SPSS Inc) was used to perform all statistical analysis.

3. RESULTS

3.1. Patients clinical characteristics

Table 1 presents patients' characteristics at presentation to the hospital. A total of 150 patients were hospitalised for COVID-19. The mean age was 46.1 years (SD: 15.3 years), and ranged between 11 and 87. Around 61.0% (n= 90) were males. Six patients (3.9%) reported working in the healthcare sector. The most common comorbidities were hypertension (28.8%, n= 42) and diabetes mellitus (DM) (26.0%, n= 38). The majority of the patients (56.0%; n= 84) were local resident. Around half of the patients (54.1%, n= 80) reported that they had a contact history with a traveller. In addition, the majority of the patients, 64.4% (n = 96) had a contact history with a COVID-19 patient. Regarding the severity of the hospitalised patients, 105 patients (70.0%) were mild, 29 (19.3%) were moderate, and 16 patients (10.7%) were severe or required ICU care. Of the 105 mild patients, around 31.3% (n= 47) were asymptomatic. Patients with comorbidities were more likely to have a severe outcome compared to other patients (p<0.05). Patients who reported a contact history with a COVID-19 patient were more likely to have mild to moderate severity of the disease (p<0.05). Mild cases were more prevalent among females, while moderate to severe and or critical were prevalent among males (**Figure 1**). For symptomatic patients, the most common symptoms at presentation were fever (49.3%, n= 72), dry cough (48.6%, n= 71), and shortness of breath (19.9%, n= 29) **(Table 2)**. Furthermore, during admission, fever and cough (28%) were the most common symptoms followed by nausea and vomiting (12%). Most of the asymptomatic patients were females (OR: 0.45 [95%CI 0.22 – 0.92]; p= 0.027). In addition, patients who reported travel history or a contact with a traveller recently were three times (OR: 3.13 [95%CI 1.52 – 6.45]; p= 0.002) and four times (OR: 4.03 [95%CI 1.84 – 8.81]; p= 0.000) more likely to be asymptomatic, respectively. Besides, patients who have reported a contact with COVID-19 patients were four times at higher risk of being symptomatic (OR: 4.50 [95%CI 1.84 – 10.99]; p= 0.001).

3.2. Radiological findings

Around half of the patients (49.7%, n= 72) had normal radiological exam at presentation. The severity of the cases was correlated with an increase in the prevalence of GGO at presentation (P=0.002). The predominant pattern of abnormality observed was ground-glass opacification (29.0%, n= 42),

peripheral (57.5%, n= 42), and (bilateral (35.3%, n= 35), which was mainly involving the lower lobes

(Figure 2). Most of the patients had stable radiological exams on follow up. Around 64.6% (n= 62)

showed progression, half of them belongs to more severe group **(Table 3).**

3.3. Recovery

Patients stayed at the hospital for a mean duration of 9.2 days (SD: 3.9). Duration of stay in hospital

ranged from two days to 23 days. At the end of the follow-up period a total of 94 patients (63.5%)

recovered and 31.8% (n= 47) improved clinically but RT-PCR results were still positive. On the other

hand, three patients (2.0%) did not fully recover and four patients (2.7%) deceased. The two patients

who had mild cases died due to other reasons. There was no statistically significant difference based

on the age regarding the recovery or whether the patient was symptomatic or asymptomatic upon

presentation to hospital (p>0.05). The majority of the patients with mild cases improved or recovered,

however, there was no statistically significant difference between cases from different severity and

recovery rate (p>0.05) **(Figure 3).**

3.4. Therapeutic management

Beside supportive care, there were three main types of therapies that were prescribed to the patients

for the management of COVID-19, this includes: a) antiviral therapy, b) antibiotics, and c)

antimalarial medications **(Table 4).**

4. DISCUSSION

To the best of our knowledge, this is the first and largest study to examine the clinical characteristics

of COVID-19 in the Middle East region. We investigated clinical, radiological, and therapeutic

characteristics of COVID-19 in a 150 hospitalised patients in Saudi Arabia. We found that around

89.0% of the cases were either mild or moderated and only 11.0% were either severe or critical. Our

finding showed that the clinical severity of COVID-19 were of a milder presentation compared to

results from China (12), Italy (13) and the United States (14, 15). These finding could be attributed to

several factors including age and other demographics differences. The mean age in our study was 46.1

years (SD: 15.3) which was younger than the age reported in other studies. Several studies have

reported poorer outcome among older population and patients with COVID-19 and comorbidities (16-

18). However, it is difficult to draw a causal inference and we urge for further studies to investigate

this association. In addition, it is important to highlight that the majority of the Saudi Arabian

population are younger than 44 years (19).

Male patients with COVID-19 were more prevalent in our study compared to females, this was also

similar to previous reports which highlight more males to be infected with COVID-19 (2, 14). These

numbers could be because of men sex-based immunological differences or it could also be because of

behavioural patterns such as smoking (20). In addition, comorbidities are more prevalent in men

which could also be a reason for this difference (21), However, more researches must focus on gender

differences and clinical outcomes with COVID-19.

Our study highlighted that around 28.8% and 26.0% of the study population had hypertension (HTN)

and DM, these results were similar to previous reports that investigated the clinical characteristics of

COVID-19 (1). Patients with DM and hypertension have an increased risk of complication of

COVID-19 including acute respiratory distress syndrome (ARDS) (22), however, the mechanism of

this remains un-investigated and it is unclear whether patients with uncontrolled blood pressure have a

poorer outcomes of COVID-19 compared to patients with controlled blood pressure. In addition,

Angiotensin-converting enzyme (ACE) inhibitors and angiotensin receptor blockers (ARBs) are two

commonly prescribed medications for the management of HTN, and since SARS-CoV-2, binds to

ACE2 in the lung, some theoretical theories have been raised about the benefits of these medications

in the treatment of COVID-19 (23).

SARS-COV2 has been described to be similar to seasonal influenza, SARS-COV and MERS, this

includes the fact that it is transmitted through respiratory droplets (24, 25). In addition, SARS-COV2

has similar symptoms to SARS-COV and MERS such as; fever, cough, and shortness of breath. This

was reported in our study and it was also in line with previous studies (6, 13), however, SARS-COV2

has a higher case fatality rate in comparison to seasonal flu (0.1%) while it is also milder in

comparison to other respiratory viruses such as SARS-COV (9.5%) and MERS (34.3%) (26). Besides

this, COVID-19 is a highly infectious pathogenic (27, 28), with some reports suggested that half of

the United Kingdom (UK) population has been infected without showing any symptoms or with

having a mild course of the disease (29). Our study demonstrated that around 31.3% of the study

sample were asymptomatic and had a mild disease. Mostly, these patients were identified through

contact tracing and were isolated in the earlier course of the disease, whether this approach have any

impact on the clinical course psychologically, this might need to be addressed in future studies. In

addition, the majority of these patients had a contact with a confirmed COVID-19 patients which may

raise concerns regarding the mechanism and the underlying inflammatory response in these patients.

More researches are encouraged to investigate the characteristics of asymptomatic patients and if

early detection and supportive treatment have a role in the clinical progression of the disease.

In our study, and unlike previous reports, nearly half of the patients presented with normal CXR, most

of them were asymptomatic or had a mild disease. Furthermore, normal CXR at presentation may

have a prognostic rule as only few numbers of those patients progressed into more severe cases. On

the other hand, presence of ground glass opacity is linked with more aggressive course. The patterns

found in abnormal exams were similar to the previously published reports and findings where

peripheral, bilateral ground glass opacification (30).

Our study highlighted that around 26.7% of the patients received antimalarial treatment and around

9.0% received antiviral treatment, these medications have been suggested to have some beneficial

effect to reduce the viral load and eliminate the disease, however, there are also uncertainties
regarding their safety (31, 32). In addition, there has been debate about their efficacy in the treatment
of COVID-19 with several trials are now in pipeline for the testing of these medications (33). To date,
there is no treatment for COVID-19, and the main approach in the management of the disease is to
provide supportive treatment and to control the symptoms including mechanical ventilator for critical
cases (34). This study has some limitations. First, the number of patients included in the study were
small. Second, the study population only included patients from a single-centre hospital in Saudi
Arabia.

5. CONCLUSION

This case series provides clinical, radiological, and therapeutic characteristics of hospitalised patients
with confirmed COVID-19 in Saudi Arabia. Our study demonstrates similar characteristics of
COVID-19 to previously reported studies worldwide.

Conflict of interest

The authors have stated explicitly that there are no conflicts of interest in connection with this article.

Authors contribution

Shabrawishi, Naser, Ghazawi and Alwafi had full access to all the data in the study and take
responsibility for the integrity of the data and the accuracy of the data analysis. Shabrawishi and
Alwafi had the original idea for this study. Shabrawishi, Naser and Alwafi contributed to the design of
the study. Obaid and Alsharif contributed to the data collection. Naser and Alwafi contributed in the
statistical analysis. Shabrawishi, Naser, Ekram and Alwafi wrote the first draft. All the authors
contributed to interpretation and edited the draft report.

Data availability statement

The data that support the findings of this study are available from the corresponding author upon reasonable
request.

References:

1.   Wang D, Hu B, Hu C, Zhu F, Liu X, Zhang J, et al. Clinical Characteristics of 138 Hospitalized
Patients With 2019 Novel Coronavirus–Infected Pneumonia in Wuhan, China. JAMA.
2020;323(11):1061-9.
2.   Guan W-j, Ni Z-y, Hu Y, Liang W-h, Ou C-q, He J-x, et al. Clinical Characteristics of Coronavirus
Disease 2019 in China. 2020.
3.   WHO. Coronavirus disease 2019 (COVID-19)Situation Report –96 2020 [Available from:
https://www.who.int/docs/default-source/coronaviruse/situation-reports/20200425-sitrep-96-
covid-19.pdf?sfvrsn=a33836bb_2.

4.   Who. Coronavirus disease (COVID-19) outbreak 2020 [Available from:
https://www.who.int/emergencies/diseases/novel-coronavirus-2019.
5.   MOFA. About Saudi Arabia 2020 [Available from:
https://www.mofa.gov.sa/sites/mofaen/aboutKingDom/Pages/KingdomGeography46466.aspx.
6.   Chen N, Zhou M, Dong X, Qu J, Gong F, Han Y, et al. Epidemiological and clinical
characteristics of 99 cases of 2019 novel coronavirus pneumonia in Wuhan, China: a descriptive
study. Lancet (London, England). 2020;395(10223):507-13.
7.   Livingston E, Bucher K. Coronavirus Disease 2019 (COVID-19) in Italy. JAMA. 2020.
8.   Chang D, Lin M, Wei L, Xie L, Zhu G, Dela Cruz CS, et al. Epidemiologic and Clinical
Characteristics of Novel Coronavirus Infections Involving 13 Patients Outside Wuhan, China. JAMA.
2020;323(11):1092-3.
9.   Ebrahim SH, Memish ZA. COVID-19: preparing for superspreader potential among Umrah
pilgrims to Saudi Arabia. The Lancet. 2020;395(10227).
10.   Zaglool DA, Khodari YA, Gazzaz ZJ, Dhafar KO, Shaker HA, Farooq MU. Prevalence of
Intestinal Parasites among Patients of Al-Noor Specialist Hospital, Makkah, Saudi Arabia. Oman Med
J. 2011;26(3):182-5.
11.   Health Mo. Al Noor Specialist Hospital 2020 [Available from:
http://nsh.med.sa/Pages/Home.aspx.
12.   Wu Z, McGoogan JM. Characteristics of and Important Lessons From the Coronavirus
Disease 2019 (COVID-19) Outbreak in China: Summary of a Report of 72314 Cases From the Chinese
Center for Disease Control and Prevention. JAMA. 2020;323(13):1239-42.
13.   Livingston E, Bucher K. Coronavirus Disease 2019 (COVID-19) in Italy. JAMA.
2020;323(14):1335-.
14.   Goyal P, Choi JJ, Pinheiro LC, Schenck EJ, Chen R, Jabri A, et al. Clinical Characteristics of
Covid-19 in New York City. 2020.
15.   Richardson S, Hirsch JS, Narasimhan M, Crawford JM, McGinn T, Davidson KW, et al.
Presenting Characteristics, Comorbidities, and Outcomes Among 5700 Patients Hospitalized With
COVID-19 in the New York City Area. JAMA. 2020.
16.   Mahase E. Covid-19: death rate is 0.66% and increases with age, study estimates.
2020;369:m1327.
17.   Verity R, Okell LC, Dorigatti I, Winskill P, Whittaker C, Imai N, et al. Estimates of the severity
of coronavirus disease 2019: a model-based analysis. The Lancet Infectious diseases. 2020.
18.   Dowd JB, Andriano L, Brazel DM, Rotondi V, Block P, Ding X, et al. Demographic science aids
in understanding the spread and fatality rates of COVID-19. 2020:202004911.
19.   Statistics GAf. Statistical Yearbook of 2016 2016 [Available from:
https://www.stats.gov.sa/en/5305.
20.   Wenham C, Smith J, Morgan R. COVID-19: the gendered impacts of the outbreak. Lancet
(London, England). 2020;395(10227):846-8.
21.   The L. The gendered dimensions of COVID-19. Lancet (London, England).
2020;395(10231):1168.
22.   Fang L, Karakiulakis G, Roth M. Are patients with hypertension and diabetes mellitus at
increased risk for COVID-19 infection? The Lancet Respiratory medicine. 2020;8(4):e21.
23.   Schiffrin EL, Flack JM, Ito S, Muntner P, Webb RC. Hypertension and COVID-19. American
Journal of Hypertension. 2020.
24.   Otter JA, Donskey C, Yezli S, Douthwaite S, Goldenberg SD, Weber DJ. Transmission of SARS
and MERS coronaviruses and influenza virus in healthcare settings: the possible role of dry surface
contamination. The Journal of hospital infection. 2016;92(3):235-50.
25.   WHO. Modes of transmission of virus causing COVID-19:implications for IPC precaution
recommendations 2020 [Available from: https://www.who.int/publications-detail/modes-of-
transmission-of-virus-causing-covid-19-implications-for-ipc-precaution-recommendations.

26.   Rajgor DD, Lee MH, Archuleta S, Bagdasarian N, Quek SC. The many estimates of the COVID-
19 case fatality rate. The Lancet Infectious diseases. 2020.
27.   Zhang S, Diao M, Yu W, Pei L, Lin Z, Chen D. Estimation of the reproductive number of novel
coronavirus (COVID-19) and the probable outbreak size on the Diamond Princess cruise ship: A data-
driven analysis. International journal of infectious diseases : IJID : official publication of the
International Society for Infectious Diseases. 2020;93:201-4.
28.   Liu Y, Gayle AA, Wilder-Smith A, Rocklöv J. The reproductive number of COVID-19 is higher
compared to SARS coronavirus. Journal of Travel Medicine. 2020;27(2).
29.   Sayburn A. Covid-19: experts question analysis suggesting half UK population has been
infected. 2020;368:m1216.
30.   Shi H, Han X, Jiang N, Cao Y, Alwalid O, Gu J, et al. Radiological findings from 81 patients with
COVID-19 pneumonia in Wuhan, China: a descriptive study. The Lancet Infectious Diseases.
2020;20(4):425-34.
31.   Ferner RE, Aronson JK. Chloroquine and hydroxychloroquine in covid-19. 2020;369:m1432.
32.   Cortegiani A, Ingoglia G, Ippolito M, Giarratano A, Einav S. A systematic review on the
efficacy and safety of chloroquine for the treatment of COVID-19. Journal of critical care. 2020.
33.   Taccone FS, Gorham J, Vincent JL. Hydroxychloroquine in the management of critically ill
patients with COVID-19: the need for an evidence base. The Lancet Respiratory medicine. 2020.
34.   NICE. COVID-19 rapid guideline: managing suspected or confirmed pneumonia in adults in
the community 2020 [Available from: https://www.nice.org.uk/guidance/ng165.

TABLES LEGENDS

Table 1: Patients demographic characteristics at presentation.

Table 2: Patient signs and symptoms at presentation and during admission.

Table 3: Radiological findings.

Table 4: Initial treatment characteristics.

FIGURES LEGENDS

Figure 1: Clinical severity stratified by gender.

Figure 2: CXR's of two different patients showing the most common abnormalities: bilateral,

peripheral ground glass opacities and consolidation

Figure 3: Recovery rates stratified by case severity

TABLES

Table 2: Patients demographic characteristics at presentation.

Demographics	All patients (n=150)	Mild cases (n= 105)	Moderate cases (n= 29)	Severe/Intensive care unit cases (n= 16)	P-value
Age (years; mean (SD))	46.1 years (15.3)	45.4 years (±16.0)	46.7 years (±12.1)	49.8 years (±15.7)	0.550
Gender					
Female No. (%)	60 (40.0)	47 (44.8)	10 (34.5)	3 (18.8)	0.112
Healthcare worker					
Yes No. (%)	6 (4.0)	6 (5.8)	0	0	0.110
Place of residency No. (%)					
Kingdom of Saudi Arabia	84 (56.0)	57 (54.3)	15 (51.7)	12 (75.0)	0.084

Other countries	66 (44.0)	48 (45.7)	14 (48.3)	4 (25.0)	0.239
Comorbidities No. (%)					
Hypertension	42 (28.8)	29 (27.6)	10 (35.7)	3 (23.1)	0.627
Diabetes mellitus	38 (26.0)	20 (19.0)	11 (39.3)	7 (53.8)	0.005**
Coronary artery disease	11 (7.5)	5 (4.8)	3 (10.7)	3 (23.1)	0.094
Renal disease	10 (6.8)	5 (4.8)	1 (3.6)	4 (30.8)	0.018*
Thyroid gland problem (hypothyroidism)	9	4 (8.2)	2 (11.1)	3 (37.5)	0.123
Asthma	4 (2.7)	3 (2.9)	0	1 (7.7)	0.306
Cancer	2 (1.4)	0	0	2 (15.4)	0.007**
CVA	1 (0.7)	1 (1.0)	0	0	0.718
COPD	1 (0.7)	0	0	1 (7.7)	0.086
CLD	1 (0.7)	0	0	1 (7.7)	0.086
Tracing history No. (%)					
Recent travel history (Yes) No. (%)	65 (43.9)	47 (45.6)	14 (48.3)	4 (25.0)	0.263
Contact with traveller (Yes) No. (%)	80 (54.1)	57 (55.3)	18 (62.1)	5 (31.3)	0.124
Contact with COVID-19 patient (Yes) No. (%)	96 (64.4)	71 (68.3)	20 (69.0)	5 (31.3)	0.013*
Outcome (n= 148) No. (%)					
Deceased	4 (2.7)	2 (1.9)	0	2 (12.5)	0.086
Improved	47 (31.8)	37 (35.2)	6 (22.2)	4 (25.0)	0.358
Not recovered	3 (2.0)	2 (1.9)	1 (3.7)	0	0.615
Recovered	94 (63.5)	64 (61.0)	20 (74.1)	10 (62.5)	0.434

347 Abbreviations; COVID-19: coronavirus disease-2019; CVA: cerebrovascular accident; COPD: chronic

348 obstructive pulmonary disease; CLD: chronic liver disease; SD: Standard deviation; No: Number (frequency)

349 **Table 2: Patient signs and symptoms at presentation and during admission.**

	Symptoms		P-vale
Variable	**At presentation** No. (%)	**During admission** No. (%)	
Fever	72 (49.3)	28 (19.2)	0.029*
Cough	71 (48.6)	28 (19.2)	0.024*
Shortness of breath	29 (19.9)	7 (4.8)	0.000***
Sore throat	24 (16.4)	2 (1.4)	0.269
Runny nose	9 (6.2)	0 (0.0)	>0.99
Sputum	5 (3.4)	1 (0.7)	0.034*
Headache	4 (2.7)	0 (0.0)	>0.99
Myalgia	4 (2.7)	1 (0.7)	0.813
Diarrhea	2 (1.4)	5 (3.4)	0.068
Nausea/vomiting	1 (0.4)	12 (8.2)	0.678
Haemoptysis	1 (0.4)	1 (0.7)	0.887
Fatigue	1 (0.4)	1 (0.7)	0.907

350 * p<0.05; **p<0.01; ***p<0.000

351 **Table 3: Radiological findings.**

Radiological findings (CXR) upon admission					
	All patients (n=150)	**Mild cases** (n= 105)	**Moderate cases** (n= 29)	**Severe/Intensive care unit cases** (n= 16)	**P-value**
Predominant finding					
Normal	72 (49.7)	62 (60.2)	7 (25.9)	3 (20.0)	0.000***
Ground glass opacity	42 (29.0)	21 (20.4)	13 (48.1)	8 (53.3)	0.002*
Consolidation	26 (17.9)	16 (15.5)	6 (22.2)	4 (26.7)	0.488
Linear atelectasis	3 (2.1)	3 (2.9)	0	0	0.354
Diffusion reticular	1 (0.7)	1 (1.0)	0	0	0.795

opacities					
Reticulation	1 (0.7)	0	1	0	0.183
Distribution within the lobe					
Central	10 (13.7)	6 (14.6)	2 (10.0)	2 (16.7)	0.833
Diffuse	21 (28.8)	14 (34.1)	4 (20.0)	3 (25.0)	0.494
Peripheral	42 (57.5)	21 (51.2)	14 (70.0)	7 (58.3)	0.378
Distribution within the lung					
Lower	24 (32.9)	12 (29.3)	6 (30.0)	6 (50.0)	0.385
Lower middle	22 (30.1)	10 (24.4)	9 (45.0)	3 (25.0)	0.236
Lower and middle and upper	10 (13.7)	8 (19.5)	2 (10.0)	0	0.089
Diffuse	10 (13.7)	6 (14.6)	2 (10.0)	2 (16.7)	0.833
Peripheral	2 (2.7)	2 (4.9)	0	0	0.309
Middle	2 (2.7)	2 (4.9)	0	0	0.309
Upper	1 (1.4)	1 (2.4)	0	0	0.559
Upper and middle	1 (1.4)	0	0	1 (8.3)	0.159
No zonal predominance	1 (1.4)	0	1 (5.0)	0	0.269
Laterality					
Bilateral	53 (35.3)	27 (25.7)	16 (55.2)	10 (62.5)	0.000***
Unilateral right	12 (16.4)	10 (24.4)	2 (10.0)	0	0.035*
Unilateral left	8 (11.0)	4 (9.8)	2 (10.0)	2 (16.7)	0.805
Progression					
Stable	62 (64.6)	49 (74.2)	11 (57.9)	2 (18.2)	0.001**
Worsen	34 (35.4)	17 (25.8)	8 (42.1)	9 (81.8)	

CXR: chest x-ray

Table 4: Initial treatment characteristics.

Treatment therapy	Frequency (%)
Antiviral therapy	
Combination of antiretroviral (lopinavir and ritonavir) and ribavirin	14 (9.3)
Antimalarial therapy	
Hydroxychloroquine	25 (16.7)
Chloroquine	15 (10.0)
Antibiotics therapy	58 (38.7)

FIGURES

356
357
358   Figure 1: Clinical severity stratified by gender.

360
361   Figure 2: CXR's of two different patients showing the most common abnormalities: bilateral,
362   peripheral ground glass opacities and consolidation
363

364
365
366   Figure 3: Recovery rates stratified by case severity