1

Estimation of the number of general anesthesia cases based on a series

of nationwide surveys on Twitter during COVID-19 in Japan: A statistical

analysis

Hiroki Daijo MD, Ph.D. [1, 4], Yosuke Fujii MD, [2, 4], Kiichi Hirota MD, Ph.D. [3,*]

[1] Department of Anesthesia, Takarazuka Daiichi Hospital, Takakazuka, Hyogo 665-

0832 Japan

[2] Department of Anesthesia, Otsu City Hospital, Otsu, Shiga 520-0804 Japan

[3] Department of Human Stress Response Science, Institute of Biomedical Science,

Kansai Medical University, Hirakata, Osaka 573-1010 Japan.

[4] These authors contributed equally to this work

*corresponding author

Kiichi Hirota

Department of Human Stress Response Science, Institute of Biomedical Science,

Kansai Medical University, Hirakata, Osaka, Japan

phone: +81-72-804-2526

email: hif1@mac.com / khirota-kyt@umin.ac.jp

Keywords: COVID-19; twitter; social network service; general anesthesia; statistical

estimation; Japan

Abstract

Coronavirus disease (COVID-19), caused by severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2), was first reported in Wuhan, Hubei, China, and has since spread to more than 200 other countries at the time of writing. In light of this situation, the Japanese Government declared a state of emergency in seven regions of Japan on the April 7, 2020 under the provisions of the law. The medical care delivery system has been under pressure. Although various surgical societies have published guidelines on which to base their surgical decisions, it is not clear how general anesthesia has been performed and will be performed in Japan. Social networks such as Twitter are becoming a part of society, as various information is accumulated on the web. One of the services provided by Twitter is a voting function—Twitter Polls—through which anonymous surveys can be conducted. We analyzed the results of a series of surveys on Twitter on the status of operating restrictions using a mathematical model, and public data provided by the Japanese Government were used to estimate the current and future changes in the number of general anesthesia performed in Japan. The number of general anesthesia was estimated as $1.91*10^4$ cases per week on May 16, 2020, approximately a half reduction.

Background

Coronavirus (COVID-19), caused by severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2), was first reported in Wuhan, Hubei, China, and has since spread to more than 200 other countries around the world at the time of writing [1]. In light of this situation, the Japanese Government declared a state of emergency in seven regions of Japan on April 7, 2020 under the provisions of the law. This declaration was extended to the entire nation on the April 17, 2020. The incidence of pneumonia from COVID-19 is considerably higher than from seasonal influenza, and the number of cases with no identifiable route of infection has increased rapidly [2, 3]. The medical care delivery

system has been under pressure [4]. Emphasis should be placed on maintaining medical care systems. Although various surgical societies have published guidelines on which to base their surgical decisions [5, 6], it is not clear how general anesthesia has been performed and will be performed in Japan.

Social networks such as Twitter are becoming a part of our society, as various information is being accumulated on the web [7]. Twitter (Twitter, Inc.) is headquartered in San Francisco, California, United States, and was launched in July, 2006. One of the services provided is a voting function called Twitter Polls. Using this function, anonymous surveys can be conducted on Twitter [8].

We analyzed the results of a series of surveys conducted on Twitter on the status of operating restrictions using a mathematical model, and public data provided by the Japanese Government were used to estimate the current and future changes in the number of general anesthesia carried out in Japan.

Methods

Twitter Surveys

Using the Twitter account @dajhiroki, Twitter Polls was used to conduct eight 24-hour surveys, which were held approximately one week apart from each other, from March 13, 2020 to May 16, 2020. During the period, the spread of COVID-19 had become a social problem in Japan.

The surveys used the same wording throughout the period, in the form of a choice of one of the following four options:

1. No surgical restrictions

2. Partial restrictions (more than half of the usual)

3. Extensive restrictions (less than half of the usual)

4. No scheduled surgery

79   **Transition of responses to the survey**

80   Let the number of responses to the survey be denoted as $A_{t,*} = \{a_{t,1}, a_{t,2}, a_{t,3}, a_{t,4}\}$,

81    $t = \{1, \dots, T\}$ and the proportion of responses to the survey be the simplex $\theta_{t,i}$, where

82    $\sum_{i=1}^{4} \theta_{t,i} = 1$, $T = 8$. The transition probability from the ith answer to the $(i + 1)$th

83   answer is defined as $g_{i \to i+1}$ (Figure 1). The transition matrix denoted as follows

84
$$Q = \begin{pmatrix} g_{1\to1} & g_{1\to2} & 0 & 0 \\ g_{2\to1} & g_{2\to2} & g_{2\to3} & 0 \\ 0 & g_{3\to2} & g_{3\to3} & g_{3\to4} \\ 0 & 0 & g_{4\to3} & g_{4\to4} \end{pmatrix}$$

86   and the proportion of responses to the questionnaire transited from θ_t to θ_{t+1}

87   by

88
$$\theta_{t+1} = Q^T \theta_t$$

89   Parameters were inferred through likelihood function $L(\Theta)$ as

90
$$L(\Theta) = \sum_{t=1}^{T} \left\{ \frac{(a_{t,1} + a_{t,2} + a_{t,3} + a_{t,4})!}{a_{t,1}! \, a_{t,2}! \, a_{t,3}! \, a_{t,4}!} \prod_{i=1}^{4} \theta_{t,i}^{a_{t,i}} \right\}$$

where Θ was the set of parameters.

Figure 1: Transition scheme of the proportion of surveys. (1: green) no surgical restrictions, (2: yellow)

partial restrictions, (3: orange) extensive restrictions, and (4: red) no scheduled surgery. $g_{i \to i+1}$ is the

transition probability from answer i to answer $i + 1$.

Estimation of the number of general anesthesia performed in Japan during

COVID-19

The restriction rate of the number of general anesthesia, p_i, $i = \{1, 2, 3, 4\}$ was

assumed for each response of the surveys: (1) no surgical restrictions were almost the

same as the past statistics, (2) partial restrictions was about 70% restriction rate

compared to the past statistics, (3) extensive restrictions was about 40% restriction rate

compared to the past statistics, and (4) no scheduled surgery was about 10% restriction

rate compared to the past statistics. Under such assumption, restriction rate for each

hospital h at week t, $p_{t,i,h}$ was generated from probability distribution as follows;

(1) for no surgical restrictive hospitals, $p_{t,1,h}$ was generated from uniform distribution,

U, ranging from 0.95 to 1.05.

109
$$p_{t,1,h} \sim U(0.95, 1.05).$$

110 (2-4) for restrictive management hospitals, $p_{t,i,h}$, $i = \{2, 3, 4\}$, was generated from beta

111 distribution, $Beta(\alpha, \beta)$. Their parameters $(\alpha_{t,i,h}, \beta_{t,i,h})$, $i = \{2, 3, 4\}$, were defined so

112 that the mean was 0.7, 0.4, 0.1 and the variance was 0.005, respectively (figure 2).

113
$$p_{t,i,h} \sim Beta(\alpha_{t,i,h}, \beta_{t,i,h}), \ i = \{2, 3, 4\}.$$

Figure 2: Distribution of the degree of restriction for each response group. The mean percentages of the

restrictions were generated from probability distributions. In particular, no surgical restrictions (green)

were generated from the uniform distribution $U(0.95, 1.05)$, partial restrictions (yellow), extensive

restrictions (orange), and no scheduled surgery (red) were generated from the beta distribution

$Beta(\alpha, \beta)$. The set of parameters for beta distribution, (α, β), were defined so that the respective means

were 0.7, 0.4, and 0.1, and the variances were 0.005.

The number of general anesthesia performed at hospital h in the tth week, $R_{h,t}$, was

generated as follows:

$$R_{h,t} \sim \begin{cases} round(N_h p_{t,1,h}) & if\ i = 1 \\ Binomial(p_{t,i,h}, N_h) & otherwise \end{cases},$$

where N_h is the reference number of the general anesthesia performed at each hospital

(data available from e-stat), $round$ is a rounding function, and $Binomial(p, N)$ is a

binomial sampling function that performs N Bernoulli trials with probability p.

The answer to the survey for each hospital at $t = 1$ was randomly allocated to the ith

answer according to $\theta_{1,i}$ and its status transitioned to the $(i + 1)$th answer by $g_{i \to i+1}$.

For sensitivity analysis, two other scenarios were considered. An optimistic scenario

assumed relatively high performance in operating rooms, with an average of 80% for

"2. Partial restrictions," 50% for "3. Extensive restrictions," and 25% for "4. No

scheduled surgery." A pessimistic scenario assumed extremely low performance, with

an average of 50% for "2. Partial restrictions," 30% for "3. Extensive restrictions," and

3% for "4. No scheduled surgery."

Analysis and estimation were performed using Markov chain Monte Carlo methods

using the rstan package (version 2.19.3) within the R programing language (version

3.3.4). Point estimates and their statistical uncertainty were presented as posterior

medians and their 95% credible intervals (CrIs). The data were obtained from the Japan

official statistics portal site (e-stat), a public database in Japan [https://www.e-

stat.go.jp/stat-search/file-download?statInfId=000031565245&fileKind=0], and were

preprocessed for analysis. The code is available online

[https://github.com/yfujii08/covid19generalanesthesia] and is available upon request

from the corresponding author.

Results

From March 13, 2020 to May 2, 2020, survey responses were solicited

approximately weekly via a Twitter account (@dajhiroki) in Japanese language, with

the number of responses received ranging from 47 to 288 (Supplementary Information

S1).

The results of the surveys are presented in Table 1. Prevalence of COVID-19 in

Japan during the period of the surveys is presented in Table 2

date of survey	2020/3/13	2020/3/20	2020/3/27	2020/4/6	2020/4/10	2020/4/17	2020/4/24	2020/5/2	2020/5/9	2020/5/16
number of votes	112	47	73	160	221	232	288	184	93	113

1. No surgical restrictions	89%	77%	82%	64%	46%	27%	25%	29%	41%	39%
2. Partially restricted	5%	17%	7%	23%	38%	38%	38%	40%	37%	44%
3. Extensively restricted	4%	2%	4%	7%	13%	27%	30%	23%	19%	12%
4. No scheduled surgery	2%	4%	7%	6%	3%	8%	7%	8%	3%	5%

155 Table 1: Results of the Twitter surveys

	2020/3/13	2020/3/20	2020/3/27	2020/4/6	2020/4/10	2020/4/17	2020/4/24	2020/5/2	2020/5/9	2020/5/1
tested positive cases (case within 7 days)	659(326)	928(233)	1349(427)	3569(1208)	5246(2425)	9027(3326)	12240(3242)	14383(1689)	15581(708)	16112(531
active cases	520	681	920	2921	4461	7884	10469	11558	6250	378
serious cases	32	50	56	79	117	207	263	324	267	23

Table 2: Prevalence of COVID-19 in Japan during the period of the surveys [9]

From the public database [https://www.e-stat.go.jp/stat-search/file-

download?statInfId=000031565245&fileKind=0], there were 3,501 hospitals that were

registered as facilities that provide general anesthesia. Of the 3,311 hospitals that

reported the number of general anesthesia, 1,989 performed more than 100 general

anesthesia.

From the survey conducted in the first week (March 13, 2020), 89.2% had no surgical

restrictions, but this proportion decreased to 24.2% in the survey conducted in the

seventh week (April , 27 2020) and slightly recovered to 38.2% (May 16, 2020). The

proportions of partial restrictions, extensive restrictions, and no scheduled surgeries

were, 5.2%, 3.9%, and 1.8% (March 13, 2020), 37.8%, 30.4%, and 7.5% (April 27,

2020), and 43.8%, 11.7%, and 5.3% (May 16, 2020), respectively (Figure 3).

The number of general anesthesia performed decreased as the restrictions increased
(Figure 4 and Table 3). Before COVID-19, $4.45 * 10^4$ cases of general anesthesia were
performed per week in Japan. The number of general anesthesia (10^4 per week unit)
was estimated to be 4.18 [95% CrI: 3.96, 4.31] in the first week survey (March 13,
2020) and 1.91 [1.28, 2.61] in the tenth week survey (May 16, 2020) (Table 3).

Figure 3: Transition of the proportion of responses to the survey. The transition probability g was fitted
to the data $t = \{1, 2, ..., T\}$, $T = 10$. The median proportions are shown with 95% credible intervals.

The number of general anesthesia performed decreased as the restrictions increased
(Figure 4 and Table 3). Before COVID-19, $4.45 * 10^4$ cases of general anesthesia were
performed per week in Japan. The number of general anesthesia (10^4 per week unit) was
estimated to be 4.18 [95% CrI: 3.96, 4.31] in the first week survey (March 13, 2020)
and 1.91 [1.28, 2.61] in the tenth week survey (May 16, 2020) (Table 3).

Figure 4: Estimation of the number of general anesthesia in Japan from the survey results. The green violinplot shows the estimated number of general anesthesia performed during the period of the surveys, and the yellow violinplot shows the estimated number of general anesthesia after the period of the surveys. The violinplots show the distribution of the estimated number of general anesthesia performed at 1,989 hospitals in Japan.

	2020/3/13	2020/3/20	2020/3/27	2020/4/6	2020/4/10	2020/4/17	2020/4/24	2020/5/2	2020/5/9	2020/5/16
2.50%	4.16	3.94	3.71	3.49	3.27	3.05	2.85	2.65	1.53	1.28
50%	4.21	4.00	3.78	3.57	3.36	3.16	2.96	2.77	2.06	1.91
97.50%	4.26	4.06	3.85	3.65	3.45	3.27	3.08	2.90	2.73	2.61

Table 3: Estimated number of general anesthesia (10^4 cases per week) in Japan during

the period of the surveys.

Sensitivity analysis showed that the number of general anesthesia performed decreased according to the intensity of the restrictions. From the first week survey (March 13, 2020), the optimistic scenario had a median of $4.24*10^4$ /week, while the pessimistic

scenario had a median of 4.11. From the tenth week survey (May 2, 2020), the

optimistic scenario median was 2.33, while that of the pessimistic scenario was 1.47,

respectively.

Figure 5: Estimation of the number of general anesthesia in Japan from the results of the surveys for the

three scenarios as a sensitivity analysis. The vertical bars show the 95% credible intervals.

Discussion

It is difficult to adequately control for the level of limitation of usual surgical care

during the expansion of COVID-19. It is determined by adherence to official guidelines,

restrictions on medical supplies and medical staff, and the level of need for surgery.

Still, because circumstances can change quickly, it is often determined by reference to

the extent of restrictions at a medical facility other than the home facility.

The staged approach—recommended by the Centers for Medicare and Medicaid

Services (CMS) and the American College of Surgeons (ACS)—is used as a guide to

determine how to perform surgery in situations where the preservation of ventilators

and personal protective equipment is necessary, and the ICU has been compromised—

or is expected to be compromised soon.

The number of respondents to our surveys increased over time, and it is thought that

the number of respondents is dependent on their training in knowing the status of

operational limitations. By using a series of surveys conducted through Twitter Polls,

we were able to quickly determine the extent of the restrictions on surgery at medical

facilities across the country. By periodically soliciting responses to the same survey on

Twitter, we were able to estimate the extent to which operations would be restricted

nationally—over time.

Our methodology in this study is an estimation method that uses mathematical

analysis. In this study, we used public data from a database operated by the Japanese

Government. The number of hospitals accredited by the Japanese Society of

Anesthesiologists (JSA) is about 1,400. However, in this study, 1,989 hospitals that

perform at least 100 surgeries per year under general anesthesia were used. The number

of operations performed at each hospital varies—mainly according to the size of the

hospital. On the other hand, since neither the size of the hospitals to which the survey

respondents belonged nor the changes in the number of operations were known, we

used mathematical analysis to calculate the number of operations.

This study has several limitations.

In this study, we used a database of the number of general anesthetic procedures. If

the number of surgeries performed under local anesthetic was increasing due to

COVID-19, we cannot deny the possibility of discrepancies with the actual number of

operations.

The number and type of survey responses is a potential limitation.

It is possible that the number of anesthesiologists using Twitter was not sufficient,

and there may be some kind of bias in terms of the responses from those that do use it.

Responses were low initially, with 47 in the early stages of the period of the

surveys, but increased to 288 in the latter half of the period of the surveys. The growing

penetration of the surveys and increasing familiarity with the issue of operating room

restrictions may have contributed to an increase in participation, but this may have been

biased.

In this study, we calculated the number of surgeries as 95% to 105% for the criteria:

"1. No surgical restrictions," 70% for "2. Partial restrictions," 40% for "3. Extensive

restrictions," and 10% for "4. No scheduled surgery." In the sensitivity analysis, two

scenarios were assumed, but the number of general anesthesia performed could be

approximately 1.6-fold difference following the period of the surveys. Because it is

difficult to estimate the accurate the extent of restrictions, intuitive values were adopted.

It needs to be compared with the descriptive statistics after era of COVID-19.

Conclusions

In this study, we conducted a series of surveys using Twitter Polls, and performed a

mathematical analysis of the results using a public database on the number of general

anesthetic procedures to estimate the changes in the number of surgeries carried out in

Japan.

Acknowledgment

This work was supported by a research grant from the Kansai Medical University

(KMU) research consortium to K.H., the branding program as a world-leading research

university on intractable immune and allergic diseases from MEXT Japan. We would

like to thank Editage (www. editage.jp) for English language editing.

Author contributions

H.D, Y.F, and K.H. conceived and designed the study. Y.F performed the analysis.

H.D, Y.F, and K.H. prepared figures and/or tables and wrote the paper. All authors read

and approved the final manuscript.

Competing interests

The authors declare no competing interests.

Additional information

Supplementary Information S1

References

1.   Zhu N, Zhang D, Wang W, Li X, Yang B, Song J, et al. A Novel Coronavirus

from Patients with Pneumonia in China, 2019. N Engl J Med. 2020;382:727-33; doi:

10.1056/NEJMoa2001017.

2.   Guan WJ, Ni ZY, Hu Y, Liang WH, Ou CQ, He JX, et al. Clinical

Characteristics of Coronavirus Disease 2019 in China. N Engl J Med. 2020;382:1708-

20; doi: 10.1056/NEJMoa2002032.

3.   Meselson M. Droplets and Aerosols in the Transmission of SARS-CoV-2. N

Engl J Med. 2020; doi: 10.1056/NEJMc2009324.

4.   Zarzaur BL, Stahl CC, Greenberg JA, Savage SA, Minter RM. Blueprint for

Restructuring a Department of Surgery in Concert With the Health Care System During

a Pandemic: The University of Wisconsin Experience. JAMA Surg. 2020; doi:

10.1001/jamasurg.2020.1386.

5.   Surgeons ACo. COVID-19: Elective Case Triage Guidelines for Surgical Care

2020. Available from: https://www.facs.org/covid-19/clinical-guidance/elective-case.

6.   The Centers for Medicare & Medicaid Services C. Non-Emergent, Elective

Medical Services, and Treatment Recommendations 2020. Available from:

https://www.cms.gov/files/document/cms-non-emergent-elective-medical-

recommendations.pdf.

7.   Aramaki E, Maskawa S, Morita M. Twitter Catches The Flu: Detecting

Influenza Epidemics using Twitter. Proceedings of the Conference on Empirical

Methods in Natural Language Processing. 2011:1568–76.

8.   Sang E, Bos J. Predicting the 2011 dutch senate election results with twitter.

Proceedings of the Workshop on Semantic Analysis in Social Media. 2012:53–60.

9.   Japan MoHaLo. Situation update from March 06 to May 02, 2020, on

coronavirus diseases 2019. 2020. Available from:

https://www.mhlw.go.jp/stf/newpage_11163.html (in Japanese).