

1

2 Estimation of the number of general anesthesia cases based on a series
3 of nationwide surveys on Twitter during COVID-19 in Japan: A statistical
4 analysis

5

6 Hiroki Daijo MD, Ph.D. ^{1,4}, Yosuke Fujii MD, ^{2,4}, Kiichi Hirota MD, Ph.D. ^{3,*}

7 ¹ Department of Anesthesia, Takarazuka Daiichi Hospital, Takakazuka, Hyogo 665-
8 0832 Japan

9 ² Department of Anesthesia, Otsu City Hospital, Otsu, Shiga 520-0804 Japan

10 ³ Department of Human Stress Response Science, Institute of Biomedical Science,
11 Kansai Medical University, Hirakata, Osaka 573-1010 Japan.

12 ⁴ These authors contributed equally to this work

13

14 *corresponding author

15 Kiichi Hirota

16 Department of Human Stress Response Science, Institute of Biomedical Science,
17 Kansai Medical University, Hirakata, Osaka, Japan

18 phone: +81-72-804-2526

19 email: hifl@mac.com / khirota-kyt@umin.ac.jp

20

21 Keywords: COVID-19; twitter; social network service; general anesthesia; statistical
22 estimation; Japan

23

24

NOTE: This preprint reports new research that has not been certified by peer review and should not be used to guide clinical practice.

25 **Abstract**

26 Coronavirus disease (COVID-19), caused by severe acute respiratory syndrome
27 coronavirus 2 (SARS-CoV-2), was first reported in Wuhan, Hubei, China, and has since
28 spread to more than 200 other countries at the time of writing. In light of this situation,
29 the Japanese Government declared a state of emergency in seven regions of Japan on
30 the April 7, 2020 under the provisions of the law. The medical care delivery system has
31 been under pressure. Although various surgical societies have published guidelines on
32 which to base their surgical decisions, it is not clear how general anesthesia has been
33 performed and will be performed in Japan. Social networks such as Twitter are
34 becoming a part of society, as various information is accumulated on the web. One of
35 the services provided by Twitter is a voting function—Twitter Polls—through which
36 anonymous surveys can be conducted. We analyzed the results of a series of surveys on
37 Twitter on the status of operating restrictions using a mathematical model, and public
38 data provided by the Japanese Government were used to estimate the current and future
39 changes in the number of general anesthesia performed in Japan. The number of general
40 anesthesia was estimated as 1.91×10^4 cases per week on May 16, 2020, approximately a
41 half reduction.

42

43 **Background**

44 Coronavirus (COVID-19), caused by severe acute respiratory syndrome coronavirus
45 2 (SARS-CoV-2), was first reported in Wuhan, Hubei, China, and has since spread to
46 more than 200 other countries around the world at the time of writing [1]. In light of this
47 situation, the Japanese Government declared a state of emergency in seven regions of
48 Japan on April 7, 2020 under the provisions of the law. This declaration was extended
49 to the entire nation on the April 17, 2020. The incidence of pneumonia from COVID-19
50 is considerably higher than from seasonal influenza, and the number of cases with no
51 identifiable route of infection has increased rapidly [2, 3]. The medical care delivery

52 system has been under pressure [4]. Emphasis should be placed on maintaining medical
53 care systems. Although various surgical societies have published guidelines on which to
54 base their surgical decisions [5, 6], it is not clear how general anesthesia has been
55 performed and will be performed in Japan.

56 Social networks such as Twitter are becoming a part of our society, as various
57 information is being accumulated on the web [7]. Twitter (Twitter, Inc.) is
58 headquartered in San Francisco, California, United States, and was launched in July,
59 2006. One of the services provided is a voting function called Twitter Polls. Using this
60 function, anonymous surveys can be conducted on Twitter [8].

61 We analyzed the results of a series of surveys conducted on Twitter on the status of
62 operating restrictions using a mathematical model, and public data provided by the
63 Japanese Government were used to estimate the current and future changes in the
64 number of general anesthesia carried out in Japan.

65

66 **Methods**

67 **Twitter Surveys**

68 Using the Twitter account @dajhiroki, Twitter Polls was used to conduct eight 24-
69 hour surveys, which were held approximately one week apart from each other, from
70 March 13, 2020 to May 16, 2020. During the period, the spread of COVID-19 had
71 become a social problem in Japan.

72 The surveys used the same wording throughout the period, in the form of a choice of
73 one of the following four options:

- 74 1. No surgical restrictions
- 75 2. Partial restrictions (more than half of the usual)
- 76 3. Extensive restrictions (less than half of the usual)
- 77 4. No scheduled surgery

78

79 **Transition of responses to the survey**

80 Let the number of responses to the survey be denoted as $A_{t,*} = \{a_{t,1}, a_{t,2}, a_{t,3}, a_{t,4}\}$,
 81 $t = \{1, \dots, T\}$ and the proportion of responses to the survey be the simplex $\theta_{t,i}$, where
 82 $\sum_{i=1}^4 \theta_{t,i} = 1$, $T = 8$. The transition probability from the i th answer to the $(i + 1)$ th
 83 answer is defined as $g_{i \rightarrow i+1}$ (Figure 1). The transition matrix denoted as follows

84
$$Q = \begin{pmatrix} g_{1 \rightarrow 1} & g_{1 \rightarrow 2} & 0 & 0 \\ g_{2 \rightarrow 1} & g_{2 \rightarrow 2} & g_{2 \rightarrow 3} & 0 \\ 0 & g_{3 \rightarrow 2} & g_{3 \rightarrow 3} & g_{3 \rightarrow 4} \\ 0 & 0 & g_{4 \rightarrow 3} & g_{4 \rightarrow 4} \end{pmatrix}$$

85
 86 and the proportion of responses to the questionnaire transited from θ_t to θ_{t+1}
 87 by


88
$$\theta_{t+1} = Q^T \theta_t$$

89 Parameters were inferred through likelihood function $L(\theta)$ as

90
$$L(\theta) = \sum_{t=1}^T \left\{ \frac{(a_{t,1} + a_{t,2} + a_{t,3} + a_{t,4})!}{a_{t,1}! a_{t,2}! a_{t,3}! a_{t,4}!} \prod_{i=1}^4 \theta_{t,i}^{a_{t,i}} \right\}$$

91 where θ was the set of parameters.

92



93

94 Figure 1: Transition scheme of the proportion of surveys. (1: green) no surgical restrictions, (2: yellow)
 95 partial restrictions, (3: orange) extensive restrictions, and (4: red) no scheduled surgery. $g_{i \rightarrow i+1}$ is the
 96 transition probability from answer i to answer $i + 1$.

97

98 **Estimation of the number of general anesthesia performed in Japan during**
 99 **COVID-19**


100 The restriction rate of the number of general anesthesia, p_i , $i = \{1, 2, 3, 4\}$ was
101 assumed for each response of the surveys: (1) no surgical restrictions were almost the
102 same as the past statistics, (2) partial restrictions was about 70% restriction rate
103 compared to the past statistics, (3) extensive restrictions was about 40% restriction rate
104 compared to the past statistics, and (4) no scheduled surgery was about 10% restriction
105 rate compared to the past statistics. Under such assumption, restriction rate for each
106 hospital h at week t , $p_{t,i,h}$ was generated from probability distribution as follows;
107 (1) for no surgical restrictive hospitals, $p_{t,1,h}$ was generated from uniform distribution,
108 U , ranging from 0.95 to 1.05.

$$109 \quad p_{t,1,h} \sim U(0.95, 1.05).$$

110 (2-4) for restrictive management hospitals, $p_{t,i,h}$, $i = \{2, 3, 4\}$, was generated from beta
111 distribution, $Beta(\alpha, \beta)$. Their parameters $(\alpha_{t,i,h}, \beta_{t,i,h})$, $i = \{2, 3, 4\}$, were defined so
112 that the mean was 0.7, 0.4, 0.1 and the variance was 0.005, respectively (figure 2).

$$113 \quad p_{t,i,h} \sim Beta(\alpha_{t,i,h}, \beta_{t,i,h}), i = \{2, 3, 4\}.$$

114



115

116 Figure 2: Distribution of the degree of restriction for each response group. The mean percentages of the
 117 restrictions were generated from probability distributions. In particular, no surgical restrictions (green)
 118 were generated from the uniform distribution $U(0.95, 1.05)$, partial restrictions (yellow), extensive
 119 restrictions (orange), and no scheduled surgery (red) were generated from the beta distribution
 120 $Beta(\alpha, \beta)$. The set of parameters for beta distribution, (α, β) , were defined so that the respective means
 121 were 0.7, 0.4, and 0.1, and the variances were 0.005.

122

123 The number of general anesthesia performed at hospital h in the t th week, $R_{h,t}$, was
 124 generated as follows:

125
$$R_{h,t} \sim \begin{cases} \text{round}(N_h p_{t,1,h}) & \text{if } i = 1 \\ \text{Binomial}(p_{t,i,h}, N_h) & \text{otherwise} \end{cases},$$

126 where N_h is the reference number of the general anesthesia performed at each hospital
 127 (data available from e-stat), round is a rounding function, and $\text{Binomial}(p, N)$ is a
 128 binomial sampling function that performs N Bernoulli trials with probability p .

129 The answer to the survey for each hospital at $t = 1$ was randomly allocated to the i th
130 answer according to $\theta_{1,i}$ and its status transitioned to the $(i + 1)$ th answer by $g_{i \rightarrow i+1}$.
131 For sensitivity analysis, two other scenarios were considered. An optimistic scenario
132 assumed relatively high performance in operating rooms, with an average of 80% for
133 “2. Partial restrictions,” 50% for “3. Extensive restrictions,” and 25% for “4. No
134 scheduled surgery.” A pessimistic scenario assumed extremely low performance, with
135 an average of 50% for “2. Partial restrictions,” 30% for “3. Extensive restrictions,” and
136 3% for “4. No scheduled surgery.”

137 Analysis and estimation were performed using Markov chain Monte Carlo methods
138 using the rstan package (version 2.19.3) within the R programming language (version
139 3.3.4). Point estimates and their statistical uncertainty were presented as posterior
140 medians and their 95% credible intervals (CrIs). The data were obtained from the Japan
141 official statistics portal site (e-stat), a public database in Japan [[https://www.e-](https://www.e-stat.go.jp/stat-search/file-download?statInfId=000031565245&fileKind=0)
142 [stat.go.jp/stat-search/file-download?statInfId=000031565245&fileKind=0](https://www.e-stat.go.jp/stat-search/file-download?statInfId=000031565245&fileKind=0)], and were
143 preprocessed for analysis. The code is available online
144 [<https://github.com/yfujii08/covid19generalanesthesia>] and is available upon request
145 from the corresponding author.

146

147 Results

148 From March 13, 2020 to May 2, 2020, survey responses were solicited
149 approximately weekly via a Twitter account (@dajhiroki) in Japanese language, with
150 the number of responses received ranging from 47 to 288 (Supplementary Information
151 S1).

152 The results of the surveys are presented in Table 1. Prevalence of COVID-19 in
153 Japan during the period of the surveys is presented in Table 2

date of survey	2020/3/13	2020/3/20	2020/3/27	2020/4/6	2020/4/10	2020/4/17	2020/4/24	2020/5/2	2020/5/9	2020/5/16
number of votes	112	47	73	160	221	232	288	184	93	113

1. No surgical restrictions	89%	77%	82%	64%	46%	27%	25%	29%	41%	39%
2. Partially restricted	5%	17%	7%	23%	38%	38%	38%	40%	37%	44%
3. Extensively restricted	4%	2%	4%	7%	13%	27%	30%	23%	19%	12%
4. No scheduled surgery	2%	4%	7%	6%	3%	8%	7%	8%	3%	5%

154

155 Table 1: Results of the Twitter surveys

156

	2020/3/13	2020/3/20	2020/3/27	2020/4/6	2020/4/10	2020/4/17	2020/4/24	2020/5/2	2020/5/9	2020/5/16
tested positive cases (case within 7 days)	659(326)	928(233)	1349(427)	3569(1208)	5246(2425)	9027(3326)	12240(3242)	14383(1689)	15581(708)	16112(531)
active cases	520	681	920	2921	4461	7884	10469	11558	6250	3784
serious cases	32	50	56	79	117	207	263	324	267	231

157


158 Table 2: Prevalence of COVID-19 in Japan during the period of the surveys [9]

159

160 From the public database [[https://www.e-stat.go.jp/stat-search/file-](https://www.e-stat.go.jp/stat-search/file-download?statInfId=000031565245&fileKind=0)
 161 [download?statInfId=000031565245&fileKind=0](https://www.e-stat.go.jp/stat-search/file-download?statInfId=000031565245&fileKind=0)], there were 3,501 hospitals that were
 162 registered as facilities that provide general anesthesia. Of the 3,311 hospitals that
 163 reported the number of general anesthesia, 1,989 performed more than 100 general
 164 anesthesia.

165 From the survey conducted in the first week (March 13, 2020), 89.2% had no surgical
 166 restrictions, but this proportion decreased to 24.2% in the survey conducted in the
 167 seventh week (April, 27 2020) and slightly recovered to 38.2% (May 16, 2020). The
 168 proportions of partial restrictions, extensive restrictions, and no scheduled surgeries
 169 were, 5.2%, 3.9%, and 1.8% (March 13, 2020), 37.8%, 30.4%, and 7.5% (April 27,
 170 2020), and 43.8%, 11.7%, and 5.3% (May 16, 2020), respectively (Figure 3).


171 The number of general anesthesia performed decreased as the restrictions increased
172 (Figure 4 and Table 3). Before COVID-19, 4.45×10^4 cases of general anesthesia were
173 performed per week in Japan. The number of general anesthesia (10^4 per week unit)
174 was estimated to be 4.18 [95% CrI: 3.96, 4.31] in the first week survey (March 13,
175 2020) and 1.91 [1.28, 2.61] in the tenth week survey (May 16, 2020) (Table 3).



176
177 Figure 3: Transition of the proportion of responses to the survey. The transition probability g was fitted
178 to the data $t = \{1, 2, \dots, T\}$, $T = 10$. The median proportions are shown with 95% credible intervals.

179

180 The number of general anesthesia performed decreased as the restrictions increased
181 (Figure 4 and Table 3). Before COVID-19, 4.45×10^4 cases of general anesthesia were
182 performed per week in Japan. The number of general anesthesia (10^4 per week unit) was
183 estimated to be 4.18 [95% CrI: 3.96, 4.31] in the first week survey (March 13, 2020)
184 and 1.91 [1.28, 2.61] in the tenth week survey (May 16, 2020) (Table 3).



185

186 Figure 4: Estimation of the number of general anesthesia in Japan from the survey results. The green
 187 violinplot shows the estimated number of general anesthesia performed during the period of the surveys,
 188 and the yellow violinplot shows the estimated number of general anesthesia after the period of the
 189 surveys. The violinplots show the distribution of the estimated number of general anesthesia performed at
 190 1,989 hospitals in Japan.

191

	2020/3/13	2020/3/20	2020/3/27	2020/4/6	2020/4/10	2020/4/17	2020/4/24	2020/5/2	2020/5/9	2020/5/16
2.50%	4.16	3.94	3.71	3.49	3.27	3.05	2.85	2.65	1.53	1.28
50%	4.21	4.00	3.78	3.57	3.36	3.16	2.96	2.77	2.06	1.91
97.50%	4.26	4.06	3.85	3.65	3.45	3.27	3.08	2.90	2.73	2.61


192

193 Table 3: Estimated number of general anesthesia (10^4 cases per week) in Japan during
 194 the period of the surveys.

195

196 Sensitivity analysis showed that the number of general anesthesia performed decreased
 197 according to the intensity of the restrictions. From the first week survey (March 13,
 198 2020), the optimistic scenario had a median of 4.24×10^4 /week, while the pessimistic

199 scenario had a median of 4.11. From the tenth week survey (May 2, 2020), the
200 optimistic scenario median was 2.33, while that of the pessimistic scenario was 1.47,
201 respectively.



202

203 Figure 5: Estimation of the number of general anesthesia in Japan from the results of the surveys for the
204 three scenarios as a sensitivity analysis. The vertical bars show the 95% credible intervals.

205

206 Discussion

207 It is difficult to adequately control for the level of limitation of usual surgical care
208 during the expansion of COVID-19. It is determined by adherence to official guidelines,
209 restrictions on medical supplies and medical staff, and the level of need for surgery.
210 Still, because circumstances can change quickly, it is often determined by reference to
211 the extent of restrictions at a medical facility other than the home facility.

212 The staged approach—recommended by the Centers for Medicare and Medicaid
213 Services (CMS) and the American College of Surgeons (ACS)—is used as a guide to
214 determine how to perform surgery in situations where the preservation of ventilators
215 and personal protective equipment is necessary, and the ICU has been compromised—
216 or is expected to be compromised soon.

217 The number of respondents to our surveys increased over time, and it is thought that
218 the number of respondents is dependent on their training in knowing the status of
219 operational limitations. By using a series of surveys conducted through Twitter Polls,
220 we were able to quickly determine the extent of the restrictions on surgery at medical
221 facilities across the country. By periodically soliciting responses to the same survey on
222 Twitter, we were able to estimate the extent to which operations would be restricted
223 nationally—over time.

224 Our methodology in this study is an estimation method that uses mathematical
225 analysis. In this study, we used public data from a database operated by the Japanese
226 Government. The number of hospitals accredited by the Japanese Society of
227 Anesthesiologists (JSA) is about 1,400. However, in this study, 1,989 hospitals that
228 perform at least 100 surgeries per year under general anesthesia were used. The number
229 of operations performed at each hospital varies—mainly according to the size of the
230 hospital. On the other hand, since neither the size of the hospitals to which the survey
231 respondents belonged nor the changes in the number of operations were known, we
232 used mathematical analysis to calculate the number of operations.

233 This study has several limitations.

234 In this study, we used a database of the number of general anesthetic procedures. If
235 the number of surgeries performed under local anesthetic was increasing due to
236 COVID-19, we cannot deny the possibility of discrepancies with the actual number of
237 operations.

238 The number and type of survey responses is a potential limitation.

239 It is possible that the number of anesthesiologists using Twitter was not sufficient,
240 and there may be some kind of bias in terms of the responses from those that do use it.

241 Responses were low initially, with 47 in the early stages of the period of the
242 surveys, but increased to 288 in the latter half of the period of the surveys. The growing
243 penetration of the surveys and increasing familiarity with the issue of operating room

244 restrictions may have contributed to an increase in participation, but this may have been
245 biased.

246 In this study, we calculated the number of surgeries as 95% to 105% for the criteria:
247 "1. No surgical restrictions," 70% for "2. Partial restrictions," 40% for "3. Extensive
248 restrictions," and 10% for "4. No scheduled surgery." In the sensitivity analysis, two
249 scenarios were assumed, but the number of general anesthesia performed could be
250 approximately 1.6-fold difference following the period of the surveys. Because it is
251 difficult to estimate the accurate the extent of restrictions, intuitive values were adopted.
252 It needs to be compared with the descriptive statistics after era of COVID-19.

253

254 **Conclusions**

255 In this study, we conducted a series of surveys using Twitter Polls, and performed a
256 mathematical analysis of the results using a public database on the number of general
257 anesthetic procedures to estimate the changes in the number of surgeries carried out in
258 Japan.

259

260 **Acknowledgment**

261 This work was supported by a research grant from the Kansai Medical University
262 (KMU) research consortium to K.H., the branding program as a world-leading research
263 university on intractable immune and allergic diseases from MEXT Japan. We would
264 like to thank Editage ([www. editage.jp](http://www.editage.jp)) for English language editing.

265

266 **Author contributions**

267 H.D, Y.F, and K.H. conceived and designed the study. Y.F performed the analysis.
268 H.D, Y.F, and K.H. prepared figures and/or tables and wrote the paper. All authors read
269 and approved the final manuscript.

270

271 **Competing interests**

272 The authors declare no competing interests.

273

274 **Additional information**

275 Supplementary Information S1

276

277 **References**

278

- 279 1. Zhu N, Zhang D, Wang W, Li X, Yang B, Song J, et al. A Novel Coronavirus
280 from Patients with Pneumonia in China, 2019. *N Engl J Med.* 2020;382:727-33; doi:
281 10.1056/NEJMoa2001017.
- 282 2. Guan WJ, Ni ZY, Hu Y, Liang WH, Ou CQ, He JX, et al. Clinical
283 Characteristics of Coronavirus Disease 2019 in China. *N Engl J Med.* 2020;382:1708-
284 20; doi: 10.1056/NEJMoa2002032.
- 285 3. Meselson M. Droplets and Aerosols in the Transmission of SARS-CoV-2. *N*
286 *Engl J Med.* 2020; doi: 10.1056/NEJMc2009324.
- 287 4. Zarzaur BL, Stahl CC, Greenberg JA, Savage SA, Minter RM. Blueprint for
288 Restructuring a Department of Surgery in Concert With the Health Care System During
289 a Pandemic: The University of Wisconsin Experience. *JAMA Surg.* 2020; doi:
290 10.1001/jamasurg.2020.1386.
- 291 5. Surgeons ACo. COVID-19: Elective Case Triage Guidelines for Surgical Care
292 2020. Available from: <https://www.facs.org/covid-19/clinical-guidance/elective-case>.
- 293 6. The Centers for Medicare & Medicaid Services C. Non-Emergent, Elective
294 Medical Services, and Treatment Recommendations 2020. Available from:
295 [https://www.cms.gov/files/document/cms-non-emergent-elective-medical-](https://www.cms.gov/files/document/cms-non-emergent-elective-medical-recommendations.pdf)
296 [recommendations.pdf](https://www.cms.gov/files/document/cms-non-emergent-elective-medical-recommendations.pdf).
- 297 7. Aramaki E, Maskawa S, Morita M. Twitter Catches The Flu: Detecting

- 298 Influenza Epidemics using Twitter. Proceedings of the Conference on Empirical
299 Methods in Natural Language Processing. 2011:1568–76.
- 300 8. Sang E, Bos J. Predicting the 2011 dutch senate election results with twitter.
301 Proceedings of the Workshop on Semantic Analysis in Social Media. 2012:53–60.
- 302 9. Japan MoHaLo. Situation update from March 06 to May 02, 2020, on
303 coronavirus diseases 2019. 2020. Available from:
304 https://www.mhlw.go.jp/stf/newpage_11163.html (in Japanese).
305
306
307