Full Title: Correlation between blood and CSF compartment cytokines and chemokines in

subjects with cryptococcal meningitis.

Short Title: Compartment specific cytokine immune environments during infection

Elizabeth C Okafor[1], Katherine H Hullsiek[2], Darlisha A Williams[1,3], James E Scriven[4], Joshua

Rhein[1,3], Henry W Nabeta[3], Abdu K Musubire[3], Radha Rajasingham[1], Conrad Muzoora[5],

Charlotte Schutz[4], Graeme Meintjes[4], David B Meya[1,3], David R Boulware[1].

[1] Department of Medicine, University of Minnesota, Minneapolis, MN, USA.

[2] School of Public Health, University of Minnesota, Minneapolis, MN, USA

[3] Infectious Diseases Institute, Makerere University, Kampala, Uganda.

[4] Wellcome Centre for Infectious Diseases Research in Africa, Institute of Infectious Disease

and Molecular Medicine, and Department of Medicine, University of Cape Town, South Africa.

[5] Department of Medicine, Mbarara University of Science and Technology, Mbarara, Uganda.

Corresponding Author:

Elizabeth Okafor

Microbiology Research Facility 4-103

689 23 rd Ave SE

Minneapolis, MN 5455 USA

Email: okafo039@umn.edu

Alternate: David Boulware boulw001@umn.edu

ABSTRACT

Background:

Though peripheral blood is a crucial sample to study immunology, it is unclear whether the immune environment in the peripheral vasculature correlates with that at the end-organ site of infection. Utilizing cryptococcal meningitis as a model, we investigated the correlation between serum and cerebrospinal fluid biomarkers over time.

Methodology and Principal Findings:

We analyzed the cerebrospinal fluid and serum of 160 subjects presenting with first episode cryptococcal meningitis for soluble cytokines and chemokines measured by Luminex assay. Specimens were collected at meningitis diagnosis, 1-week, and 2-weeks post cryptococcal diagnosis. We compared paired samples by Spearman correlation and the p-value was set at <0.01. Of the 21 analytes tested at baseline, there was no correlation detected between nearly all analytes. A weak negative correlation was found between serum and cerebrospinal fluid levels of interferon-gamma (Rho= -0.214, p= .007) and interleukin- 4 (Rho= -0.232, p=. 003). There was no correlation at 1-week post cryptococcal diagnosis. However, at 2-weeks post diagnosis, there was a weak positive correlation between levels of granulocyte-macrophage colony-stimulating factor (Rho= 0.25, p= .007) between serum and cerebrospinal fluid. No cytokine or chemokine showed consistent correlation overtime.

Conclusion and Significance:

Based on our analysis of 21 biomarkers, serum and cerebrospinal fluid immune responses do not correlate. There appears to be a distinct immune environment in terms of soluble biomarkers in the vasculature versus end-organ site of infection. While this is a model of HIV-related cryptococcal meningitis, we postulate that assuming the blood compartment is representative of the immune function at the end-organ site of infection may not be appropriate.

Author Summary:

In subjects with localized infections, peripheral blood is often used as a surrogate to

investigate the immune environment at the end-organ site of infection. Sampling the end-organ

site of infection is not routinely performed. The current project serves to fill in the gap of

knowledge regarding whether the peripheral blood and end-organ site of infection immune

environments correlate. Due to therapeutic lumbar punctures in cryptococcal meningitis, we are

able to readily sample the cerebrospinal fluid to investigate compartment specific immune

responses by quantifying cytokine and chemokine levels. We determined that there was no

correlation between peripheral blood and cerebrospinal fluid cytokine levels. The findings from

this works serve to caution the use of peripheral blood as a surrogate for compartment specific

immune responses. Though our findings are solely from subjects with cryptococcal meningitis,

more research should be performed in other compartment specific infections such as

pneumonia, osteomyelitis, or hepatitis for validation. Underestimating or overestimating the

activation of the immune response at the end-organ site may impact clinical care and prognosis,

especially if immunomodulating agents are being considered as therapy; thus, determining

whether peripheral blood and end-organ site of infection immune environments correlate is

imperative.

Word Count: Abstract - 260 words; Author Summary - 194; Manuscript – 1529

INTRODUCTION:

Immunologic investigations are critical to understanding the pathogenesis of infectious disease. Often human or animal subjects have their blood drawn to assess cell phenotype, function, or soluble cytokines or chemokines in order to characterize the immune response to infection. Blood is a convenient and available compartment to analyze. However, it is unclear if analysis of cells trafficking in the peripheral vasculature reflects the actual immune response at the end organ site compartment (e.g. CSF in meningitis, lung in pneumonia). This is important because the end organ site of infection is often the primary site from which initial or disease-specific signs and symptoms manifest. Of even more concern is the possibility of unknowingly overestimating or underestimating the immune response based on analysis of immune cells in the vasculature.

Analysis of site-specific immune responses in humans is difficult but not impossible. Cryptococcal meningitis is one model which allows investigation of the compartment-specific immune responses. In cryptococcal meningitis, frequent therapeutic lumbar punctures to control intracranial pressure provide a large volume of CSF over the course of treatment. Analysis of CSF provides information about the immune responses in the central nervous system (CNS). Yet, cryptococcal meningitis is not solely a localized CNS infection, but a systemic infection whereby inhaled cryptococcal spores germinate in the lung and yeast cells disseminate via the blood throughout the body (1, 2). Peripheral blood mononuclear cells, serum, and plasma provide information about immune responses in the peripheral vascular compartment.

In cryptococcosis, but more broadly in infectious disease, the correlation between the immune response in the peripheral vasculature versus immune response in the end-organ compartment is poorly characterized and should be elucidated. We aim to determine the

correlation between 21 peripheral blood and CSF cytokines and chemokines in a cohort of

subjects with first episode cryptococcosis. Utilizing three distinct time points, we analyzed the

correlation between peripheral blood and CSF over time. The results from this analysis serves

to fill the gap in knowledge regarding the correlation between immune responses, particularly

soluble cytokines and chemokines, in the vasculature and tissue compartments utilizing

cryptococcal meningitis as a representative model.

METHODS

110   Subjects with first episode of cryptococcal meningitis were originally enrolled in the

Cryptococcal Optimal ART Timing (COAT) trial from November 2010 to April 2012 to determine

the optimal time to initiate antiretrovirals after cryptococcal meningitis (3). The COAT trial was

conducted at: Mulago Hospital, Kampala, Uganda; Mbarara University of Science and

Technology, Mbarara, Uganda; and the former G.F. Jooste Hospital in Cape Town, South

Africa. All subjects completed written informed consent and the institutional review board from

each organization approved the study protocol. Details regarding this study trial have been

previously described (Clinicaltrials.gov Identifier No: NCT01075152) (3).

119   We collected peripheral blood and CSF samples at cryptococcal meningitis diagnosis,

trial randomization at 1-week, and 2-weeks post cryptococcal meningitis diagnosis for cytokines

and chemokine analysis. Serum and CSF were analyzed in a targeted multiplex panel for 19

cytokines and chemokines using a Luminex multiplex cytokine panel (Bio-Rad, Hercules, CA)

(4), following the manufacturer's instructions being run at 1:2 dilution taking the average of two

replicates. We also measured CD14 and CD163 by ELISA (R&D Systems, Minneapolis, MN).

We have previously reported the longitudinal CSF changes by trial arm (4).

As secondary analysis, we assessed the correlation, after log transformation of raw

readout, between the intra-subject matched serum and CSF using non-parametric Spearman

correlation. Data at each time point represents all subjects included in the previous study

irrespective of trial arm randomization. Changes in cohort size represents fluctuations in

specimen availability, most often due to subjects who declined therapeutic lumbar punctures or

subjects who unfortunately died. Significance was set at a p-value of 0.01 to adjust for multiple

comparisons in a post-hoc exploratory analysis.

RESULTS

Of 177 ART-naïve subjects with first episode of cryptococcal meningitis enrolled into the

COAT trial, matching CSF and serum existed for 160 subjects at baseline, 105 subjects at 1-

week, and 115 subjects at 2-weeks. Missing specimens were most often due to lack of CSF

collection on the same day as serum collection as subjects could refuse therapeutic lumbar

punctures. Clinical and demographic data comparing COAT trial arms (early versus late ART

initiation) have been previously described (4). Of the160 subjects included, 79 subjects (49%)

were female. The median age was 35 (interquartile range (IQR) 28.5 to 40) years with a median

CD4 count of 25 (IQR 10 to 74) cells/μL determined at 1-week post diagnosis.

We assessed whether there was a correlation between immune biomarkers in CSF

versus serum for 21 analytes irrespective of study arm randomization. At time of diagnosis,

none of the 21 immune biomarkers had a positive correlation by Spearman's non-parametric

test **(Table 1)**. Two analytes had statistically significant but weak negative correlations; the type

1 T-helper cell (Th_1) cytokine interferon-gamma (IFN-γ, Rho= -0.214, p=.007) and the type 2 T-

helper cell cytokine (Th_2) interleukin-4 (IL-4, Rho= -0.232, p=.003). While statistically significant,

the correlations were relatively weak and inverse for both IFN-γ and IL-4 **(Figure 1)**, meaning as

blood levels increased, CSF levels decreased.

Of the 19 biomarkers analyzed among 105 paired samples at 1-week post cryptococcal

meningitis diagnosis, no cytokines or chemokines were statistically correlated. Of the 21

biomarkers analyzed at 2-weeks post diagnosis among 115 paired samples, a weak positive

correlation existed between serum and CSF levels of granulocyte-macrophage colony-

stimulating factor (GM-CSF, Rho=0.25, p=.007). Across the three time points as subjects

continued standard antifungal therapy, there were no immune biomarkers that were consistently

positively or negatively correlated between blood and CSF (**Table 2**). Additionally, analysis of

rate of change over time, through slope calculation across all three time points, demonstrated

no significant positive or negative correlation between blood and CSF (**Table 2**).

DISCUSSION

We hypothesized that immune biomarkers would have a general positive correlation

between serum and CSF. Surprisingly, we found no consistent correlation between soluble

immune biomarkers in the peripheral vasculature versus at the site of infection in the CSF. The

cytokine environment was strikingly different, even though cryptococcosis is a disseminated

infection throughout the vasculature, lungs, and CNS (2, 5, 6). Additionally, no single analyte

had a significant correlation pattern over the course of the first two weeks of standard antifungal

therapy. The results from this focused analysis provide evidence that peripheral blood analytes

are not necessarily similar to CSF analytes. This builds upon our previous findings which

demonstrated that lymphocyte populations and activation levels in CSF are distinct from

peripheral blood (7).

There is a large body of published literature comparing human peripheral blood and CSF

cytokines and chemokines across various infectious etiologies of meningitis. Comparatively,

little literature has been published regarding the correlation within the same individuals' between

human tissue compartments and peripheral blood analytes in infectious etiologies of meningitis

(8) or other conditions. Additionally, prior to this study, there was no literature characterizing

CSF and blood analyte correlations over time in subjects with cryptococcal meningitis.

Employing cryptococcal meningitis as a model of compartmentalization, we raise a

fundamental question of whether the immunology observed in peripheral blood represents what

occurs at the end-organ site of infection. This is particularly important as the infectious disease

field continues to consider the potential benefits and risks of immunomodulating agent adjuvant

therapy, such as administration of proinflammatory cytokines and/or monoclonal antibodies to

inhibit anti-inflammatory cytokines or exhaustion receptor engagement. Therefore, it is crucial to

truly understand the immune environment at the end-organ site of infection.

The effector immune response at the site of infection appears to be very different from

that of cells trafficking in blood. Previous studies on cryptococcal meningitis immune

reconstitution inflammatory syndrome documented differential expression of leukocyte

phenotype when comparing peripheral blood to the CSF (7). We doubt these findings are

unique to cryptococcosis, for example as demonstrated by differential gene expression when

comparing pericardial fluid to peripheral blood in subjects with pericardial tuberculosis (9). There

is also the potential impact of tissue niche on leukocyte activation and cytokine environment.

Research on compartmentalization in infection is limited and thus characterizing the immune

response in peripheral blood vs end-organ compartments should be rigorously assessed in

other localized infections.

Our study is restricted to immunocompromised persons with cryptococcosis, thus

whether the lack of correlation between tissue-specific and peripheral vasculature compartment

immune responses is universal, requires future study. Though 160 subjects were included in the

baseline analysis, our data are limited by the absence of 35% and 30% of study participants'

specimens at 1 and 2 weeks, respectively. However, we postulate that more specimens would

not have changed the generally poor strength of the correlations. The strength of correlations for

all analytes was generally weak with Rho correlation coefficient ranging from -0.23 to +0.25.

Thousands of specimens may have been able to generate a statistically significant p-value, but

the correlations were weak. Additionally, a more stringent and formal p-value calculation, (for

example Bonferroni correction or Benjamini-Hochberg procedure) would not have changed the

overall weak correlation.

In conclusion, we demonstrate that cytokines and chemokines released upon leukocyte

activation do not correlate between serum and CSF in cryptococcal meningitis. This

demonstrates that, though readily available, utilizing peripheral blood as an approximation for

the immune status in tissue compartments, such as the CNS, is potentially unreliable.

Acknowledgements: We thank the medical officers, nurses, and laboratory staff who were a

part of the COAT Trial Study Team. Additionally, we express our deepest gratitude to all study

participants and their families, healthcare professionals who provided care for study

participants, and laboratory staff at all study sites.

Disclaimer: The findings from this project have not been previously presented at any

conferences and has not been previously published.

Potential Conflict of Interest: The authors have declared that no competing interests exist.

References

1.	Coelho C, Bocca AL, Casadevall A. The Intracellular Life of Cryptococcus neoformans. Annu Rev Pathol. 2014;9:219-38.

2.	Denham ST, Brown JCS. Mechanisms of Pulmonary Escape and Dissemination by *Cryptococcus neoformans*. Journal of Fungi. 2018;4(25).

3.	Boulware DR, Meya DB, Muzoora C, Rolfes MA, Huppler Hullsiek K, Musubire A, et al. Timing of antiretroviral therapy after diagnosis of cryptococcal meningitis. N Engl J Med. 2014;370(26):2487-98.

4.	Scriven JE, Rhein J, Hullsiek KH, von Hohenberg M, Linder G, Rolfes MA, et al. Early ART after cryptococcal meningitis is associated with cerebrospinal fluid pleocytosis and macrophage activation in a multisite randomized trial. J Infect Dis. 2015;212(5):769-78.

5.	Vanherp L, Ristani A, Poelmans J, Hillen A, Lagrou K, Janbon G, et al. Sensitive bioluminescence imaging of fungal dissemination to the brain in mouse models of cryptococcosis. Dis Model Mech. 2019;12(6):dmm039123.

6.	Nicola AM, Robertson EJ, Albuquerque P, Derengowski LdS, Casadevalla A. Nonlytic Exocytosis of Cryptococcus neoformans from Macrophages Occurs In Vivo and Is Influenced by Phagosomal pH. mBio. 2011;2(4):e00167-11.

7.	Meya DB, Okurut S, Zziwa G, Rolfes MA, Kelsey M, Cose S, et al. Cellular immune activation in cerebrospinal fluid from ugandans with cryptococcal meningitis and immune reconstitution inflammatory syndrome. J Infect Dis. 2015;211(10):1597-606.

8.	Fassbender K, Schminke U, Ries S, Ragoschke A, Kischka U, Fatar M, et al. Endothelial-derived adhesion molecules in bacterial meningitis: association to cytokine release and intrathecal leukocyte-recruitment. J Neuroimmunol. 1997;74(1-2):130-4.

9.	Matthews K, Deffur A, Ntsekhe M, Syed F, Russell JBW, Tibazarwa K, et al. A Compartmentalized Profibrotic Immune Response Characterizes Pericardial Tuberculosis, Irrespective of HIV-1 Infection. Am J Respir Crit Care Med. 2015;192(12):1518-21.

273 **Table 1: Correlation of Paired CSF and Serum Analytes at Baseline, 1-Week, and 2-Weeks**

274 **post Cryptococcal Meningitis Diagnosis.**

	At CM diagnosis			1-week post diagnosis			2-weeks post CM diagnosis		
Biomarker	**N**	**Rho**	**P-value**	**N**	**Rho**	**P-value**	**N**	**Rho**	**P-value**
Innate Immune response									
IL-6	160	0.021	0.79	105	-0.025	0.80	115	-0.228	0.01
IL-1β	160	-0.042	0.60	105	-0.122	0.22	115	0.052	0.58
IL-8	160	-0.034	0.67	105	0.141	0.15	115	0.035	0.71
G-CSF	160	-0.003	0.97	105	0.1	0.31	115	0.087	0.36
GM-CSF	160	0.033	0.68	105	0.16	0.10	115	0.25	0.007*
MCP-1	160	0.133	0.09	105	0.216	0.03	115	-0.117	0.21
MIP-1α	75	0.01	0.93	5	0.4	0.51	93	-0.07	0.51
MIP-1β	160	0.117	0.14	105	0.114	0.25	115	-0.057	0.55
CD14	87	0.078	0.47	n	N/A	N/A	83	0.187	0.09
CD163	87	-0.249	0.02	n	N/A	N/A	83	-0.099	0.38
Adaptive Immune response									
IFN-γ	160	-0.214	0.007*	105	0.089	0.37	115	0.198	0.03
TNF-α	160	0.077	0.33	105	0.038	0.70	115	0.078	0.41
IL-12	160	0.09	0.26	105	-0.153	0.12	115	-0.005	0.96
IL-2	160	-0.025	0.76	105	-0.041	0.68	115	0.22	0.02
IL-7	160	0.038	0.64	105	0.11	0.26	115	-0.004	0.96
IL-10	160	0.021	0.80	105	0.045	0.65	115	-0.025	0.80
IL-5	160	-0.046	0.56	105	0.099	0.32	115	0.004	0.97
IL-4	160	-0.232	0.003*	105	0.107	0.28	115	-0.064	0.49
IL-13	160	0.056	0.48	105	-0.004	0.97	115	-0.057	0.54
IL-17	160	-0.008	0.92	105	0.128	0.19	115	0.012	0.90
Hemostasis									
VEGF	111	-0.049	0.61	62	0.062	0.63	115	-0.002	0.99

*P value of less than 0.01.

Interleukin (IL), Interferon (IFN), Tumor- necrosis factor (TNF), Granulocyte-colony stimulating factor (G-CSF),

Granulocyte-macrophage colony stimulating factor (GM-CSF), Monocyte chemoattractant protein-1/CCL2 (MCP-1),

Macrophage inflammatory protein (MIP), and Vascular Endothelial Growth Factor (VEGF)

Figure 1: **Scatterplot of Interferon-Gamma (IFN-γ) and Interleukin-4 (IL-4) at Cryptococcal**

Meningitis in CSF Versus Serum.

IFN-γ is a Th_1 cytokine which is protective in cryptococcosis causing classical activation of macrophages to enhance

killing of yeast. IL-4 is Th_2 cytokine which is not protective in mouse models of cryptococcosis. Both cytokines have a

weak, inverse correlation between levels observed in serum versus CSF. Levels are in pg/mL.

Table 2. Correlation of CSF and Serum Cytokines Pooled Across All Three Time Points at Baseline, 1-Week, and 2-Weeks post Cryptococcal Meningitis Diagnosis.

BIOMARKER	N	Rho	P-value	N	Rho	P-value
	Combined Correlation			*Rate of Change Over Time*		
Innate Immune response						
IL-6	403	-0.07	0.16	160	0.151	0.06
IL-1β	403	0.029	0.57	160	0.070	0.38
IL-8	403	0.029	0.56	160	0.132	0.10
G-CSF	403	0.067	0.18	160	0.086	0.28
GM-CSF	403	0.147	0.003*	160	-0.030	0.70
MCP-1	403	0.073	0.15	160	0.111	0.16
MIP-1α	214	-0.127	0.06	118	0.061	0.51
MIP-1β	403	0.10	0.05	160	-0.119	0.13
CD14	312	0.095	0.09	128	0.180	0.04
CD163	312	-0.051	0.37	128	-0.143	0.11
Adaptive Immune response						
IFN-γ	403	0.119	0.02	160	0.120	0.13
TNF-α	403	0.102	0.04	160	0.096	0.23
IL-12	403	-0.021	0.67	160	-0.003	0.97
IL-2	403	0.089	0.08	160	-0.057	0.48
IL-7	403	0.036	0.47	160	-0.032	0.69
IL-10	403	-0.062	0.21	160	0.068	0.40
IL-5	403	0.041	0.41	160	0.059	0.46
IL-4	403	0.082	0.10	160	-0.063	0.43
IL-13	403	-0.093	0.06	160	0.194	0.014
IL-17	403	0.066	0.19	160	-0.082	0.30
				160		
Hemostasis						
VEGF	307	0.051	0.37	147	-0.059	0.48

*P value of less than 0.01.

Interleukin (IL), Interferon (IFN), Tumor- necrosis factor (TNF), Granulocyte-colony stimulating factor (G-CSF), Granulocyte-macrophage colony stimulating factor (GM-CSF), Monocyte chemoattractant protein-1/CCL2 (MCP-1), Macrophage inflammatory protein (MIP), and Vascular Endothelial Growth Factor (VEGF)