Healthcare workers preparedness for COVID-19 pandemic in the occupied Palestinian territory: a cross-sectional survey

Authors: Osaid Alser, MD, MSc(Oxon)[1*]; Heba Alghoul, MD[2]; Zahra Alkhateeb, BSc, MS[3]; Ayah Hamdan, BSc[4]; Loai Albaraqouni, MD, PhD[5]; Kiran Saini, BM BCh(Oxon)[6]

Affiliation:

1. Ministry of Health, Gaza Strip, occupied Palestinian territory and OxPal Medlink, UK.
2. Faculty of Medicine, Islamic University of Gaza, occupied Palestinian territory.
3. Vertex Pharmaceuticals, Boston, USA.
4. Harvard T.H. Chan School of Public Health, Boston, USA.
5. Institute for Evidence-Based Healthcare, Faculty of Health Sciences & Medicine, Bond University, Australia
6. Medical Sciences Division, University of Oxford and OxPal Medlink, UK.

*Corresponding author: Dr Osaid Alser, MD, MSc (Oxon)
165 Cambridge St, Suite 810, Massachusetts General Hospital, Harvard Medical School, Boston, MA 02114, USA. E-mail: osaidalserr@hotmail.com

Abstract:

Background: The coronavirus disease 19 (COVID-19) pandemic threatens to overwhelm the capacity of the vulnerable healthcare system in the occupied Palestinian territory (oPt). Sufficient training of healthcare workers (HCWs) in how to manage COVID-19 and the provision of personal protective equipment (PPE) to enable them to do so will be key tools in allowing oPt to mount a credible response to the crisis.

Methods: A cross-sectional study was conducted using a validated online questionnaire. Data collection occurred between March 30, 2020 and April 12, 2020. The primary outcomes was PPE provision and the secondary outcome was HCWs preparedness for the COVID-19 pandemic.

Results: Of 138 respondents, only 38 HCWs (27.5%) always had access to facemasks when needed and 15 (10.9%) for isolation gowns. The vast majority of HCWs did not find eye protection (n=128, 92.8%), N95 respirators (n=132, 95.7%), and face shields (n=127, 92%) always available. Compared to HCWs in West Bank, those in the Gaza Strip were significantly less likely to have access to alcohol sanitizers (p=0.026) and gloves (p <0.001). On average, governmental hospitals were significantly less likely to have all appropriate PPE measures than non-governmental institutions (p = 0.001). As for preparedness, only 16 (11.6%) surveyed felt confident in dealing with a potential COVID-19 case. With 57 (41.3%) having received any COVID-19 related training and 57 (41.3%) not having a local hospital protocol.

Conclusion: HCWs in oPt are underprepared and severely lacking adequate PPE provision. The lack of local protocols, and training has left HCWs confidence exceedingly low. The lack of PPE provision will exacerbate spread of COVID-19 and deepen the crisis, whilst putting HCWs at risk.

Introduction

With the ongoing coronavirus disease 2019 (COVID-19) pandemic, the humanitarian and healthcare crisis in low-to-middle income countries (LMICs) such as the occupied Palestinian territory (oPt) is expected to be amplified and this will further cripple the healthcare system. As of May 7, 2020, the World Health Organization (WHO) has recorded 547 confirmed cases of COVID-19 in the oPt; 527 in the West Bank and 20 in the Gaza Strip – with 4 fatalities.(1)

The United Nations Relief and Works Agency (UNRWA) has been unable to support Palestinians' COVID-19 response needs at their full capacity at the consequence of funding cut and legal restrictions that were in place prior to the pandemic.(2) Multiple COVID-19 testing sites serving Palestinians in East Jerusalem have been closed by the Israeli authorities.(3) The West Bank is particularly vulnerable due to checkpoint closures, halt of the transportation of patients to hospitals, and redistribution of clinical supplies. The Gaza Strip is one of the most densely populated places on earth with 2 million inhabitants, mostly refugees, live in 365 sq. km^2, allowing for an accelerated spread of disease should a COVID-19 outbreak manifest.(4) Other LMICs in the Middle East and Africa have also reported scarcity of personal protective equipment (PPE) for front line healthcare workers (HCWs).(5, 6)

We hypothesize that (HCWs) in the oPt are largely underprepared to address COVID-19 related needs of the Palestinian population in both the West Bank and Gaza Strip. Shortages of PPE pose a serious threat to COVID-19 containment in the oPt. It is also expected that HCWs in the oPt have likely received insufficient training on how to address spread and containment of COVID-19; institutions themselves may not have yet been equipped to draw up or implement preventative or management protocols. To the best of our knowledge, there have been no studies evaluating the preparedness of the HCWs to face COVID-19 pandemic. In this study, we aim to evaluate the availability of PPE and the level of preparedness among the HCWs in the oPt.

Methods

Study design, setting and data collection
We conducted a cross-sectional study using an online survey tool. Our survey (*Supplementary material*) was modified from two validated questionnaires; the first was utilized during the H1N1 influenza pandemic (7) and the second one was the Personnel, Infrastructure, Procedures, Equipment and Supplies (PIPES) surgical capacity assessment tool. (8) Our modified questionnaire consisted of 22 questions divided into 3 different sections (respondent and healthcare facility characteristics, availability of PPE and HCWs preparedness). Availability of PPE and HCWs preparedness were assessed on a 5-point Likert scale. The questionnaire was distributed to HCWs in the oPt through convenient sampling between March 30, 2020 and April 12, 2020. E-mail lists for participants in an educational link (OxPal) and social media (Facebook, Twitter, and LinkedIn) groups of HCWs in oPt were used to disseminate the questionnaire. Participants were required to sign in to limit the number of responses to one per respondent.

Study outcomes
The primary outcomes assessed were availability of PPE and HCWs preparedness in oPt in the era of COVID-19 pandemic. The secondary outcome was to assess the differences between Gaza Strip and West Bank, and between governmental and non-governmental in oPt in terms of availability of PPE and HCWs preparedness to face the COVID-19 pandemic.

Statistical analysis
Respondent characteristics were summarized using descriptive statistics. For continuous data, mean and standard deviation (SD) were used to report normally distributed data, while median and interquartile ranges (IQR) were used for non-normally distributed data. For categorical data, results were summarized as counts (n) and percentages (cumulative incidence). Univariate analysis (chi-squared and Fisher's Exact tests) was also used to compare participants' profession, geographical location, and responses to questions related to the availability of PPE and HCWs preparedness for the COVID-19 pandemic. Likert scale variables were converted from 5-point to binary variables for univariate analysis. For example, often available, sometimes available, rarely availably and never available were grouped together as 'not always available'. Strongly agree and moderately agree were grouped into 'agree' variable compared to 'neutral' and 'disagree' categories. Missing data were considered missing completely at random, therefore we performed complete case analysis. All statistical analyses were performed using IBM Corp. Released 2019. IBM SPSS Statistics for Windows, Version 26.0. Armonk, NY: IBM Corp.

Results

Of 140 completed surveys, two were excluded from the study as they were either working outside the oPt or in a non-medical profession.

HCWs and their healthcare facility characteristics
Of 138 HCWs included in the study, 97 respondents (70.3%) were from Gaza Strip and 41 (29.7%) were from the West Bank. The median (IQR) age was 28 (24-35) years with a range from 19 to 57 years old. 85 respondents (61.6 %) were males. Exactly half of respondents were medical doctors, with approximately 35 (25.4%) in nursing and the remaining quarter in physiotherapy, dentistry, or another health-related profession. 20 (14.5%) of the respondents worked in emergency medicine and 19 (13.8%) in surgery, 14 (10.1%) in primary care and 13 (9.4%) in internal medicine. With regards to place of work, 63 (45.7%) of the respondents worked in a tertiary hospital, 29 (21%) in a secondary facility and 43 (31%) in a primary healthcare center or clinic. One respondent worked in a COVID-19 isolation center. 98 (71%) worked in a governmental institution operated by the Ministry of Health, 26 (18.8%) worked in a private hospital and 13 (9.4%) in a non-governmental organization (NGO) or mission-based place of care (Table 1).

Availability of PPE and HCWs preparedness in terms of infection control training
Only 67 (48.6%) and 71 (51.4%) of HCWs surveyed indicated that they always had alcohol-based sanitizer and gloves available in their institutions, respectively. Only 38 (27.5%) of respondents indicated that regular face masks were always available when needed, and just over 15 (10.9%) of respondents reported that isolation gowns were always available in their

institutions. Over 128 (92.8%), 132 (95.7%), and 127 (92%) of respondents indicated that eye protection, N95 respirators, and face shields were not always available to them at their institutions, respectively. 57 (41.3%) of HCWs surveyed indicated that their hospital did not provide a local protocol for the management of COVID-19. Only 57 (41.3%) of respondents had received any COVID-19 related training courses by the time of survey administration. 16 (11.6%) of HCSw surveyed agreed with the statement of feeling confident or well-prepared to deal with a potential COVID-19 case (Table 2).

Univariate analysis comparing Gaza Strip and West Bank in terms of availability of PPE and HCWs preparedness in terms of infection control training
Compared to the West Bank, respondents from the Gaza Strip reported significantly greater lack of alcohol-based hand sanitizers (p=0.03) and gloves (p<0.001), but no meaningful differences were observed between regions on other PPE or infection control readiness (Table 3).

Univariate analysis comparing governmental and non-governmental hospitals in terms of availability of PPE and HCWs preparedness in terms of infection control training
On average, governmental hospitals run by the MOH were also reported by respondents to be significantly lacking in sanitizer, gloves, facemasks, eye protection, and face shields compared to non-governmental institutions (p<0.05) (Table 4).

Discussion

Our study demonstrates that the availability of PPE in both Gaza and the West Bank is insufficient to support the COVID-19 response needs of the oPt. Alcohol-based hand sanitizers, gloves, face masks, eye protection, isolation gowns, N95 respirators and face shields were reported to be inconsistently available, despite being internationally recommended as critical equipment needed for protecting health care workers from infection.(9) Governmental hospitals, as opposed to non-governmental settings, appear to be particularly lacking in equipment. Lessons from prior outbreaks have underlined the importance of PPE in infection control.(10) Recommendations from the WHO suggest the inadequate supply of infection prevention and control measures is vital to address immediately, with assistance from international partners if necessary.(11) The WHO specifically mentioned supplies needed to implement recommended protocols, such as PPE, being a key resource to all national authorities currently not producing sufficient volumes themselves. Suggestions for other methods of procurement, conservation and management of PPE have been extensively covered in the literature during the pandemic.(12) Many of these suggestions may not be viable in the geopolitical and economic context in which oPt operates. However, methods such as governmental coordination of all PPE supply, extending or creating new supply through 3D printing all provide viable means of blunting the dearth of PPE in oPt currently.(13, 14)

Our study showed that most HCWs surveyed did not receive adequate training on local protocol or measures to address COVID-19 spread from an institutional perspective. Comparing the preparedness of HCWs in oPt to those around the world, will be a vital element of the debrief from this pandemic and important in developing strategies to ensure the oPt have protocols in place for future public health crises. The lack of current data makes this comparison impossible,

currently. In previous pandemics, clinicians in other countries have been substantially more confident in their clinical ability to manage infected patients than what our results reflect; for example, Chinese ICU HCWs during the 2009 H1N1 pandemic were substantially more confident in their preparedness.(15) This may partly be due to a far greater provision of PPE amongst these workers, that permits greater clinical confidence.

Our study has some important strengths. To our knowledge, this study represents the first attempt to assess the availability of PPE in oPt and the preparedness of HCWs to face the COVID-19 pandemic. We provided a comprehensive evaluation of most PPE described in the literature and used clinically. Participants were well-represented across gender, geographic region, department/specialty, level of training, profession and type of health care facility.

Potential limitations of this study include small sample size, which may impact generalizability to the greater population of Palestinians. Another weakness of our study was the failure to elicit whether the lack of appropriate PPE was one of the driving factors in reducing HCW confidence in their preparedness. This would then imply attempts to target increasing PPE provision could both protect HCW and improve clinical confidence in managing COVID-19 patients. Potential selection bias arises due to sampling method. Most study participants were recruited from social media posts and emails to the networks of the researchers involved, which may limit some of the study's generalizability. However, other studies have demonstrated the viability of social media recruitment and snowball sampling to access difficult to reach populations.(16) Additionally, participants were asked to report on their individual experiences and thus may or may not be wholly representative of the institutions in which they are employed. The cross-sectional nature of this study is also by definition unable to take into account any changes in equipment or training preparedness over time and is only representative of the point-in-time data were collected. These limitations were acknowledged by the authors during study enrolment due to the need to publish findings within the international community in a time-sensitive manner and address the gap in literature regarding COVID-19's unique impact on the population in the oPt.

Low-to-middle income countries (LMICs) are particularly vulnerable to the spread of disease because they often grapple with detrimental resource and financial constraints that existed prior to the spread of pandemic. OPt and other LMICs often not only lack proper infrastructure and resources, but also have to navigate restrictions on movement, travel and transportation of essential supplies. In this global pandemic, procurement of adequate supply of PPE and the development of necessary protocols specific to the unique needs and challenges of the region are urgently needed.

Ethical approval: Ethical approval was not needed as the study did not involve patients or animals. However, participants consented to share their responses for research purposes.

Funding: None

Conflict of interest: We declare no conflict of interest.

References

1.   World Health Organization (WHO). Coronavirus disease (COVID-19) Situation Update 27, issued May 7, 2020. The occupied Palestinian territory 2020 [May 7, 2020]. Available from: https://who18.createsend.com/campaigns/reports/viewCampaign.aspx?d=j&c=99FA4938D049E3A8&ID=C3BC22C4771C92C22540EF23F30FEDED&temp=False&tx=0&source=Report&fbclid=IwAR2X6rkDzLsRcnMAjoF9sTL6M5585MrudomM8go2nWQzqRloE41MBbzFS8k.
2.   Monitor ME. Israel restricts UNRWA coronavirus efforts in Jerusalem refugee camps 2020 [April 29, 2020

]. Available from: https://www.middleeastmonitor.com/20200416-israel-restricts-unrwa-coronavirus-efforts-in-jerusalem-refugee-camps/.
3.   Israel Shuts Palestinian Coronavirus Testing Clinic in East Jerusalem 2020 [April 29, 2020]. Available from: https://www.haaretz.com/israel-news/.premium-israeli-police-raid-palestinian-coronavirus-testing-clinic-in-east-jerusalem-1.8767788.
4.   United Nations Country Team in the occupied Palestinian territory: Gaza ten years later 2020 [updated April 29, 2020. Available from: https://unsco.unmissions.org/sites/default/files/gaza_10_years_later_-_11_july_2017.pdf.
5.   Shortage of personal protective equipment endangering health workers worldwide 2020 [May 7, 2020]. Available from: https://www.who.int/news-room/detail/03-03-2020-shortage-of-personal-protective-equipment-endangering-health-workers-worldwide.
6.   Vijayakumar V. Personal Protection Prior to Preoperative Assessment—Little more an anaesthesiologist can do to prevent SARS-CoV-2 transmission and COVID-19 infection. Ain-Shams Journal of Anesthesiology. 2020;12(1):13.
7.   Daugherty EL, Perl Tm Fau - Needham DM, Needham Dm Fau - Rubinson L, Rubinson L Fau - Bilderback A, Bilderback A Fau - Rand CS, Rand CS. The use of personal protective equipment for control of influenza among critical care clinicians: A survey study. (1530-0293 (Electronic)).
8.   Personnel, Infrastructure, Procedures, Equipment and Supplies (PIPES) Surgical Assessment tool 2020 [April 29, 2020]. Available from: http://www.adamkushnermd.com/files/PIPES_tool_103111.pdf.
9.   Centers for Disease Control and Prevention (CDC). Strategies to Optimize the Supply of PPE and Equipment 2020 [May 8, 2020]. Available from: https://www.cdc.gov/coronavirus/2019-ncov/hcp/ppe-strategy/index.html.
10.   Fischer WA, 2nd, Weber D, Wohl DA. Personal Protective Equipment: Protecting Health Care Providers in an Ebola Outbreak. (1879-114X (Electronic)).
11.   2019 Novel Coronavirus (2019-nCoV): Strategic prepardness and response plan 2019 [May 3, 2020]. Available from: https://www.who.int/docs/default-source/coronaviruse/srp-04022020.pdf.
12.   Livingston E, Desai A, Berkwits M. Sourcing Personal Protective Equipment During the COVID-19 Pandemic. JAMA. 2020.
13.   Qarawi ATAN, Sze Jia, et al Awareness and Preparedness of Hospital Staff against Novel Coronavirus (COVID-2019): A Global Survey - Study Protocol. Elsevier BV. 2020.
14.   Canadian doctor who works in Gaza makes 3D-printed face shields for COVID-19 2020 [May 8, 2020]. Available from: https://www.cbc.ca/radio/asithappens/as-it-happens-thursday-edition-1.5502954/canadian-doctor-who-works-in-gaza-makes-3d-printed-face-shields-for-covid-19-1.5502964.

15.   Hu X, Zhang Z, Li N, Liu D, Zhang L, He W, et al. Self-reported use of personal protective equipment among Chinese critical care clinicians during 2009 H1N1 influenza pandemic. PloS one. 2012;7(9):e44723-e.
16.   Dusek G, Yurova Y, Ruppel C. Using Social Media and Targeted Snowball Sampling to Survey a Hard-to-reach Population: A Case Study. International Journal of Doctoral Studies. 2015;10.

Table 1. Healthcare workers and their healthcare facility characteristics

Parameter	Value
Age, median (IQR) - [range]	28 (24, 35) - [19-57]
Male, n (%)	85 (61.6)
Regional Location, n (%)	
Gaza Strip	97 (70.3)
West Bank (including East Jerusalem)	41 (29.7)
Profession, n (%)	
Medicine	69 (50)
Nursing	35 (25.4)
Physiotherapy	13 (9.4)
Dentistry	6 (4.3)
Other*	15 (10.9)
Level of training, n (%)	
Student	23 (16.7)
Trainee	23 (16.7)
Non-specialized	69 (50.0)
Specialized	23 (16.7)
Department / Specialty, n (%)	
Emergency medicine	20 (14.5)
Surgery (including sub-specialties)	19 (13.8)
Family medicine (primary care)	14 (10.1)
Internal medicine (including sub-specialties)	13 (9.4)
Pediatrics	10 (7.2)
Dentistry	8 (5.8)
Anesthesiology/critical care	7 (5.1)
Obstetrics/Gynecology	7 (5.1)
Radiology	6 (4.3)
Other*	13 (9.4)
Not applicable (e.g. students)	21 (15.2)
Type of healthcare facility (I), n (%)	
Primary healthcare center/ clinic	43 (31.2)
Secondary hospital	29 (21)
Tertiary (referral) hospital	63 (45.7)
Isolation Center	1 (0.7)
Not applicable	2 (1.4)
Type of healthcare facility (II), n (%)	
Governmental (Ministry of Health)	98 (71)
Private	26 (18.8)
Non-governmental (NGO) or Mission	13 (9.4)
Not applicable	1 (0.7)

*Includes: Optometry, medical secretary, medical laboratory, pharmacy, radiography.

Table 2. Availability of PPE and preparedness of healthcare workers in oPt for COVID-19 pandemic

Parameter	n (%)
Alcohol sanitizer	
Always available	67 (48.6)
Not always available	71 (51.4)
Gloves	
Always available	71 (51.4)
Not always available	67 (48.6)
Facemask	
Always available	38 (27.5)
Not always available	100 (72.5)
N95 respirator	
Always available	6 (4.3)
Not always available	132 (95.7)
Isolation gowns	
Always available	15 (10.9)
Not always available	123 (89.1)
Eye protection (goggles/glasses)	
Always available	10 (7.2)
Not always available	128 (92.8)
Face shields	
Always available	11 (8)
Not always available	127 (92)
All protective measures (above)	
Always available	15 (10.9)
Not always available	123 (89.1)
Receipt of any COVID-19 training course (e.g. infection control)	
Yes	57 (41.3)
No	81 (58.7)
Hospital has provided local protocol for the management of COVID-19	
Yes	81 (58.7)
No	57 (41.3)
I feel confident / well prepared dealing with a potential COVID-19 case	
Agree	16 (11.6)
Neutral / Disagree	122 (88.4)

Table 3. Univariate analysis comparing Gaza Strip and West Bank in terms of healthcare workers preparedness and personal protective equipment (PPE) availability

Parameter	Gaza Strip, n (%) N=97	West Bank, n (%) N=41	*p*-value
Alcohol sanitizer			
Always available	41 (42.3)	26 (63.4)	0.026
Not always available	56 (57.7)	15 (36.6)	
Gloves			
Always available	40 (41.2)	31 (75.6)	<0.001
Not always available	57 (58.8)	10 (24.4)	
Facemask			
Always available	24 (24.7)	14 (34.1)	0.299
Not always available	73 (75.3)	27 (65.9)	
N95 respirator			
Always available	5 (5.2)	1 (2.4)	0.669
Not always available	92 (94.8)	40 (97.6)	
Isolation gowns			
Always available	10 (10.3)	87 (212.2)	0.769
Not always available	5 (5.2)	36 (87.8)	
Eye protection			
Always available	8 (8.2)	2 (4.9)	0.723
Not always available	89 (91.8)	39 (95.1)	
Face shields			
Always available	10 (10.3)	1 (2.4)	0.174
Not always available	87 (89.7)	40 (97.6)	
All PPE any other measures (above)			
Always available	12 (12.4)	3 (7.3)	0.552
Not always available	85 (87.6)	38 (92.7)	
Receipt of any COVID-19 training course (e.g. infection control)			
Yes	44 (45.4)	13 (31.7)	0.185
No	53 (54.6)	28 (68.3)	
Hospital has provided local protocol for the management of COVID-19			
Yes	58 (59.8)	23 (56.1)	0.709
No	39 (40.2)	18 (43.9)	
I feel confident / well prepared dealing with a potential COVID-19 case			
Agree	13 (13.4)	3 (7.3)	<0.001
Neutral/disagree	84 (86.6)	38 (92.7)	

Table 4. Univariate analysis comparing Governmental and non-governmental hospitals in oPt in terms of healthcare workers preparedness and personal protective equipment (PPE) availability

Parameter	Governmental hospitals, n (%) N=98	Non-governmental hospital, n (%) N=39	*p*-value
Alcohol sanitizer			
Always available	35 (35.7)	32 (82.1)	<0.001
Not always available	63 (64.3)	7 (17.9)	
Gloves			
Always available	38 (38.8)	32 (82.1)	<0.001
Not always available	60 (61.2)	7 (17.9)	
Facemask			
Always available	15 (15.3)	23 (59.0)	<0.001
Not always available	83 (84.7)	16 (41.0)	
N95 respirator			
Always available	3 (3.1)	3 (7.7)	0.352
Not always available	95 (96.9)	36 (92.3)	
Isolation gowns			
Always available	8 (8.2)	7 (17.9)	0.129
Not always available	90 (91.8)	32 (82.1)	
Eye protection			
Always available	2 (2.0)	8 (20.5)	0.001
Not always available	96 (98.0)	31 (79.5)	
Face shields			
Always available	4 (4.1)	7 (17.9)	0.012
Not always available	94 (95.9)	32 (82.1)	
All PPE any other measures (above)			
Always available	5 (5.1)	10 (25.6)	0.001
Not always available	93 (94.9)	29 (74.4)	
Receipt of any COVID-19 training course (e.g. infection control)			
Yes	37 (37.8)	19 (48.7)	0.254
No	61 (62.2)	20 (51.3)	
Hospital has provided local protocol for the management of COVID-19			
Yes	57 (58.2)	23 (59.0)	1
No	41 (41.8)	16 (41.0)	
I feel confident / well prepared dealing with a potential COVID-19 case			
Agree	10 (10.2)	6 (15.4)	0.556
Neutral / Disagree	88 (89.8)	33 (84.6)	

Supplementary Table. Healthcare workers preparedness and availability of personal protective equipment in occupied Palestinian territory to face the COVID-19 pandemic

Section I: participant demographics and healthcare facility characteristics		
Age (years):		
Gender:	a.	Male
	b.	Female
Profession:	a.	Medicine
	b.	Nursing
	c.	Other (please specify)
Level of training:	a.	Clinical medical student
	b.	Nursing student
	c.	Intern doctor
	d.	Non-specialized doctor
	e.	Resident
	f.	Registered nurse/staff nurse
	g.	Nurse practitioner
	h.	Specialist doctor
	i.	Consultant doctor
	j.	Other (please specify)
Department/specialty:	a.	Internal medicine (including sub-specialties)
	b.	Surgery (including sub-specialties)
	c.	Emergency medicine
	d.	Anesthesiology/critical care
	e.	Family medicine (primary care)
	f.	Pediatrics
	g.	Obstetrics/Gynecology
	h.	Radiology
	i.	Psychiatry
	j.	Non-applicable
	k.	Other (please specify)
Place of work (city/governorate):	a.	Jerusalem
	b.	Hebron
	c.	Bethlehem
	d.	Ramallah and al-Bireh
	e.	Jenin
	f.	Jericho
	g.	Qalqilya
	h.	Tulkarm

	i. Tubas
	j. Salfit
	k. Nablus
	l. North Gaza
	m. Gaza
	n. Deir al-Balah (includes all cities in middle of Gaza Strip)
	o. Khan Younis
	p. Rafah

Healthcare facility name (in English):	
Type of healthcare facility - I	a. Primary healthcare center
	b. Secondary (district) hospital
	c. Tertiary (referral) hospital
	d. Other
Type of healthcare facility - II	a. Governmental (Ministry of Health)
	b. Private
	c. Non-governmental (NGO) or Mission
	d. Other

Hospital beds (total number)

Section II: Availability of personal protective equipment (PPE) and other protective measures

Does your healthcare facility provide/have the following?

Alcohol based hand sanitizer?	Always available
	Often available
	Sometimes available
	Rarely available
	Never available
Gloves?	Always available
	Often available
	Sometimes available
	Rarely available
	Never available
Regular facemasks?	Always available
	Often available
	Sometimes available
	Rarely available
	Never available
N95 respirators?	Always available
	Often available
	Sometimes available

	Rarely available
	Never available
Isolation gowns?	Always available
	Often available
	Sometimes available
	Rarely available
	Never available
Eye protection (goggles/glasses)?	Always available
	Often available
	Sometimes available
	Rarely available
	Never available
Face shields?	Always available
	Often available
	Sometimes available
	Rarely available
	Never available
All of the above protective measures are readily available in your healthcare facility?	Always available
	Often available
	Sometimes available
	Rarely available
	Never available
	Never available
Section III: Healthcare workers preparedness for COVID-19 pandemic	
Have you received any Covid-19 related training course? (e.g. infection control)	Yes
	No
Does your hospital provide a local protocol for management of COVID-19?	Yes
	No
I feel confident/well-prepared dealing with a potential COVID-19 case	Strongly agree
	Moderately agree
	Neutral
	Moderately disagree
	Strongly disagree

Any comments?