1
2
3
4   **Time is of the essence: impact of delays on effectiveness of contact tracing**

5   **for COVID-19, a modelling study**

7   Prof. Mirjam E. Kretzschmar[1]*, PhD, Ganna Rozhnova[1,2], PhD, Martin Bootsma[1,3], PhD,

8   Michiel van Boven[1], PhD, Prof. Janneke van de Wijgert[1,4], PhD, Prof. Marc Bonten[1,5], MD

9
10
[1] Julius Center for Health Sciences and Primary Care, University Medical Center Utrecht,

Utrecht University, Utrecht, the Netherlands

[2] BioISI – Biosystems & Integrative Sciences Institute, Faculdade de Ciências, Universidade

de Lisboa, Lisboa, Portugal

[3] Mathematical Institute, Utrecht University, Utrecht, The Netherlands

[4] Institute of Infection and Global Health, University of Liverpool, Liverpool, UK

[5] Department of Medical Microbiology, University Medical Center Utrecht, Utrecht

University, the Netherlands

19
20
21
22 *corresponding author
23
24 Mirjam E. Kretzschmar

25 Julius Center for Health Sciences and Primary Care

26 University Medical Center Utrecht, Utrecht

27 Email: m.e.e.kretzschmar@umcutrecht.nl

28 Phone: +31 88 75 687 61

Summary

Background

With confirmed cases of COVID-19 declining in many countries, lockdown measures are

gradually being lifted. However, even if most social distancing measures are continued,

other public health measures will be needed to control the epidemic. Contact tracing via

conventional methods or mobile app technology is central to control strategies during de-

escalation of social distancing. We aimed to identify key factors for a contact tracing

strategy (CTS) to be successful.

Methods

We evaluated the impact of timeliness and completeness in various steps of a CTS using a

stochastic mathematical model with explicit time delays between time of infection and

symptom onset, and between symptom onset, diagnosis by testing, and isolation (testing

delay). The model also includes tracing of close contacts (e.g. household members) and

casual contacts, followed by testing regardless of symptoms and isolation if positive, with

different delays (tracing delay) and coverages (tracing coverage). We computed effective

reproduction numbers of a CTS (R_{cts}) for a population with social distancing measures and

various scenarios for isolation of index cases and tracing and quarantine of its contacts.

Findings

For the best-case scenario (testing and tracing delays of 0 days and tracing coverage of

80%), and assuming that around 40% of transmission occur before symptom onset, the

model predicts that the effective reproduction number of 1.2 (with social distancing only)

will be reduced to 0.8 by adding contact tracing. A testing delay of 2 days requires tracing

delay to be at most 1 day, or tracing coverage to be at least 80% to keep R_{cts} below 1. With a

testing/isolation delay of 3 days, even the most efficient CTS cannot reach R_{cts} values below

1. The effect of minimizing tracing delay (e.g., with app-based technology) declines with

decreasing coverage of app use, but app-based tracing alone remains more effective than

conventional tracing alone even with 20% coverage. The proportion of transmissions per

index case that can be prevented depends on testing and tracing delays, and ranges from up

to 80% in the best-case scenario (testing and tracing delays of 0 days) to 42% with a 3-day

testing delay and 18% with a 5-day testing delay.

Interpretation

In our model, minimizing testing delay had the largest impact on reducing onward

transmissions. Optimizing testing and tracing coverage and minimizing tracing delays, for

instance with app-based technology, further enhanced CTS effectiveness, with a potential to

prevent up to 80% of all transmissions. Access to testing should therefore be optimized, and

mobile app technology may reduce delays in the CTS process and optimize contact tracing

coverage.

Funding: ZonMw projects 91216062 and 10430022010001, Fundação para a Ciência e a Tecnologia project
reference 131_596787873, and EU H2020 grant RECOVER (H2020-101003589).

Research in context

Evidence before this study

We searched PubMed, bioRxiv, and medRxiv for articles published in English from January 1, 2020, to June 20, 2020, with the following keywords: ("2019-nCoV" OR "novel coronavirus" OR "COVID-19" OR "SARS-CoV-2") AND "contact tracing" AND "model*". Population-level modelling studies of severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) have suggested that isolation and tracing alone might not be sufficient to control outbreaks and additional measures might be required. However, few studies have focused on the effects of lifting individual measures once the first wave of the epidemic has been controlled. Lifting measures must be accompanied by effective contact tracing strategies (CTS) in order to keep the effective reproduction number below 1. A detailed analysis, with special emphasis on the effects of time delays in testing of index patients and tracing of contacts, has not been done.

Added value of this study

We performed a systematic analysis of the various steps required in the process of testing and diagnosing an index case as well as tracing and isolating possible secondary cases of the index case. We then used a stochastic transmission model which makes a distinction between close contacts (e.g. household members) and casual contacts to assess which steps and (possible) delays are crucial in determining the effectiveness of CTS. We evaluated how delays and the level of contact tracing coverage influence the effective reproduction number, and how fast CTS needs to be to keep the reproduction number below 1. We also analyzed what proportion of onward transmission can be prevented for

short delays and high contact tracing coverage. Assuming that around 40% of transmission

occurs before symptom onset, we found that keeping the time between symptom onset and

testing and isolation of an index case short (<3 days) is imperative for a successful CTS. This

implies that the process leading from symptom onset to receiving a positive test should be

minimized by providing sufficient and easily accessible testing facilities. In addition, reducing

contact-tracing delays also helps to keep the reproduction number below 1.

Implications of all the available evidence

Our analyses highlight that CTS will only contribute to containment of COVID-19 if it can be

organised in a way that time delays in the process from symptom onset to isolation of the

index case and his/her contacts are very short. The process of conventional contact tracing

should be reviewed and streamlined, while mobile app technology may offer a tool for

gaining speed in the process. Reducing delay in testing subjects for SARS-CoV-2 should be a

key objective of CTS.

Introduction

As the first wave of the SARS-CoV-2 has reached its peak of cases in many countries, societies are preparing so-called exit-strategies from the COVID-19 lockdown, while still successfully controlling transmission. Contact tracing, in combination with quarantine and potentially testing of the contacts, is considered a key component in a phase when lockdown measures are gradually lifted[1-8]. Contact tracing is an intervention, where an index case with confirmed infection is asked to provide information about contact persons, who were at risk of acquiring infection from the index case within a given time period before the positive test result. These contact persons are then traced and informed about their risk, quarantined, and tested if eligible for testing according to national testing guidelines. This requires upscaling of conventional contact tracing capacity. The potential of mobile apps to support contact tracing is widely discussed and such technology has been used in several Asian countries. Although these countries have successfully reduced case numbers, no causal relationship between use of app technology and epidemic control has yet been shown[9-14]. Many uncertainties remain on the optimal process of contact tracing with conventional methods and/or mobile applications, on the timing of testing for current or past infection, and on the required coverage of contact tracing needed.

Modelling studies have demonstrated how mobile applications can increase effectiveness of contact tracing, compared to conventional approaches for contact tracing, but effectiveness depends on what proportion of the population will use the app consistently for a sufficiently long period of time[9]. Modelling studies have predicted that contact tracing alone cannot control an outbreak if tracing coverage is too low[2,15]. What tracing coverage is needed depends on how much transmission occurs before symptom onset, and on the details of the tracing process.

In previous work, we have investigated the impact of timeliness and completeness of case

reporting for the effectiveness of surveillance and interventions[16,17], and we quantified the

timeliness of contact tracing of infected passengers during an airline flight for the 2009

pandemic influenza[18]. In all of these studies, the timing of various steps in the monitoring and

intervention chain emerged as one of the key factors for effectiveness of a public health

response. Usually, there are identifiable delays in the response chain that may be critical to

the overall effectiveness of a strategy.

Here we analyze in detail the process chain of identifying index cases by symptom-reporting

followed by testing, and subsequent contact tracing, with the aim to inform policy makers on

the relative importance of key steps in the process. We use a mathematical model that reflects

the various steps and delays in the contact tracing process to quantify the impact of delays on

the effective reproduction number and the fraction of onward transmission prevented per

diagnosed index case[5,19].

Methods

Time delays in contact tracing

Our starting point is an assumed effective reproduction number (R_e) for COVID-19 of around

1, describing a situation with "social distancing but measures lifted to some extent". We then

quantify the relative contribution of the individual components of a contact tracing strategy

(CTS) required to bring and maintain the effective reproduction number with CTS (R_{CTS}) to a

value below 1. For simplicity we do not include transmission in healthcare settings, as in

healthcare settings like nursing homes, which can be viewed as closed populations, other

interventions are more appropriate.

We break down the process of contact tracing in two steps (Figure 1; Supplementary

Information Table S1).

• An index case acquires infection (at time T_0), then after a short latent period becomes

infectious (at time T_1), and finally symptomatic (at time T_2), which is here defined as

"being eligible for testing". Subsequently a proportion of all symptomatic subjects gets

tested and diagnosed (at time T_3). The time between T_2 and T_3 is called the "testing

delay" ($D_1 = T_3 - T_2$), and may vary between 0 and 7 days, and in this period individuals

might self-quarantine. We refer to the proportion of all symptomatically infected cases

that are tested as testing coverage and vary it from 20% to 80% in increments of 20%.

After being diagnosed, we assume index cases are isolated with no further transmission.

• The second step is tracing contacts of the index case, which occurs at time T_4. A fraction

of those contacts will be found and tested. We assumed that all contacts, regardless of

symptoms, are offered testing, and that those testing negative do not spread. Those who

are found infected will be isolated, with effectiveness ranging from 0% to 100%. We

assume that contacts in isolation do not spread. The time between T_3 and T_4 is the

"tracing delay" ($D_2 = T_4 - T_3$), which may range from 0 (for instance with app

technology) to 3 days (with conventional approaches). In this step, tracing coverage is

defined as the proportion of contacts detected, which either depends on the capacity of

conventional approaches (ranging from 20% to 80% in increments of 20%) or on the

fraction of the population using suitable app technology for screening (ranging from 20%

to 100% in increments of 20%). We did not consider hybrid approaches of combined

conventional and app-based CTS.

The best-case scenario we consider is that persons eligible for testing are immediately tested

(coverage 80%) with a very fast test result (test-delay 0 days) and immediate isolation when

testing positive, followed by immediate tracing (trace delay 0 days) of all contacts, that

immediately adhere to isolation measures (coverage 80%). We consider more realistic

scenarios where testing and tracing are suboptimal, e.g. a conventional CTS, and we vary

these parameters separately in a sensitivity analysis (see Supplementary Information).

Impact on effectiveness on population level

To analyse the impact of these time delays on the effectiveness of contact tracing we use a

model introduced by Kretzschmar et al[19], which was adapted for SARS-CoV-2[5]. The

stochastic model describes an epidemic as a branching process with progression through

latent infection and infectious period in time steps of 1 day. Infectivity and probability of

symptom onset per day of the infectious period, and numbers of contacts per day were fitted

to distributions taken from published data. [20-24] We distinguish between close contacts (e.g.

household contacts, but also other high-risk contacts) and casual contacts, which differ in the

risk of acquiring infection from the index case. Also, the time required for tracing and

isolating infected contacts and the coverage of tracing may differ between these types of

contacts and between different CTS (i.e., conventional contact tracing versus mobile app

supported contact tracing). We assume that isolation is perfect, i.e. that isolated persons do

not transmit any longer, and that all traced infected contacts are isolated, regardless of

whether they develop symptoms or not. The model allows for explicit computation of the

basic reproduction number R_0, the effective reproduction number under social-distancing

interventions R_e, and the effective reproduction number with CTS (R_{cts}). Reproduction

numbers were calculated as expectations, and distributions of individual reproduction

numbers were simulated. The model was coded in Mathematica 12.1. For details, see the

Supplementary Information.

Parameter settings

We assumed that without social distancing individuals have on average 4 close contacts per

day and around 9 casual contacts per day, with stochastic variability. The distributions were

fitted to data from the Polymod study for the Netherlands[23]. Transmission probability per

contact for close contacts was taken to be 4 times higher than for casual contacts.

Symptomatic and asymptomatic cases were assumed to be equally infectious. Overall, the

transmission probability was calibrated to a basic reproduction number of $R_0 = 2.5$. For

social distancing, we assumed that close contacts were reduced by 40% and casual contacts

by 70%. The resulting effective reproduction number was $R_e = 1.2$.

Uncertainty of model outcomes

We considered uncertainty due to stochastic variability, and uncertainty due to possible

variation in parameter estimates. We dealt with stochastic variability by computing

individual reproduction numbers for 1000 individuals for all scenarios, and plotted their

distributions as boxplots. Parameter uncertainty was explored by performing simulations

using hypercube sampling for transmission probabilities and probabilities of symptom onset

per day of the infectious period (Supplementary Information).

Scenarios modelled

We analyzed the impact of various testing and tracing delays and tracing coverage on the

effective reproduction number R_{cts} while keeping the testing coverage at 80%. For

comparison, we also considered the strategy where symptomatic individuals get tested and

isolated, without subsequent tracing (R_{iso}). We varied the testing delay D_1 between 0 and 7

days, the tracing delay D_2 between 0 and 3 days, and tracing coverages between 0% and

100% in increments of 20%. For conventional contact tracing, we assumed that coverage is

higher for close contacts than for casual contacts.

We then compared the effectiveness of conventional CTS alone with a scenario in which

mobile app technology is used for alerting subjects to be tested and for tracing contacts.

Differences between these strategies were taken as follows. The testing delay (D_1) is reduced

with app technology. We assumed a conventional CTS setting in which symptomatic

individuals need to decide to seek health care to get tested, and we assumed that with app

technology symptomatic persons get alerted and can be tested without health care

interference. For conventional CTS we assumed suboptimal coverage in identifying contacts

from the week before diagnosis due to recall bias, especially for casual contacts. For CTS

with mobile app technology we assume 60% and 80% tracing coverage of the contacts of

subjects using app technology. We show also results for 100% coverage, although

realistically more than 80% is not feasible, because not all contacts may be correctly

identified and compliance with isolation of those tested positive may not be perfect. We

assume that tracing goes back for 7 days before the positive test result. The exact parameter

values for this comparison are shown in Table 1.

Next, we quantified the impact of coverage of testing and app use on the effectiveness of

CTS. We varied the percentage of app users in the population between 20% and 100% in

increments of 20%. We first considered the situation that testing is provided for 80% of

persons with symptoms independent of app use, and app use only influences the fraction of

contacts that are traced. Alternatively, assumed that only app users are tested (i.e. testing

coverage varies between 20% and 100% in increments of 20%), and coverage of tracing also

depends on fraction of app use. In all cases, a contact person could only be traced if both the

index case and the contact person were app users, i.e. the probability of a contact being traced

is given by the square of the proportion of app users.

Finally, we quantified the fraction of transmissions of an index person that can be prevented,

and the contribution to the fraction prevented from isolation and from tracing contacts with

decreasing delays. The number of onward transmissions of an index case is by definition

described by the effective reproduction number of the realized scenario. Therefore, the

difference of reproduction numbers between two intervention scenarios under the condition

that an index case is diagnosed, describes the fraction of onward transmissions prevented. For

contact persons, this is the fraction of the total infectivity that lies after the time of isolation,

i.e. the part of infectiousness that is prevented by contact tracing. In other words, a contact

person who is detected and isolated before the start of their infectious period is a fully

prevented transmission, while a contact person who is only traced and identified after 70% of

their infectivity has passed, is counted as 0.3 of a prevented onward transmission.

Role of the funding source

Results

In the best-case scenario, if 80% of infectious persons that develop symptoms are tested and

isolated within 1 day after symptom onset the effective reproduction number R_e is expected

to decline from 1.2 to R_{iso} = 1.0, without contact tracing (Figure 2). Contact tracing may

further decrease the reproduction number to R_{cts}=0.8 in the best case scenario. In the best

case scenario – a testing delay of 0 days, a tracing delay of 0 days, and a tracing coverage of

80%, the additional reduction of R_{cts} predicted by the model is 33%. Yet, with a testing delay

of 2 days, tracing delay should be at most 1 day, or tracing coverage should be at least 80% to

keep R_{cts} below 1. In these scenarios, the reduction of R_{cts} compared to the best-case scenario

is estimated at 17% (Supplementary Information Figure S4). With a testing delay of more

than 3 days, even perfect contact tracing cannot bring R_{cts} values below 1.

We assumed that conventional CTS has longer tracing delay and lower tracing coverage than

CTS based on app technology which results in marked differences in R_{cts} for the whole range

of testing delay (Figure 2A). With conventional CTS, R_{cts} would remain above 1, if the

testing delay exceeds 0 days, whereas contact tracing based on app technology could still

keep R_{cts} below 1, as long as testing and tracing coverage would be at least 80%, or if testing

delay is 1 day and tracing coverage 60%. If the testing delay reaches 5 days or more, app

technology adds little effectiveness to conventional CTS or just isolating symptomatic cases.

The reductions of R_e (based on social distancing) achieved by isolation of symptomatic cases

only, conventional CTS, and mobile app-based CTS are shown in figure 3A. For isolation

only and for conventional CTS we assumed a delay of 4 days between symptom onset and

isolation of the index case. The relative reductions are independent of the level of R_e, similar

reductions are seen for R_0, i.e. in a situation without social distancing (Supplementary

Information). Conventional CTS, even if applied for all infected subjects with symptoms, is

27% less effective than mobile app-based CTS alone, due to longer tracing delays and lower

tracing coverage. Figure 3B shows the distributions of individual reproduction numbers for

the testing delays assumed in Table 1, i.e. 4 days for isolation and conventional CT, and 0

days for app-based CTS. Only for app-based CTS the means of the individual reproduction

numbers are below 1.

The effectiveness of app-based technology declines with lower fractions of persons using it

(Figure 4). Yet, app-based tracing on its own remains more effective than conventional

tracing alone, even with 20% coverage, due to its inherent speed. Even with low coverage

there is a reduction of R_e, due to fast tracing of a small part of the population. Depending on

R_e, such an approach might be sufficient to reduce R_{cts} to levels below 1. This can be seen in

the distributions of the individual reproduction numbers (Figure 4B and 4D), where in 4B the

means of the distributions are below 1 for 40% and more app use, while in 4D this is the case

for 60% and above.

323
324
325   In Table 2, we quantified proportions of transmissions per index case that can be prevented

326   depending on testing delay, stratified by of isolation of index cases and tracing delays. In the

327   best-case scenario (testing and tracing delay being 0 days) around 80% of transmissions can

328   be prevented if tracing coverage is 80%.

329
330
331
Discussion

Using a mathematical model that describes the different steps of CTS for COVID-19 we have

quantified the relevance of delays and coverage proportions for controlling SARS-CoV-2

transmission. We conclude that reducing the testing delay, i.e. shortening the time between

symptom onset and positive test result (assuming immediate isolation), is the most important

factor for improving CTS effectiveness. Reducing the tracing delay, i.e. shortening the time

of contact tracing (assuming immediate testing and isolation if positive), may further enhance

CTS effectiveness. Yet this additional effect rapidly declines with increasing testing delay.

The effectiveness of app-based CTS declines with lower app use coverage, but it remains

more effective than conventional contact tracing even with lower coverage, due to its

inherent speed. CTS therefore has the potential to control virus transmission, and to enable

alleviation of other control measures, but only if all delays are maximally reduced. It should

be noted that we simulated two CTS systems (conventional CTS with testing and tracing

delays and app-based CTS without delays) and ignored hybrid approaches. At present, most

European countries are using conventional CTS, but are attempting to reduce delays (for

example, by improving testing and tracing capacity and by removing testing barriers), and are

piloting or planning the addition of app-based contact-tracing. Such hybrid CTS systems

would fall somewhere between the fully conventional and app-based scenarios described in

this paper.

Several factors can reduce CTS effectiveness, such as large proportions of cases who remain

asymptomatic or are otherwise not diagnosed, and large proportions of contacts who cannot

be traced. App-based technology could increase the proportion of tracable contacts, because

it does not rely on recall of names and contact details, but this would require the participation

of a substantial proportion of the population. App use acceptance may be hampered by

privacy concerns and other ethical considerations, which limit its acceptance.   Also, app use

needs to continue over a long time period, requiring sustained adherence by app users. Low

participation does not render CTS useless, however, because it could help to locally

extinguish clusters before they grow larger. In addition, every measure that lowers the

effective reproduction number, even if it is already below 1, will lower the cumulative case

number and speed up the time until elimination of the virus from the population.

The strength of our approach is that it explicitly takes many details of the contact tracing

process into account, such that the key factors can be identified. A limitation of our approach

is that it does not take population age-structure into account, which may influence the

proportion of asymptomatic cases and mobile app use coverage. Also, the willingness of a

case to self-isolate depends on age and social norms, may depend on socio-economic status,

and is affected by perceived benefit of isolation in relation to perceived risk of the infection

to others[25]. We also excluded other heterogeneities while assuming homogeneous mixing[26,27],

and assumed homogeneously distributed use of app technology for different coverage levels.

Clustering of non-users may have consequences for overall effectiveness of CTS, similar to

clustering of non-vaccinated persons. Furthermore, we ignored that a sizeable portion of

transmissions may be acquired nosocomially when population prevalence is still low. [28] The

model also ignores that some contacts of the index case may have self-quarantined with

symptoms before they are traced by CTS, which lowers the benefits of CTS.

Our results add to results from other modelling studies, which showed that CTS can be an

effective intervention if tracing coverage is high and if the process is fast[2,15]. A determining

factor is the proportion of transmissions occurring before symptom onset, which determines

the urgency of tracing and isolating contacts as fast as possible. Our study showed in detail

what the role is of each step in the CTS process in making it successful. Our model differs in

that it makes a distinction between close and casual contacts, and that we consider scenarios

for conventional CTS and mobile app-based CTS characterized by specific delays and

coverages.

Our finding of the crucial importance of the first step of CTS, establishing a diagnosis in

cases with symptoms, has important consequences. It requires an infrastructure for testing,

that allows persons with symptoms to be tested, preferably, within one day of symptom onset.

Studies have demonstrated that viral shedding in the respiratory tract is highest at the start of

symptoms[29,30], so early testing will also increase the sensitivity of this approach. To further

enhance effectiveness, as many infectious persons as possible need to be tested regardless of

symptoms, which requires a low threshold for testing. As the clinical symptoms of COVID-

19 are mostly mild and heterogeneous, many persons should be eligible for testing, resulting

in a large proportion of negative test results. Future work should determine the optimal

balance between the proportion of test-negatives and the effectiveness of CTS.

Our findings also provide strong support to optimize contact tracing. In the Netherlands, CTS

was based on establishing contact between an index case and a public health officer, followed

by an interview after which contacts are traced. This procedure is labor intensive, time

consuming, prone to recall bias, incomplete (anonymous contacts cannot be traced), and

usually takes several days. Optimizing this process by improving testing and tracing capacity,

removing testing barriers, and by adding app-based and/or other digital technologies to

minimize tracing delay are needed to establish optimal control of transmission. These

improvements are currently being implemented or considered. Overall, our findings suggest

that optimized CTS, with short delays and high coverage for testing and tracing could

substantially reduce the reproduction number, which would allow alleviation of more

stringent control measures.

Data availability

The Mathematica code used for the analysis are available on Github under

https://github.com/mirjamkretzschmar/ContacttracingModel

Contributors

415   MEK and MB conceived the study. MK designed and programmed the model, and produced

416   output. MvB, MCJB, and GR helped with the analysis and literature research. JvdW

417   contributed to data interpretation and writing. All authors interpreted the results, contributed

418   to writing the manuscript, and approved the final version for submission.

Declaration of interests

421   We declare no competing interests.

Acknowledgements

424   MEK received funding from ZonMw projects number 91216062 and number 10430022010001. GR received
425   funding from Fundação para a Ciência e a Tecnologia project reference 131_596787873. MB received funding
426   from EU H2020 grant RECOVER (H2020-101003589).
427
428   We thank Patricia Bruijning-Verhagen and Hans Heesterbeek for useful discussions. This
429   work forms part of RECOVER (*Rapid European COVID-19 Emergency Response research*).
430   RECOVER is funded by the European Union's Horizon 2020 research and innovation
431   programme under grant agreement No 101003589.
432

References

1.   Fraser C, Riley S, Anderson RM, Ferguson NM. Factors that make an infectious

disease outbreak controllable. Proc Natl Acad Sci U S A 2004; 101(16): 6146-51.

2.   Hellewell J, Abbott S, Gimma A, et al. Feasibility of controlling COVID-19 outbreaks

by isolation of cases and contacts. Lancet Glob Health 2020.

3.   Keeling MJ, Hollingsworth TD, Read JM. The Efficacy of Contact Tracing for the

Containment of the 2019 Novel Coronavirus (COVID-19). medRxiv 2020:

2020.02.14.20023036.

4.   Klinkenberg D, Fraser C, Heesterbeek H. The effectiveness of contact tracing in

emerging epidemics. PLoS One 2006; 1: e12.

5.   Kretzschmar ME, Rozhnova G, van Boven M. Isolation and contact tracing can tip

the scale to containment of COVID-19 in populations with social distancing. medRxiv 2020:

2020.03.10.20033738.

6.   Muller J, Kretzschmar M, Dietz K. Contact tracing in stochastic and deterministic

epidemic models. Math Biosci 2000; 164(1): 39-64.

7.   Tang B, Wang X, Li Q, et al. Estimation of the Transmission Risk of the 2019-nCoV

and Its Implication for Public Health Interventions. J Clin Med 2020; 9(2).

8.   Salathe M, Althaus CL, Neher R, et al. COVID-19 epidemic in Switzerland: on the

importance of testing, contact tracing and isolation. Swiss Med Wkly 2020; 150: w20225.

9.   Ferretti L, Wymant C, Kendall M, et al. Quantifying SARS-CoV-2 transmission

suggests epidemic control with digital contact tracing. Science 2020.

10.   Kamel Boulos MN, Geraghty EM. Geographical tracking and mapping of

coronavirus disease COVID-19/severe acute respiratory syndrome coronavirus 2 (SARS-CoV-

2) epidemic and associated events around the world: how 21st century GIS technologies are

supporting the global fight against outbreaks and epidemics. Int J Health Geogr 2020; 19(1):

8.

11.   Parker MJ, Fraser C, Abeler-Dorner L, Bonsall D. Ethics of instantaneous contract

tracing using mobile phone apps in the control of the COVID-19 pandemic. J Med Ethics

2020.

12.   Nguyen THD, Vu DC. Summary of the COVID-19 outbreak in Vietnam - Lessons and

suggestions. Travel Med Infect Dis 2020: 101651.

13.   Ruan L, Wen M, Zeng Q, et al. New measures for COVID-19 response: a lesson from

the Wenzhou experience. Clin Infect Dis 2020.

14.   Show evidence that apps for COVID-19 contact-tracing are secure and effective.

Nature 2020; 580(7805): 563.

15.   Peak CM, Kahn R, Grad YH, et al. Individual quarantine versus active monitoring of

contacts for the mitigation of COVID-19: a modelling study. Lancet Infect Dis 2020.

16.   Bonacic Marinovic A, Swaan C, van Steenbergen J, Kretzschmar M. Quantifying

reporting timeliness to improve outbreak control. Emerg Infect Dis 2015; 21(2): 209-16.

17.   Bonacic Marinovic AA, Koopmans M, Dittrich S, et al. Speed versus coverage trade

off in targeted interventions during an outbreak. Epidemics 2014; 8: 28-40.

18.   Swaan CM, Appels R, Kretzschmar ME, van Steenbergen JE. Timeliness of contact

tracing among flight passengers for influenza A/H1N1 2009. BMC Infect Dis 2011; 11: 355.

19.   Kretzschmar M, van den Hof S, Wallinga J, van Wijngaarden J. Ring vaccination and

smallpox control. Emerg Infect Dis 2004; 10(5): 832-41.

20.   Bi Q, Wu Y, Mei S, et al. Epidemiology and Transmission of COVID-19 in Shenzhen

China: Analysis of 391 cases and 1,286 of their close contacts. medRxiv 2020:

2020.03.03.20028423.

21.   Chan JF, Yuan S, Kok KH, et al. A familial cluster of pneumonia associated with the

2019 novel coronavirus indicating person-to-person transmission: a study of a family cluster.

Lancet 2020; 395(10223): 514-23.

22.   Chen N, Zhou M, Dong X, et al. Epidemiological and clinical characteristics of 99

cases of 2019 novel coronavirus pneumonia in Wuhan, China: a descriptive study. Lancet

2020; 395(10223): 507-13.

23.   Mossong J, Hens N, Jit M, et al. Social contacts and mixing patterns relevant to the

spread of infectious diseases. PLoS Med 2008; 5(3): e74.

24.   Backer JA, Klinkenberg D, Wallinga J. Incubation period of 2019 novel coronavirus

(2019-nCoV) infections among travellers from Wuhan, China, 20-28 January 2020. Euro

Surveill 2020; 25(5).

25.   Webster RK, Brooks SK, Smith LE, Woodland L, Wessely S, Rubin GJ. How to

improve adherence with quarantine: rapid review of the evidence. Public Health 2020; 182:

163-9.

26.   Dowd JB, Andriano L, Brazel DM, et al. Demographic science aids in understanding

the spread and fatality rates of COVID-19. Proc Natl Acad Sci U S A 2020; 117(18): 9696-8.

27.   Prem K, Cook AR, Jit M. Projecting social contact matrices in 152 countries using

contact surveys and demographic data. PLoS Comput Biol 2017; 13(9): e1005697.

28.   Wang D, Hu B, Hu C, et al. Clinical Characteristics of 138 Hospitalized Patients With

2019 Novel Coronavirus-Infected Pneumonia in Wuhan, China. JAMA 2020.

29.   He X, Lau EHY, Wu P, et al. Temporal dynamics in viral shedding and

transmissibility of COVID-19. Nat Med 2020; 26(5): 672-5.

30.   Kucirka LM, Lauer SA, Laeyendecker O, Boon D, Lessler J. Variation in False-

Negative Rate of Reverse Transcriptase Polymerase Chain Reaction-Based SARS-CoV-2 Tests

by Time Since Exposure. Ann Intern Med 2020.

511 **Table 1: Comparison Conventional CT and Mobile app CT**

	Conventional CT	**Mobile app CT**
Testing coverage	80%	20%, 40%, 60%, 80%, 100%
Testing delay (D_1), assuming immediate isolation when testing positive	4 days	0 day
Time to trace close contacts (D_2)	3 days	0 day
Time to trace other contacts, assuming testing and isolation of those who test positive	3 days	0 day
Tracing coverage close contacts	80%	20%, 40%, 60%, 80%, 100%
Tracing coverage casual contacts	50%	20%, 40%, 60%, 80%, 100%
Time traced back	7 days	7 days

515 **Table 2.** Percentage of onward transmissions prevented per diagnosed index case for various
516 interventions: only isolation of the index case (left column) or isolation of the index case with tracing
517 and isolation of 80% of infected contacts (columns 2-5).
518
519

Delay D_1 (days)	Isolation only	Contact tracing Delay D_2 (days)			
		3	2	1	0
0	50.4	62.4	67.8	73.9	79.9
1	35.7	47.3	53.4	60.7	68.5
2	23.4	33.0	38.9	46.5	55.4
3	14.2	21.0	26.0	32.9	41.8
4	7.8	11.9	15.7	21.4	29.1
5	3.8	5.9	8.4	12.5	18.4
6	1.6	2.4	3.8	6.4	10.4

521 **Figure 1: Schematic of the contact tracing process and its time delays.**
522
523
524
525

527
528

Figure 2: Comparison of a conventional and mobile app CTS. For parameter values, see table 1.
We assumed that testing coverage is 80% for the conventional CTS and 60%, 80%, and 100% for the
mobile app CTS. For mobile app CTS it is assumed that the tracing coverage equals the testing rate,
i.e. it is 60%, 80%, and 100%, respectively. Expected reproduction numbers are shown as a function
of testing delay D_1.

Figure 3: The reduction of the effective reproduction number for various CTS. (A) The
reproduction number with CTS, R_{cts}, is shown as a percentage of the reproduction number where only
social distancing is implemented (R_e). For the isolation scenario and conventional tracing scenario we
assumed that there is a delay of 4 days between symptom onset and isolation of the index case. For
the mobile app CTS, testing delay was assumed to be 0 days. Testing coverage was assumed to be
80% in the isolation and conventional CT scenarios; app use prevalence was assumed to be 60%,
80%, and 100% in the mobile app CTS. (B) Distributions of individual reproduction numbers for
1000 individuals and the same scenarios as in (A).

552
553 (A)   **(B)**
554

555
556

559 **Figure 4: The impact of mobile app use on R_{cts} for varying levels of app use.** In 4A and 4B, we
560 assume that there is also testing of those who do not use the mobile app, so app use only is used for
561 tracing contacts. In 4C and 4D, only app users, who develop symptoms, are tested. Panels A and C
562 show percentage reductions of R_e achieved by the CTS; panels B and D show the impact of various
563 CTS on distributions of individuals reproduction numbers.
564
565

566 (A)
567

568
569
570
5 5 4
5
575
576 (C)
577
578
579