1
2
3
4 **Time is of the essence: impact of delays on effectiveness of contact tracing**

5 **for COVID-19**

7 Mirjam Kretzschmar[1]*, Ganna Rozhnova[1,2], Martin Bootsma[1,3], Michiel van Boven[1], Janneke

8 van de Wijgert[1,4], Marc Bonten[1,5]

9
10
[1] Julius Center for Health Sciences and Primary Care, University Medical Center Utrecht,

Utrecht University, Utrecht, the Netherlands

[2] BioISI – Biosystems & Integrative Sciences Institute, Faculdade de Ciências, Universidade

de Lisboa, Lisboa, Portugal

[3] Mathematical Institute, Utrecht University, Utrecht, The Netherlands

[4] Institute of Infection and Global Health, University of Liverpool, Liverpool, UK

[5] Department of Medical Microbiology, University Medical Center Utrecht, Utrecht

University, the Netherlands

19
20
21
22 *corresponding author
23
24 Mirjam E. Kretzschmar

25 Julius Center for Health Sciences and Primary Care

26 University Medical Center Utrecht, Utrecht

27 Email: m.e.e.kretzschmar@umcutrecht.nl

28 Phone: +31 88 75 687 61

29
30
Summary

Background

With confirmed cases of COVID-19 declining in many countries, lockdown measures are

gradually being lifted. However, even if most social distancing measures are continued,

other public health measures will be needed to control the epidemic. Contact tracing either

via conventional methods or via mobile app technology is central to control strategies

during de-escalation of social distancing. It is therefore essential to identify key factors for a

contact tracing strategy (CTS) to be successful.

Methods

We evaluated the impact of timeliness and completeness in various steps of a CTS using a

stochastic mathematical model with explicit time delays between time of infection,

symptom onset, diagnosis by testing, and isolation. The model also includes tracing of close

contacts (e.g. household members) and casual contacts with different delays and coverages.

We computed effective reproduction numbers of a CTS (R_{cts}) for a population with social

distancing measures and various scenarios for isolation of index cases and tracing and

quarantine of its contacts.

Findings

In the best-case scenario (testing and tracing delays of 0 days and tracing coverage of 100%)

the effective reproduction number will be reduced with 50% from 1.2 (with social distancing

only) to 0.6 (R_{cts}) by contact tracing. A testing delay of 3 days requires tracing delay or

coverage to be at most 1 day or at least 80% to keep R_{cts} below 1, with the R_{cts} reduction

being 15% and 17%, respectively. With a testing delay of 4 days, even the most efficient CTS

cannot reach R_{cts} values below 1. The effect of minimizing tracing delay (e.g., with app-

based technology) declines with declining coverage of app use, but app-based tracing

remains more effective than conventional contact tracing even with 20% coverage. The

proportion of transmissions per index case that can be prevented depending on testing and

tracing delay and isolation of index cases ranges from above 80% in the best-case scenario

(testing and tracing delays of 0 days) to 40% and 17% with testing delays of 3 and 5 days,

respectively.

Interpretation

Minimizing testing delay is of key importance for the effectiveness of CTS. Optimizing testing

and tracing coverage and minimizing tracing delays, for instance with app-based technology

further enhances effectiveness of CTS, with a potential to prevent up to 80% of all

transmissions. The process of conventional contact tracing should be reviewed and

streamlined, while mobile app technology may offer a tool for gaining speed in the process.

73   **Funding:** ZonMw project number 91216062

Research in context

Evidence before this study

As of 8 May 2020, the novel coronavirus (SARS-CoV-2) has spread globally and has caused more than 263,000 confirmed deaths of COVID-19 worldwide. In the absence of effective medicines and vaccines, many countries have implemented strict measures of social distancing, thereby reducing transmission and bringing the epidemic under control. For lifting these measures, adequate tools are needed to deal with possible newly arising transmission clusters. Strategies including isolation of confirmed and suspected cases, and identification and quarantining of their contacts are considered a key part of the response during de-escalation of social distancing. As a substantial portion of transmission may occur before the onset of symptoms and before cases can be isolated, it is unclear how successful contact tracing strategies (CTS) can be in reducing onward transmission.

Added value of this study

We performed a systematic analysis of the various steps required in the process of testing and diagnosing an index case as well as tracing and isolation possible secondary cases of the index case. We then used a stochastic transmission model which makes a distinction between close contacts (e.g. household members) and casual contacts to assess which steps and (possible) delays are crucial in determining the effectiveness of CTS. We evaluated how delays and the level of contact tracing coverage influence the effective reproduction number, and how fast CTS needs to be to keep the reproduction number below 1. We also analyzed what proportion of onward transmission can be prevented for short delays and high contact tracing coverage. Assuming that around 40% of transmission

occurs before symptom onset, we found that keeping the time between symptom onset and

isolation of an index case short (<3 days) is imperative for a successful CTS. This implies that

the process leading from symptom onset to receiving a positive test should be minimized by

providing sufficient and easily accessible testing facilities. In addition, reducing contact-

tracing delays also helps to keep the reproduction number below 1.

Implications of all the available evidence

Our analyses highlight that CTS will only contribute to containment of COVID-19 if it can be

organised in a way that time delays in the process from symptom onset to isolation of the

index case and his/her contacts are very short. The process of conventional contact tracing

should be reviewed and streamlined, while mobile app technology may offer a tool for

gaining speed in the process.

Introduction

As the first wave of the SARS-CoV-2 has reached its peak of cases in many countries, societies are preparing so-called exit-strategies from the COVID-19 lockdown, while still successfully controlling transmission. Contact tracing, in combination with testing and quarantine or isolation of the contacts, is considered a key component in a phase when lockdown measures are gradually lifted[1-8] This requires upscaling of conventional contact tracing capacity. The potential of mobile apps to support contact tracing is widely discussed and such technology has been used in several Asian countries that have successfully reduced case numbers[9-14]. Yet, many uncertainties remain on the optimal process of contact tracing with conventional methods and/or mobile applications, on the timing of testing for current or past infection, and on the required coverage of contact tracing needed. As a result, predicting the effects of contact tracing, and predicting whether and at which level of virus circulation contact tracing can sufficiently control remaining transmission is difficult.

Modelling studies have demonstrated how mobile applications can increase effectiveness of contact tracing, compared to conventional approaches for contact tracing, but effectiveness depends on what proportion of the population will use the app consistently and for a sufficiently long period of time[9].

In previous work, we have investigated the impact of timeliness and completeness of case reporting for the effectiveness of surveillance and interventions[15-17], and we quantified the timeliness of contact tracing of infected passengers during an airline flight for the 2009 pandemic influenza[18]. In all of these studies, the timing of various steps in the monitoring

and intervention chain emerged as one of the key factors for effectiveness of a public health

response. Usually, there are identifiable delays in the response chain that may be critical to

the overall effectiveness of a strategy.

Here we analyze in detail the process chain of identifying index cases by symptom-reporting

followed by testing, and subsequent contact tracing, with the aim to inform policy makers

on the relative importance of key steps in the process. We use a mathematical model that

reflects the various steps and delays in the test and contact tracing process to quantify the

impact of delays on the effective reproduction number and the fraction of onward

transmission prevented per diagnosed index case[5,19].

147 Time delays in contact tracing

Our starting point is an assumed effective reproduction number (R_e) for COVID-19 of around

1, describing a situation with "social distancing but measures lifted to some extent". We

then quantify the relative contribution of the individual components of a contact trace

strategy (CTS) required to bring and maintain the effective reproduction number with CTS

(R_{CTS}) to a value below 1. For simplicity we do not include transmission in healthcare

settings.

We break down the process of contact tracing in two different steps (Table 1 and Figure 1).

• An index case acquires infection (at time T_0), then after a short latent period becomes

infectious (at time T_1), and finally symptomatic (at time T_2), which is here defined as

"being eligible for testing". Subsequently a proportion of all symptomatic subjects gets

tested and diagnosed (at time T_3). The time between T_2 and T_3 is called the "testing

delay" ($D_1 = T_3 - T_2$), and may vary between 0 and 5 days, and in this period individuals

might self-quarantine. We refer to the proportion of all symptomatically infected

subjects that is tested as testing coverage and vary it from 20% to 100%. After being

diagnosed, we assume index cases are quarantined with no further transmission.

• The second step is tracing contacts of the index, which occurs at time T_4. A fraction of

those contacts will be quarantined, with effectiveness ranging from 0%-100%. For

simplicity we assume that contacts in quarantine do not spread. The time between T_3

and T_4 is the "tracing delay" ($D_2 = T_4 - T_3$), which may range from 0 (for instance with app

technology) to 4 days (with conventional approaches). In this step, tracing coverage is

defined as the proportion of contacts detected, which either depends on the capacity of

conventional approaches (ranging from 40% to 80%) or on the fraction of the population

using suitable app technology for screening (ranging from 40% to 80%).

Figure 1: Schematic of the contact tracing process and its time delays.

173
174

T0   time of infection of index case
T1   onset of infectiousness
T2   symptom onset
T3   time of positive diagnosis
T4   time of tracing and quarantining contacts

D1 = T3 – T2 delay to diagnosis
D2 = T4 – T3 delay in tracing contacts

Coverage of contact tracing:
percentage of contacts found
and isolated

175
176
177

Table 1: Time delays in the test and contact tracing process (see also Figure 1).

Time	Event	Comments	Model implementation
T_0	the time of infection of the index case	Not observed	Start of the latent period, which lasts 1-3 days.
T_1	Time the index case becomes infectious	Proportion of pre-symptomatic transmission may range from 0% to 40% of all transmissions	After 1-3 days after infection, the infectious stage starts, which lasts 10 days with variable infectiousness. About 40% of transmission takes place in the first 2 days of infectiousness[25].
T_2	Time that the index (case) becomes	T_0 until T_2 reflects the time window in which	The incubation period in the model is taken in agreement with published literature[21].

		symptomatic, and eligible for testing	prevention is not possible with CTS	
T_3		Time that index (case) is tested positive	T_2 until T_3 is the testing delay, which may range from 0-5 days. The proportion being tested varies from 0-100%. During this period we expect subjects to self-quarantine, with effectiveness ranging from 0%-100%	After a testing delay D_1 after symptom onset, an individual receives a positive test result and gets isolated. If an individual self isolates immediately, $D_1=0$. After isolation, no transmission takes place.
T_4		Time that contacts of index case are traced and quarantined.	T_3 until T_4 is the tracing delay, which may range from 0 (for instance with app technology) to 4 days (with current GGD approach). Here we can also vary the proportion with short post-test-delay (those with apps) and not.	After a tracing delay D_2, contacts of the index case are traced and isolated. D_2 and the tracing coverage (proportion of contacts found and isolated) may differ between close and casual contacts. If household contact self-isolate immediately with the index case, it means that $D_2=0$ and coverage 100% for close contacts. For simplicity we assume that contacts in quarantine do not spread.

The best-case scenario is that all eligible for testing are immediately tested (coverage 100%)

with a very fast test result (test-delay 1 day), followed by immediate tracing (trace delay 0

days) of all contacts (coverage 100%), that immediately adhere to quarantine measures.

More realistic scenarios include testing and tracing delays, with suboptimal testing and

tracing coverages and suboptimal adherence to quarantining and testing.

Impact on effectiveness on population level

To analyse the impact of these time delays on the effectiveness of contact tracing we use a

model first described in Kretzschmar et al[19], which was recently adapted for SARS-CoV-2[5].

The stochastic model describes an epidemic in its early phase as a branching process.

Starting from a small set of initially infected individuals, the model calculates the numbers

of latently infected persons, infectious persons, and persons that are diagnosed and isolated

in time steps of one day. Latent infection, infectivity during the infectious period, and daily

contact rates are quantified using distributions taken from published data.[20-24] We

distinguish between close contacts (e.g. household contacts, but also other high-risk

contacts) and casual contacts, which differ in the risk of acquiring infection from the index

case. Also, the time required for tracing and quarantining contacts and the coverage of

tracing may differ between these types of contacts and between different CTS (i.e.,

conventional contact tracing versus mobile app supported contact tracing). Intervention

effectiveness is determined by the daily probability of an index case being diagnosed by

testing during the infectious period, and depends on various delays in the process of tracing

household and non-household contacts, respectively, and on the proportions of contacts

that can be traced and isolated (see Figure 1). We assume that isolation is perfect, i.e. that

isolated persons do not transmit any longer. The model is described by a set of difference

equations, and allows for explicit computation of the basic reproduction number R_0, the

effective reproduction number under social-distancing interventions R_e and the effective

reproduction number with CTS (R_{cts}). The model was coded in Mathematica 12.1.

Parameter settings

We assumed that without social distancing individuals have on average 4 close contacts per

day and around 9 casual contacts per day, with certain stochastic variability. The

distributions were fitted to data from the Polymod study[23]. Transmission probability per

contact for close contacts was taken to be 4 times higher than for casual contacts.

Symptomatic and asymptomatic cases were assumed to be equally infectious. Overall, the

transmission probability was calibrated to a basic reproduction number of R_0 = 2.5. For the

social distancing, we assumed that close contacts were reduced by 40% and casual contacts

by 70%. The resulting effective reproduction number was R_e = 1.2. Without further

interventions, the doubling time of the epidemic would be around 19 days.

Scenarios modelled

We analyzed the impact of various testing and tracing delays and tracing coverage on the

effective reproduction number R_{cts} while keeping the testing coverage at 100%. For

comparison, we also considered the strategy where symptomatic individuals get tested and

isolated, without subsequent tracing (R_{iso}). We varied the testing delay D_1 between 0 and 7

days, the tracing delay D_2 between 0 and 3 days, and tracing coverages between 0% and

100%. Tracing delays and coverages were allowed to differ between close contacts and

casual contacts.

We then compared the effectiveness of conventional CTS with a scenario that reflects

mobile app technology for alerting subjects to be tested and for tracing contacts.

Differences between these strategies were taken as follows. The testing delay (D_1) is

reduced with app technology. With conventional CTS symptomatic individuals need to

decide to seek health care to get tested, and we assume that with app technology

symptomatic subjects get alerted and can be tested without health care interference, for

instance in specific test facilities for app users. For conventional CTS we assume suboptimal

coverage in identifying contacts from the week before diagnosis by testing due to recall

bias, especially for casual contacts. For CTS with mobile app technology we assume 100%

tracing coverage of the proportion of subjects using app technology. For simplicity we

assume 100% compliance with quarantining. We assume that tracing goes back for 7 days

before the positive test result. The exact parameter values for this comparison are shown in

Table 2.

Next, we quantified the impact of coverage of testing and app use on the effectiveness of

CTS. We varied the percentage of app users in the population between 20% and 80%. We

first considered the situation that testing is provided for 100% of persons with symptoms

independent of app use, and app use only influences the fraction of contacts that are

traced. Alternatively, we considered the situation that only app users with symptoms are

tested (i.e. testing coverage varies between 20% and 80%) and coverage of tracing also

depends on fraction of app use, i.e. varies as the testing coverage.

Finally, we quantified the fraction of transmissions of an index person that can be

prevented, and the contribution to the fraction prevented from isolation and from tracing

contacts with decreasing delays. The number of onward transmissions of an index case is by

definition described by the effective reproduction number of the realized scenario.

Therefore, the difference of reproduction numbers between two intervention scenarios

under the condition that an index case is diagnosed, will describe the fraction of onward

transmissions prevented. For contact persons, this is the fraction of the total infectivity that

lies after the time of isolation, i.e. the part of infectiousness that is prevented by contact

tracing. In other words, a contact person who is detected and isolated before the start of his

infectious period is a fully prevented transmission, while a contact person who is only traced

and identified after 70% of his infectivity has passed, is counted as 0.3 of a prevented

onward transmission.

Table 2: Comparison Conventional CT and Mobile app CT

	Conventional CT	Mobile app CT
Testing coverage	100%	100%
Time to (self)-isolation (D_1)	4 days	0 day
Time to trace close contacts (D_2)	3 days	0 day
Time to trace other contacts	3 days	0 day
Tracing coverage close contacts	80%	100%
Tracing coverage casual contacts	50%	100%
Time traced back	7 days	7 days

Results

In the best-case scenario, if all infectious persons that develop symptoms are tested and

isolated within 1 day after symptom onset the effective reproduction number R_e will decline

from 1.2 to R_{iso} = 0.97, without contact tracing (Figure 2). Contact tracing will further

decrease the reproduction number to R_{cts}=0.6 in the best case. In the optimal scenario – a

testing delay of 0 days and a tracing delay of 0 days and a tracing coverage of 100%, the

additional reduction of R_{cts} is 50%. Yet, with a diagnosis delay of 3 days, tracing delay or

tracing coverage should be at most 1 day or at least 80% to keep R_{cts} below 1. In these

scenarios the reduction of R_{cts} compared to the best-case scenario is 15% and 17%. With a

testing delay of 4 days, even the most efficient contact tracing cannot reach R_{cts} values

below 1.

Figure 2: Impact of contact tracing on the effective reproduction number depending on various delays and tracing coverages. In these analyses, 100% of those who develop symptoms get tested. For comparison the reproduction number R_{iso} with only isolation of index cases without contact tracing is plotted (green). (A) Influence of varying tracing delay D_1 on the x-axis. The curves plotted in blue show varying tracing delays D_2; (B) Here the tracing coverage is varied in the curves plotted in blue, while there is assumed to be no delay in tracing the contacts.

We assumed that conventional CTS has longer tracing delay and lower tracing coverage than

CTS based on app technology which results in marked differences in R_{cts} for the whole range

of testing delay (Figure 3). With conventional CTS, R_{cts} would remain above 1, if the testing

delay exceeds 2 days, whereas contact tracing based on app technology could still keep R_{cts}

below 1, as long as testing and tracing coverage would be at least 80%. If the testing delay

reaches 5 days or more, app technology adds little effectiveness to conventional CTS or just

isolating symptomatic cases.

306 **Figure 3: Comparison of a conventional and mobile app CTS.** For parameter values, see table 2. We
307 assumed that ascertainment is 100% for the conventional CTS and 100% and 80% for the mobile app
308 CTS.

The reductions of R_e (based on social distancing) achieved by isolation only, conventional

CTS, and mobile app-based CTS is shown in figure 4. For isolation only and for conventional

CTS we assumed a delay of 4 days between symptom onset and isolation of the index case.

The relative reductions are independent of the level of R_e, as there is a linear relationship

between the various reproduction numbers. Conventional CTS, even if applied for all

infected subjects with symptoms is 45% less effective than mobile app-based CTS, due to

longer tracing delays and lower tracing coverage.

The effectiveness of app-based technology declines with lower fractions of persons using it

(Figure 5). Yet, it remains more effective than conventional contact tracing even with 20%

coverage, due to its inherent speed. In Figure 5a we assume that all symptomatic persons

get tested, and then vary coverage of app use. In Figure 5b, we assume that only app users

who develop symptoms get diagnosed, and that only app users get traced and isolated.

Even with low coverage there is a reduction of R_e, due to fast tracing of a small part of te

population. Depending on R_e, such an approach might be sufficient to reduce R_{cts} to levels

below 1.

In Figure 6, we quantified proportions of transmissions per index case that can be prevented

depending on testing delay, as well as the contributions of isolation of index cases and

tracing of contacts. In the best-case scenario (testing and tracing delay being 0 days) more

than 80% of transmissions can be prevented if coverage of infected persons being tested is

100%.

339
340
341

Figure 4: The reduction of the effective reproduction number for various CTS. The reproduction number with CTS, R_{cts}, is shown as a percentage of the reproduction number where only social distancing is implemented (R_e). For the isolation scenario and conventional tracing scenario we assumed that there is a delay of 4 days between symptom onset and isolation of the index case.

Figure 5: The impact of mobile app use on R_{cts} for varying levels of app use. In (A), we assume that there is also testing of those who do not use the mobile app, so app use only is used for tracing contacts. In (B), only app users, who develop symptoms, are tested.

(A)

364 (B)

376 **Figure 6: The fraction of onward transmissions prevented by isolation of the index case and**
377 **his/her infected contacts.** The fraction prevented by contact tracing increases with decreasing
378 tracing delay.

379
380

Discussion and conclusions

Using a mathematical model that describes the different steps of the CTS for COVID-19 we have quantified the relevance of delays and coverage proportions for controlling transmission of SARS-CoV-2. Based on these analyses we conclude that reducing the testing delay, i.e. shortening the time between symptom onset and test positivity, is the most crucial step. Reducing the tracing delay, i.e. shortening the time of contact tracing, may further enhance the effectiveness of CTS. Yet this additional effect rapidly declines with increasing testing delay. Naturally, the effectiveness of CTS increases when proportions of index subjects detected and contacts traced increase as well. CTS has huge potential to control virus transmission, and thus to alleviate other control measures, but only if all delays are maximally reduced.

There are several obvious factors that can reduce the effectiveness of CTS, such as a large proportion of infectious subjects that remain asymptomatic or are otherwise not ascertained and a large proportion of contacts that cannot be traced. The latter implies that the potential benefits of using app-based technology for contact tracing requires participation of a substantial proportion of the population. Also, app use needs to continue over a long time period, so required continued adherence of app users. Low proportions of participation do not render CTS useless, however, because it could help to locally extinguish clusters before they grow larger. Also, for this purpose, the timeliness and completeness of CTS in local populations should be high to make it successful.

The strength of the approach is that it explicitly takes many details of the contact tracing

process into account, such that the key factors can be identified. A limitation of our

approach is that it does not take population age-structure into account, which may

influence the proportion of asymptomatic cases and the mobile app use coverage. Also, the

willingness of an index case or contact person to self-isolate may be different in different

age groups. We have also assumed homogeneous mixing of the population, and

homogeneous distributed use of app technology for the different coverage levels. Yet,

clustering of non-users may have consequences for overall effectiveness of CTS, similar to

clustering of non-vaccinated subjects. This is an important aspect to be addressed in

subsequent work. The model also ignores that some contacts of the index case may have

symptoms before they are traced by CTS. As these contacts may already self-isolate, this

lowers the benefits of contact tracing.

Our finding of the crucial importance of the first step of CTS, establishing a diagnosis in

subjects with symptoms, has important consequences. It requires an infrastructure for

testing, that allows subjects with symptoms to be tested, preferably, within one day of

symptom onset. Studies have demonstrated that viral shedding in the respiratory tract is

highest at the start of symptoms[25], so early testing will also increase the sensitivity of this

approach. To further enhance effectiveness, as many infectious subjects need to be tested,

which requires a low threshold for testing. As the clinical symptoms of COVID-19 are mostly

mild and heterogeneous, many subjects should be eligible for testing, resulting in a large

proportion of subjects with negative test results. Future work should determine the optimal

balance between the proportion of test-negatives and the effectiveness of CTS. In our

country, testing of ambulatory subjects is coordinated by the public health services and

general practitioners. That infrastructure may introduce a considerable delay in testing. To

optimize the effectiveness of CTS a different infrastructure with direct access of

symptomatic subjects to testing facilities should be considered. Finally, laboratories should

be prepared to deliver high-throughput rapid testing.

Our findings also provide strong support to optimize contact tracing. In our country this is

now based on establishing a contact between public health officers and index patients,

followed by an interview after which contacts are traced. This procedure is labor intensive,

time consuming, prone to recall bias and usually takes several days. Optimizing this process

with app technology, or any other method achieving the same goal of minimizing tracing

delay, will be needed to establish optimal control of transmission. An important advantage

of app-based technology is the possibility of performing multiple step tracing, as not only

the first-line contacts can be traced, but also their (second-line) contacts and so on.

Naturally, the number of contacts than rapidly increases, which increases the number of

both correctly and unnecessarily quarantined subjects. Further work will focus on finding an

optimal balance for this aspect. In fact, our findings suggest that optimized CTS, with short

delays and high coverage for testing and tracing could reduce the reproduction number by

50%, which would allow alleviation of most of the currently implemented control measures.

Acknowledgements

We thank Patricia Bruijning-Verhagen and Hans Heesterbeek for useful discussions.

References

1.   Fraser C, Riley S, Anderson RM, Ferguson NM. Factors that make an infectious

disease outbreak controllable. *Proc Natl Acad Sci U S A* 2004; **101**(16): 6146-51.

2.   Hellewell J, Abbott S, Gimma A, et al. Feasibility of controlling COVID-19 outbreaks

by isolation of cases and contacts. *Lancet Glob Health* 2020.

3.   Keeling MJ, Hollingsworth TD, Read JM. The Efficacy of Contact Tracing for the

Containment of the 2019 Novel Coronavirus (COVID-19). *medRxiv* 2020:

2020.02.14.20023036.

4.   Klinkenberg D, Fraser C, Heesterbeek H. The effectiveness of contact tracing in

emerging epidemics. *PLoS One* 2006; **1**: e12.

5.   Kretzschmar ME, Rozhnova G, van Boven M. Isolation and contact tracing can tip the

scale to containment of COVID-19 in populations with social distancing. *medRxiv* 2020:

2020.03.10.20033738.

6.   Muller J, Kretzschmar M, Dietz K. Contact tracing in stochastic and deterministic

epidemic models. *Math Biosci* 2000; **164**(1): 39-64.

7.   Tang B, Wang X, Li Q, et al. Estimation of the Transmission Risk of the 2019-nCoV and

Its Implication for Public Health Interventions. *J Clin Med* 2020; **9**(2).

8.   Salathe M, Althaus CL, Neher R, et al. COVID-19 epidemic in Switzerland: on the

importance of testing, contact tracing and isolation. *Swiss Med Wkly* 2020; **150**: w20225.

9.   Ferretti L, Wymant C, Kendall M, et al. Quantifying SARS-CoV-2 transmission suggests

epidemic control with digital contact tracing. *Science* 2020.

10.   Kamel Boulos MN, Geraghty EM. Geographical tracking and mapping of coronavirus

disease COVID-19/severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) epidemic

and associated events around the world: how 21st century GIS technologies are supporting

the global fight against outbreaks and epidemics. *Int J Health Geogr* 2020; **19**(1): 8.

11.   Parker MJ, Fraser C, Abeler-Dorner L, Bonsall D. Ethics of instantaneous contract

tracing using mobile phone apps in the control of the COVID-19 pandemic. *J Med Ethics*

2020.

12.   Nguyen THD, Vu DC. Summary of the COVID-19 outbreak in Vietnam - Lessons and

suggestions. *Travel Med Infect Dis* 2020: 101651.

13.   Ruan L, Wen M, Zeng Q, et al. New measures for COVID-19 response: a lesson from

the Wenzhou experience. *Clin Infect Dis* 2020.

14.   Show evidence that apps for COVID-19 contact-tracing are secure and effective.

Nature 2020; **580**(7805): 563.

15.   Bonacic Marinovic A, Swaan C, van Steenbergen J, Kretzschmar M. Quantifying

reporting timeliness to improve outbreak control. *Emerg Infect Dis* 2015; **21**(2): 209-16.

16.   Bonacic Marinovic AA, Koopmans M, Dittrich S, et al. Speed versus coverage trade

off in targeted interventions during an outbreak. *Epidemics* 2014; **8**: 28-40.

17.   Reijn E, Swaan CM, Kretzschmar ME, van Steenbergen JE. Analysis of timeliness of

infectious disease reporting in the Netherlands. *BMC Public Health* 2011; **11**: 409.

18.   Swaan CM, Appels R, Kretzschmar ME, van Steenbergen JE. Timeliness of contact

tracing among flight passengers for influenza A/H1N1 2009. *BMC Infect Dis* 2011; **11**: 355.

19.   Kretzschmar M, van den Hof S, Wallinga J, van Wijngaarden J. Ring vaccination and

smallpox control. *Emerg Infect Dis* 2004; **10**(5): 832-41.

20.   Bi Q, Wu Y, Mei S, et al. Epidemiology and Transmission of COVID-19 in Shenzhen

China: Analysis of 391 cases and 1,286 of their close contacts. *medRxiv* 2020:

2020.03.03.20028423.

21.   Chan JF, Yuan S, Kok KH, et al. A familial cluster of pneumonia associated with the

2019 novel coronavirus indicating person-to-person transmission: a study of a family cluster.

Lancet 2020; **395**(10223): 514-23.

22.   Chen N, Zhou M, Dong X, et al. Epidemiological and clinical characteristics of 99

cases of 2019 novel coronavirus pneumonia in Wuhan, China: a descriptive study. *Lancet*

2020; **395**(10223): 507-13.

23.   Mossong J, Hens N, Jit M, et al. Social contacts and mixing patterns relevant to the

spread of infectious diseases. *PLoS Med* 2008; **5**(3): e74.

24.   Backer JA, Klinkenberg D, Wallinga J. Incubation period of 2019 novel coronavirus

(2019-nCoV) infections among travellers from Wuhan, China, 20-28 January 2020. *Euro*

Surveill 2020; **25**(5).

25.   He X, Lau EHY, Wu P, Deng X, Wang J, Hao X, Lau YC, Wong JY, Guan Y, Tan X, Mo X,

Chen Y, Liao B, Chen W, Hu F, Zhang Q, Zhong M, Wu Y, Zhao L, Zhang F, Cowling BJ, Li F,

Leung GM. Temporal dynamics in viral shedding and transmissibility of COVID-19. *Nat Med*.

2020 Apr 15. [Epub ahead of print]