The confounded crude case-fatality rates for COVID-19 hide more than they reveal - a comparison of age-specific and age-adjusted rates between six countries

Manfred S Green MBChB,PhD[1], Victoria Peer MSc[1], Dorit Nitzan MD, MPH[2]

Affiliation

1. School of Public Health, University of Haifa, Israel
2. World Health Organization, European Region, Copenhagen, Denmark

Correspondence:

School of Public Health

University of Haifa,

Abba Khoushy 199,

Haifa 3498838,

Israel.

E-mail: manfred.s.green@gmail.com

Abstract

Background

The reported crude case-fatality rates (CFRs) vary widely between countries. The serious limitations of using crude COVID-19 CFRs for comparisons between countries have been addressed in the literature but are still either overlooked or misunderstood, both in the scientific literature and in the media. In this paper we examined the extent to which age distribution of the cases is responsible for the differences in CFRs between countries.

Methods

Data on COVID-19 were extracted from the reports of individual countries. Overall and age-specific CFRs were available for six countries. The CFRs by country were adjusted for age using the direct method, using the combined age-specific number of cases of all six countries as the standard population.

Findings

The age distribution of the cases varied widely between countries. The crude CFRs varied between 1.6% and 11%. The differences in the age-specific CFRs were much smaller and the age-adjusted rates were much closer than the crude rates. The ratio of the crude CFR for the country with the highest to that with the lowest, was reduced substantially from 7.4 to 2.3 for the age-adjusted rates.

Conclusions

The age structure of the cases dramatically impacts on the differences in the crude CFRs between countries. Adjusting for age substantially reduces this variation. Other factors such as the differences in the definition of the denominators, the definition of a case and the standard of healthcare are likely to account for much of the residual variation. It is misleading to compare the crude COVID-19 CFRs between countries and should be avoided. At the very least, comparisons should be based on age-specific and age-adjusted rates.

Key words: COVID-19, case-fatality rates, age-specific rates, age-adjusted rates, confounding

No funding was provided for this study. The authors declare no conflict of interests

Background

The novel coronavirus SARS-2-CoV 2019 was first identified in Wuhan in China in early December 2019. On January 30, 2020, the World Health Organization declared the disease caused by the virus, a Public Health Emergency of International Concern (PHEIC) (1) and on March 11, 2020, it was declared as a pandemic. By May, 2020, almost all countries had been affected with more than three million cases of COVID-19 reported. Early estimates indicated that the average proportion of deaths among the diagnosed cases, defined as the case-fatality rate (CFR), was around 2.0% (2). However, subsequently, the reported crudeCOVID-19 CFRs varied widely between countries (3,4). A related concept is the infection fatality rate (IFR) which includes asymptomatic cases in the denominator, which need to be identified by screening tests. Since the IFR is rarely available for COVID-19, in this paper we consider only the CFR. The limitations of comparing crude CFRs in general, has been evaluated previously (5). The strong positive association between the COVID-19 CFR and age has been demonstrated both as in observational studies (6,7) and in a model-based analysis (8), particularly over the age of 40.

The interpretation of the CFR depends on the context in which its used. In a single cohort, such as a country, it can indicate the severity of the disease at a single point in time. It can also be used to assess trends in the impact of changes in health care over time. In the current context of the COVID-19 pandemic, CFRs are commonly compared between countries, which sometime leads to speculation on differences in healthcare. This has frequently led to suggestions that the differences in the CFRs somehow reflect the successes and failures of the different countries to control the pandemic. For diseases in general and for COVID-19 in particular, a number of factors could impact on the both the numerator and denominator of the CFR. Examples are possible misclassification of causes of death and the variations in the definition of cases in the denominator. In this paper, we examined the contribution of the age distribution of the cases when comparing the COVID-19 CFRs between different countries. We focused our analyses on six countries, with widely varying COVID-19 CFRs.

Methods

We studied published crude and age-adjusted CFRs in cohorts of cases of COVID-19 in six countries, with varying periods of follow-up. The countries chosen were based on the accessibility of the data at the beginning of May, 2020. The data for each country were not necessarily updated to the time of the study, and the CFR's may have changed over time.

Age-specific data on the cases and deaths were available for six countries: Italy (9), Spain (10), Sweden (11), China (12), S Korea (13) and Israel (unpublished data). All published data on cases and deaths by ten-year age groups (0-9, 10-19, …..80+). The outcome variable was defined as the crude CFR defined as the number of deaths divided by the number of reported cases. The exposure variable was the individual countries. Age-group was considered as a confounding variable. Age-adjustment was carried by the direct method, using the distribution of the combined cases of all six countries by age group, as the standard population. 95% confidence intervals were computed for each age-adjusted rate, using WinPepi (PEPI-for-Windows).

Ethical considerations

Open access aggregative and anonymous data were used and there was no need for ethics committee approval.

Results

The age-specific number of cases, number of deaths and the crude CFRs by country are given in Table 1. In general, the crude overall CFR's varied widely between and 1.6% to 11.9% for Italy. The age distribution of the cases is shown in Figure 1. They distributions vary markedly. The distribution of the cases for Israel and South Korea are heavily weighted in the 20-39 age group, China has a more balanced distribution and Italy and Sweden are heavily weighted in the over 70 age groups.

Table 1. Age-specific case-fatality rate (CFR) from COVID-19 for six countries

Age	China		Israel		Italy		S Korea		Spain		Sweden	
	Cases/died	CFR%	Cases/died	CFR%	Cases/died	CFR%	Cases/died	CFR%	Cases/died	CFR%	Cases/died	CFR%
0-9	416 0	0	838 0	0	589 0	0	112 0	0	288 1	0.3	64 0	0.0
10-19	549 1	0.2	2028	0	766 0	0	513 0	0	593 1	0.2	170 0	0.0
20-29	3619 7	0.2	3299 0	0	3830 2	0.1	2630 0	0	5477 11	0.2	873 3	0.3
30-39	7600 18	1.2	1952 2	0.1	6523 20	0.3	1002 1	0.1	10518 24	0.2	1110 4	0.4
40-49	8571 38	0.4	1792 1	0.0	12084 89	0.7	1297 1	0.1	16451 66	0.2	1518 10	0.7
50-59	10008 130	1.3	1752 3	0.2	18678 369	2.0	1812 10	0.6	20342 206	1.0	2026 41	2.0
60-69	8583 309	3.6	1364 14	1.1	16395 1162	7.1	1218 21	1.7	18095 634	3.4	1655 95	5.7
70-79	3918 312	8.0	761 41	4.0	17464 3456	19.8	640 45	7.0	17297 1848	10.7	1683 301	17.9
80+	1408 208	14.8	541 110	13.5	17759 4923	27.7	437 80	18.3	20085 4325	21.5	2822 759	26.9
Total	44672 1023	2.3	14327 230	1.6	124352 14589	11.9	10728 242	2.3	204377 15674	7.7	18633 2194	11.8

Figure 1. Cases distribution by age for six countries (China, Italy, S Korea, Spain, Israel and Sweden)

Figure 2. Age specific case-fatality rates for six countries – China, Italy, S Korea, Spain, Israel and Sweden

The age-specific CFRs are shown in Figure 2. While there are still marked differences in the age-specific CFRs, the trend of steeply increasing CFRs in the oldest age groups is evident. The crude and age-adjusted CFRs are compared in Table 2. The crude CFRs varied from 1.6% for Israel to 11.9% for Italy and the ratios of the crude CFRs compared with lowest

CFR, varied between 1.4 and 7.4. The age-adjusted CFRs varied between 4.8% for China to 10.8% for Italy and the ratios of the age-adjusted CFRs for each country compared with the lowest CFR adjusted varied between 1.2 to 2.3. For example, the ratio of the CFR for Italy to Israel dropped from 7.4 to 1.7. Figure 3 shows the marked reduction in the differences between the crude and age-adjusted CFRs for the six countries.

Table 2. Crude and age standardized case-fatality rates (CFR) for COVID-19 in six countries

Country	Crude CFR %	Age-adjusted CFR rate % (95% CI)	Ratio to lowest crude CFR	Ratio to lowest age-adjusted CFR
Israel	1.6	6.4 (5.6-7.1)	1.0	1.3
Korea	2.3	6.5 (5.7-7.3)	1.4	1.4
China	2.3	4.8 (4.4-5.2)	1.4	1.0
Spain	7.7	6.9 (6.8-7.0)	4.8	1.4
Sweden	11.8	10.6 (10.2-11.0)	7.4	2.2
Italy	11.9	10.8 (10.6-11.0)	7.4	2.3

Figure 3. Crude and age-adjusted case-fatality rates by country for six countries

Discussion

In our study, we examined crude, age-specific and age-adjusted CFRs for COVID-19 in six countries, with widely varying crude CFRs. The trends in the age-specific CFRs were remarkably similar in the six countries, with the CFR's increasing steeply in those over 70. After adjusting for age, the marked differences in the crude CFRs were substantially reduced. These findings demonstrate the importance of accounting for age when comparing rates in general and CFRs in particular. The results of this study are strengthened by the use of national data or large datasets from a number of countries, with considerable differences in the extent of the pandemic in each country. It should be stressed that the age distribution of the cases and not the age distribution of the total population in each country cases that was used to compute the age-adjusted CFRs. This was previously demonstrated in a study of three countries (7).

In addition to the age distribution of the cases, the use of the CFR for comparisons between countries has other important limitations. Selection bias is clearly present when calculating the denominator on the basis of reported cases. As mentioned, the CFR must be distinguished from the IFR, which includes asymptomatic cases identified by deliberate or incidental screening with diagnostic tests. In addition, if only those with more severe symptoms are tested this will affect the denominator of the CFR and will depend on the testing strategy of each country. If more mild cases are identified, this is likely to reduce the CFR. Since there is a lag time between the reporting of the case and the death which can occur up to weeks later. In the country reports, cases and deaths are usually reported at the same time, so the cases in the denominator are usually an overestimate of the true denominator which should be the number of cases reported sometime earlier (14,15). This will have a more dramatic effect when the number of cases are rising rapidly. Selection bias may also affect the numerator if only deaths occurring in hospital are reported. Information bias can be present inboth the numerator and denominator of the CFR. The definition of the cases due to may be biased due to the variability of the sensitivity and specificity of the diagnosis of COVID-19. Information bias in the numerator can occur when the cause of death is coded. This could be particularly problematic in elderly people with multiple co-morbidities.

However, while these factors will affect the calculation of the true CFR, the purpose of this paper is to demonstrate the dramatic effect of confounding by the age distribution of the cases when using crude CFRs for country comparisons. This was shown in an earlier paper when

comparing three countries, and we have extended it to a comparison of six countries with widely different CFRs. In this study, the age structures of the population of the six countries included varied markedly. The percentage of the population 65 and over is 11% in Israel, in Italy 23% in Italy, S Korea 14%, in Spain 19%, in Sweden 20% and in China 11%. However, the main impact of confounding by age was due to the differences in the age distribution of the cases. This was largely due to the specific circumstances of exposure. For example, most of the cases in Italy occurred in an area of a particularly old population (3). In some countries, many of the cases were medical personnel, a large number of whom were relatively young women (16). In South Korea, a large percentage of the cases were young women associated with a specific religious group (17). In Germany, many of the cases were relatively young people returning from skiing holidays in Austria and Italy (18). In Israel, the largest outbreaks occurred in the ultra-orthodox Jewish community, where the number of children per family is much higher than in the general population. Other factors affecting the age distribution of the cases, depended on the frequency of outbreaks in homes for the elderly (19).

The results of this study once again demonstrate the pitfalls of comparing unadjusted rates. The assumption that differences between countries in testing policies or standard of treatment accounted for the wide discrepancies in CFRs, is not well-founded. This does not mean that there are no differences. For example, it is possible that where the health services were overloaded, younger patients were more likely to be admitted to intensive care units with better chances of survival. Clearly, the data are incomplete and other factors affecting CFRs such as case definitions, use of different denominators, underlying health conditions and the standard of health services are likely to play important roles. In order to assess the impact of these factors, age-specific and age-adjusted CFRs must be used.

Conclusions

In addition to the selection and information biases inherent in computing CFRs, the age structure of the cases dramatically impacts on the differences in the crude CFRs between countries. Failure to account for this source of confounding markedly distorts the country comparisons. The substantial reduction in the differences in the age-adjusted CFRs suggest that differences in the standard of healthcare between these countries may not play as important a role in affecting the death rates, as some have hypothesized. Crude COVID-19 CFRs have no real use for between country comparisons and should be avoided. In general,

for comparisons between groups and countries, age-adjusted CFRs can be used, but age-specific COVID-19 CFRs are generally far more meaningful.

Acknowledgements

We express our appreciation to the official institutions of all countries for the providing their data on Covid-19.

Funding

No funding was provided for the study

Conflicts of interests

The authors declare that they have no conflicts of interests.

References

1. WHO: Available at: https://www.who.int/emergencies/diseases/novel-coronavirus-2019/events-as-they-happen. Accessed on1 May 2020.

2. Coronavirus disease 2019 (COVID-19): situation report - 36. Geneva: World Health Organization, February 25, 2020. Available at: https://www .who .int/ docs/ default - source/coronavirus/ situation-reports/20200225 -sitrep-36-covid-19.pdf?sfvrsn=2791b4e0_2. Accessed on 20 April 2020

3. Onder G, Rezza G, Brusaferro S. Case-fatality rate and characteristics of patients dying in relation to COVID-19 in Italy. JAMA. 2020

4. Yang S, Cao P, Du P, Wu Z, Zhuang Z, Yang L, et al. Early estimation of the case fatality rate of COVID-19 in mainland China: a data-driven analysis. Ann Transl Med. 2020 Feb;8(4):128. doi: 10.21037/atm.2020.02.66.

5. Lipsitch M, Donnelly CA, Fraser C, Blake IM, Cori A, Dorigatti I, et al. Potential Biases in Estimating Absolute and Relative Case-Fatality Risks during Outbreaks. PLoS Negl Trop Dis. 2015 Jul 16;9(7):e0003846. doi: 10.1371/journal.pntd.0003846. eCollection 2015.

6. Mahase E. Covid-19: death rate is 0.66% and increases with age, study estimates. BMJ 2020;369:m1327.

7. Kim DD, Goel A. Estimating case fatality rates of COVID-19. Lancet Infect Dis. 2020 Mar 31:S1473-3099(20)30234-6. doi:

8. Verity R, Okell LC, Dorigatti I, Winskill P, Whittaker C, Imai N, et al. Estimates of the severity of coronavirus disease 2019: a model-based analysis. The Lancet Infectious Diseases. 2020.

9. Epidemia COVID-19. Aggiornamento nazionale. 6 aprile 2020. Available at: https://www.epicentro.iss.it/coronavirus/bollettino/Bollettino-sorveglianza-integrata-COVID-19_6-aprile-2020.pdf. Accessed on 25 April 2020.

10. Centro de Coordinación de Alertas y Emergencias Sanitarias. Available at: https://www.mscbs.gob.es/profesionales/saludPublica/ccayes/alertasActual/nCov-China/documentos/Actualizacion_88_COVID-19.pdf. Accessed on 1 May 2020.

11. Statista. Available at: https://www.statista.com/topics/6267/coronavirus-covid-19-in-sweden/. Accessed on 20 April 2020.

12. Zhonghua Liu Xing Bing Xue Za Zhi. The epidemiological characteristics of an outbreak of 2019 novel coronavirus diseases (COVID-19) in China. Novel Coronavirus Pneumonia Emergency Response Epidemiology Team. 2020;41:145-151.

13. Korea Centers for Disease Control (CDC). Available at: https://www.cdc.go.kr/board/board.es?mid=a30501000000&bid=0031&list_no=367045&act=view. Accessed on 20 April 2020

14. Baud D, Qi X, Nielsen-Saines K, Musso D, Pomar L, Favre G. Real estimates of mortality following COVID-19 infection. 1. Lancet Infect Dis. 2020 Mar 12:S1473-3099(20)30195-X. doi: 10.1016/S1473-3099(20)30195-X.

15. Lipsitch M. Estimating case fatality rates of COVID-19. 1. Lancet Infect Dis. 2020 Mar 31:S1473-3099(20)30245-0. doi: 10.1016/S1473-3099(20)30245-0.

16. Kang L, Ma S, Chen M, Yang J, Wang Y, Li R, Yao L, Bai H, Cai Z, Xiang Yang B, Hu S, Zhang K, Wang G, Ma C, Liu Z. Impact on mental health and perceptions of psychological care among medical and nursing staff in Wuhan during the 2019 novel coronavirus disease outbreak: A cross-sectional study.Brain Behav Immun. 2020. pii: S0889-1591(20)30348-2.

17. Korean Society of Infectious Diseases; Korean Society of Pediatric Infectious Diseases; Korean Society of Epidemiology; Korean Society for Antimicrobial Therapy; Korean Society for Healthcare-associated Infection Control and Prevention; Korea Centers for Disease Control and Prevention. Report on the Epidemiological Features of Coronavirus Disease 2019 (COVID-19) Outbreak in the Republic of Korea from January 19 to March 2, 2020. J Korean Med Sci. 2020;35(10):e112.

18. A German Exception? Why the Country's Coronavirus Death Rate Is Low. New York Times, Published April 4, 2020, updated April 6, 2020.

19. Available at: https://www.euronews.com/2020/04/20/coronavirus-and-the-elderly-what-s-life-inside-a-french-retirement-home-amid-covid-19-pand.