

Medical overuse in the Iranian healthcare system: a systematic scoping review and practical recommendations for decreasing medical overuse during unexpected COVID-19 pandemic opportunity

Mohammad Zakaria Pezeshki¹, Ali Janati², Morteza Arab-Zozani^{3,4}

1. Social Determinants of Health Research Center, Department of Community and Family Medicine, Tabriz Medical School, Tabriz University of Medical Sciences, Tabriz, Iran
2. Iranian Center of Excellence in Health Management, Department of Health Services Management, School of Management and Medical Informatics, Tabriz University of Medical Sciences, Tabriz, Iran
3. Social Determinants of Health Research Center, Birjand University of Medical Sciences, Birjand, Iran
4. Student Research Committee, Tabriz University of Medical Sciences, Tabriz, Iran

Corresponding author; Morteza Arab-Zozani

Tel: (+98) 9153317843

Email address: arab.hta@gmail.com

Funding Sources: This work was supported by Tabriz University of Medical Sciences [grant number 5/d/633456, 15 January 2018; IR.TBZMED.REC.1396.908].

Running title: Medical overuse in Iran

Abstract

Background: Overuse of medical care is a major problem across health systems as well as in Iran. By our knowledge, this is the first scoping review in which medical overuse in the Iranian healthcare system was investigated.

Objective: To perform an inclusive search for original studies that report medical overuse in the Iranian healthcare system.

Methods: A systematic search of the literature conducted in bibliographic databases including PubMed, Embase, Scopus, Web of Sciences, Cochrane and Scientific Information Database using a comprehensive search strategy without time limit until the end of 2018, accomplished by reference tracking, author contacting and expert consultation to identify studies on the overuse of medical care.

Results: We reviewed 4124 published articles based on predetermined inclusion criteria. The author's consensus included a total of 40 articles. Of these, 31 were in English and 9 in Farsi, published between 19975-2018. The result categorized into two distinct clinical areas: treatment (17 articles), and diagnostic (23 articles) services. Almost all of the studies only described the magnitude of unnecessary overuse. Unnecessary overuse of Antibiotics, MRI, and CT-scan were the most reported topics. The ranges of their overuse proportion were as follows; Antibiotic (31 to 97%); MRI (33 to 88%), and CT-scan (19 to 50%).

Conclusions: Our review showed, even so, the magnitude of unnecessary overuse of medical services is high but there are only a few interventional studies in clinical and administrative level for finding effective methods for decreasing these unnecessary services. Researchers should be encouraged to conducting interventional studies. We suggest the ministry of health to use the golden opportunity of COVID-19 epidemic for designing Iran national policy and

action plan for controlling and preventing unnecessary health care services and including a section for “Interventional Research” in the action plan.

Keywords: Medical Overuse; Healthcare system; Iran

Introduction

Medical overuse means services that "are more harmful than beneficial, does not seem to increase the quality and quantity of life, impose excessive cost on the patients and their healthcare system, has low quality and if the patient has enough information, he or she will not ask for it" (1, 2). Overuse can take place in different area including medication, test or procedure (3). Recent studies have shown that overuse of tests and treatments can lead to serious consequences on patients in six domain including physical, psychological, social, financial, treatment burden, and dissatisfaction with health care (4). Medical overuse can delay access to the goals of health systems-improved health, responsiveness, financial risk protection and efficiency-by increasing cost and decreasing quality of medical care (5).

Given the fact that medical overuse is one of the problems that can make system performances problematic, its identification is of great importance for a health system (6). In addition, overuse in medical care is one of the obstacles to achieving universal health coverage (UHC) (7). If we want to achieve better UHC, we need to be able to manage costs, and one of the most important tasks in this direction is to reduce overuse in medical services (8, 9).

Over the years, many efforts have been made to identify overuse in medical care across health systems around the world (10, 11). In Iran, as in many other countries, there is little evidence about the amount of medical overuse in the health care system (1). So, the identification of medical overuse is an essential issue for the Iranian healthcare system and

also helps health policymakers, health and medical managers, researchers, general practitioners, patients and their families to cope with harms, costs and quality of services. Ultimately, identifying overuse of medical care can bring our health system to balance in the right use of services. Thus, the objectives of this systematic scoping review were 1) to review the literature on the overuse of medical care, (2) to identify the areas in which the overuse of medical care take placed (3) to determine the rate of overuse of medical care in the Iranian healthcare system and its drivers and (4) to identify the interventional studies in clinical and administrative level for decreasing the rate of overuse

Methods

We conducted a systematic scoping review on medical overuse in accordance with the five stages outlined in the Arksey and O'Malley framework (12). The review protocol was registered in PROSPERO before starting our study (registration no. CRD42017075481).

Stage 1: Identifying research questions

The following questions guided this scoping review of medical overuse in the Iranian healthcare system: How many literatures have dealt with this issue in the Iranian healthcare system? Which area does medical overuse occur? What is the rate of medical overuse in the Iranian healthcare system? What are the drivers and preventive ways of medical overuse in the Iranian healthcare system?

Stage 2: Identifying relevant studies

All original articles that investigated the overuse in medical care were included in the study. Of these, only studies were included that has addressed overuse in the Iranian healthcare system. All the included studies were limited in English and Farsi languages. Articles were excluded if the researchers did not have access to the Full-text.

We searched six databases including PubMed, Web of Science, Embase, Scopus, Cochrane and Scientific Information Database (SID) without time limit until the end of 2018. We also contacted the authors of included studies and use reference tracking to get the articles we probably did not find in the search. We used a set of Medical Subject Headings (MeSH) terms and free term to maximize the sensitivity of the search. For more information on search strategy, see study protocol at: <https://bmjopen.bmj.com/content/8/4/e020355#ref-17> (1).

Stages 3 and 4: Study selection and data charting process

After the search was completed, duplicate records were removed. Then, two reviewer screened the records based on title, abstract and full text, and extracted the data about authors, publication year, type of study, study population, type of service, clinical area, and overuse rate or range. . These set of checklists designed for various types of studies and are available online (<http://joannabriggs.org/research/critical-appraisal-tools.html>). All potential disagreements in each level of study were resolved by consensus with a third researcher.

Stage 5: Summarizing results

We categorized the results of the included studies based on publication year, clinical area, type of service (diagnostic tests, therapeutic procedures and medications), and range or rate of overuse.

Results

Initially, a total of 4179 records were screened. After removing duplicates, 3023 records were considered for eligibility. Of these, 40 studies were included. The study selection process is outlined in the Preferred Reporting Items for Systematic Reviews and Meta-analysis (PRISMA) diagram (Figure 1).




Figure 1: study flow diagram

Most studies were published in English (77.5%). Included studies published between 1975 and 2018. Most studies were published in 2014 (25%), 2012 (15%) and 2011 (12.5%). Also, in term of design, 37 were cross-sectional, and three RCT studies. The summary characteristics of included studies are shown in Tables 1.

Table 1: The summary characteristics of the included studies

Authors	Year	Language	Study Design	Sample Size
Ahmadi et al, (13)	2006	Fa	Cross-Sectional	400
Ahmadizar et al, (14)	2011	En	Cross-Sectional	4456755
Amidi et al, (15)	1975	En	Cross-Sectional	40
Alavi et al, (16)	2014	En	Cross-Sectional	410
Alavi et al, (17)	2014	En	Cross-Sectional	8586

Azami et al, (18)	2006	Fa	Cross-Sectional	150
Ahmadi et al, (19)	2014	En	Cross-Sectional	1309
Alizadeh et al, (20)	2012	En	RCT	200
Bijani et al, (21)	2014	En	Cross-Sectional	1543
Hatam et al, (22)	2011	En	Cross-Sectional	1000
Khakhshour et al, (23)	2011	Fa	Cross-Sectional	292
Refahi et al, (24)	2016	En	Cross-Sectional	115
Sobhani et al, (25)	2001	Fa	Cross-Sectional	4750
Saadat et al, (26)	2008	En	Cross-Sectional	1650
Sadeghi et al, (27)	2015	Fa	Cross-Sectional	2105
Taghizadeh et al, (28)	2013	En	Cross-Sectional	234
Zargar et al, (29)	2014	En	Cross-Sectional	400
Vessal et al, (30)	2011	En	Cross-Sectional	155
Bakhit et al, (31)	2014	En	Cross-Sectional	270
Bayani et al, (32)	2014	Fa	Cross-Sectional	400
Bilehjani et al, (33)	2017	En	Cross-Sectional	620
Davoodian et al, (34)	2012	En	Cross-Sectional	206
Eini et al, (35)	2012	Fa	Cross-Sectional	100
Fard et al, (36)	2001	En	Cross-Sectional	279
Forouzanfar et al, (37)	2014	En	Cross-Sectional	2607
Ghadimi et al, (38)	2011	En	Cross-Sectional	2041
Ghaffarpassand et al, (39)	2011	En	Cross-Sectional	1679
Ghazizadeh et al, (40)	2009	En	RCT	60
Jame et al, (41)	2014	En	Cross-Sectional	400
Moussavi et al, (42)	2015	En	RCT	100
Khaji et al, (43)	2006	En	Cross-Sectional	1209
Meidani et al, (44)	2017	Fa	Cross-Sectional	361
Meidani et al, (45)	2016	En	Cross-Sectional	384
Memari et al, (46)	2012	En	Cross-Sectional	345

Moghimi et al, (47)	2008	Fa	Cross-Sectional	1220
Mohammadi et al, (48)	2016	En	Cross-Sectional	279
Mokhtari et al, (49)	2014	En	Cross-Sectional	1219
Nikbakhsh et al, (50)	2010	En	Cross-Sectional	498
Raji et al, (51)	2018	En	Cross-Sectional	112
Paydar et al, (52)	2012	En	Cross-Sectional	5091

The result categorized onto two distinct clinical area: treatment (17 articles), and diagnostic (23 articles) area. Unnecessary overuse of Antibiotics, MRI and CT-scan were the most reported topics. The ranges of their overuse proportion were as follow: Antibiotic (31 to 97%); MRI (33 to 88%), and CT-scan (19 to 50%). Among the studies in the area of treatment, the most studied were antibiotics (7 studies, 41.1%), and three studies (17.6%) did not compare the results with any other standard. Also, among the studies in the area of diagnosis, the most studied were related to MRI (4 studies, 17.3%), and CT (4 studies, 17.3%), and three studies (13%) did not compare the results with any other standard. For more details see Table 2 and 3.

Table 2: The characteristics of included studies in treatment area

Authors	Clinical Domain	Subject	The ranges of overuse proportion (%)	Standard
Amidi et al,	Treatment	Antibiotic	92.5	Clinical guideline
Khakhshour et al,	Treatment	Antibiotic	70	Clinical guideline
Alavi et al,	Treatment	Antibiotic	41.4	Clinical guideline
Vessal et al,	Treatment	Antibiotic	31.6	American Society of Hospital Pharmacists (ASHP) guidelines
Hatam et al,	Treatment	Antibiotic	98	ASHP
Alavi et al,	Treatment	Antibiotic	44	Mangram's guideline

Taghizadeh et al,	Treatment	Antibiotic	61	NR
Ahmadizar et al,	Treatment	Drug-Drug Interaction	0.77	National guideline
Bijani et al,	Treatment	Polypharmacy	16.5- 35.1	National guideline
Sobhani et al,	Treatment	Polypharmacy	88	NR
Ahmadi et al,	Treatment	Polypharmacy	39.6	National guideline
Azami et al,	Treatment	Blood transfusion	26.8	Scientific criteria
Ahmadi et al,	Treatment	Intravenous Acetaminophen	NR	NR
Alizadeh et al,	Treatment	Heparin	NR	Heparinization protocol
Ghadimi et al,	Treatment	Prescribing pattern	30	Beers criteria & WHO indicators
Ghazizadeh et al,	Treatment	GnRH antagonist	NR	Study protocol
Memari et al,	Treatment	Psychotropic medication	80	DSM-IV-TR criteria

NR; Not Reported

Table 3: The characteristics of included studies in diagnostic area

Authors	Clinical Domain	Subject	The ranges of overuse proportion (%)	Standard
Refahi et al,	Diagnostic	MRI	45.2	Local guideline
Zargar et al,	Diagnostic	MRI	46.5	Clinical guideline
Saadat et al,	Diagnostic	MRI	82.8	Clinical guideline
Sadeghi et al,	Diagnostic	MRI	76	Clinical guideline
Bakhit et al,	Diagnostic	Diagnosis of dizziness	NR	Clinical guideline
Bayani et al,	Diagnostic	Clinical diagnosis and candida culture	NR	Clinical guideline
Bilehjani et al,	Diagnostic	Erythrocyte Sedimentation Rate (ESR)	NR	NR

Davoodian et al,	Diagnostic	Urinary catheters	20.6	NR
Eini et al,	Diagnostic	Antibacterial therapy	97	Clinical guideline
Fard et al,	Diagnostic	Venous duplex ultrasonography (VDUS)	NR	Scientific criteria
Forouzanfar et al,	Diagnostic	Chest X-ray (CXR)	7.5	Thoracic Injury Rule out Criteria (TIRC)
Ghaffarpasand et al,	Diagnostic	Radiography	NR	ATLS protocol
Jame et al,	Diagnostic	Computed tomography	19.8-51.6	Glasgow coma score
Moussavi et al,	Diagnostic	Computed tomography	NR	Glasgow coma score
Khaji et al,	Diagnostic	Computed tomography	66.5	Glasgow coma score
Meidani et al,	Diagnostic	Computed tomography	14.1	ACR criteria
Meidani et al,	Diagnostic	Laboratory test	26.4	ACR criteria
Moghimi et al,	Diagnostic	Preclinical test	1.3-9.6	NR
Mohammadi et al,	Diagnostic	MRI	33	NICE and AHRQ guidelines
Mokhtari et al,	Diagnostic	Venous thromboembolism (VTE) prophylaxis	NR	ACCP guidelines
Nikbakhsh et al,	Diagnostic	electrocardiogram (ECG)	77.3	American Society of Anesthesiologists status (ASA) criteria
Rajier al,	Diagnostic	Pulmonary CT angiography	NR	Geneva score and Wells' criteria
Paydar et al,	Diagnostic	Routine chest radiography for stable blunt trauma	19.8	ATLS

NR; Not Reported

Discussion

The aim of this systematic scoping review was to perform an inclusive search for original studies that report medical overuse in the Iranian healthcare system. Finally 40 original studies were included in our study, of which 17 article related to treatment area and 23 article related to diagnostic area. Antibiotics and MRI were the most reported issues in each category where overuse has been reported.

As Table 2 and Table 3 shows the majority of studies only have focused on the magnitude of unnecessary diagnostic and treatment services. There are only few interventional studies regarding diagnostic and treatment services. Also there is not any study regarding unnecessary clinical preventive services like unnecessary check up and also unnecessary public health services. Unfortunately as the Tables shows there is not any study at regional or national level that clarify the drivers of unnecessary services in Iran and how to address them. To address the shortage of study regarding the interventions for decreasing overuse rate in Iran we already conducted a qualitative research at national level to clarify the drivers of overuse and strategies for controlling these drivers in Iran. In this qualitative study we did interview with 21 well respected old hand policy makers and researchers of Iran. After analyzing the interview, our study showed that the main drivers of unnecessary overuse in the Iranian healthcare system are physician, patient, organizational, socio-cultural, market, and mass media factors. Also, a Policy Delphi analysis as part of our national study and based on the key informant's opinion (53, 54), showed that the main interventions for decreasing unnecessary overuse of medical services include; implementing strategic purchasing, active engaging of insurance companies, promoting payment system, use of clinical practice guideline in decision making, and increasing political commitment and reducing conflicts of interest. We are going to publish the results of our study in detail.

COVID-19 pandemic has created a golden opportunity for addressing the drivers of unnecessary overuse of medical services by countries because of the three main reasons : 1)

There is shortage of health care resources for controlling COVID-19 pandemic and unnecessary services waste the resources 2) Overuses of health care services unnecessarily expose the patients and healthy individuals to the virus in outpatient clinics and hospitals, 3) Overuse of medications may suppress the immune response and predispose people to COVID-19 infection. Our preliminary search shows that COVID-19 pandemic has decreased the use of several clinical interventions in countries (55) for example screening tests (56), admission and hospitalization (57), and elective surgeries (58, 59). Considerable proportions of these clinical interventions are unnecessary. We suggest Iranian ministry of health to use the golden opportunity of COVID-19 pandemic to develop national policy and action plans for controlling and preventing unnecessary health care services in Iran. These policies will facilitate the controlling of COVID-19 epidemic and preventing underuse of necessary services during COVID-19 epidemic and after the end of epidemic.

Conclusion

Our systematic review shows even so the magnitude of unnecessary overuse of medical services is high but there are only few interventional studies in clinical and administrative level for finding effective methods for decreasing these unnecessary services. Researchers should be encouraged for conducting such researches. It is necessary to be included a section for “Interventional Research” in the action plans we suggest to the ministry of health for controlling and preventing unnecessary health care services in Iran.

References

1. Arab-Zozani M, Pezeshki MZ, Khodayari-Zarnaq R, Janati AJBo. Medical overuse in the Iranian healthcare system: a systematic review protocol. 2018;8(4):e020355.
2. Brownlee S, Chalkidou K, Doust J, Elshaug AG, Glasziou P, Heath I, et al. Evidence for overuse of medical services around the world. 2017;390(10090):156-68.
3. Morgan DJ, Dhruva SS, Coon ER, Wright SM, Korenstein DJJIm. 2017 update on medical overuse: a systematic review. 2018;178(1):110-5.

4. Korenstein D, Chimonas S, Barrow B, Keyhani S, Troy A, Lipitz-Snyderman A. Development of a conceptual map of negative consequences for patients of overuse of medical tests and treatments. *JAMA Internal Medicine*. 2018.
5. Korenstein D, Falk R, Howell EA, Bishop T, Keyhani S, Ahoim. Overuse of health care services in the United States: an understudied problem. 2012;172(2):171-8.
6. Morgan DJ, Leppin A, Smith CD, Korenstein DJ, John. A practical framework for understanding and reducing medical overuse: conceptualizing overuse through the patient-clinician interaction. 2017;12(5):346.
7. Kleinert S, Horton RJTL. From universal health coverage to right care for health. 2017;390(10090):101-2.
8. Berwick DMJTL. Avoiding overuse—the next quality frontier. 2017;390(10090):102-4.
9. Elshaug AG, Rosenthal MB, Lavis JN, Brownlee S, Schmidt H, Nagpal S, et al. Levers for addressing medical underuse and overuse: achieving high-value health care. 2017;390(10090):191-202.
10. Rosenberg A, Agiro A, Gottlieb M, Barron J, Brady P, Liu Y, et al. Early trends among seven recommendations from the Choosing Wisely campaign. 2015;175(12):1913-20.
11. Rumball-Smith J, Shekelle PG, Bates DWJJ. Using the electronic health record to understand and minimize overuse. 2017;317(3):257-8.
12. Arksey H, O'Malley L. Scoping studies: towards a methodological framework. *International journal of social research methodology*. 2005;8(1):19-32.
13. Ahmadi B, Alimohamadian M, Mahmoodi M. Polypharmacy among older adults in Tehran. *Tehran University Medical Journal TUMS Publications*. 2006;64(9):65-71.
14. Ahmadizar F, Soleymani F, Abdollahi M. Study of drug-drug interactions in prescriptions of general practitioners and specialists in Iran 2007-2009. *Iranian Journal of Pharmaceutical Research*. 2011;10(4):921-31.
15. Amidi S, Solter S, Rashidian B, Zokaian AR, Razmjoian F. Antibiotic use and abuse among physicians in private practice in Shiraz, Iran. *Medical care*. 1975;13(4):341-5.
16. Alavi SM, Roozbeh F, Behmanesh F, Alavi L. Antibiotics use patterns for surgical prophylaxis site infection in different surgical wards of a teaching hospital in Ahvaz, Iran. *Jundishapur Journal of Microbiology*. 2014;7(11).
17. Alavi SM, Sharifi M, Eghtesad M. Bacterial sepsis: Challenges of diagnosis and treatment in a teaching hospital Southwest of Iran. *Jundishapur Journal of Microbiology*. 2014;7(3).
18. Azami A, Amani F, Khorasani S. Unnecessary blood and blood products transfusion to patients admitted to Ardabil Dr Fatemi Hospital, 2003. *Journal of Ardabil University of Medical Sciences*. 2006;6(4):345-50.
19. Ahmadi K, Hashemian AM, Pishbin E, Sharif-Alhoseini M, Rahimi-Movaghar V. Impact of intravenous acetaminophen therapy on the necessity of cervical spine imaging in patients with cervical spine trauma. *Chinese journal of traumatology = Zhonghua chuang shang za zhi*. 2014;17(4):204-7.
20. Alizadeh A, Yazdi AH, Kafi M, Rad MA, Moradi M, Emkanjoo Z. Predictors of local venous complications resulting from electrophysiological procedures. *Cardiology journal*. 2012;19(1):15-9.
21. Bijani A, Roshan ARH, Yazdanpour S, Hosseini SR. Are older women likely to use medicines than older men? (Results from AHAP study). *Caspian Journal of Internal Medicine*. 2014;5(2):77-81.
22. Hatam N, Askarian M, Moravveji AR, Assadian O. Economic burden of inappropriate antibiotic use for prophylactic purpose in Shiraz, Iran. *Iranian Red Crescent Medical Journal*. 2011;13(4):234-8.
23. Khakhshour A, Taherpour M, Khorashadi Zadeh F, Maddi I, Nojomi S, Shaikhi Z. Unnecessary antibiotic therapy in children with diarrhea may cause antibiotic resistance. *Journal of North Khorasan University of Medical Sciences*. 2011;3(2):29-33.

24. Refahi S, Kachooei AR, Farsadpour M, Shahrayeri R, Goudarzian M, Taleghani YM, et al. IS PRESCRIPTION OF KNEE MRI ACCORDING TO STANDARD CLINICAL GUIDELINE? *Acta Medica Mediterranea*. 2016;32:1207-11.
25. Sobhani A, Shodjai H. Prevalence of polypharmacy and correlations with sex, age and drug regimen in insurance prescription. 2001.
26. Saadat S, Ghodsi SM, Firouznia K, Etminan M, Goudarzi K, Naieni KH. Overuse or underuse of MRI scanners in private radiology centers in Tehran. *International journal of technology assessment in health care*. 2008;24(3):277-81.
27. Sadeghi A, Keshavarz K, Ahmadzadeh MS, Yousefi A. Survey of Appropriate Use of Magnetic Resonance Imaging Services Provided in Shahid Chamran Hospital of Shiraz. *Journal of health research in community*. 2015;1(3):33-40.
28. Taghizadeh S, Haghdoost M, Mashrabi O, Zeynalikhasraghi Z. Antibiotic usage in intensive care units of Tabriz Imam Reza hospital, 2011. *American Journal of Infectious Diseases*. 2013;9(4):123-8.
29. Zargar BJS, Sari AA, Majdzadeh R, Rashidian A, Arab M, Rahmani H. The extent of inappropriate use of magnetic resonance imaging in low back pain and its contributory factors. *International Journal of Preventive Medicine*. 2014;5(8):1029-36.
30. Vessal G, Namazi S, Davarpanah MA, Foroughinia F. Evaluation of prophylactic antibiotic administration at the surgical ward of a major referral hospital, Islamic republic of Iran. *Eastern Mediterranean Health Journal*. 2011;17(8):663-7.
31. Bakhit M, Heidarian A, Ehsani S, Delphi M, Latifi SM. Clinical assessment of dizzy patients: the necessity and role of diagnostic tests. *Global journal of health science*. 2014;6(3):194-9.
32. Bayani M, Asghar Sefidgar SA, Basirat Z, Haji Ahmadi M, Asgharzadeh S, Haddad G, et al. Association of clinical symptoms and laboratory results in diagnoses of *Candida* vaginitis. *Journal of Babol University of Medical Sciences*. 2014;16(1):50-5.
33. Bilehjani E, Fakhari S, Farzin H, Yaghoubi A, Mirinazhad M, Shadvar K, et al. The correlation between preoperative erythrocyte sedimentation rate and postoperative outcome in adult cardiac surgery. *International Journal of General Medicine*. 2017;10:15-21.
34. Davoodian P, Nematee M, Sheikhvatan M. Inappropriate use of urinary catheters and its common complications in different hospital wards. *Saudi journal of kidney diseases and transplantation : an official publication of the Saudi Center for Organ Transplantation, Saudi Arabia*. 2012;23(1):63-7.
35. Eini P, Mobaien A, Agha Sharif M. Importance of diagnostic laboratory methods of beta hemolytic streptococcus group A in comparison with clinical findings in the diagnosis of streptococcal sore throat and unnecessary antibacterial therapy. *ISMJ*. 2012;15(1):59-68.
36. Fard MN, Mostaan M, Anaraki M. Utility of lower-extremity duplex sonography in patients with venous thromboembolism. *Journal of Clinical Ultrasound*. 2001;29(2):92-8.
37. Forouzanfar MM, Safari S, Niazazari M, Baratloo A, Hashemi B, Hatamabadi HR, et al. Clinical decision rule to prevent unnecessary chest X-ray in patients with blunt multiple traumas. *Emergency Medicine Australasia*. 2014;26(6):561-6.
38. Ghadimi H, Esmaily HM, Wahlstrom R. General practitioners' prescribing patterns for the elderly in a province of Iran. *Pharmacoepidemiology and drug safety*. 2011;20(5):482-7.
39. Ghaffarpasand F, Paydar S, Foroughi M, Saberi A, Abbasi H, Karimi AA, et al. Role of cervical spine radiography in the initial evaluation of stable high-energy blunt trauma patients. *Journal of Orthopaedic Science*. 2011;16(5):498-502.
40. Ghazizadeh S, Pourmatroud E, Shariat M, Masomi M, Bagheri M. Study of Positive and Negative Consequences of Using GnRH Antagonist in Intrauterine Insemination Cycles. *International Journal of Fertility & Sterility*. 2009;3(2):56-61.
41. Jame SZB, Majdzadeh R, Sari AA, Rashidian A, Arab M, Rahmani H. Indications and Overuse of Computed Tomography in Minor Head Trauma. *Iranian Red Crescent Medical Journal*. 2014;16(5).

42. Moussavi N, Davoodabadi AH, Atoof F, Razi SE, Behnampour M, Talari HR. Routine Chest Computed Tomography and Patient Outcome in Blunt Trauma. *Archives of Trauma Research*. 2015;4(2).
43. Khaji A, Eftekhari B, Karbakhsh M, Ardalan KM. The use of head CT scanning in mild head injury. *Neurosciences*. 2006;11(4):248-51.
44. Meidani Z, Moosavi G, Hamidian Y, Farzandipour M, Aliasgharzadeh A, Nazemi Bidgoli Z. Assessing the Appropriate Utilization of CT Scan in Triage Units. *Hospital*. 2017;16(2):27-35.
45. Meidani Z, Farzandipour M, Farrokhian A, Haghighat M. A review on laboratory tests' utilization: A trigger for cutting costs and quality improvement in health care settings. *Medical Journal of the Islamic Republic of Iran*. 2016;30(1).
46. Memari AH, Ziaee V, Beygi S, Moshayedi P, Mirfazeli FS. Overuse of psychotropic medications among children and adolescents with autism spectrum disorders: perspective from a developing country. *Research in developmental disabilities*. 2012;33(2):563-9.
47. Moghimi M, Marashi Sa, Salehian M, Motavalizadeh K. Unnecessary paraclinical tests before elective surgeries. *Journal of Iranian Society Anesthesiology and Intensive Care*. 2008;30(62):33-70.
48. Mohammadi N, Farahmand F, Kharazi HH, Mojdehipanah H, Karampour H, Nojomi M. Appropriateness of physicians' lumbosacral MRI requests in private and public centers in Tehran, Iran. *Medical Journal of the Islamic Republic of Iran*. 2016;30(1):1-7.
49. Mokhtari M, Attarian H, Norouzi M, Koucheh M, Kashani BS, Sirati F, et al. Venous thromboembolism risk assessment, prophylaxis practices and interventions for its improvement (AVAIL-ME Extension Project, Iran). *Thrombosis Research*. 2014;133(4):567-73.
50. Nikbakhsh N, Alijanpoor E, Adabi F. Preoperative medical evaluation in elective surgery versus standard criteria. *Caspian Journal of Internal Medicine*. 2010;1(2):53-7.
51. Raji H, Javadmoosavi SA, Dastoorpoor M, Mohamadipour Z, Mousavi Ghanavati SP. Overuse and underuse of pulmonary CT angiography in patients with suspected pulmonary embolism. *Medical Journal of the Islamic Republic of Iran*. 2018;32:3-.
52. Paydar S, Johari HG, Ghaffarpasand F, Shahidian D, Dehbozorgi A, Ziaeeian B, et al. The role of routine chest radiography in initial evaluation of stable blunt trauma patients. *American Journal of Emergency Medicine*. 2012;30(1):1-4.
53. Arab-Zozani M, Janati A, Zakaria Pezeshki M, Khodayari-Zarnaq R. Policy package for preventing overuse and underuse of health care services in the Iranian health care system: A study protocol. *Medical Journal of the Islamic Republic of Iran*. 2019;33:86-.
54. Arab-Zozani M, Pezeshki MZ, Janati A, Khodayari-zarnaq R. 6 Medical overuse: how to define, detect and prevent? *BMJ Evidence-Based Medicine*. 2019;24(Suppl 2):A10-A.
55. Emanuel L, Navathe A. Will 2020 Be the Year That Medicine Was Saved? *The New York Times*. 2020;<https://www.nytimes.com/2020/04/14/opinion/coronavirus-hospitals.html>.
56. Bewley S. Things should never be the same again in the screening world. *The BMJ* 2020;<https://blogs.bmj.com/bmj/2020/04/14/susan-bewley-things-should-never-be-the-same-again-in-the-screening-world/>.
57. Arab-Zozani M, Pezeshki MZ, Khodayari-Zarnaq R, Janati A. Inappropriate Rate of Admission and Hospitalization in the Iranian Hospitals: A Systematic Review and Meta-Analysis. *Value in Health Regional Issues*. 2020;21:105-12.
58. Elshaug A, Duckett S. Hospitals have stopped unnecessary elective surgeries – and shouldn't restart them after the pandemic. *The Conversation*. 2020;<https://theconversation.com/hospitals-have-stopped-unnecessary-elective-surgeries-and-shouldnt-restart-them-after-the-pandemic-136259>.
59. Prasad S. An Unexpected Opportunity Presented by the COVID-19 Pandemic. The clear health. 2020;<https://www.realclearhealth.com/articles/2020/04/03/an-unexpected-opportunity-presented-by-the-covid-19-pandemic-111011.html>.

