DNA methylation mediates the effect of cocaine use on HIV severity

Chang Shu[1,2], Amy C. Justice[2,3], Xinyu Zhang[1,2], Zuoheng Wang[4], Dana B. Hancock[5], Eric O.

Johnson[5,6], Ke Xu[1,2]

1. Department of Psychiatry, Yale School of Medicine, New Haven, CT, USA

2. Connecticut Veteran Healthcare System, West Haven, CT, USA

3. Department of Internal Medicine, Yale School of Medicine, New Haven, CT, USA

4. Department of Biostatistics, Yale School of Public Health, New Haven, CT, USA

5. GenOmics, Bioinformatics, and Translational Research Center, Biostatistics and

9   Epidemiology Division, RTI International, Research Triangle Park, NC, USA

6. Fellow Program, RTI International, Research Triangle Park, NC, USA.

Keywords: Cocaine use, HIV severity, mortality, DNA methylation, mediation effect

Running title: DNA methylation mediates cocaine exposure on HIV

Word count:

Abstract: 276; Text: 4,785

Figures: 4

Tables: 5

Corresponding Author

Ke Xu, MD., Ph.D.

Associate Professor of Psychiatry

Yale School of Medicine

950 Campbell Ave, West Haven, CT

Email: ke.xu@yale.edu

Tel: 203-932-5711 x7430

Abstract

Background: Cocaine use accelerates human immunodeficiency virus (HIV) progression and worsens HIV outcomes. We assessed whether DNA methylation in blood mediates the association between cocaine use and HIV severity in a veteran population.

Methods: We analyzed 1,435 HIV-positive participants from the Veterans Aging Cohort Study Biomarker Cohort (VACS-BC). HIV severity was measured by the Veteran Aging Cohort Study (VACS) index. We assessed the effect of cocaine use on VACS index and mortality among the HIV-positive participants. We selected candidate mediators that were associated with both persistent cocaine use and VACS index by epigenome-wide association (EWA) scans at a liberal p-value cutoff of 0.001. Mediation analysis of the candidate CpG sites between cocaine's effect and the VACS index was conducted, and the joint mediation effect of multiple CpGs was estimated. A two-step epigenetic Mendelian randomization (MR) analysis was conducted as validation.

Results: More frequent cocaine use was significantly associated with a higher VACS index (β=1.00, p=2.7E-04), and cocaine use increased the risk of 10-year mortality (hazard ratio=1.10, p=0.011) with adjustment for confounding factors. Fifteen candidate mediator CpGs were selected from the EWA scan. Twelve of these CpGs showed significant mediation effects, with each explaining 11.3%-29.5% of the variation. The mediation effects for 3 of the 12 CpGs were validated by the two-step epigenetic MR analysis. The joint mediation effect of the 12 CpGs accounted for 47.2% of cocaine's effect on HIV severity. Genes harboring these 12 CpGs are involved in the antiviral response (*IFIT3*, *IFITM1*, *NLRC5*, *PLSCR1*, *PARP9*) and HIV progression (*CX3CR1*, *MX1*).

Conclusions: We identified 12 DNA methylation CpG sites that appear to play a mediation role in the association between cocaine use and HIV severity.

Introduction

Cocaine use is common among persons with chronic human immunodeficiency virus (HIV)

infection, with prevalence estimates for current or recent use ranging from 5% to 30% [1-6],

compared with 2% in the US general population [7]. Previous studies have shown that cocaine

use accelerated HIV progression [8-11]. However, the biological mechanism of cocaine's effect

on HIV outcomes remains largely unknown. Some studies have suggested that cocaine use

may worsen HIV outcomes due to poor adherence to antiretroviral therapy (ART) among HIV-

positive participants [2, 12]. Other studies have demonstrated that cocaine's adverse effect on

HIV outcomes is independent of ART [10, 11, 13-15], supporting the hypothesis that cocaine

exposure may lead to long-lasting pathophysiological changes in the immune system that

worsen HIV outcomes.

DNA methylation (DNAm) is an important mechanism associated with many environmental

exposures such as smoking, alcohol, and drug misuse [16-26] and diseases such as cancer,

diabetes and cardiovascular diseases [27-32]. Our previous study showed that two DNAm CpG

sites in *NLRC5* were differentially methylated between HIV-positive and HIV-negative

participants in peripheral blood [33]. DNAm may play an important mediation role linking

environmental exposure and disease outcomes [34-40]. Environmental exposure such as

substance use or toxicants can directly or indirectly affect DNA methylatransferases, causing

global or site-specific DNAm changes that may lead to disease [41]. A recent study reported

that DNAm sites in *PIM3* (energy metabolism) and *ABCG1* (lipid metabolism) mediated the

association between prenatal famine exposure and long-term metabolic outcomes [38]. Another

study reported the mediation effect of cg05575921 (*AHRR*) between smoking and the risk of

bladder cancer among postmenopausal women [42].

Previous studies have shown that the use of cocaine enhances HIV-1 replication and

undermines immune function by dysregulating gene expression on HIV-1 entry co-receptors,

enhancing HIV-1 cellular toxicity and dysregulating interleukins (IL) in the host [43, 44]. Cocaine

use increases the release of cytokines in immune cells and alters cytokine profile in HIV-

infected individuals [45, 46]. Specifically, cocaine use was positively associated with IL-4 and IL-

10 [47], which likely worsens HIV severity and disease progression. Epigenetic mechanisms

may play a role in cocaine's effect on the HIV severity because cocaine exposure has been

showed to increase the expression of Methyl CpG binding protein 2 (MeCP2) expression [48] as

well as *DNMT3A* and *DNMT3B* expressions [49] in the animal brains. In a well-matched, case-

control human pilot study, cocaine use alters DNA methylation profile in blood [50]. It is

plausible that cocaine use may lead to DNAm changes in immune response genes and gene

expression changes in cytokine gene family, which further affects HIV progression. Thus, we

hypothesized that DNAm may mediate the effect of cocaine exposure on HIV severity.

In this study, we first validated previous findings by examining cocaine's adverse effect on HIV

severity and mortality. We further conducted mediation analyses to assess the mediation role of

DNAm sites (or CpGs) on cocaine's effect on HIV severity using the Veteran Aging Cohort

Study Biomarker Cohort (VACS-BC, n=1,435). To assess how sensitive our results are to the

violation of model assumptions and validate our findings using a different approach, we

performed the sensitivity analysis [51] and the two-step epigenetic Mendelian randomization

(MR) analysis that used genetic variants as instrumental variables to assess the mediation role

of DNAm between cocaine use and HIV severity [52]. Our results provide new insights for the

role of DNAm on how cocaine affects HIV severity.

Methods

Study samples: VACS is a prospective cohort study of veterans designed to study substance

use and HIV-related outcomes with patient surveys, electronic medical records and

biospecimen data [53]. A baseline survey was conducted at enrollment [53]. The follow-up

survey of 5 visits occurred at approximately one-year intervals [53]. Blood samples were

collected in the middle of follow-up for a subset of participants in the cohort (VACS-BC) [54]. A

total of 1,435 HIV-positive participants from the VACS-BC were used to examine cocaine's

effect on mortality and HIV severity, and a subset of participants (n=875) with DNAm data

available were used for mediation analyses (**Figure 1**). Demographic and clinical information of

baseline samples and a subset of the samples at the time of blood collection are summarized in

Table 1.

Assessment of cocaine use: The timeline of cocaine use assessment for each analysis is

illustrated in **Figure 1**. Information on cocaine use status was self-reported through telephone

interviews for a total of 5 visits. We defined the "persistent cocaine use" group as self-reported

cocaine use across all 5 visits and the "no cocaine use" group as self-reported no cocaine use

across all 5 visits. This definition led to a subset of samples with 265 persistent cocaine users

and 202 nonusers for the mediation analyses to eliminate the inconsistent response across 5

visits and examine the effect of long-term cocaine exposure on DNAm and HIV severity.

The frequency of cocaine use was also assessed at baseline (**Figure 1**). Each participant was

asked "how often in the past year have you used cocaine or crack?", from which cocaine

frequency of use was coded as an ordinal variable as follows: 0=never tried; 1= no use in the

last year; 2=less than once a month; 3=1-3 times a month; 4=1-3 times a week; 5=4 or more

times a week.

Assessment of mortality and HIV severity: The timeline of HIV severity measurement and

survival information for each analysis is shown in **Figure 1**. Mortality and survival year

information were based on medical records. The VACS index was used as a measure of HIV

severity [55-58] and was obtained at each visit and at the time of blood collection (**Figure 1**).

The VACS index was calculated by summing preassigned points for age, routinely monitored

indicators of HIV disease (CD4 count and HIV-1 RNA) and other general indicators of organ

system injury [55]. A high VACS index corresponds to worsened HIV outcomes, and the VACS

index is positively associated with increased mortality [59]. The VACS index and DNAm profiling

were measured at the same time for the selection of candidate mediator CpGs, and the average

VACS index after blood collection was used for mediation analyses (**Figure 1**).

DNA methylation profiling and quality control: DNA samples were extracted from blood for a

subset of 875 HIV-positive participants (**Figure 1**). DNAm was profiled using two different

methylation arrays, with 475 samples profiled by the Infinium Human Methylation 450K

BeadChip (HM450K, Illumina Inc., CA, USA) and 400 samples later profiled by the Infinium

Human Methylation EPIC BeadChip (EPIC, Illumina Inc., CA, USA) [54]. DNA samples were

randomly selected for each methylation array regardless of cocaine use status or other clinical

demographic variables.

The quality control (QC) for samples measured by each array was conducted separately using

the same pipeline as previously described [60] by the R package *minfi* [61]. After QC, a total of

408,583 CpGs measured by both the HM450k and EPIC array remained for analysis. Six cell

type proportions (CD4+ T cells, CD8+ T cells, NK T cells, B cells, monocytes and granulocytes)

were estimated for each participant using the established method [62]. Negative control probes

were designed to capture background signals in Illumina arrays, and negative control principal

components (PCs) were extracted by *minfi* to control for background noise [61]. Batch effect

removal was conducted by *combat* after QC [63].

Genotyping and quality control: The 1,177 samples were genotyped using the

Illumina HumanOmniExpress Beadchip and imputed for 18,960,156 single nucleotide

polymorphisms (SNPs). *IMUPTE2* (ver 2.3.2) was used for imputation with the reference of

1000 genome phase 3 [64]. QC was conducted using *plink* (ver 1.90b21) [65]. SNPs and

samples with low call rate less than 0.05 were removed. The Hardy-Weinberg equilibrium test

cutoff was set to 1E-06. SNPs with minor allele frequency less than 0.01 were filtered.

Statistical analysis

Cocaine survival analysis among HIV-positive participants at baseline: Survival analysis

was conducted using baseline information among 1,435 HIV-positive participants with cocaine

use frequency (0-5) and other covariates (**Figure 1**). Kaplan-Meier analyses on 10-year follow-

up among HIV-positive and HIV-negative participants by cocaine use frequency (0-5) at

baseline were conducted, and the Kaplan-Meier curves were plotted by using the R package

survminer [66]. A test on ordered differences of Kaplan-Meier curves by cocaine use frequency

was conducted by *survminer* [66].

To adjust for confounding factors, a Cox proportional-hazards model was used to assess the

hazard ratio of baseline cocaine use frequency (0-5) on mortality during the follow-up using the

R package *survival* [67]. The following model was used to calculate the adjusted hazard ratio

among HIV-positive participants:

$$h(t) = h_0(t)\exp\left(\beta_1 cocaine\ use\ frequency\ +\ \beta_2\ sex\ +\ \beta_3\ baseline\ age\ +\ \beta_4\ race \right.$$
$$\left. +\ \beta_5 \log_{10}(viral\ load) + \beta_6\ CD4\ count\ +\ \beta_7\ antiviral\ medication\ adherence\right)$$

173   ***Association between cocaine use frequency and HIV severity among HIV-positive***

174   ***participants at baseline:*** This analysis was conducted using baseline information on cocaine

175   use frequency (0☐5), the VACS index and other covariates (**Figure 1**). The following linear

176   regression model was performed to test the association of cocaine use frequency and HIV

177   severity, adjusting for confounders as shown in the following model:

$$HIV\ disease\ severity$$
$$= \beta_1 cocaine\ use\ frequency\ +\ \beta_2\ sex\ +\ \beta_3\ age\ +\ \beta_4\ race$$
$$+\ \beta_5\ \log_{10}(viral\ load) + \beta_6\ \ CD4\ count\ +\ \beta_7\ antiviral\ medication\ adherence$$

Selection of candidate CpGs by epigenome-wide association (EWA) of persistent cocaine

use and HIV severity: To select candidate CpGs for mediation analysis, we conducted two

separate EWAs, one for persistent cocaine use and the other for HIV severity (**Figure 1**). Each

EWA model adjusted for sex, baseline age, race, smoking, self-reported antiviral medication

adherence, white blood cell count, estimated cell type proportions and negative control PCs. We

used the linear regression model with methylation as dependent variable for EWA as described

previously [33, 60, 68]. Since CD4+ T cell count is one component of the VACS index, to avoid

overrepresented CpGs associated with CD4+ T cells in the EWA results, we extracted the top

1,000 CD4+ T cell type relevant CpGs based on data from *FlowSorted.Blood.450k* [69]. The top

2 PCs that in total account for > 80% variation of the 1,000 CD4+ T cell CpGs were used as

covariates in the VACS index EWA model. CpGs with p<0.001 in both EWAs for persistent

cocaine use and HIV severity were selected as candidate CpGs for mediation analyses. A

liberal selection threshold was arbitrarily set to make sure there would be a sufficient number of

candidate CpGs for the mediation analysis. To limit confounding by use of other substances, we

tested the association of each candidate CpG site with alcohol use, cannabis use, and opioid

use based on self-reported data. Alcohol use was assessed by using 3 items of Alcohol Use

Diagnosis Identification Test- consumption (AUDIT-C). Cannabis and opioid uses were

assessed by asking the same questions as for cocaine use, described earlier.

Single-site mediation analysis and joint mediation analysis: The selected candidate CpGs

were assessed as potential mediators of the association between persistent cocaine use and

HIV severity among HIV-positive participants (n=467). We performed single-site mediation

analysis using the mediation method as previously described [51] and the R package *mediation*

[70]. Here, we used the average VACS index after DNAm profiling to ensure the temporality of

our mediation hypothesis that DNAm measurement preceded the HIV severity measurement. In

our mediation model, we adjusted for sex, age, race, smoking, self-reported antiviral medication

adherence, white blood cell count, and estimated cell type proportions as confounding factors.

We used M to represent the candidate CpGs (mediator), X to represent persistent cocaine use

status (exposure), Y to represent the average VACS index after blood collection (outcome), and

C_i to represent k confounding variables (sex, age, race, smoking, self-report antiviral medication

adherence, white blood cell count, estimated CD8 T cells, granulocytes, NK cells, B cells, and

monocytes). The mediator model $f(M|X, C)$ examined the association between persistent

cocaine use and CpGs:

$$f(M|X, C) = \beta_0\, X\; +\; \sum_{i=1}^{k} \beta_i\, C_i$$

213   The outcome model $f(Y|X, M, C)$ examined both the direct effect of persistent cocaine use on

214   VACS index and the mediation effect by CpG:

$$f(Y|X, M, C) = \alpha_0\, X\; +\; \alpha_1\, M + \sum_{i=1}^{k} \alpha_{i+1}\, C_i$$

Thus, the mediation effect, or the average causal mediation effect (ACME) of CpG M is $\alpha_1\beta_0$,

the total effect is $\alpha_0 + \alpha_1\beta_0$, and the proportion mediated is $\alpha_1\beta_0/(\alpha_0 + \alpha_1\beta_0)$. The confidence

interval and p-value were estimated by bootstrapping 1,000,000 iterations.

To assess the robustness of the results if the sequential ignorability assumption was violated,

we conducted a sensitivity analysis developed by Imai et al. [51] using the R package *mediation*

[70]. Sequential ignorability consists of two assumptions: (a) conditional on the covariates C_i,

the exposure X is independent of all potential values of the outcome Y and mediator M; and (b)

the observed mediator M is independent of all potential outcomes Y given the observed

exposure X and covariates C_i. The sensitivity parameter ρ was calculated on a grid of 0.05 and

he ρ at which ACME = 0 was calculated. For each mediator, sensitivity plots were illustrated to

show the estimated ACME and their 95% confidence interval as a function of ρ (**Figure S2**). If

the ρ at which ACME = 0 is close to 0, it indicates that the mediation analysis was sensitive to

violation of the sequential ignorability assumption.

The joint mediation analysis of all significant mediator CpGs were conducted as previously

described [71]. The mediator model $f\left(M_j \middle| X, C\right)$ for multiple mediators M_j $(M_1, M_2, \dots, M_n)$ is:

$$f\left(M_j \middle| X, C\right) = \beta_{0j}\, X \ + \ \sum_{i=1}^{k} \beta_{ij}\, C_i$$

232   The outcome model $f\left(Y \middle| X, M_1, \dots, M_n, C\right)$ is:

$$f\left(Y \middle| X, M_1, \dots, M_n, C\right) = \alpha_0\, X \ + \ \sum_{j=1}^{n} \alpha_j M_j + \sum_{i=1}^{k} \alpha_{i+n}\, C_i$$

The joint mediation effect of CpGs $M_1, \dots, M_j$ is $\sum_{j=1}^{n} \alpha_j \beta_{0j}$, the total effect is $\alpha_0 + \sum_{j=1}^{n} \alpha_j \beta_{0j}$, and

the proportion mediated is $\sum_{j=1}^{n} \alpha_j \beta_{0j} / (\alpha_0 + \sum_{j=1}^{n} \alpha_j \beta_{0j})$. The confidence interval and p-value

were estimated by bootstrapping 1,000,000 iterations.

Two-step epigenetic Mendelian randomization of cocaine and HIV severity: To evaluate

whether the results from the mediation analysis were influenced by reverse causation or

unmeasured confounding, we conducted a two-step epigenetic MR analysis [52] (n=1,177) on

cocaine use, candidate mediator CpGs, and HIV severity using the inverse-variance weighted

(IVW) method by R package *MendelianRandomization* [72].

In step 1, we conducted a two-sample MR on the effect of cocaine use on candidate CpGs

(n=1,177). Based on a recent meta-analysis of a cocaine dependence genome-wide association

study (GWAS) [73], 8 SNPs genotyped in our samples pruned at linkage disequilibrium (LD)

r^2<0.1 by the R package *LDlinkR* [74] were used as instrumental variables (p<1E-05) (**Table S4**).

We tested the associations between the 8 SNPs and the candidate CpGs, adjusting for age, sex,

race and 5 ancestry PCs using linear regression model in our sample (n=1,177). Based on

these summary statistics, we conducted MR using the IVW method to evaluate the effect of

cocaine use on candidate CpGs.

In step 2, we conducted a one-sample MR on the effect of candidate CpGs on HIV severity

(n=1,177). Here, *cis-* methylation quantitative trait loci (meQTLs) were used as instrumental

variables. *cis*-meQTLs were defined by the distance between a candidate CpG and a SNP

within 1 Mb. A linear regression analysis was performed to identify *cis*-meQTLs, adjusted for

age, sex, race and 5 ancestry PCs. For each candidate CpG, *cis*-meQTLs with p<0.01 after

pruning (LD r^2<0.1 using 1000 genome African ancestry samples as references [75]) were used

as instrumental variables in the MR analysis (**Table S4**) by the R package *LDlinkR* [74].

Association between each *cis*-meQTL and HIV severity was assessed by linear regression,

adjusting for age, sex and 5 ancestry PCs. Similar to the first step, we conducted an MR using

the IVW method to evaluate the effect of candidate CpGs on HIV severity.

Results

Cocaine use affects HIV severity and mortality among HIV-positive participants

We found that among HIV-positive participants, higher cocaine use frequency was associated

with increased mortality (p=0.008, **Figure 2a**). This difference was not found among HIV-

negative participants (p= 0.180, **Figure 2b**). Using Cox proportional-hazards model, this trend

remained significant with a hazard ratio (HR) of 1.10 (95% CI: 1.02-1.19, p=0.011), controlling

for sex, baseline age, race, viral load, CD4 count and antiviral medication adherence (**Table 2**).

A higher frequency of cocaine use at baseline was also significantly associated with a higher

VACS index (i.e., higher HIV severity, β=1.00, p= 0.00027) after adjusting for sex, age, race,

viral load, CD4 count and antiviral medication adherence (**Table 2**). To account for other drug

use, we further adjusted for baseline use of alcohol, cigarette smoking, cannabis, and opioids in

the model. Cocaine use frequency remains significantly associated with HIV severity after

adjusting for use of other substances (p=0.049). Our results suggest that cocaine use

accelerated HIV progression and increased mortality independent of antiviral medication

adherence, which is consistent with previous reports [10, 11, 13-15].

Selection of candidate DNAm sites for mediation analysis by EWA scan

The EWA scan of persistent cocaine use showed good control of inflation (λ=1.034, **Figure S1**).

A total of 497 CpGs met our candidate selection threshold (p<0.001). The top ranked CpG site,

cg22917487, was close to the epigenome-wide significance threshold with p-value of 1.69E-07.

This CpG site is located in *CX3CR1*, a gene that encodes a coreceptor for HIV-1 and leads to

rapid HIV progression (**Table S1**).

The EWA scan of the VACS index also showed good control of inflation (λ=1.116, **Figure S1**).

There were 876 CpGs that reached the candidate selection threshold (p<0.001) (**Table S2**). Of

note, 6 CpGs reached the epigenome-wide significance threshold (p<1.2E-07). These CpGs

were located near the genes involved in the viral and immune response (*PARP9, IFITM1,*

CD247, IFIT3, VASN, and *RUNX1*).

We selected candidate CpGs that were both associated with cocaine use and HIV severity

(p<0.001) by two separate EWA scans of 408,583 CpGs for mediation analysis. Fourteen CpGs

met both candidate selection thresholds. Additionally, cg22917487 in *CX3CR1* showed a strong

association with cocaine (p=1.69E-07) and a marginal association with the VACS index

(p=1.73E-03). Given its biological plausibility, this CpG was also included as a candidate

mediator for mediation analysis. Five of the top 10 VACS index EWA CpGs were selected as

candidate mediator CpGs (cg08122652, *PARP9*, p=2.30E-10; cg03038262, *IFITM1*, p=7.65E-

09; cg06188083, *IFIT3*, p=4.76E-08; cg08818207, *TAP1*, p=2.11E-07; cg26312951, *MX1*,

p=2.50E-07). Overall, a total of 15 CpGs were selected as candidates to assess their potential

mediation roles on the association between persistent cocaine use and HIV severity. Notably,

the DNAm from each of the 15 CpGs was not associated with cannabis, opioid or alcohol use

(p>0.05, **Table S3**).

Mediation analysis of candidate CpGs between persistent cocaine use and HIV severity

We examined the mediation role of DNAm between persistent cocaine use and HIV severity.

Twelve out of the 15 candidate CpGs showed significant mediation effects on the association

between persistent cocaine use and the VACS index, with p-values ranging from 1.00E-06 to

0.00307 (**Table 4**). These results remained significant after Bonferroni correction (p<0.0033).

Each CpG mediator explained between 11.3% to 29.5% of persistent cocaine use affecting HIV

severity. Notably, the direction of mediation effects among these 12 mediator CpGs were the

same. The average direct effects of cocaine on HIV severity were attenuated from 0.329 to

0.231-0.291 after adjusting for each mediator CpG. These 12 CpGs collectively mediated 47.2%

of the cocaine's effects on HIV severity by joint mediation analysis.

We also conducted a sensitivity analysis on these 15 candidate CpGs to assess the robustness

of our mediation analysis when the sequential ignorability assumption was violated [51]. The

absolute sensitivity parameters at which ACME=0 of the 12 significant mediator CpGs were

relatively higher (|ρ|≥0.15) than 3 nonsignificant CpGs (|ρ|≤0.10) (**Table 4**, **Figure S2**). Notably,

6 significant mediator CpGs had |ρ| of 0.30, indicating that these mediation effects were robust

even when the assumptions are slightly violated. The sensitivity analysis showed that our

mediation results were relatively stable.

Significant mediator CpGs are located near 11 viral and immune response genes: *MX1*, *PARP9*,

IFIT3, *IFITM1*, *NLRC5*, *EPSTI1*, *PLSCR1*, *TAP2*, *TAP1*, *CX3CR1* and *RIN2*. Five CpGs are

located on 5' gene regulatory regions, 4 CpGs on gene bodies, 2 CpGs on transcription start

sites and 1 CpG on 3' gene regulatory region. Notably, these 12 CpGs were mostly less

methylated in the persistent cocaine use group than in the no cocaine use group (**Table 3**,

Figure 3). **Figure 4** illustrates the mediation effect of cg26312951 (*MX1*), cg08122652 (*PARP9*),

cg07839457 (*NLRC5*) and cg22917487 (*CX3CR1*) on persistent cocaine use affecting HIV

severity.

Two-step epigenetic Mendelian randomization of cocaine and HIV severity

To validate our mediation results while eliminating unmeasured confounding and reverse

causation, we used the two-step epigenetic MR method [52] to test our mediation hypotheses

(n=1,177): whether cocaine use has a causal effect on candidate CpGs (step 1) and whether

candidate CpGs have causal effects on HIV severity (step 2).

In step 1, we conducted the MR analysis based on summary statistics of a meta-analysis of

GWAS on cocaine dependence [73]. The effect estimates of the association between 8 SNP

instrumental variables and 15 candidate CpG sites were obtained in our sample. Our MR

analysis showed that cocaine had significant MR estimates (p<0.05) on 4 CpGs (cg03753191,

EPSTI1; cg06188083,*IFIT3*; cg26312951,*MX1*; cg22917487,*CX3CR1*), as shown in **Table 5**.

Three of these CpGs were also among the top significant mediators in our previous mediation

analysis (**Table 4**).

In step 2, we conducted the MR analysis based on *cis*-meQTLs of the candidate CpGs and their

association with HIV severity in our sample. Seven CpGs showed significant MR estimates on

HIV severity (**Table 5**). Of note, 3 significant CpGs in the MR analysis in step 1 were also

significant in step 2.

Overall, 3 mediator CpGs discovered by the mediation analysis were validated as significant

mediators by two-step epigenetic MR analysis (cg03753191,*EPSTI1*; cg06188083,*IFIT3*;

cg26312951,*MX1*). Three CpGs without significant mediation effects in the mediation analysis

were also found to be nonsignificant in the two-step MR analysis (cg26396492,*RIN2*;

cg22385827,*C2orf67*; cg08623256).

Discussion

Our findings provide evidence that cocaine use worsens HIV severity and increases mortality

among HIV positive participants and that cocaine's adverse effects are partially mediated by

DNAm in the blood. We identified 12 CpGs that collectively accounted for a total of 47.2% of

cocaine affecting HIV severity. Three of the 12 mediator CpGs were further validated by a two-

step epigenetic MR approach, which provides supporting evidence that our mediation results

were not affected by unmeasured confounders or reverse causation. The sensitivity analysis

showed that our mediation analyses are relatively robust to slight violation of assumptions.

These 12 mediator CpGs offer new insights into the mechanisms of how cocaine use may affect

HIV outcomes by DNAm.

Methodological considerations are important for examining the mediation effect of DNAm. It is

possible that our mediation analyses could be undermined by violation of model assumptions,

reverse causation and unmeasured confounding. To address these concerns, we performed the

sensitivity analysis and two-step epigenetic MR analysis to further evaluate the mediation

effects of the 12 CpG sites. The results from the sensitivity test showed that the 12 mediator

CpGs were robust when slight violation of the sequential ignorability assumption is present. The

two-step epigenetic MR analysis confirmed 3 of 12 CpG sites as mediators of cocaine affecting

HIV severity and were not affected by reverse causation and unmeasured confounding. Of note,

the 8 SNPs used in the MR analysis showed marginal association with cocaine use, which

limited their utility as instrumental variables and may explain why 9 CpG sites did not show

significant mediation effects in two-step epigenetic MR analysis. In addition, in the mediation

analysis, the HIV severity was measured after the blood collection for DNAm profiling to assure

that the measurement of mediator precedes the measurement of outcome. Our study design

intended to match the temporality of exposure, mediator and outcome and to avoid reverse

causation. Of note, we observed discrepancy on the direction of cocaine use effect on DNA

methylation between EWA scan and step 1 MR analysis. This may happen because EWA scan

assessed association while MR evaluated the causal effect by removing reverse causality. This

difference might also be due to different ways on adjusting for confounding factors in two

models. Additionally, to assess whether cocaine use influenced cell type proportions as

reflected by DNA methylation, we conducted a MR on cocaine affecting six cell type proportions

using the same SNP instruments as used in step 1 MR. We found no significant MR estimates

across six cell types ($p > 0.1$) (**supplementary table S5**), suggesting that cocaine use does not

directly affect cell type proportions in our sample. Overall, we took various measures to make

sure our mediation results are valid and robust.

We observed that the sum of individual mediation proportion for 12 mediator CpGs exceeded

100%. An alternative approach is to test the joint mediation effect of all mediators [71]. We

found that the 12 mediator CpGs jointly accounted for 47% of the total effect (effect size=0.329)

of cocaine use on HIV severity. This finding indicates that these mediators may affect one

another or that there is an interaction effect [71]. For example, several mediator CpG sites are

near genes on the response to cytokine pathway (*PARP9*, *PLSCR1*, *CX3CR1*, *IFITM1*, *IFIT3*,

MX1 and *NLRC5*). It is possible that these CpGs may share the common biological pathway on

mediating the effect of cocaine on HIV severity.

These 12 CpGs are located in or near 11 biologically meaningful genes that were previously

reported to be involved in inflammation, HIV-1 viral replication and other pathways that play

critical roles in HIV progression. Specifically, cg06188083 on *IFIT3* mediated 28.8% of the

variation, and *IFIT3* encodes an IFN-induced antiviral protein which acts as an inhibitor of viral

processes and viral replication [76]. Another significant mediator CpG site, cg06188083, is

located near interferon gene *IFITM1*. We previously reported the hypomethylation of

cg07839457 due to HIV infection, which is located in the promoter region of *NLRC5* [33]. This

CpG site was also a significant mediator between cocaine and HIV severity in this study.

NLRC5 plays an important role in the cytokine response and antiviral immunity through its

inhibition of NF-kappa-B activation and negative regulation of type I interferon signaling

pathways [77]. The converging evidence on cg07839457 (*NLRC5*) warrants further investigation

of its role in HIV infection and progression. Another interesting CpG site, cg22917487 on

CX3CR1, showed both strong association with persistent cocaine use and a significant

mediation effect of cocaine affecting HIV severity. *CX3CR1* is involved in leukocyte adhesion

and migration and was recently identified as an HIV-1 coreceptor [78]. Some studies also

showed that genetic variants on *CX3CR1* were associated with HIV susceptibility and rapid HIV

progression to AIDS [79]. cg25114611, located in the promoter region of *FKBP5*, is also

biologically plausible, given the implication for chronic cocaine administration upregulating

FKBP5 expression in rats [80].

Cocaine use commonly cooccurred with the use of other substances, and this may confound

cocaine's effects on HIV severity and the mediation effects of CpGs between cocaine use and

HIV outcomes. However, our results show that the association between cocaine use and HIV

severity remained significant after accounting for smoking, alcohol, cannabis and opioid use.

Additionally, our cocaine use EWA model adjusted for smoking as a covariate, and the selected

candidate CpGs were not associated with alcohol, marijuana and opioid use (**Table S3**).

There are several strengths of this study. First, instead of selecting candidate mediator CpGs

based on the literature or hypotheses, we applied an unbiased epigenome-wide screening to

select CpGs associated with both cocaine use and HIV severity. Second, to limit self-reporting

bias of cocaine use, we leveraged longitudinal data in defining persistent cocaine use and no

cocaine use. We included only those participants who consistently reported cocaine use or no

cocaine use across all 5 visits for the selection of candidate CpGs and the mediation analyses.

Last, we used the average VACS index after blood collection so that DNAm measurements

(mediator) preceded HIV severity (outcome) for the mediation analyses.

One limitation of the study is that our sample size for the mediation analyses is small. However,

the strict definition of cocaine use helped reduce self-reporting bias and can potentially increase

power by comparing extreme groups. In addition, we used a less stringent criterion when

selecting candidate CpGs for mediation analysis due to limited sample size to achieve

epigenome-wide significance. To our knowledge, there are no sufficiently sized independent

cohorts for replication. Although this approach has also been adopted by previous studies [42,

81], using epigenome-wide significant CpG sites as candidate mediators may show stronger

signals in the future study with a larger sample size. Additionally, other unmeasured

confounding factors such as socioeconomic status may not be fully addressed in the mediation

model. Lastly, our samples consisted of mostly male veterans, which may limit the

generalizability of our findings.

Conclusions

We validated previous reports that the use of cocaine worsened HIV severity and increased the

risk of all-cause mortality among HIV positive participants. For the first time, this study found

that several biologically meaningful DNAm sites mediated the adverse effect of cocaine use on

HIV severity. These results merit future studies to further explore the biological mechanisms

revealed by these DNAm sites on how cocaine affects HIV disease outcomes.

Tables

Table 1: Sample characteristics in HIV-positive participants

	Baseline[*]	Follow-up (time of blood collection)[†]
Sample size	1435	875
Age [mean (sd)]	48.8 (8.1)	51.5 (7.7)
Sex		
Male	1399 (97.5%)	861 (98.4%)
Female	36 (2.5%)	14 (1.6%)
Race		
Caucasian	290 (20.2%)	84 (9.6%)
African American	964 (67.2%)	726 (83.0%)
Other	181 (12.6%)	65 (7.4%)
Persistent Cocaine use status		
Persistent cocaine use		265 (30.3%)
Non-cocaine use		202 (23.1%)
Cocaine use at baseline (%)		
have never tried	555 (38.7%)	
no use in the last year	540 (37.6%)	
less than once a month	122 (8.5%)	
1-3 times a month	103 (7.2%)	
1-3 times a week	39 (2.7%)	
≥4 times a week	76 (5.3%)	
Cigarette smoking	762 (53.1%)	521 (59.5%)
VACS index[‡]	30.0 (19.1)	
Average VACS index after blood collection[mean(sd)]		39.20 (22.4)
CD4+ count [mean (sd)]	418.7 (273.0)	440.02 (289.1)
log10 viral load [mean (sd)]	3.1 (1.2)	2.74 (1.2)
Antiviral medication adherence (%)	1117 (77.8%)	666 (76.1%)

[*]Samples at baseline for the survival analysis

[†]A subset of samples at the time of blood collection with DNA methylation measurements

[‡]VACS index: Veteran Aging Cohort Study index

Table 2: Association between cocaine use frequency and HIV severity and survival analysis among HIV-positive participants (n=1,435)

	HIV severity[*]				Mortality[†]			
	Estimate	SE	T	P value	Hazard ratio	95% CI	Z	P value
Cocaine use frequency[‡]	1.00	0.28	3.65	2.70E-04	1.10	(1.02,1.19)	2.54	1.10E-02
Sex (reference: male)	5.78	2.33	2.48	1.34E-02	0.27	(0.07,1.08)	-1.86	6.33E-02
Age[‡]	1.09	0.05	23.58	< 2E-16	1.05	(1.04,1.07)	7.35	1.90E-13
Race (reference: Caucasian)								
African American	7.30	0.94	7.73	2.00E-14	1.05	(0.80,1.39)	0.37	7.15E-01
Other	5.33	1.31	4.08	4.80E-05	0.71	(0.47,1.09)	-1.57	1.17E-01
log 10 viral load[‡]	4.72	0.36	13.11	< 2eE-16	1.07	2	1.36	1.73E-01
CD4 count[‡]	-0.03	0.00	-19.05	< 2E-16	1.00	(1.00,1.00)	-3.74	1.80E-04
Antiviral medication adherence[‡]	2.59	0.99	2.62	8.87E-03	1.02	(0.76,1.37)	0.13	8.95E-01

[*]Summary statistics from linear regression model

[†]Summary statistics from Cox proportional hazards model

[‡]Measured at baseline

Table 3: The selected candidate CpG sites by epigenome-wide association (EWA) scan on persistent cocaine use (n=467) and HIV severity (n=875)

CpG	Chr	Position	Nearest gene	Cocaine EWA effect size	Cocaine EWA p-value	VACS index[*] EWA effect size	VACS index[*] EWA p-value	Reference Gene Group	Relations to CpG Islands
cg22917487	3	39322103	CX3CR1	1.71E-02	1.69E-07	1.96E-04	1.73E-03	Body;TSS200;5 UTR;TSS1500	
cg07839457	16	57023022	NLRC5	-5.18E-02	3.16E-05	-7.08E-04	4.02E-04	TSS1500	N_Shore
cg26312951	21	42797847	MX1	-3.93E-02	3.24E-05	-7.80E-04	2.50E-07	TSS200;5UTR	N_Shore
cg06188083	10	91093005	IFIT3	-5.06E-02	3.34E-05	-9.96E-04	4.76E-08	Body	
cg08122652	3	1.22E+08	PARP9	-6.13E-02	1.10E-04	-1.26E-03	2.30E-10	5UTR;TSS1500	N_Shore
cg22385827	2	2.11E+08	C2orf67	8.10E-03	2.01E-04	1.20E-04	7.52E-04	TSS1500	Island
cg22930808	3	1.22E+08	PARP9	-6.52E-02	2.44E-04	-1.17E-03	1.52E-06	5UTR;TSS1500	N_Shore
cg06981309	3	1.46E+08	PLSCR1	-4.29E-02	2.63E-04	-5.72E-04	4.21E-04	5UTR	N_Shore
cg03753191	13	43566902	EPSTI1	-1.36E-02	2.81E-04	-2.44E-04	5.50E-05	TSS1500	S_Shore
cg00096307	2	1.07E+08	UXS1	-1.12E-02	3.59E-04	1.74E-04	5.73E-04	Body	
cg26396492	20	19915762	RIN2	-1.26E-02	4.76E-04	-2.00E-04	9.15E-04	Body	
cg22940798	6	32805554	TAP2	-1.93E-02	4.82E-04	-2.89E-04	4.72E-04	Body	N_Shore
cg08623256	5	3858275		-1.73E-02	6.96E-04	3.17E-04	7.18E-04		S_Shelf
cg08818207	6	32820355	TAP1	-2.67E-02	8.18E-04	-6.18E-04	2.11E-07	Body	N_Shore
cg03038262	11	315262	IFITM1	-2.98E-02	9.59E-04	-6.46E-04	7.65E-09	3UTR	N_Shore

[*]VACS index: Veteran Aging Cohort Study index

462
463

464   **Table 4: Mediation analyses on candidate CpGs between cocaine use and VACS index[*] (n=467)**

CpG	Chr	Position	Nearest gene	Reference Gene Group	Relations to CpG Islands	Average Causal Mediation Effect			Proportion Mediated			Average Direct Effect	Total Effect	Sensitivity analysis: Rho at which ACME = 0
						Effect estimate	95% CI	p value	Proportion Mediated	95% CI	p value			
cg26312951	21	42797847	MX1	TSS200;5UTR	N_Shore	0.097	(0.051,0.153)	1.00E-06	29.5%	(15.9%,53.0%)	1.00E-06	0.231	0.329	-0.30
cg08122652	3	122281939	PARP9	5UTR;TSS1500	N_Shore	0.092	(0.049,0.142)	1.00E-06	28.2%	(15.7%,49.2%)	1.00E-06	0.236	0.329	-0.30
cg06188083	10	91093005	IFIT3	Body		0.095	(0.052,0.145)	2.00E-06	28.8%	(16.2%,50.7%)	2.00E-06	0.234	0.329	-0.30
cg03038262	11	315262	IFITM1	3UTR	N_Shore	0.078	(0.040,0.123)	4.00E-06	23.7%	(12.4%,42.8%)	4.00E-06	0.251	0.329	-0.30
cg22930808	3	122281881	PARP9	5UTR;TSS1500	N_Shore	0.091	(0.047,0.141)	8.00E-06	27.7%	(15.1%,48.9%)	8.00E-06	0.237	0.329	-0.30
cg07839457	16	57023022	NLRC5	TSS1500	N_Shore	0.079	(0.038,0.127)	8.00E-06	24.0%	(11.0%,47.5%)	8.00E-06	0.250	0.329	-0.25
cg03753191	13	43566902	EPSTI1	TSS1500	S_Shore	0.078	(0.036,0.128)	3.90E-05	23.7%	(10.6%,46.8%)	3.90E-05	0.251	0.329	-0.25
cg06981309	3	146260954	PLSCR1	5UTR	N_Shore	0.074	(0.033,0.123)	4.70E-05	22.6%	(10.7%,41.8%)	4.70E-05	0.254	0.329	-0.25
cg22940798	6	32805554	TAP2	Body	N_Shore	0.058	(0.025,0.100)	5.70E-05	17.6%	(7.6%,35.6%)	5.70E-05	0.271	0.329	-0.20
cg08818207	6	32820355	TAP1	Body	N_Shore	0.081	(0.037,0.130)	5.80E-05	24.5%	(11.9%,44.7%)	5.90E-05	0.248	0.329	-0.30
cg22917487	3	39322103	CX3CR1	Body;TSS200;5UTR;TSS1500		0.054	(0.015,0.104)	1.81E-03	16.3%	(4.6%,34.4%)	1.81E-03	0.276	0.329	0.15
cg26396492	20	19915762	RIN2	Body		0.037	(0.009,0.073)	3.07E-03	11.3%	(2.7%,24.9%)	3.07E-03	0.291	0.329	-0.15
cg08623256	5	3858275			S_Shelf	0.009	(-0.012,0.032)	2.00E-01	0.026	(-0.038,0.108)	2.00E-01	0.321	0.329	-0.05
cg22385827	2	211036478	C2orf67	TSS1500	Island	-0.008	(-0.027,0.007)	8.28E-01	-0.023	(-0.089,0.023)	8.28E-01	0.336	0.329	-0.05
cg00096307	2	106761754	UXS1	Body		-0.027	(-0.064,0.000)	9.75E-01	-0.084	(-0.221,0.000)	9.75E-01	0.356	0.329	0.10

465   [*]VACS index: Veteran Aging Cohort Study index

Table 5: Two-step epigenetic mendelian randomization on cocaine and HIV severity (n=1,177)

CpG	Chr	Position	Nearest gene	Number of IV*	Step 1: cocaine affecting Methylation				Number of IV*	Step 2: methylation affecting HIV severity			
					Estimate	Standard Error	95% CI	p value†		Estimate	Standard Error	95% CI	p value†
cg03753191	13	43566902	*EPSTI1*	8	0.206	0.070	(0.069,0.343)	**3.27E-03**	9	-10.688	2.300	(-15.197,-6.180)	**3.38E-06**
cg06188083	10	91093005	*IFIT3*	8	0.269	0.109	(0.056,0.482)	**1.33E-02**	4	-6.756	2.602	(-11.856,-1.656)	**9.42E-03**
cg26312951	21	42797847	*MX1*	8	0.286	0.137	(0.018,0.554)	**3.68E-02**	12	-4.582	1.261	(-7.053,-2.111)	**2.79E-04**
cg22917487	3	39322103	*CX3CR1*	8	0.090	0.045	(0.001,0.179)	**4.83E-02**	7	-3.522	5.577	(-14.453,7.410)	5.28E-01
cg08122652	3	122281939	*PARP9*	8	0.259	0.141	(-0.017,0.536)	6.61E-02	8	-4.517	1.558	(-7.571,-1.463)	**3.75E-03**
cg22930808	3	122281881	*PARP9*	8	0.245	0.164	(-0.076,0.567)	1.35E-01	10	-5.245	1.135	(-7.470,-3.019)	**3.85E-06**
cg08818207	6	32820355	*TAP1*	8	0.092	0.071	(-0.048,0.231)	1.96E-01	2	-10.287	5.519	(-21.104,0.530)	6.23E-02
cg03038262	11	315262	*IFITM1*	8	0.099	0.081	(-0.059,0.257)	2.20E-01	4	-9.610	3.955	(-17.361,-1.859)	**1.51E-02**
cg22940798	6	32805554	TAP2	8	0.047	0.042	(-0.035,0.128)	2.62E-01	2	-28.774	9.027	(-46.466,-11.082)	**1.43E-03**
cg06981309	3	146260954	*PLSCR1*	8	0.107	0.097	(-0.083,0.296)	2.69E-01	4	1.502	2.983	(-4.344,7.349)	6.15E-01
cg00096307	2	106761754	*UXS1*	8	0.036	0.033	(-0.030,0.101)	2.84E-01	2	3.455	11.853	(-19.776,26.685)	7.71E-01
cg07839457	16	57023022	*NLRC5*	8	0.109	0.158	(-0.200,0.418)	4.88E-01	7	-2.961	1.552	(-6.004,0.081)	5.64E-02
cg26396492	20	19915762	*RIN2*	8	0.014	0.031	(-0.047,0.075)	6.50E-01	10	-9.992	5.226	(-20.234,0.250)	5.59E-02
cg08623256	5	3858275		8	0.009	0.029	(-0.048,0.066)	7.58E-01	1	18.246	15.315	(-11.772,48.263)	2.34E-01
cg22385827	2	211036478	*C2orf67*	8	0.025	0.125	(-0.219,0.269)	8.41E-01	3	1.054	1.855	(-2.582,4.689)	5.70E-01

*IV: instrumental variable

†p<0.05 are bolded

Figure Legends

Figure 1. Timeline of data and blood sample collection for each analysis

Figure 2. Kaplan-Meier curves of cocaine use frequency at baseline among HIV-positive

(n=1,435, **2a**) and HIV-negative (n=795, **2b**) participants. The higher frequency of cocaine use

is associated with lower survival probability among HIV-positive participants but not among HIV-

negative participants.

Figure 3. DNA methylation level of the significant CpG mediators by persistent cocaine use

status

Figure 4. Significant mediation effect of cg26312951 (*MX1*), cg08122652(*PARP9*), cg07839457

(*NLRC5*), and cg22917487 (*CX3CR1*) between persistent cocaine use and HIV severity

(p<0.0033)

Figure S1. Manhattan and quantile-quantile (QQ) plot of persistent cocaine use epigenome-

wide association (EWA) (λ=1.034) and HIV severity EWA (λ=1.116)

Figure S2. The estimated ACME and their 95% confidence interval as a function of the

sensitivity parameter ρ among 15 candidate CpGs

485
List of abbreviations

ACME: average causal mediation effect

ART: antiretroviral therapy

AUDIT-C: Alcohol Use Diagnosis Identification Test- consumption

CI: Confidence interval

DNA: Deoxyribonucleic acid

DNAm: DNA methylation

EWA: epigenome-wide association

FDR: False discovery rate

GWAS: Genome-wide association study

GO: Gene ontology

HIV: Human immunodeficiency virus

HM450K: Infinium Human Methylation 450K BeadChip

EPIC: Infinium Human Methylation EPIC BeadChip

HR: Hazard ratio

IL: interleukins

IVW: Inverse-variance weighted

LD: Linkage disequilibrium

meQTL: Methylation quantitative trait loci

MR: Mendelian randomization

NK: Natural killer

PC: Principal component

QC: Quality control

VACS: Veterans Aging Cohort Study

VACS-BC: Veteran Aging Cohort Study Biomarker Cohort

VACS index: Veterans Aging Cohort Study index

Declarations

Ethics approval and consent to participate

The study was approved by the committee of the Human Research Subject Protection at Yale

University and the Institutional Research Board Committee of the Connecticut Veteran

Healthcare System. All participants provided written consent.

Availability of data and materials

Demographic and clinical variables and DNAm data for the VACS samples were submitted to

the GEO dataset (GSE117861) and are available to the public. All codes for analysis are also

available upon a request to the corresponding author.

Competing interest

All authors declare that they have no conflict of interest.

Funding

The work was supported by the National Institute on Drug Abuse (R03DA039745,

R01DA038632, R01DA047063, R01DA047820).

Authors' contributions

CS was responsible for data analysis and manuscript preparation. ACJ provided DNA samples,

clinical data, and contributed to manuscript preparation. XZ was responsible for the

bioinformatics data processing. ZW contributed to mediation analysis and manuscript

preparation. DH and EJ contributed to the analytical approach and the manuscript preparation.

KX was responsible for the study design, study protocol, sample preparation, data analysis,

interpretation of findings, and manuscript preparation.

Acknowledgements The authors appreciate the support of the Veteran Aging Study Cohort

Biomarker Core and Yale Center of Genomic Analysis.

Reference

1. Daskalopoulou M, Rodger A, Phillips AN et al. Recreational drug use, polydrug use, and sexual behaviour in HIV-diagnosed men who have sex with men in the UK: results from the cross-sectional ASTRA study. *Lancet HIV* 1(1), e22-31 (2014).

2. Cofrancesco Jr J, Scherzer R, Tien PC et al. Illicit drug use and HIV treatment outcomes in a US cohort. *AIDS (London, England)* 22(3), 357 (2008).

3. Hatfield LA, Horvath KJ, Jacoby SM, Simon Rosser BR. Comparison of substance use and risky sexual behavior among a diverse sample of urban, HIV-positive men who have sex with men. *J Addict Dis* 28(3), 208-218 (2009).

4. Grabovac I, Meilinger M, Schalk H, Leichsenring B, Dorner TE. Prevalence and Associations of Illicit Drug and Polydrug Use in People Living with HIV in Vienna. *Scientific Reports* 8(1), 8046 (2018).

5. Wolitski RJ, Parsons JT, Gómez CA. Prevention with HIV-seropositive men who have sex with men: lessons from the Seropositive Urban Men's Study (SUMS) and the Seropositive Urban Men's Intervention Trial (SUMIT). *J Acquir Immune Defic Syndr* 37 Suppl 2 S101-109 (2004).

6. Purcell DW, Moss S, Remien RH, Woods WJ, Parsons JT. Illicit substance use, sexual risk, and HIV-positive gay and bisexual men: differences by serostatus of casual partners. *Aids* 19 Suppl 1 S37-47 (2005).

7. Key substance use and mental health indicators in the United States: Results from the 2018 National Survey on Drug Use and Health (HHS Publication No. PEP19-5068, NSDUH Series H-54). Rockville, MD: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration. Retrieved from https://www.samhsa.gov/data/

8. Webber MP, Schoenbaum EE, Gourevitch MN, Buono D, Klein RS. A prospective study of HIV disease progression in female and male drug users. *Aids* 13(2), 257-262 (1999).

9. Vittinghoff E, Hessol NA, Bacchetti P, Fusaro RE, Holmberg SD, Buchbinder SP. Cofactors for HIV disease progression in a cohort of homosexual and bisexual men. *Journal of acquired immune deficiency syndromes (1999)* 27(3), 308-314 (2001).

10. Cook JA, Burke-Miller JK, Cohen MH et al. Crack cocaine, disease progression, and mortality in a multi-center cohort of HIV-1 positive women. *AIDS (London, England)* 22(11), 1355 (2008).

11. Baum MK, Rafie C, Lai S, Sales S, Page B, Campa A. Crack-cocaine use accelerates HIV disease progression in a cohort of HIV-positive drug users. *JAIDS Journal of Acquired Immune Deficiency Syndromes* 50(1), 93-99 (2009).

12. Sharpe TT, Lee LM, Nakashima AK, Elam-Evans LD, Fleming PL. Crack cocaine use and adherence to antiretroviral treatment among HIV-infected black women. *J Community Health* 29(2), 117-127 (2004).

13. Carrico AW, Johnson MO, Morin SF et al. Stimulant use is associated with immune activation and depleted tryptophan among HIV-positive persons on anti-retroviral therapy. *Brain, behavior, and immunity* 22(8), 1257-1262 (2008).

579   14.   Dash S, Balasubramaniam M, Villalta F, Dash C, Pandhare J. Impact of cocaine abuse on
580   HIV pathogenesis. *Front Microbiol* 6 1111-1111 (2015).

581   15.   Rasbach DA, Desruisseau AJ, Kipp AM *et al*. Active cocaine use is associated with lack of
582   HIV-1 virologic suppression independent of nonadherence to antiretroviral therapy: use
583   of a rapid screening tool during routine clinic visits. *AIDS care* 25(1), 109-117 (2013).

584   16.   Breitling LP, Yang R, Korn B, Burwinkel B, Brenner H. Tobacco-smoking-related
585   differential DNA methylation: 27K discovery and replication. *The American Journal of
586   Human Genetics* 88(4), 450-457 (2011).

587   17.   Joubert BR, Håberg SE, Nilsen RM *et al*. 450K epigenome-wide scan identifies differential
588   DNA methylation in newborns related to maternal smoking during pregnancy.
589   *Environmental health perspectives* 120(10), 1425-1431 (2012).

590   18.   Lee KW, Pausova Z. Cigarette smoking and DNA methylation. *Frontiers in genetics* 4 132
591   (2013).

592   19.   Tsaprouni LG, Yang T-P, Bell J *et al*. Cigarette smoking reduces DNA methylation levels at
593   multiple genomic loci but the effect is partially reversible upon cessation. *Epigenetics*
594   9(10), 1382-1396 (2014).

595   20.   Gao X, Zhang Y, Saum K-U, Schöttker B, Breitling LP, Brenner H. Tobacco smoking and
596   smoking-related DNA methylation are associated with the development of frailty among
597   older adults. *Epigenetics* 12(2), 149-156 (2017).

598   21.   Zhang X, Hu Y, Aouizerat BE *et al*. Machine learning selected smoking-associated DNA
599   methylation signatures that predict HIV prognosis and mortality. 10(1), 155 (2018).

600   22.   Zhang R, Miao Q, Wang C *et al*. Genome-wide DNA methylation analysis in alcohol
601   dependence. *Addiction biology* 18(2), 392-403 (2013).

602   23.   Zhang H, Gelernter J. DNA methylation and alcohol use disorders: Progress and
603   challenges. *The American journal on addictions* 26(5), 502-515 (2017).

604   24.   Sharp GC, Arathimos R, Reese SE *et al*. Maternal alcohol consumption and offspring DNA
605   methylation: findings from six general population-based birth cohorts. *Epigenomics*
606   10(1), 27-42 (2018).

607   25.   Liu C, Marioni RE, Hedman ÅK *et al*. A DNA methylation biomarker of alcohol
608   consumption. *Molecular psychiatry* 23(2), 422 (2018).

609   26.   Joehanes R, Just AC, Marioni RE *et al*. Epigenetic Signatures of Cigarette Smoking. *Circ
610   Cardiovasc Genet* 9(5), 436-447 (2016).

611   27.   Feinberg AP, Koldobskiy MA, Göndör A. Epigenetic modulators, modifiers and mediators
612   in cancer aetiology and progression. *Nature Reviews Genetics* 17 284 (2016).

613   28.   Hao X, Luo H, Krawczyk M *et al*. DNA methylation markers for diagnosis and prognosis of
614   common cancers. *Proceedings of the National Academy of Sciences* 114(28), 7414 (2017).

615   29.   Michalak EM, Burr ML, Bannister AJ, Dawson MA. The roles of DNA, RNA and histone
616   methylation in ageing and cancer. *Nature Reviews Molecular Cell Biology*
617   doi:10.1038/s41580-019-0143-1 (2019).

618   30.   Davegårdh C, García-Calzón S, Bacos K, Ling C. DNA methylation in the pathogenesis of
619   type 2 diabetes in humans. *Mol Metab* 14 12-25 (2018).

620   31.   Zhong J, Agha G, Baccarelli AA. The role of DNA methylation in cardiovascular risk and
621   disease: methodological aspects, study design, and data analysis for epidemiological
622   studies. *Circulation research* 118(1), 119-131 (2016).

32.   Nakatochi M, Ichihara S, Yamamoto K *et al*. Epigenome-wide association of myocardial
infarction with DNA methylation sites at loci related to cardiovascular disease. *Clinical*
epigenetics 9(1), 54 (2017).
33.   Zhang X, Justice AC, Hu Y *et al*. Epigenome-wide differential DNA methylation between
HIV-infected and uninfected individuals. *Epigenetics* 11(10), 750-760 (2016).
34.   Liu Y, Aryee MJ, Padyukov L *et al*. Epigenome-wide association data implicate DNA
methylation as an intermediary of genetic risk in rheumatoid arthritis. *Nature*
biotechnology 31(2), 142 (2013).
35.   Bind M-A, Lepeule J, Zanobetti A *et al*. Air pollution and gene-specific methylation in the
Normative Aging Study: association, effect modification, and mediation analysis.
Epigenetics 9(3), 448-458 (2014).
36.   Cao-Lei L, Dancause KN, Elgbeili G *et al*. DNA methylation mediates the impact of
exposure to prenatal maternal stress on BMI and central adiposity in children at age
13½ years: Project Ice Storm. *Epigenetics* 10(8), 749-761 (2015).
37.   Timms JA, Relton CL, Rankin J, Strathdee G, Mckay JA. DNA methylation as a potential
mediator of environmental risks in the development of childhood acute lymphoblastic
leukemia. *Epigenomics* 8(4), 519-536 (2016).
38.   Tobi EW, Slieker RC, Luijk R *et al*. DNA methylation as a mediator of the association
between prenatal adversity and risk factors for metabolic disease in adulthood. *Science*
advances 4(1), eaao4364 (2018).
39.   Barker ED, Walton E, Cecil CA. Annual Research Review: DNA methylation as a mediator
in the association between risk exposure and child and adolescent psychopathology.
Journal of Child Psychology and Psychiatry 59(4), 303-322 (2018).
40.   Rutten BPF, Mill J. Epigenetic Mediation of Environmental Influences in Major Psychotic
Disorders. *Schizophrenia Bulletin* 35(6), 1045-1056 (2009).
41.   Ladd-Acosta C, Fallin MD. The role of epigenetics in genetic and environmental
epidemiology. *Epigenomics* 8(2), 271-283 (2016).
42.   Jordahl KM, Phipps AI, Randolph TW *et al*. Differential DNA methylation in blood as a
mediator of the association between cigarette smoking and bladder cancer risk among
postmenopausal women. *Epigenetics* doi:10.1080/15592294.2019.1631112 1-9 (2019).
43.   Shirazi J, Shah S, Sagar D *et al*. Epigenetics, drugs of abuse, and the retroviral promoter. *J*
Neuroimmune Pharmacol 8(5), 1181-1196 (2013).
44.   Dhillon NK, Williams R, Peng F *et al*. Cocaine-mediated enhancement of virus replication
in macrophages: implications for human immunodeficiency virus-associated dementia. *J*
Neurovirol 13(6), 483-495 (2007).
45.   Atluri VS, Pilakka-Kanthikeel S, Garcia G *et al*. Effect of Cocaine on HIV Infection and
Inflammasome Gene Expression Profile in HIV Infected Macrophages. *Sci Rep* 6 27864
(2016).
46.   Castro FOF, Silva JM, Dorneles GP *et al*. Distinct inflammatory profiles in HIV-infected
individuals under ART using cannabis, cocaine or cannabis plus cocaine. *AIDS*
doi:10.1097/QAD.0000000000002296 (2019).
47.   Parikh N, Dampier W, Feng R *et al*. Cocaine alters cytokine profiles in HIV-1-infected
African American individuals in the DrexelMed HIV/AIDS genetic analysis cohort. *J Acquir*
Immune Defic Syndr 66(3), 256-264 (2014).

48.   Robison AJ, Nestler EJ. Transcriptional and epigenetic mechanisms of addiction. *Nat Rev*
668   *Neurosci* 12(11), 623-637 (2011).
669   49.   Anier K, Malinovskaja K, Aonurm-Helm A, Zharkovsky A, Kalda A. DNA methylation
670   regulates cocaine-induced behavioral sensitization in mice. *Neuropsychopharmacology*
671   35(12), 2450-2461 (2010).
672   50.   Camilo C, Maschietto M, Vieira HC *et al*. Genome-wide DNA methylation profile in the
673   peripheral blood of cocaine and crack dependents. *Braz J Psychiatry* 41(6), 485-493
674   (2019).
675   51.   Imai K, Keele L, Tingley D. A general approach to causal mediation analysis.
676   *Psychological methods* 15(4), 309 (2010).
677   52.   Relton CL, Davey Smith G. Two-step epigenetic Mendelian randomization: a strategy for
678   establishing the causal role of epigenetic processes in pathways to disease. *Int J*
679   *Epidemiol* 41(1), 161-176 (2012).
680   53.   Justice AC, Dombrowski E, Conigliaro J *et al*. Veterans aging cohort study (VACS):
681   overview and description. 44(8 Suppl 2), S13 (2006).
682   54.   Veterans Aging Cohort Study. VACS Biomarker Cohort Description. (2016).
683   55.   Tate JP, Justice AC, Hughes MD *et al*. An internationally generalizable risk index for
684   mortality after one year of antiretroviral therapy. *AIDS (London, England)* 27(4), 563-572
685   (2013).
686   56.   Justice AC, Modur SP, Tate JP *et al*. Predictive accuracy of the Veterans Aging Cohort
687   Study index for mortality with HIV infection: a North American cross cohort analysis.
688   *Journal of acquired immune deficiency syndromes (1999)* 62(2), 149-163 (2013).
689   57.   Bebu I, Tate J, Rimland D *et al*. The VACS Index Predicts Mortality in a Young, Healthy
690   HIV Population Starting Highly Active Antiretroviral Therapy. *J. Acquir. Immune. Defic.*
691   *Syndr.* 65(2), 226-230 (2014).
692   58.   Brown ST, Tate JP, Kyriakides TC *et al*. The VACS index accurately predicts mortality and
693   treatment response among multi-drug resistant HIV infected patients participating in
694   the options in management with antiretrovirals (OPTIMA) study. *PLoS One* 9(3), e92606
695   (2014).
696   59.   Justice AC, Modur S, Tate JP *et al*. Predictive accuracy of the Veterans Aging Cohort
697   Study (VACS) index for mortality with HIV infection: a north American cross cohort
698   analysis. *Journal of acquired immune deficiency syndromes (1999)* 62(2), 149 (2013).
699   60.   Lehne B, Drong AW, Loh M *et al*. A coherent approach for analysis of the Illumina
700   HumanMethylation450 BeadChip improves data quality and performance in epigenome-
701   wide association studies. 16(1), 37 (2015).
702   61.   Aryee MJ, Jaffe AE, Corrada-Bravo H *et al*. Minfi: a flexible and comprehensive
703   Bioconductor package for the analysis of Infinium DNA methylation microarrays. 30(10),
704   1363-1369 (2014).
705   62.   Houseman EA, Accomando WP, Koestler DC *et al*. DNA methylation arrays as surrogate
706   measures of cell mixture distribution. *BMC Bioinformatics* 13(1), 86 (2012).
707   63.   Johnson WE, Li C, Rabinovic A. Adjusting batch effects in microarray expression data
708   using empirical Bayes methods. *Biostatistics* 8(1), 118-127 (2007).

64.   Howie BN, Donnelly P, Marchini J. A flexible and accurate genotype imputation method
for the next generation of genome-wide association studies. *PLoS Genet* 5(6), e1000529
(2009).
65.   Purcell S, Neale B, Todd-Brown K *et al*. PLINK: a tool set for whole-genome association
and population-based linkage analyses. *The American journal of human genetics* 81(3),
559-575 (2007).
66.   Kassambara A, Kosinski M, Biecek P. survminer: Drawing Survival Curves using'ggplot2'.
R package version 0.3 1 (2017).
67.   Therneau T. A Package for Survival Analysis in S. version 2.38. (2015).
68.   Zhang X, Hu Y, Justice AC *et al*. DNA methylation signatures of illicit drug injection and
hepatitis C are associated with HIV frailty. *Nature Communications* 8(1), 2243 (2017).
69.   Jaffe A. Flowsorted. blood. 450k: illumina humanmethylation data on sorted blood cell
populations. *R Package Version* 1(0), (2015).
70.   Tingley D, Yamamoto T, Hirose K, Keele L, Imai K. Mediation: R package for causal
mediation analysis. (2014).
71.   Vanderweele T, Vansteelandt S. Mediation analysis with multiple mediators.
Epidemiologic methods 2(1), 95-115 (2014).
72.   Yavorska OO, Burgess S. MendelianRandomization: an R package for performing
Mendelian randomization analyses using summarized data. *International Journal of*
Epidemiology 46(6), 1734-1739 (2017).
73.   Cabana-Domínguez J, Shivalikanjli A, Fernàndez-Castillo N, Cormand B. Genome-wide
association meta-analysis of cocaine dependence: Shared genetics with comorbid
conditions. *Progress in Neuro-Psychopharmacology and Biological Psychiatry* 94 109667
(2019).
74.   Myers TA, Chanock SJ, Machiela MJ. LDlinkR: An R Package for Rapidly Calculating
Linkage Disequilibrium Statistics in Diverse Populations. *Frontiers in Genetics* 11 157
(2020).
75.   Auton A, Abecasis GR, Altshuler DM *et al*. A global reference for human genetic variation.
Nature 526(7571), 68-74 (2015).
76.   Schmeisser H, Mejido J, Balinsky CA *et al*. Identification of alpha interferon-induced
genes associated with antiviral activity in Daudi cells and characterization of IFIT3 as a
novel antiviral gene. *J Virol* 84(20), 10671-10680 (2010).
77.   Cui J, Zhu L, Xia X *et al*. NLRC5 negatively regulates the NF-kappaB and type I interferon
signaling pathways. *Cell* 141(3), 483-496 (2010).
78.   Garin A, Tarantino N, Faure S *et al*. Two novel fully functional isoforms of CX3CR1 are
potent HIV coreceptors. *J Immunol* 171(10), 5305-5312 (2003).
79.   Faure S, Meyer L, Costagliola D *et al*. Rapid progression to AIDS in HIV+ individuals with a
structural variant of the chemokine receptor CX3CR1. *Science* 287(5461), 2274-2277
(2000).
80.   Connelly KL, Unterwald EM. Chronic cocaine administration upregulates FKBP5 in the
extended amygdala of male and female rats. *Drug Alcohol Depend* 199 101-105 (2019).
81.   Chu SH, Loucks EB, Kelsey KT *et al*. Sex-specific epigenetic mediators between early life
social disadvantage and adulthood BMI. *Epigenomics* 10(6), 707-722 (2018).
752

a. HIV-Positive

b. HIV-Negative