

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35

TITLE

Patterns of COVID-19 related excess mortality in the municipalities of Northern Italy

AUTHORS

Dino Gibertoni, Kadjo Yves Cedric Adja, Davide Golinelli, Chiara Reno, Luca Regazzi, Maria Pia Fantini

AUTHORS' AFFILIATION

Dino Gibertoni, PhD, Department of Biomedical and Neuromotor Sciences, Alma Mater Studiorum, University of Bologna, Italy.

Kadjo Yves Cedric Adja, MD, MSc, Department of Biomedical and Neuromotor Sciences, Alma Mater Studiorum, University of Bologna, Italy.

Davide Golinelli, MD, MSc, MPH, Research Fellow in Public Health and Epidemiology at the Department of Biomedical and Neuromotor Sciences, Alma Mater Studiorum, University of Bologna, Italy.

Chiara Reno, MD, MSc, Department of Biomedical and Neuromotor Sciences, Alma Mater Studiorum, University of Bologna, Italy.

Luca Regazzi, Alma Mater Studiorum, Medical University of Bologna, Italy.

Maria Pia Fantini, MD, MSc Department of Biomedical and Neuromotor Sciences, Alma Mater Studiorum, University of Bologna, Italy.

CORRESPONDING AUTHOR

Name: Kadjo Yves Cedric
Last name: Adja
Address: Department of Biomedical and Neuromotor Sciences, Alma Mater Studiorum, University of Bologna. Via San Giacomo 12, 40126, Bologna, Italy
E-mail: adjayvescedric@gmail.com

1 **DECLARATIONS**

2 **Conflicts of interest**

3 Declarations of interest: none.

4 **Availability of data and material**

5 Data are publicly available.

6 **Ethics approval**

7 Not applicable.

8

9 **Patterns of COVID-19 related excess mortality in the municipalities of**
10 **Northern Italy**

11

12 **Abstract**

13 The Coronavirus Disease 2019 (COVID-19) spatial distribution in Italy is inhomogeneous, because
14 of its ways of spreading from the initial hotspots. The impact of COVID-19 on mortality has been
15 described at the regional level, while less is known about mortality in demographic subgroups
16 within municipalities.

17 We aimed to describe the excess mortality (EM) due to COVID-19 in the three most affected Italian
18 regions, by estimating EM in subgroups defined by gender and age classes within each
19 municipality from February 23 to March 31, 2020.

20 EM varied widely among municipalities even within the same region; it was similar between
21 genders for the ≥ 75 age group, while in the other age groups it was higher in males. Thus, nearby
22 municipalities may show a different mortality burden despite being under common regional health
23 policies, possibly as a result of policies adopted both at the regional and at the municipality level.

24

25 **Keywords**

26 Excess Mortality, COVID-19, Italy, Municipalities, Regions.

27

28 **Funding**

29 This research did not receive any specific grant from funding agencies in the public, commercial, or
30 not-for-profit sectors.

31

32

33

34

35

36

37

38

39

40

1 1. Introduction

2 As of May 10, 2020, 4,051,431 confirmed cases and 279,734 deaths of Coronavirus Disease 2019
3 (COVID-19), a respiratory infectious disease caused by the Severe Acute Respiratory Syndrome -
4 Coronavirus 2 (SARS-CoV-2), were reported worldwide, affecting 187 countries (Center for
5 Systems Science and Engineering (CSSE) at Johns Hopkins University (JHU), 2020). Among
6 Western countries, Italy was the first affected. The first case of local transmission of COVID-19
7 was confirmed in a 38-year-old man in the municipality of Codogno (Lombardy region) on February
8 20, 2020 (Zangrillo et al., 2020), while on February 21, a resident of Vo', a small town near Padua
9 (Veneto region), died of pneumonia due to SARS-CoV-2 infection (Crisanti and Cassone, 2020;
10 Lavezzo et al., 2020). On February 23, the Italian government created two red zones confining ten
11 towns including Codogno and the town of Vo', quarantining 50,000 people.

12 Despite these containment measures, SARS-CoV-2 rapidly spread not only in the Lombardy and
13 Veneto region, but also in some areas of the bordering Emilia Romagna region. It became clear
14 that, without appropriate measures, SARS-CoV-2 would have spread dangerously all over Italy.
15 This prospect convinced the government to lockdown the whole country on March 10, 2020, when
16 confirmed cases were 10,149 and 631 deaths (Italian Civil Protection Department, 2020). At first,
17 commercial activities were closed, but some productive ones were allowed to stay open. On March
18 22, the government closed all productive activities, except for those necessary to produce
19 essential goods. As of May 10, 2020, official cases in Italy are 219.070 (Italian Civil Protection
20 Department, 2020), while deaths from COVID-19 reached 30.560. The COVID-19 had an
21 extremely varied distribution in Italy, because of its ways of spreading and of geographical
22 differences in population density, distribution of healthcare facilities, presence of care and nursing
23 homes and locally defined strategies to contain/mitigate its spreading. As a matter of fact, in Italy,
24 regions must provide the so called "Essential Levels of Care" (i.e. the core benefit package and
25 standard of health services), set by the central government, through the Regional Health Service
26 (RHS), but they are responsible and autonomous for the local organization and delivery of health
27 care (Lenzi et al., 2013). In each region, the municipalities followed the general indications given
28 by the Regional Government, but with different scaling up strategies at the local level.

29 Deaths from COVID-19 as reported by official data may be underestimated because of various
30 factors, among which the classification of cause of death when the deceased suffered from other
31 pre-existing conditions and the absence of post-mortem testing among those who have died before
32 they could be swabbed. For this reason, the most reliable method to estimate COVID-19 related
33 mortality to date is to calculate the excess all-cause mortality occurred during the COVID-19
34 outbreak compared to all-cause mortality in the same period in the previous years.
35 The Italian National Institute of Statistics (ISTAT), called on by researchers and local
36 administrators, promptly released and regularly updated the data of confirmed deaths (Italian
37 Institute for National Statistics (ISTAT), 2020b). The first studies (Colombo and Impicciatore, 2020;
38 Mancino, 2020) performed on these data reported higher excess mortality mainly in the
39 municipalities closer to the SARS-CoV-2 outbreak epicenters, such as those in Lombardy. Most of
40 these studies (Bucci et al., 2020; Buonanno et al., 2020; Modi et al., 2020) aimed to infer excess
41 mortality at the regional or national level, but all of them had to acknowledge the caveats implied
42 by relying on incomplete data to draw such predictions. Other studies, utilizing data provided by
43 the heat waves surveillance systems collected only in a limited number of municipalities (Davoli et
44 al., 2020), provided conflicting results, without showing a particular excess of mortality.

45 Importantly, excess mortality is not evenly distributed over the national territory and at the regional
46 and municipality level (Davoli et al., 2020; Italian Institute for National Statistics (ISTAT), 2020b;
47 Italian Institute for National Statistics (ISTAT) and Italian Superior Institute of Health, 2020;
48 Mancino, 2020; ScienzaInRete, 2020). In fact, some municipalities seem to be unaffected by the
49 epidemic, with a number of deaths equal to or even lower than that related to the same period of
50 previous years, while in others the number of deaths was estimated between 6 to 10 times higher.

51 Therefore, the aim of our study is to describe the spatial and demographic distribution of excess
52 mortality due to COVID-19 in the three most affected Italian regions, by estimating excess mortality
53 in subgroups defined by gender and age classes within each municipality with available data.

1 **2. Methods**

2 We conducted a longitudinal study using the data provided by the Italian Institute of Statistics
3 (ISTAT) on May 4, 2020, which include the number of deaths available for 6,866 Italian
4 municipalities (87.0% out of the 7,904 municipalities) (Italian Institute for National Statistics
5 (ISTAT), 2020c) divided by gender and age classes in the period from January 1 to March 31 of
6 the years 2015-2020. The municipalities with available data were 1,439/1,506 (95.8%) in
7 Lombardy, 490/563 (87.0%) in Veneto and 295/328 (89.9%) in Emilia-Romagna.

8 Excess mortality (EM) was estimated for each municipality in six subgroups defined by gender and
9 three age classes consistent with data provided by ISTAT (0-64, 65-74 and ≥ 75 years) in the time
10 period from February 23, 2020 to March 31, 2020, corresponding to the EPidemic Period (EPP). In
11 each subgroup a simple linear regression was conducted on the observed deaths from 2015 to
12 2019; the constant and slope obtained were used to estimate by extrapolation the expected
13 number of deaths in 2020. The ratio of observed to expected death is the relative mortality (RM)
14 experienced by each subgroup and the corresponding excess mortality is obtained by subtracting
15 1 from the RM and multiplying this value by 100. To avoid obtaining a negative expected number
16 of deaths, which might happen in the smallest municipalities, we set to 1 each prediction ≤ 0 .
17 Moreover, taking into account that 2020 is a leap year we corrected estimates by dividing for a
18 coefficient of 1.022.

19 To summarize the impact of COVID-19, we reported for each age and gender subgroup how many
20 municipalities showed an excess mortality (separating those under +50%, which could include also
21 not anomalous increases of mortality, from those over +50%) and how many showed a lower
22 mortality in the EPP.

23 Lastly, relative mortality ratios were applied to choropleth maps (Pisati, 2007) of the three Italian
24 regions considered, in order to visually identify the areas that suffered the greatest burden of
25 COVID-19 mortality during the epidemic and the municipalities where a reduction in mortality was
26 observed. We drew side by side regional maps of two age groups (0-64, ≥ 75 years) to visually
27 check the RM differences.

28 All analyses were carried out using Stata v.15.1 (StataCorp; College Station, TX, USA).
29

30

31

32

33

34

35

36

37

38

39

40

41

42

43

44

45

1
2
3
4
5
6
7
8
9
10
11
12
13
14

3. Results

Relative mortality during the COVID-19 outbreak exhibited a very composite pattern among the three examined regions, and between age classes and genders (Table 1). Generally, Lombardy experienced the highest burden of mortality, and Veneto the lowest. In the three regions, females displayed an excess mortality only among those aged ≥ 75 years. Males in age groups 65-74 and ≥ 75 had excess mortality in the three regions, except for the age group 65-74 in Veneto. People aged ≥ 75 years had the highest excess mortality. In Lombardy, this excess was +170% for males and +146% for females, in Emilia-Romagna +123% for males and +104% for females; in Veneto +35% for males and +37% for females. While among people aged ≥ 75 , EM was quite similar between males and females, in the other two age classes it was higher in males. For instance, in the 65-74 age group the EM of males was almost twofold compared to females in Lombardy and Emilia Romagna.

	EPP		
	median	mean	maximum
LOMBARDIA (n=1439) (23.02.2020 to 31.03.2020)			
0-64y Females	0	0.43	7
0-64y Males	0.29	0.87	10
65-74y Females	0	0.71	9
65-74y Males	1	1.57	22
≥ 75 y Females	1.5	2.46	36
≥ 75 y Males	2	2.70	47
VENETO (n=490) (23.02.2020 to 31.03.2020)			
0-64y Females	0	0.36	3
0-64y Males	0	0.52	5
65-74y Females	0	0.39	5
65-74y Males	0	0.64	6
≥ 75 y Females	1	1.37	11
≥ 75 y Males	1	1.35	13
EMILIA-ROMAGNA (n=295) (23.02.2020 to 31.03.2020)			
0-64y Females	0	0.47	5
0-64y Males	1	1.02	26
65-74y Females	0.5	0.82	6
65-74y Males	1	1.38	8
≥ 75 y Females	1.43	2.04	14
≥ 75 y Males	1.53	2.23	14

15
16
17
18
19
20
21
22
23

Table 1. Relative mortality in municipalities (age and gender subgroups) during the COVID-19 epidemic. Median, mean and maximum values are reported for each subgroup.

1
2
3
4
5
6
7
8
9

3.1. Municipality level

In Lombardy, males aged ≥ 75 experienced an EM in 60.3% of municipalities, compared to the previous five years, while in the remaining 39.7% no EM was observed (Table 2). In Emilia-Romagna the proportion of municipalities with EM was similar to that of Lombardy (64.4% of municipalities, males ≥ 75 years) while in Veneto it was lower (40.6%). These figures were similar for females aged ≥ 75 . As illustrated in Table 2, in most of the municipalities no EM was found for males and females in the age groups 0-64 and 65-74.

LOMBARDIA (n=1439)	Low mortality	Excess mortality		Total
		EM \leq 50%	EM $>$ 50%	
0-64y Females, n(%)	1052(78.7)	11(0.8)	273(20.4)	1336(100.0)
0-64y Males, n(%)	898(67.2)	13(1.0)	425(31.8)	1335(100.0)
65-74y Females, n(%)	1043(76.5)	6(0.4)	315(23.1)	1364(100.0)
65-74y Males, n(%)	774(56.7)	21(1.5)	569(41.7)	1364(100.0)
≥ 75 y Females, n(%)	607(42.2)	107(7.4)	725(50.4)	1439(100.0)
≥ 75 y Males, n(%)	572(39.7)	73(5.1)	794(55.2)	1439(100.0)
Total (%)	59.7	2.8	37.5	100.0

VENETO (n=490)	Low mortality	Excess mortality		Total
		EM \leq 50%	EM $>$ 50%	
0-64y Females, n(%)	402(84.4)	0(0)	74(15.6)	476(100.0)
0-64y Males, n(%)	376(79.0)	7(1.5)	93(19.5)	476(100.0)
65-74y Females, n(%)	413(86.9)	0(0)	62(13.1)	475(100.0)
65-74y Males, n(%)	374(78.7)	10(2.1)	91(19.2)	475(100.0)
≥ 75 y Females, n(%)	277(56.5)	59(12.0)	154(31.4)	490(100.0)
≥ 75 y Males, n(%)	291(59.4)	48(9.8)	151(30.8)	490(100.0)
Total (%)	74.0	4.3	21.7	100.0

EMILIA-ROMAGNA (n=295)	Low mortality	Excess mortality		Total
		EM \leq 50%	EM $>$ 50%	
0-64y Females, n(%)	240(83.0)	3(1.0)	46(15.9)	289(100.0)
0-64y Males, n(%)	207(71.6)	7(2.4)	75(26.0)	289(100.0)
65-74y Females, n(%)	210(72.4)	3(1.0)	77(26.6)	290(100.0)
65-74y Males, n(%)	175(60.3)	11(3.8)	104(35.9)	290(100.0)
≥ 75 y Females, n(%)	108(36.6)	56(19.0)	131(44.4)	295(100.0)
≥ 75 y Males, n(%)	105(35.6)	42(14.2)	148(50.2)	295(100.0)
Total (%)	59.8	7.0	33.2	100.0

10
11
12
13
14
15
16
17


Table 2. Cross-classification of municipalities (age and gender subgroups) with low or excess mortality during the COVID-19 outbreak. The number (and percentage) of municipalities for each subgroup is reported.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22

Concerning individual municipalities (Supplementary materials), the highest EM was estimated in Brembate di Sopra (10 km from Bergamo, in Lombardy) in males aged ≥ 75 (+4,600%), with 47 deaths occurred against 1 expected.


In the major towns, Parma (Emilia-Romagna) displayed a +633% EM in males aged 65-74; Bergamo (Lombardy) +551% EM in males aged ≥ 75 , +409% EM in males aged 65-74 and +368% EM in females aged ≥ 75 ; Piacenza (Emilia-Romagna) had +491% EM in males ≥ 75 . The regional capitals Venezia (Veneto) and Bologna (Emilia-Romagna) experienced EM rates generally below +50%. On the contrary, Milano (Lombardy) had an EM of +166% in males aged 65-74 and an EM between +56% and +76% in the other groups except for females aged 0-64 with +4%.

Examination of the Lombardy map for men aged ≥ 75 (Figure 1) confirmed that the municipalities of Bergamo, Brescia, Cremona and Lodi provinces, corresponding to the central and eastern part of the region, experienced the highest EM. The “green” municipalities with a mortality reduction are mainly located in the peripheral areas of the region which are more distant and less connected to the epidemic epicenters. In men aged 0-64 this pattern is less visible, because most municipalities did not experience an EM for this age group. Differently from Lombardy, a geographical pattern of epidemic spread for males aged ≥ 75 is not clearly visible in Veneto (Figure 2), as municipalities with EM seem to be scattered in different areas, intertwined with “green zones”. Moreover, in men aged 0-64 very few municipalities displayed an EM. The western Verona province, bordering Lombardy, was not the most hit by mortality within the region. From these maps we can argue that Veneto successfully isolated itself from the potential epidemic flow coming from the nearby region.



23
24
25
26
27
28
29
30


Figure 1. Relative mortality in males aged 0-64 (left) and ≥ 75 years (right) in Lombardy region. Municipalities are depicted with different colors according to the magnitude of their excess mortality. Green refers to municipalities with a relative mortality ≤ 1 , that is when observed deaths are lower than or equal to the expected deaths. When the relative mortality is > 1 , increasing saturation of red is used, to reflect increasing values of relative mortality. Municipalities depicted in white are those for which no data was released by ISTAT.




1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27

Figure 2. Relative mortality in males aged 0-64 (left) and ≥ 75 years (right) in Veneto region. Municipalities are depicted with different colors according to the magnitude of their excess mortality. Green refers to municipalities with a relative mortality ≤ 1 , that is when observed deaths are lower than or equal to the expected deaths. When the relative mortality is > 1 , increasing saturation of red is used, to reflect increasing values of relative mortality. Municipalities depicted in white are those for which no data was released by ISTAT.

1 As for the Emilia-Romagna region (Figure 3), it is evident for both age groups an EM north-south
2 gradient from the area bordering the Lodi and Cremona provinces of Lombardy, following the
3 trajectory of the main regional route that connects almost all the major regional centers. Moreover,
4 one local hotspot of COVID-19 can be seen in the east of the region, spreading from the
5 municipality of Medicina, and another local critical area might be noticed in the southernmost
6 municipalities, bordering the Pesaro province of the Marche region, where another known hotspot
7 was located. Similarly to Lombardy and Veneto, EM in the ≥ 75 age group was remarkably higher
8 than in the 0-64 age group.



Source: Istat



Source: Istat

9
10

11 **Figure 3.** Relative mortality in males aged 0-64 (above) and ≥ 75 years (below) in Emilia-Romagna region.
12 Municipalities are depicted with different colors according to the magnitude of their excess mortality. Green
13 refers to municipalities with a relative mortality ≤ 1 , that is when observed deaths are lower than or equal to
14 the expected deaths. When the relative mortality is > 1 , increasing saturation of red is used, to reflect
15 increasing values of relative mortality. Municipalities depicted in white are those for which no data was
16 released by ISTAT.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51

Comparing the epidemic period with the first five weeks of the year, in the large majority of municipalities of Lombardy and Emilia-Romagna, and to a lesser extent in Veneto, mortality in 2020 before the COVID-19 outbreak was lower than expected (data not shown). This trend has been observed in many countries worldwide (EuroMOMO, 2020).

4. Discussion

In this study we found that excess mortality due to COVID-19 in Italy has an inhomogeneous spatial and demographic distribution in the most affected regions of Lombardy, Emilia-Romagna and Veneto, and within each region between the different municipalities.

Our study highlighted that COVID-19 related EM was similar between males and females for the ≥ 75 age group, while in the other age groups it was higher in males. Whether SARS-CoV-2 has a higher rate of severe clinical manifestations and mortality among older people will be clarified when more studies on the characteristics of patients will be available. However increasing age and pre-existing conditions such as diabetes, obesity and cardiovascular diseases could play an important role (Onder et al., 2020; Italian Superior Institute of Health, 2020).

The spread of COVID-19 epidemic clearly originated from specific areas and initially affected the surrounding municipalities. The epidemic was spread by individuals living in a community, and circulated across the regions through the main commercial and commuting routes, as suggested by other analyses (Adnkronos, 2020; Gloria, 2020; Golinelli et al., 2020; Sebastiani, 2020). Some areas distant from the principal epicenters most likely benefited from the lockdown measures enforced nationwide before the epidemic spread could massively reach them.

The different RHS organizations must be taken into account when analyzing the responses to the epidemic. Lombardy has a Regional Health System based on a strong hospital-centered approach, with many excellences in secondary and tertiary care. The Veneto region has a RHS with a solid primary care level; Emilia-Romagna is somewhere in between, but its health care system is more similar to the one in Veneto (Cicchetti et al., 2020). When SARS-CoV-2 started spreading in the town of Vo', the RHS in Veneto and the policies adopted at the regional and local level allowed the region to quickly react. Adopting mainly a community based approach, a multitude of actions were taken at once: the municipality of Vo' was immediately declared a red zone, massive testing (even if at the beginning of the outbreak test capacity is usually low (Ruan, 2020), isolation of cases, rapid contact tracing. In Vo', all people, also the asymptomatic, were tested (Crisanti and Cassone, 2020; Lavezzo et al., 2020). Importantly, Veneto largely relied on home care assistance, limiting hospital admissions to the most severe cases and started early testing of the healthcare workers operating in the community and in the hospitals. In Emilia-Romagna, the community-based approach was used to contrast the epidemic as well, but Piacenza, where the first outbreak of the region was registered, originating from the nearby hotspot of Codogno, still remains one of the most affected municipalities in Italy. Probably the virus had already extensively spread when the first case was found, and the community services were not equipped to promptly react, being overwhelmed by the high volumes of contemporary cases. Also, it could be that the municipality was not quickly declared a red zone, when there was the suspicion that the number of cases still undetected might have been dangerously high. Nevertheless, the RHS was able to adapt and started being more effective in mitigating the spread of the virus, following the same approach used by Veneto, particularly leaning towards home care assistance and relying on general practitioners to implement an active surveillance system among their patients with phone calls to monitor their symptoms.

On the contrary, the organizational structure of the RHS in Lombardy that promotes a hospital centered approach at the expenses of the community-based services, may have contributed to exacerbate the criticalities presented by COVID-19. Thereto, the region did not address the outbreak adopting the same plan in municipality experiencing similar circumstances: while the

1 towns around Lodi were quickly declared red zones when the first cases were identified, thus
2 containing the spread of the virus, the outbreak of the virus in a hospital in the province of
3 Bergamo, didn't lead to the establishment of a confinement, that was implemented only when the
4 national lockdown was enforced (De Luca, 2020). This contributed to make the Bergamo area one
5 of the most hit areas in Europe. Especially at the beginning of the epidemic, most people were
6 admitted to hospitals. The dramatic flow of patients saturated the intensive care units, forcing
7 doctors to decide who could go forward (Rosenbaum, 2020). In the hospital setting, the virus was
8 spread not only by patients, but also by healthcare workers, who could not always rely on
9 appropriate personal protective equipment (PPE) risking their lives while doing their job.

10 Analyzing the deaths' toll in Lombardy, Veneto and Emilia-Romagna, we argue that the best way
11 to address epidemics is through proactive medicine at a community-based level, opposite to
12 reactive medicine at a hospital level. Moreover, to foster discussions and more in depth analysis
13 on the EM due to COVID-19 infections, we hypothesized several factors that could have equally
14 affected the infections and mortality in the three regions: household transmission, which led to the
15 most vulnerable being exposed when a family member got the infection; congestion of the
16 healthcare system; late diagnosis of the healthcare workers for COVID-19; limited number of
17 executed swabs; inadequate contact tracing; public transportation; inadequate community based
18 interventions.

19 At the regional level, the difference in adopted strategies certainly reflects different RHS's
20 organization and operational approaches. In tackling the epidemic, the Veneto and Lombardy
21 regions adopted two opposite strategies regarding, for example, the recommendations on testing.
22 As of March 31, Veneto performed 21.6 swabs per 1,000 population (4,905,854 inhabitants) and,
23 whenever possible, tested also asymptomatic people; Lombardy (10,060,574 inhabitants)
24 performed 11.3 swabs per 1,000 population, testing mainly patients with more severe clinical
25 symptoms, as instructed by the Italian Ministry of Health in that moment. On this topic, Emilia
26 Romagna had a more similar approach to Lombardy, performing 12.2 swabs per 1,000 population
27 on a total population of 4,459,477 inhabitants (Italian Civil Protection Department, 2020; Italian
28 Institute for National Statistics (ISTAT), 2020a; Onder et al., 2020).

29 At the local level, the reason of higher COVID-19 mortality in some specific municipalities can be
30 debated, as several different causes other than the distance from the hotspots or a possible bias in
31 estimation or some casual local factors might be involved. Possible explanations for a marked
32 difference in excess mortality between close municipalities could be the presence of gathering
33 places (i.e. sport venues, cinemas), or of some industrial/commercial activities, care and nursing
34 homes or of particular events that may have taken place before the lockdown. In the municipality
35 of Medicina (Emilia Romagna), a gathering at a bowls club was the cause of the rapid spread of
36 the virus; almost all the people present got infected, and all but one died. Medicina immediately
37 became a red zone (Regione Emilia-Romagna, 2020), successfully narrowing the afflicted area.
38 This demonstrates that it is important to contrast the spread of the virus at the smallest possible
39 level and in the most accurate possible way.

40 Several municipalities showed a lower increase of COVID-19 related deaths or even a reduced
41 mortality. This can be likely attributed to a smaller incidence of deaths for causes such as
42 accidents or occupational injuries, favored by the lockdown. On the other hand, the high mortality
43 figures especially for people aged ≥ 75 may also have been favored by the lower incidence of
44 influenza observed this year in Italy, compared to last year, which left more elderly people exposed
45 to COVID-19 (Bella, 2020).

46 Our study suggests that the difference in the response to the epidemic can be the result of policies
47 adopted both at the regional and at the municipality level. Most of the studies conducted on this
48 epidemic (Bucci et al., 2020; Buonanno et al., 2020; Modi et al., 2020; ScienzaInRete, 2020) have
49 tried to explain its course considering the regional policies, as so did we, but, as an added value, in
50 the present study we explored also the pattern of the epidemic within regions and noticed that, net
51 of regional policies, even nearby municipalities showed different excess mortality rates.
52 Accordingly, the causes of such differences should be sought not only at the national and the
53 regional policies, but also in local factors which cannot be brought to light if data are analyzed as
54 regional or higher aggregates. Therefore, focusing on the regional and municipality level could be

1 more helpful than a regional level analysis alone and we advocate that our approach may help to
2 shed some light on the areas where the death toll was highest and thus trigger further
3 investigations on some specific/local dynamics of the outbreak.

4 Our study presents some caveats. First, it is based on incomplete territorial coverage, therefore
5 our conclusions could have been partially different if data for all municipalities was available.
6 However, our data cover >87% of Italian municipalities, which is a relevant coverage. We used
7 linear regression to derive a mortality trend from 5 years, which could not always be adequate as it
8 may be leveraged by anomalous mortality figures in the first or in the last year of the time interval
9 used for prediction. By using linear regression, we intended to capture a trend in mortality where it
10 exists, and in these cases its prediction is more accurate than the projection of the average
11 mortality observed in the previous five years. In addition, we reduced potential confounding by
12 estimating excess mortality within subgroups defined by age class and gender. Nevertheless, our
13 results should be considered as provisional estimates until official data of verified quality covering
14 all municipalities will be released and made available to researchers for further investigations.
15

16 **5. Conclusions**

17 Our approach could be used to generate or to confirm hypotheses regarding the flow of epidemics
18 starting from the main originating sites, and to detect affected loci that are distant from these initial
19 affected areas. Analyses are fundamental at the regional and subregional level. Identifying the
20 municipalities where the mortality burden was higher, and the pathways used by the virus to
21 spread may help to concentrate efforts in understanding the reasons why this happened and to
22 identify the frailest areas in case of the occurrence of a second epidemic outbreak. Whether higher
23 excess mortality depended on lack of preparedness, or on a particularly old age structure, or on
24 inefficacy in the enforcement of containment measures, this should lead to different types of
25 interventions in the preparation of a possible second wave of COVID-19.

26 The phase of coexistence with the virus will be as challenging as the phase of the outbreak. An
27 adequate organization of the health services is imperative.

28 Our study demonstrates that nearby municipalities within each region may show highly different
29 mortality levels, despite being under common regional health policies, therefore further studies are
30 necessary to analyze more in depth the local determinants of COVID-19 spread.
31
32
33

34 **6. References**

35 Adnkronos, 2020. Il matematico: 'Virus ha viaggiato in autostrada sui tir' [online], 12 April. URL
36 [https://www.adnkronos.com/fatti/cronaca/2020/04/12/matematico-virus-viaggiato-autostrada-](https://www.adnkronos.com/fatti/cronaca/2020/04/12/matematico-virus-viaggiato-autostrada-sui-tir_D0v4m2iPjmBghbr1hskyqJ.html)
37 [sui-tir_D0v4m2iPjmBghbr1hskyqJ.html](https://www.adnkronos.com/fatti/cronaca/2020/04/12/matematico-virus-viaggiato-autostrada-sui-tir_D0v4m2iPjmBghbr1hskyqJ.html) (accessed 5.3.20).

38 Bella, A., 2020. Rapporto Epidemiologico Influnet - Stagione Influenzale 2019-2020 - Settimana
39 2020-14.

40 Bucci, E., Leuzzi, L., Marinari, E., Parisi, G., Ricci-Tersenghi, F., 2020. Un'analisi dei dati ISTAT sui
41 decessi legati all'epidemia Covid-19 in Italia: verso una stima del numero di morti dirette ed
42 indirette, anche grazie allo sbilanciamento di genere [online]. URL
43 [https://www.scienzainrete.it/articolo/verso-stima-di-morti-dirette-e-indirette-covid/enrico-bucci-](https://www.scienzainrete.it/articolo/verso-stima-di-morti-dirette-e-indirette-covid/enrico-bucci-luca-leuzzi-enzo-marinari)
44 [luca-leuzzi-enzo-marinari](https://www.scienzainrete.it/articolo/verso-stima-di-morti-dirette-e-indirette-covid/enrico-bucci-luca-leuzzi-enzo-marinari) (accessed 4.24.20).

45 Buonanno, P., Galletta, S., Puca, M., 2020. Estimating the severity of COVID-19: evidence from the
46 Italian epicenter? SSRN Electron. J. <https://doi.org/10.2139/ssrn.3567093>

47 Center for Systems Science and Engineering (CSSE) at Johns Hopkins University (JHU), 2020.
48 COVID-19 Dashboard [online]. URL <https://coronavirus.jhu.edu/map.html> (accessed 5.3.20).

- 1 Cicchetti, A., Damiani, G., Specchia, M.L., Basile, M., Di Bidino, R., Di Brino, E., Di Paolo, M.G., Di
2 Pilla, A., Ferrara, F.M., Giorgio, L., Riccardi, M.T., Rumi, F., Tattoli, A., 2020. Analisi dei
3 modelli organizzativi di risposta al Covid-19. Focus su Lombardia, Veneto, Emilia-Romagna,
4 Piemonte e Lazio. Rome.
- 5 Colombo, A.D., Impicciatore, R., 2020. La crescita della mortalità ai tempi del Covid-19, Cattaneo
6 Institute. Bologna.
- 7 Crisanti, A., Cassone, A., 2020. In one Italian town, we showed mass testing could eradicate the
8 coronavirus [online], 20 March. The Guardian. URL
9 [https://www.theguardian.com/commentisfree/2020/mar/20/eradicated-coronavirus-mass-](https://www.theguardian.com/commentisfree/2020/mar/20/eradicated-coronavirus-mass-testing-covid-19-italy-vo)
10 [testing-covid-19-italy-vo](https://www.theguardian.com/commentisfree/2020/mar/20/eradicated-coronavirus-mass-testing-covid-19-italy-vo)
- 11 Davoli, M., Sario, M. De, Michelozzi, P., Noccioli, F., Rossi, P., Scortichini, M., 2020. Andamento
12 della Mortalità Giornaliera (SiSMG) nelle città italiane in relazione all'epidemia di Covid-19,
13 Secondo Rapporto 1 febbraio - 28 marzo, Epidemiology Department of Latium's Regional
14 Health Service. Rome.
- 15 De Luca, D.M., 2020. Il disastro in Val Seriana [online], 1 April. il Post. URL
16 <https://www.ilpost.it/2020/04/01/disastro-alzano-lombardo-nembro/>
- 17 EuroMOMO, 2020. Maps and Graphs [online]. URL <https://www.euromomo.eu/graphs-and-maps>
18 (accessed 5.3.20).
- 19 Gloria, S., 2020. 'È l'epidemia della A21': così il virus si è spostato lungo l'autostrada da Brescia a
20 Torino [online], 23 April. URL [https://www.fanpage.it/attualita/e-lepidemia-della-a21-cosi-il-](https://www.fanpage.it/attualita/e-lepidemia-della-a21-cosi-il-virus-si-e-spostato-lungo-lautostrada-da-brescia-a-torino/)
21 [virus-si-e-spostato-lungo-lautostrada-da-brescia-a-torino/](https://www.fanpage.it/attualita/e-lepidemia-della-a21-cosi-il-virus-si-e-spostato-lungo-lautostrada-da-brescia-a-torino/) (accessed 5.3.20).
- 22 Golinelli, D., Fantini, M.P., Maietti, E., 2020. COVID-19 in Italy: is the virus running through an
23 ancient Roman road? Preprints 2020040513. <https://doi.org/10.20944/preprints202004.0513.v1>
- 24 Italian Civil Protection Department, 2020. Official Italian Data about COVID-19 [online]. URL
25 <https://github.com/pcm-dpc/COVID-19> (accessed 4.26.20).
- 26 Italian Institute for National Statistics (ISTAT), 2020a. Official Italian Data about the resident
27 population [online]. URL http://dati.istat.it/Index.aspx?DataSetCode=DCIS_POPSTRCIT1
28 (accessed 5.10.20).
- 29 Italian Institute for National Statistics (ISTAT), 2020b. Official Data about 2015-2020 mortality in
30 Italy [online]. URL <https://www.istat.it/it/archivio/240401> (accessed 5.3.20).
- 31 Italian Institute for National Statistics (ISTAT), 2020c. Official Italian Data about the Nomenclature of
32 Territorial Units for Statistics [online]. URL <https://www.istat.it/it/archivio/6789> (accessed
33 5.3.20).
- 34 Italian Institute for National Statistics (ISTAT), Italian Superior Institute of Health, 2020. Impatto
35 dell'epidemia COVID-19 sulla mortalità totale della popolazione residente - Primo Trimestre
36 2020. Rome.
- 37 Italian Superior Institute of Health, 2020. Caratteristiche dei pazienti deceduti positivi a COVID-19 in
38 Italia - 9 Aprile 2020. Rome.
- 39 Lavezzo, E., Franchin, E., Ciavarella, C., Cuomo-Dannenburg, G., Barzon, L., Vecchio, C. Del,
40 Rossi, L., Manganelli, R., Loregian, A., Navarin, N., Abate, D., Sciro, M., Merigliano, S.,
41 Decanale, E., Vanuzzo, M.C., Saluzzo, F., Onelia, F., Pacenti, M., Parisi, S., Carretta, G.,
42 Donato, D., Flor, L., Cocchio, S., Masi, G., Sperduti, A., Cattarino, L., Salvador, R., Gaythorpe,
43 K.A.M., Team, I.C.L.C.-19 R., Brazzale, A.R., Toppo, S., Trevisan, M., Baldo, V., Donnelly,

- 1 C.A., Ferguson, N.M., Dorigatti, I., Crisanti, A., 2020. Suppression of COVID-19 outbreak in the
2 municipality of Vo, Italy. medRxiv 2020.04.17.20053157.
3 <https://doi.org/10.1101/2020.04.17.20053157>
- 4 Lenzi, J., Rucci, P., Franchino, G., Domenighetti, G., Damiani, G., Fantini, M.P., 2013. Regional and
5 gender variation in mortality amenable to health care services in Italy. J. Hosp. Adm. 2, 28.
6 <https://doi.org/10.5430/jha.v2n3p28>
- 7 Mancino, D., 2020. Coronavirus, a marzo anche a Milano la mortalità è raddoppiata [online], 3 April.
8 Il Sole 24 Ore. URL <https://www.infodata.ilsole24ore.com/2020/04/03/coronavirus-marzo->
9 [anche-milano-la-mortalita-raddoppiata/](https://www.infodata.ilsole24ore.com/2020/04/03/coronavirus-marzo-)
- 10 Modi, C., Böhm, V., Ferraro, S., Stein, G., Seljak, U., 2020. Total COVID-19 Mortality in Italy:
11 Excess Mortality and Age Dependence through Time-Series Analysis. medRxiv
12 2020.04.15.20067074. <https://doi.org/https://doi.org/10.1101/2020.04.15.20067074>
- 13 Onder, G., Rezza, G., Brusaferro, S., 2020. Case-Fatality Rate and Characteristics of Patients
14 Dying in Relation to COVID-19 in Italy. JAMA. <https://doi.org/10.1001/jama.2020.4683>
- 15 Pisati, M., 2007. SPMAP: Stata module to visualize spatial data, Statistical Software Components,
16 Boston College Department of Economics.
- 17 Regione Emilia-Romagna, 2020. Coronavirus, chiuso il comune di Medicina (Bo): contagio troppo
18 elevato, non è possibile uscire e entrare Ordinanza [online], 16 March. URL
19 <https://www.regione.emilia-romagna.it/notizie/2020/marzo/coronavirus-chiuso-il-comune-di->
20 [medicina](https://www.regione.emilia-romagna.it/notizie/2020/marzo/coronavirus-chiuso-il-comune-di-) (accessed 5.3.20).
- 21 Rosenbaum, L., 2020. Facing Covid-19 in Italy — Ethics, Logistics, and Therapeutics on the
22 Epidemic's Front Line. N. Engl. J. Med. <https://doi.org/10.1056/NEJMp2005492>
- 23 Ruan, S., 2020. Likelihood of survival of coronavirus disease 2019. Lancet Infect. Dis.
24 [https://doi.org/10.1016/S1473-3099\(20\)30257-7](https://doi.org/10.1016/S1473-3099(20)30257-7)
- 25 ScienzaInRete, 2020. Mortalità in Italia (1 gennaio - 15 aprile 2020) [online]. URL
26 <https://www.scienzainrete.it/articolo/dati-covid-19/2020-04->
27 [19?utm_source=phplist1435&utm_medium=email&utm_content=HTML&utm_campaign=Crona](https://www.scienzainrete.it/articolo/dati-covid-19/2020-04-)
28 [che+della+ricerca+%23134](https://www.scienzainrete.it/articolo/dati-covid-19/2020-04-) (accessed 5.10.20).
- 29 Sebastiani, G., 2020. Il coronavirus ha viaggiato in autostrada? [online]. URL
30 <https://www.scienzainrete.it/articolo/coronavirus-ha-viaggiato-autostrada/giovanni->
31 [sebastiani/2020-04-09](https://www.scienzainrete.it/articolo/coronavirus-ha-viaggiato-autostrada/giovanni-) (accessed 5.3.20).
- 32 Zangrillo, A., Beretta, L., Silvani, P., Colombo, S., Scandroglio, A.M., Dell'Acqua, A., Fominskiy, E.,
33 Landoni, G., Monti, G., Azzolini, M.L., Monaco, F., Oriani, A., Belletti, A., Sartorelli, M.,
34 Pallanch, O., Saleh, O., Sartini, C., Nardelli, P., Lombardi, G., Morselli, F., Scquizzato, T.,
35 Frontera, A., Ruggeri, A., Scotti, R., Assanelli, A., Dagna, L., Rovere-Querini, P., Castagna, A.,
36 Scarpellini, P., Di Napoli, D., Ambrosio, A., Ciceri, F., Tresoldi, M., 2020. Fast reshaping of
37 intensive care unit facilities in a large metropolitan hospital in Milan, Italy: facing the COVID-19
38 pandemic emergency. Crit. care Resusc. J. Australas. Acad. Crit. Care Med.