

## Pregnancy and Neonatal Outcomes in SARS-CoV-2 Infection: a systematic review

Reem S Chamseddine<sup>1</sup>, Farah Wahbeh<sup>1</sup>, Frank Chervenak<sup>5</sup>, Laurent J Salomon<sup>6</sup>, Baderledeen Ahmed<sup>2,3</sup>, Arash Rafii<sup>1,4</sup>

<sup>1</sup> Genetic Intelligence Laboratory, Weill Cornell Medicine-Qatar, Doha, Qatar

<sup>2</sup> Fetal Maternal Center, Doha, Qatar

<sup>3</sup> Qatar University, Doha, Qatar

<sup>4</sup> Department of Genetic Medicine and Obstetrics and Gynecology, Weill Cornell Medicine, New York, USA

<sup>5</sup> Department of Obstetrics and Gynecology, Zucker School of Medicine at Hofstra/Northwell--Lenox Hill Hospital, New York, USA

<sup>6</sup> Maternité, médecine, chirurgie et imagerie fœtales, Hôpital Universitaire Necker Enfants Malades, Université de Paris, Paris, France and EA FETUS, affiliate of Institute IMAGINE, Paris, France

Corresponding Author: Dr. Arash Rafii ([jat2021@qatar-med.cornell.edu](mailto:jat2021@qatar-med.cornell.edu)).

Department of Genetic Medicine and Obstetrics and Gynecology, Weill Cornell Medical College, Genetic Intelligence laboratory Weill Cornell Medicine College in Qatar, Qatar-Foundation PO: 24144, Doha, Qatar. Phone: +974 3313 5828. Fax: +974 4492 8422.

### Abstract

*With the emergence of SARS-CoV-2 and its rapid spread, concerns regarding its effects on pregnancy outcomes have been growing. We reviewed 164 pregnancies complicated by maternal SARS-CoV-2 infection across 20 studies. The most common clinical presentations were fever (57.9%), cough (35.4%), fatigue (15.2%), and dyspnea (12.2%). Only 2.4% of patients developed respiratory distress. Of all patients, 84.5% delivered via Cesarean section, with a 23.9% rate of maternal gestational complications, 20.3% rate of preterm delivery, and a concerning 2.3% rate of stillbirth delivery. Relative to known viral infections, the prognosis for pregnant women with SARS-CoV-2 is good, even in the absence of specific antiviral treatment. However, neonates and acute patients, especially those with gestational or pre-existing co-morbidities, must be actively managed to prevent severe outcomes.*

## **Introduction**

The emergence of SARS-CoV-2 as a novel infection in late December 2019 poses unique challenges to healthcare systems and practitioners. Chief among them is the management for pregnant women who are infected with SARS-CoV-2. Pregnant women are prone to a range of fetal and maternal complications that could impact the outcome of any concurrent infection.<sup>1</sup> The first trimester bears a major risk for miscarriage and fetal developmental abnormalities. The late second and third trimesters carry an increased likelihood for the development of maternal conditions, such as gestational diabetes and hypertensive disorders, which contribute to maternal morbidity and premature birth.

It is well known that infectious pneumonia is a common cause of morbidity and mortality in pregnant women due to several physiological factors such as lower lung volumes and increased oxygen consumption.<sup>2,3</sup> Indeed, a quarter of pregnancies complicated by pneumonia are estimated to require critical care hospitalization and ventilation support.<sup>4</sup> Pregnancy-induced changes to immunity, such as reduction in cell-mediated cytotoxicity, diminished lymphokine response and reduction in lymphocyte proliferative response, further add to the prognostic picture of pneumonia in pregnant women.<sup>3</sup> Hence, the advent of the SARS-CoV-2 pandemic could raise new questions about perinatal and obstetric management. Here, we review 164 pregnancies reported in the literature across 20 studies.

## **Methods**

We searched PubMed using the terms “pregnancy,” “pregnant,” “prenatal,” “Covid-19,” “SARS-CoV2”, “miscarriage,” “fetal anomalies” and “complications.” We limited our search to all published studies between December 20th, 2019 to April 18th, 2020 and added a filter for English-language articles. Our search retrieved 84 papers. We carried out the same search on MedRxiv, which yielded 27 results. We excluded reviews, meta-reviews, letters to the editor, and guidelines that are specific to a region. We then reviewed the results for relevance and included the articles that reported on SARS-CoV-2 in pregnant patients. This search strategy resulted in 19 case reports and articles from the primary literature that we included in this review. We added one study outside the aforementioned date range due to its novelty and relevance to the subject at hand. We extracted

all information about patient characteristics, symptoms, laboratory results, and imaging studies. We also gathered the relevant data about the pregnancy course such as delivery method, gestational complications, and neonatal outcomes.

## **Results**

Results from individual studies are described and summarized in Table 1 and Table 2 below.

### ***Maternal Symptoms***

With regards to prepartum symptoms, the most common symptoms were fever (57.9%) followed by cough (35.4%), fatigue (15.2%) and dyspnea (12.2%). Less common symptoms included diarrhea and gastrointestinal symptoms, sore throat, nasal congestion and runny nose, muscle aches, rash, headache, hoarseness, myalgia, high blood pressure, tachycardia, and cholecystitis. Only 4.3% of all patients were asymptomatic prior to delivery. Two patients who were asymptomatic prior to delivery became symptomatic post-partum, both requiring ICU admission.

The majority of the studies included patients in the third trimester only with the exception of four studies which also included patients in the second trimester. Most of these studies reported the number of patients experiencing each symptom without specifying the GA, so it is difficult to determine whether patients in earlier trimesters differed in their clinical presentations from those in later trimesters. D. Liu and colleagues (2020) reported 15 patients who ranged in GA from 12 to 38 weeks, but it was not clear whether the patients in the earlier trimester experienced different symptoms or complications.<sup>5</sup> Nie (2020) included 33 patients in total which ranged in GA from 24 to 36 weeks.<sup>6</sup> No ICU admissions or mortalities were reported, and the study concluded that pregnant women were not at an increased risk for severe illness or mortality. Based on the findings reported by Y. Liu et al. (2020) and Nie (2020), patients' clinical presentations do not appear to differ based on the gestational age of pregnancy.

Among the reported postpartum symptoms, the most common one was fever (24.7%). In one patient who was admitted with worsening hypertension and a history of diabetes mellitus and asthma, postpartum symptoms included low oxygen saturation, tachycardia, dyspnea, diaphoresis, and cough that progressed to respiratory failure.<sup>7</sup> Gidlof et al. (2020) also reported low oxygen

saturation postpartum.<sup>8</sup> Among 13 patients, Y. Liu and colleagues (2020) reported that one patient developed Multi Organ Dysfunction Syndrome (MODS) including Acute Respiratory Distress Syndrome (ARDS).<sup>9</sup> Karami et al. (2020) also reported a case of pre-partum respiratory distress, which progressed to multi-organ failure post-partum and unfortunately in the eventual death of the patient and the fetus.<sup>10</sup> Thus, out of 164 pregnant women with SARS-CoV-2, 4 (2.44%) developed respiratory distress.

### ***Laboratory Findings***

The most common laboratory abnormalities seen in pregnant women infected with SARS-CoV-2 were elevated C-reactive protein (CRP), elevated neutrophils, and leukocytosis. In the studies that reported laboratory findings, 114 pregnant women were included and among them 50.9% had elevated CRP, 44.7% had lymphopenia, 34% had elevated neutrophils, and 16.7% had leukocytosis. Among 24 pregnant women with SARS-CoV-2, Liu H reported that 58.6% of them had low or normal leukocyte levels.

Other laboratory abnormalities included low platelets, elevated D-dimer, abnormal liver enzymes, decreased albumin, low hemoglobin, elevated alkaline phosphate, decreased eosinophils, elevated uric acid, elevated ferritin, pancytopenia, prolonged APTT, low fibrinogen, and progressive coagulopathy.

### ***Chest X-ray (CXR)***

Most studies reported imaging abnormalities in either chest X-ray (CXR), computed tomography (CT), or both. In terms of CXR, Y. Li et al. (2020) (n=1) reported scattered multiple patchy infiltrates in both lungs, Lee et al. (2020) (n=1) reported left lower/middle lobe consolidation and increased vascular marking, while Breslin et al. (2020) (n=1) reported mild pulmonary vascular congestion with no consolidation or effusion<sup>7,11,12</sup> All three women were of similar GA (mean 36 weeks).

### ***Computed Tomography (CT) Scan***

Typical signs of viral pneumonia were frequently reported, with most patients having bilateral lung involvement. CT findings did not show signs of worsening pneumonia postpartum. However, with disease progression, Zhu et al. (2020) reported that lesions merged into strips and D. Liu et al.

(2020) observed a paving pattern and consolidation.<sup>5,13</sup> Ground glass opacities (GGO) were reported in the majority of studies. Mixed GGO with consolidation were more common among pregnant SARS-CoV-2 patients in comparison to non-pregnant SARS-CoV-2 patients.<sup>5</sup> Lee and colleagues (2020) also reported GGO with consolidation.<sup>11</sup> In addition to multiple bilateral consolidations, bilateral pleural effusion were observed.<sup>14</sup> Given the prevalence of respiratory symptoms in pregnant women infected with SARS-CoV-2, lung ultrasound can a valuable tool in confirming diagnosis of SARS-CoV-2 infection as well as monitoring disease progression.<sup>15</sup> Lung ultrasound eliminates the radiation risk and avoids the incidence of false-negative results associated with other diagnostic tests. As outlined by Moro and colleagues (2020), specific guidelines can be followed when performing lung ultrasound for pregnant women infected with SARS-CoV-2.<sup>15</sup>

### ***Maternal and Obstetric Complications***

There have been several reports of uncomplicated term delivery in SARS-CoV-2 positive women.<sup>11,16</sup> Of 110 deliveries reported, 93 were performed as C-sections, including 10 that were performed as emergency surgeries. The rate of C-sections is thus 84.5% of all deliveries, significantly higher than the expected rate of approximately 15% for pregnancies in the general population.<sup>17,18</sup> This rate of C-section is also higher than that in pregnancies complicated by viral Influenza infection, where the likelihood of C-section did not increase relative to the general population.<sup>19</sup>

Several case reports highlight the frequency of C-section deliveries compared to vaginal deliveries, whether in healthy-appearing patients or patients presenting with fever.<sup>13,20</sup> The indications for C-section varied and were not consistently reported across studies. In certain hospitals, the choice of C-section was determined by obstetric factors.<sup>21</sup> In others, C-section was performed proactively due to maternal SARS-CoV-2 infection. Presumably, C-sections minimize cross-infection and reduce maternal exertion during labor, as per certain recommendations.<sup>22</sup> Due to the uneven distribution of C-sections as opposed to vaginal delivery, it is not possible to comment on differences in neonatal and maternal outcomes between vaginal delivery and C-sections in SARS-CoV-2 pregnancies. Importantly, there are no contraindications for vaginal delivery as evidenced by the uncomplicated vaginal delivery in at least five patients.<sup>23 24</sup> It is not clear from this review if the high Cesarean section rate is warranted.

Out of 163 pregnancies reviewed, we looked at the following gestational complications: gestational diabetes, preeclampsia or gestational hypertension, and placental complications such as premature rupture or placenta previa. The rate of complications was at 23.93%, composed of gestational diabetes (7.99%), preeclampsia or gestational hypertension (5.52%), and placental complications (6.13%). At least 6 additional patients (3.68%) had complications, but it is not possible to ascertain the specific condition due to the method of reporting in the primary literature.

### ***Fetal and Neonatal Outcomes***

The 110 pregnancies followed in the literature resulted in 128 newborns. Out of 128 neonates, 26 (20.31%) were delivered preterm, before 36 weeks of gestation. Again, it is not consistently clear whether early delivery was induced in light of obstetric indications or maternal SARS-CoV-2 infection. The vast majority of neonates for whom an Apgar score was reported (n=68) scored in the normal range, at least 8 at minute 1 and minute 5, with the exception of three neonates. Three newborns (4.41%) had a concerning Apgar score of <7 at minute 1.

Eight fetuses (6.35%) exhibited fetal distress, which was the indication for preterm delivery in some cases. Postnatally, the rate of respiratory distress or abnormal lung findings among the neonates was at 7.94%, divided as follows: 6 neonates had shortness of breath, 2 had pneumonia-like presentation on imaging, and two had respiratory distress including one case of ARDS. All cases resolved with antibiotics without complication. Additionally, three newborns presented with fever (2.34%).

Of the 128 fetuses, four had very poor outcomes: three deliveries resulted in stillbirth, including one fetus who was born to a mother who passed away soon after delivery.<sup>9,10,25</sup> One neonate developed refractory shock and multiple organ failure and passed on day 8 of life.<sup>13</sup> Although no fetus is thought to have died as a result of fetal infection, Baud et al. (2020) suggest that placental inflammation in the setting of the virus contributes to fetal distress, even when the fetal is not infected.<sup>25</sup> Notably, several case studies have ruled out vertical transmission of SARS-CoV-2. This is largely supported by our review. Out of the neonates tested for SARS-CoV-2 infection (n=44), 3 tested positive for the infection through nasopharyngeal swab (6.81%). Two of the cases in question raised concerns about potential vertical transmission, since the mother and the obstetric team practiced protection and isolation measures to prevent infection during the delivery process

and immediately afterwards.<sup>6,14</sup> However, because of the asymptomatic presentation of many infected individuals, it is possible that the neonates had acquired SARS-CoV-2 in the hospital after birth.

Certain studies went beyond testing the newborn for evidence of vertical transmission. Gidlof et al. (2020) tested breast milk and maternal vaginal secretion of a SARS-CoV-2 positive mother.<sup>8</sup> Y. Li et al. (2020) tested the breast milk, amniotic fluid, umbilical cord blood and placenta, maternal serum, urine, and feces of a SARS-CoV-2 positive mother.<sup>9</sup> Fan (2020) tested breast milk, maternal vaginal secretion, placental tissue, umbilical cord blood, amniotic fluid and maternal serum in two pregnancies.<sup>20</sup>

### ***Breastfeeding***

Collected breast milk samples from the four aforementioned pregnancies were negative for SARS-CoV-2. These results, in addition to several guidelines such as those published by the ACOG and the RCOG, converge on the idea that SARS-CoV-2 is not carried through breast milk.<sup>26,27</sup> However, the underlying risk of neonatal exposure to maternal respiratory droplets during the breastfeeding process persists. Overall, the many benefits of breastfeeding do make breastfeeding safe, as long as SARS-CoV-2 positive mothers take reasonable precautions while in close contact with the newborn.

### **Discussion**

According to Liu et al., pregnancy and delivery did not aggravate the severity of SARS-CoV-2 in the patients studied.<sup>5</sup> However, as one expects, the variability in the severity of infection and pneumonia presentation in different patients makes such a statement difficult to prove.

Pregnancy, in particular the third trimester, is associated with changes in lung physiology, such as decreases in ERV and FRC volumes and increases in respiratory resistance.<sup>28</sup> A prime example is the fact that pregnant women infected with the H1N1 virus during the 2009 epidemic were at a significantly higher risk of complications and hospitalization than non-pregnant women.<sup>29,30</sup> Hence, an increased rate of respiratory complications, in addition to maternal and neonatal complications, could be expected in pregnant patients with SARS-CoV-2 infection. The absence of widespread respiratory complications is surprising and suggests that a larger number of patients will help better understand the natural history of SARS-CoV-2 infections during pregnancy.

It is certainly true that, at least from the available reports, that the mortality risk in SARS-CoV-2 pregnant women is extremely low. There exists only one case of maternal mortality and three other cases of patients developing respiratory distress.<sup>7-9</sup> At the time of writing, there exists a single report of a (late) miscarriage due to SARS-CoV-2 infection, and no reports of termination due to early congenital defects. In the 2002-2004 SARS epidemic, miscarriage rates were reported to be as high as 50% for infected women in the first trimester.<sup>31</sup> Similarly, infection with the seasonal influenza virus is associated with higher miscarriage and maternal mortality rates than the non-infected population.<sup>1,29</sup> It should be noted however that miscarriage could be due to the maternal response to the infection, rather than a direct placental effect induced by the infection.<sup>32</sup> In the case of SARS-CoV-2, it is highly likely that the higher rate of miscarriage could be attributed to fever. In contrast to these viral infections, the outcomes for SARS-CoV-2 infected pregnant women in the first trimester are reassuring.

With regards to fetal and neonatal outcomes, pneumonias during pregnancies are associated with an increased risk of preterm birth compared to the general population.<sup>33</sup> Our findings are consistent with this observation. Fortunately, premature neonates fared well and had comparable Apgar scores to the full-term neonates included in the analysis. Unfortunately, the rate of stillbirth among this group was 2.34%, significantly higher than the rate among the general population. Intrauterine growth restriction (IUGR) was an expected finding in this group due to its prevalence in prior SARS infections during pregnancy.<sup>34</sup> However, there was little reporting on IUGR among fetuses in SARS-CoV-2 positive pregnancies.

Although our review includes a high number of patients, it is limited by the heterogeneity of the studies' primary outcomes and reporting methods. Secondly, the availability of few reports of severe outcomes makes it difficult to explain the pathophysiology of stillbirths and maternal morbidity in the setting of SARS-CoV-2. As more case reports become available, sub-group analyses will enable stronger understanding of the impact of the virus on maternal and fetal health.

## References

1. Mertz D, Geraci J, Winkup J, Gessner BD, Ortiz JR, Loeb M. Pregnancy as a risk factor for severe outcomes from influenza virus infection: A systematic review and meta-analysis of observational studies. *Vaccine*. 2017;35(4):521-528. doi:10.1016/j.vaccine.2016.12.012
2. Berkowitz K, LaSala A. Risk factors associated with the increasing prevalence of pneumonia during pregnancy. *Am J Obstet Gynecol*. 1990. doi:10.1016/0002-9378(90)91109-P
3. Brito V, Niederman MS. Pneumonia Complicating Pregnancy. *Clin Chest Med*. 2011;32(1):121-132. doi:10.1016/j.ccm.2010.10.004
4. Madinger NE, Greenspoon JS, Ellrodt AG. Pneumonia during pregnancy: Has modern technology improved maternal and fetal outcome? *Am J Obstet Gynecol*. 1989. doi:10.1016/0002-9378(89)90373-6
5. Liu D, Li L, Wu X, et al. Pregnancy and Perinatal Outcomes of Women With Coronavirus Disease (COVID-19) Pneumonia: A Preliminary Analysis. *Am J Roentgenol*. March 2020:1-6. doi:10.2214/ajr.20.23072
6. Nie. Clinical features and the maternal and neonatal outcomes of pregnant women with coronavirus disease 2019. doi:10.1101/2020.03.22.20041061
7. Breslin N, Baptiste C, Miller R, et al. COVID-19 in pregnancy: early lessons. *Am J Obstet Gynecol MFM*. March 2020:100111. doi:10.1016/j.ajogmf.2020.100111
8. Gidlöf S, Savchenko J, Brune T, Josefsson H. COVID-19 in pregnancy with comorbidities: More liberal testing strategy is needed. *Acta Obstet Gynecol Scand*. April 2020. doi:10.1111/aogs.13862
9. Liu Y, Chen H, Tang K, Guo Y. Clinical manifestations and outcome of SARS-CoV-2 infection during pregnancy. *J Infect*. March 2020. doi:10.1016/j.jinf.2020.02.028
10. Karami P, Naghavi M, Feyzi A, et al. Mortality of a pregnant patient diagnosed with COVID-19: A case report with clinical, radiological, and histopathological findings. *Travel Med Infect Dis*. 2020. doi:10.1016/j.tmaid.2020.101665
11. Lee DH, Lee J, Kim E, Woo K, Park HY, An J. Emergency cesarean section on severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) confirmed patient. *Korean J Anesthesiol*. March 2020. doi:10.4097/kja.20116
12. Li Y, Zhao R, Zheng S, et al. Lack of Vertical Transmission of Severe Acute Respiratory Syndrome Coronavirus 2, China. *Emerg Infect Dis*. 2020;26(6). doi:10.3201/eid2606.200287
13. Zhu H, Wang L, Fang C, et al. Clinical analysis of 10 neonates born to mothers with 2019-nCoV pneumonia. *Transl Pediatr*. 2020. doi:10.21037/tp.2020.02.06
14. Alzamora MC, Paredes T, Caceres D, Webb CM, Valdez LM, La Rosa M. Severe COVID-19 during Pregnancy and Possible Vertical Transmission. *Am J Perinatol*. April 2020. doi:10.1055/s-0040-1710050
15. Moro F, Buonsenso D, Moruzzi MC, et al. How to perform lung ultrasound in pregnant women with suspected COVID-19 infection. *Ultrasound Obstet Gynecol*. March 2020. doi:10.1002/uog.22028
16. Yu N, Li W, Kang Q, et al. Clinical features and obstetric and neonatal outcomes of pregnant patients with COVID-19 in Wuhan, China: a retrospective, single-centre, descriptive study. *Lancet Infect Dis*. March 2020. doi:10.1016/s1473-3099(20)30176-6
17. Librero J, Peiró S, Belda A, Calabuig J. Porcentaje de cesáreas en mujeres de bajo riesgo: Un indicador útil para comparar hospitales que atienden partos con riesgos diferentes. *Rev*

- Esp Salud Publica*. 2014;88(3):315-326. doi:10.4321/S1135-57272014000300002
18. Betran AP, Torloni MR, Zhang J, et al. What is the optimal rate of caesarean section at population level? A systematic review of ecologic studies. *Reprod Health*. 2015;12(1). doi:10.1186/s12978-015-0043-6
  19. He J, Liu ZW, Lu YP, et al. A Systematic Review and Meta-Analysis of Influenza A Virus Infection during Pregnancy Associated with an Increased Risk for Stillbirth and Low Birth Weight. *Kidney Blood Press Res*. 2017. doi:10.1159/000477221
  20. Fan C, Lei D, Fang C, et al. Perinatal Transmission of COVID-19 Associated SARS-CoV-2: Should We Worry? [published online ahead of print, 2020 Mar 17]. *Clin Infect Dis*. 2020;ciaa226. doi:10.1093/cid/ciaa226
  21. Chen S, Liao E, Shao Y. Clinical analysis of pregnant women with 2019 novel coronavirus pneumonia. *J Med Virol*. March 2020. doi:10.1002/jmv.25789
  22. Qi H, Luo X, Zheng Y, et al. Safe Delivery for COVID-19 Infected Pregnancies. *BJOG An Int J Obstet Gynaecol*. March 2020. doi:10.1111/1471-0528.16231
  23. Chen H. Clinical Characteristics and Intrauterine Vertical Transmission Potential of COVID-19 Infection in Nine Pregnant Women: A Retrospective Review of Medical Records - PubMed. [https://pubmed.ncbi.nlm.nih.gov/32151335/?from\\_term=pregnancy+OR+prenatal+AND+covid-19+OR+SARS-CoV2&from\\_filter=language.english&from\\_filter=years.2019-2020&from\\_pos=5](https://pubmed.ncbi.nlm.nih.gov/32151335/?from_term=pregnancy+OR+prenatal+AND+covid-19+OR+SARS-CoV2&from_filter=language.english&from_filter=years.2019-2020&from_pos=5). Accessed April 10, 2020.
  24. Li N, Han L, 2^ M, et al. Maternal and neonatal outcomes of pregnant women with COVID-19 pneumonia: a case-control study. doi:10.1101/2020.03.10.20033605
  25. Baud D, Greub G, Favre G, et al. Second-Trimester Miscarriage in a Pregnant Woman With SARS-CoV-2 Infection. *JAMA*. April 2020. doi:10.1001/jama.2020.7233
  26. Gynecologists. AC of O and. No Title. Novel Coronavirus 2019 (COVID-19). Washington, DC.
  27. No Title. Royal College of Obstetricians and Gynaecologists. Coronavirus (COVID-19) Infection in Pregnancy. London, UK: 2020.
  28. LoMauro A, Aliverti A. Respiratory physiology of pregnancy. *Breathe*. 2015;11(4):297-301. doi:10.1183/20734735.008615
  29. Jamieson DJ, Honein MA, Rasmussen SA, et al. H1N1 2009 influenza virus infection during pregnancy in the USA. *Lancet*. 2009;374(9688):451-458. doi:10.1016/S0140-6736(09)61304-0
  30. Creanga AA, Johnson TF, Graitcer SB, et al. Severity of 2009 pandemic influenza a (H1N1) virus infection in pregnant women. *Obstet Gynecol*. 2010;115(4):717-726. doi:10.1097/AOG.0b013e3181d57947
  31. Wong SF, Chow KM, Leung TN, et al. Pregnancy and perinatal outcomes of women with severe acute respiratory syndrome. *Am J Obstet Gynecol*. 2004;191(1):292-297. doi:10.1016/j.ajog.2003.11.019
  32. Giakoumelou S, Wheelhouse N, Cuschieri K, Entrican G, Howie SEM, Horne AW. The role of infection in miscarriage. *Hum Reprod Update*. 2016. doi:10.1093/humupd/dmv041
  33. Goodnight WH, Soper DE. Pneumonia in pregnancy. *Crit Care Med*. 2005;33(Supplement):S390-S397. doi:10.1097/01.CCM.0000182483.24836.66
  34. Ng PC, Leung C, Chiu WK, Wong SF, Hon EKL. SARS in newborns and children. In: *Biology of the Neonate*. Vol 85. Biol Neonate; 2004:293-298. doi:10.1159/000078174

## Legends

**Table 1:** Patient Characteristics and Symptoms in SARS-CoV-2 positive pregnancies

**Table 2:** Summary of Gestational Complications and Delivery Outcomes in SARS-CoV-2 positive pregnancies

**Table 1:** Patient Characteristics and Symptoms in SARS-CoV-2 positive pregnancies

Study	Number of patients	Maternal age (years)	Gestational age (weeks)	Pre-existing comorbidities	Maternal symptoms prepartum	Maternal symptoms postpartum	Imaging (CXR, CT)	Laboratory findings
Alzamora, 2020	1	41	33	Admitted due to resp insufficiency. PMH included diabetes mellitus (DM), and history of C-sections	Fever, malaise, fatigue, worsening dyspnea, tachycardia, tachypnea, metabolic acidosis	-	No CXR abnormalities  CT: multiple bilateral consolidations and bilateral pleural effusion	Elevated CRP, elevated ferritin, elevated D dimer, pancytopenia, lymphopenia Low Hb, WBCs, platelets, and creatinine.
Baud, 2020	1	28	19	Obesity  Admitted due to severe uterine contractions and a 5-cm dilated cervix at 19 weeks	Fever, fatigue, myalgia, dry cough, pain with swallowing, diarrhea	-	Physical examination did not show signs of pneumonia	
Breslin, 2020	2	38, 33	37 (both)	Patient 1 admitted for poorly-controlled type 2 DM and intrahepatic cholestasis of pregnancy	Negative for fever, cough, dyspnea, or sore throat	Patient 1: Fever  Patient 2: Cough that progressed to respiratory distress, fever, tachycardia,	Patient 2 CXR: mild pulmonary vascular congestion with no consolidation or effusion	-

				Patient 2 admitted for labor induction due to worsening chronic hypertension. PMH included DM and asthma		88% O2 saturation, dyspnea, diaphoresis		
Chen H, 2020	9	Mean 30 (Range 26-40)	36-39	Not remarkable	Fever (7), cough (4), myalgia (3), sore throat (2), and malaise (2), GI symptoms (1), dyspnea (1)	Fever (6)	CT: patchy ground glass shadows (8)	Elevated CRP (6), lymphopenia (5), and increased ALT and AST (3)
Chen S, 2020	5	Mean 28.8 (Range 25-31)	38-41	-	Cough (2), runny nose (1)	Fever (5)	CT: ground glass opacities bilaterally (3) or unilaterally (2)	Elevated CRP, decreased albumin, elevated alkaline phosphatase
Fan, 2020	2	34, 29	36, 37	-	Nasal congestion (2), fever (2), abdominal rash (1), sore throat (1), chills (1)	-	CT patient 1: patchy consolidation in both lungs CT patient 2: multiple patchy infiltrates on left lung	Lymphopenia (2)

Study	Number of patients	Maternal age (years)	Gestational age (weeks)	Pre-existing comorbidities	Maternal symptoms prepartum	Maternal symptoms postpartum	Imaging (CXR, CT)	Laboratory findings
Gidlof, 2020	1	34	36	Referred due to hypertension and proteinuria. Dichorionic twin pregnancy	Headache, hoarseness, malaise, high BP, photophobia, brisk patellar reflexes	87% O2 saturation	CT: typical signs of SARS-CoV-2 pneumonia	Elevated uric acid
Karami, 2020	1	27	30+3	None	Fever, cough, myalgia, Respiratory Distress	Multi Organ Failure and maternal death	CXR: faint bilateral patchy opacities CT: faint subpleural ground-glass opacities with pleural thickening	Leukopenia, lymphopenia, Elevated CPR, creatinine, and LDH
Lee, 2020	1	28	36+2	-	Fever, cough, sore throat. Nausea and low BP following spinal anesthesia	-	CXR: left lower/middle lobe consolidation and increased vascular marking CT: multifocal peribronchial ground glass appearance and consolidation in left lower lobe	Normal CRP, elevated erythrocyte sedimentation rate, low Hb

Li N, 2020	16**	Mean: 30.9 (Range 26-37)	33-40	Hypertension, hepatitis B infection, , polycystic ovary syndrome	Fever (4)	Fever (8)	Typical images of pneumonia bilaterally (10) and unilaterally (7)	Leukopenia, low neutrophils, low CRP, and low ALT
Li Y, 2020	1	30	35	-	Dry cough, chills, or dyspnea	-	CXR: scattered multiple patchy infiltrates in both lungs	Slightly abnormal
Liu D, 2020	15	23-40 (mean 32)	12-38	Mitral and tricuspid valve replacement	Fever (13), cough (9), sore throat (1), dyspnea (1), myalgia (3), fatigue (4), diarrhea (1)	Fever (1)	CT: ground-glass opacity (GGO). consolidations reported with disease progression	Lymphocytopenia (12), elevated CRP (10)
Liu H, 2020	41***	Mean: 30	22-40+5	Hepatitis B (1)	Fever (16), cough (15), dyspnea (5), fatigue(5), loss of appetite (3)	Fever (14)	CT: Mixed GGO with consolidation	Elevated CRP (27), lymphopenia (25), leukocytosis (17), elevated neutrophils (34)
Liu W, 2020	3	34, 34, 30	40, 38+4, 39+5	Patient 1: hypothyroidism	Fever (2), cough (2)	-	CT: abnormal in all 3 patients	Leukocytosis (2), elevated neutrophils (2), lymphocytosis (1), low

				and epiglottic cysts Patient 2: none Patient 3: gestational diabetes				eosinophils (1), low platelets (1)
Study	Number of patients	Maternal age (years)	Gestational age (weeks)	Pre-existing comorbidities	Maternal symptoms prepartum	Maternal symptoms postpartum	Imaging (CXR, CT)	Laboratory findings
Liu Y, 2020	13	Mean: 29.7 (Range 22-36)	2 women less than 28 GA (25, 27). 11 in third trimester (32 - 36)	None	Fever (10), cough (2), fatigue (4), dyspnea (3), sore throat (1)	1 patient had MODS including ARDS	-	-
Nie, 2020	33	Mean: 30.5 (Range 24-36)	3 women in 2nd trimester (17, 20, 26), 30 women in 3rd trimester	15 women had pre-existing chronic disorders	Fever (21), dry cough(13), fatigue(7), dyspnea (7)  Less common: diarrhea, muscle ache, sore throat, chest pain,	Fever (5)	CT: bilateral or unilateral pneumonia, GGO	-

					stomachache 4 asymptomatic			
Vlachodimitropoulou, 2020	2	40, 23	35+3, 35+2	Patient 1: neutropenia, mild respiratory infections Patient 2: not reported	Pt 1: fever, cough, tachycardia Pt 2: fever, cough	-	Pt 1: normal CXR Pt 2: not reported	Pt 1: progressive thrombocytopenia, declining fibrinogen, rising APTT, elevated CRP, low neutrophil count, lymphocytopenia, elevated D dimer  Pt 2: thrombocytopenia, prolonged APTT, transaminitis, progressive coagulopathy, elevated CRP, lymphocytopenia, elevated D dimer
Wen, 2020	1	31	30	-	Mild diarrhea	Recovered and discharged prior to delivery	CT: patchy consolidation and GGO	Not reported
Yu, 2020	7	Mean: 32 (Range 29 - 34)	39+1 (37 - 41+2)	2 patients had chronic diseases (hypothyroidism and polycystic ovary syndrome) 3	Fever (6), cough (1), dyspnea (1), diarrhea (1)	-	CT: bilateral pneumonia (6), unilateral pneumonia (1)	Elevated CRP (7), elevated neutrophils (5), lymphopenia (5), low platelets (2), elevated D-

								dimer (7), abnormal liver enzymes (2)
Zhu, 2020	9	Mean: 30 (Range 25-35)	31-39	-	Fever (8), cough (4), diarrhea (1), sore throat (1), cholecystitis (1)	-	CT: typical of viral pneumonia. Lesions merged into strips as disease progressed	-
<b>Total n</b>	<b>164</b>							

\*Mixed diagnosis: 25 clinically-diagnosed and 16 laboratory-diagnosed

\*\*Compared with non-SARS-CoV-2 pneumonia pregnant women

\*\*\*Compared with 14 non-pregnant patients

- in Table 1 is used to refer to information that was not reported in the publication

**Table 2:** Summary of Gestational Complications and Delivery Outcomes in SARS-CoV-2 positive pregnancies

Study	Number of patients	Delivery Method	Gestational Complications	Neonatal Outcomes
Alzamora, 2020	1	1 C-Section due to maternal respiratory compromised status	None	1/1 had Apgars 6,8 1/1 neonate positive for SARS-CoV-2 infection on hour 16 of life
Baud, 2020	1	1 vaginal delivery (at 19 weeks due to cervical dilation and severe contractions)	Miscarriage with abnormal placental histopathology suggestive of a fetal inflammatory response	1 stillbirth Placenta tested positive for SARS-CoV-2 infection
Breslin, 2020	2	2 C-sections due to obstetric indication	None	2/2 neonates negative for SARS-CoV-2 Infection
Chen H, 2020	9	9 C-sections due to uncertainty about the risk of intrapartum mother-to-child transmission by vaginal delivery	1 gestational hypertension 1 pre-eclampsia 2 premature rupture	4 preterm neonates 2 fetal distress
Chen S, 2020	5	3 vaginal deliveries 2 C-sections, including 1 emergency C-section due to fetal tachycardia	2 gestational diabetes 1 preeclampsia	5/5 had Apgars of 10,10
Fan, 2020	2	2 C-sections, one of which due to	-	2/2 had Apgars 9,10

		persistent fevers		1 fever with diffuse haziness in both lungs fields on CT scan  2/2 show no evidence of transmission in newborn, maternal serum, placental tissue, umbilical cord blood, amniotic fluid, maternal vaginal swab, and mother's breast milk
Gidlof, 2020	1	1 emergency C-section, twin pregnancy	1 gestational diabetes	2/2 (twins) had Apgars of 9,10  1/2 had breathing problems soon after birth that resolved with CPAP.
Karami, 2020	1	1 vaginal delivery due to maternal body movement under sedation following maternal respiratory distress	Severe systolic dysfunction with reduced ejection fraction, multi-organ failure, and death	1/1 had Apgars of 0,0 1 stillbirth
Lee, 2020	1	1 Emergency C-section due to obstetric indication	None	1/1 had Apgars 9, 10  1/1 neonate negative for SARS-CoV-2 Infection
Li N, 2020	16	2 vaginal deliveries 14 C-sections	3 gestational diabetes 3 gestational hypertension 1 preeclampsia 1 placenta previa 1 placental bleeding 1 premature rupture	5 preterm neonates
Li Y, 2020	1	1 emergency C-section	-	1/1 newborn without complications

Study	Number of patients	Delivery Method	Gestational Complications	Neonatal Outcomes
Liu D, 2020	15	1 vaginal delivery 10 C-sections	1 Gestational diabetes 1 complete placenta previa	1/1 neonate negative for SARS-Cov2 infection. 1/1 shows no evidence for maternal serum, urine, feces, amniotic fluid, umbilical cord blood and placenta, and breast milk samples transmission.  11/11 had Apgar scores higher than 8
Liu H, 2020	41	No delivery data reported	4 gestational diabetes 3 gestational hypertension	16 neonates reported on have no abnormalities
Liu W, 2020	3	1 vaginal delivery 2 C-sections due to SARS-CoV-2	1 gestational diabetes 1 chorioamnionitis	1 Chronic fetal distress in utero
Liu Y, 2020	13	1 vaginal delivery 10 C-sections, including 5 emergency procedures due to fetal distress, premature rupture of the membrane and stillbirth	1 premature rupture	1 stillbirth 6/10 preterm (<36 weeks of gestation) 9/10 had Apgars 10
Nie, 2020	33	5 vaginal deliveries 22 C-sections	The rate of obstetrical complications was reported at 22.2%	10/28 preterm 1/28 neonate developed ARDS (then recovered)  27/27 had Apgars range 8-19, Apgar 9-10 at minute 5

				1/26 newborns tested positive for SARS-CoV-2
Vlachodimitr opoulou, 2020	2	2 C-sections, including 1 emergency C - section, due to maternal coagulopathy and fetal heart rate problem,	1 gestational diabetes	1/2 had Apgars 9, 10 1/2 had Apgars 4,2,7
Yu, 2020	7	7 C-sections	-	1/3 positive for SARS-COV2 Infection, with no evidence of vertical transmission; All 7 neonates had Apgar scores of no lower than 8 at 1 minute, no lower than 9 at 5 min.
Zhu, 2020	9	2 vaginal deliveries 7 C-sections	-	6/10 preterm  8/9 had Apgars 8-10 at minute 1 and 5. 1/9 had Apgars 7,8.  9/9 negative for SARS-Cov-2  6/10 had shortness of breath, 2/10 had fevers  1 neonate death (born at GA 34 weeks, refractory shock and MOF eight days later.)
<b>Total</b>	<b>163 pregnancies</b> <b>*</b>	<b>110 deliveries</b>	<b>39 total counts of complications (including overlapping patients)</b>	<b>128 newborns**</b>

\*The number of studies in Table 2 excludes one study that was included in Table 1. That study did not provide any information on delivery outcome, neonatal outcome, or gestational complications.

\*\*The number of newborns is different from the number of pregnancies due to certain studies not following the delivery to birth, and due to twin pregnancies.

- in Table 2 is used to refer to information that was not reported in the publication