1   # Anti-Spike, anti-Nucleocapsid and neutralizing antibodies in
2   # SARS-CoV-2 hospitalized patients and asymptomatic carriers
3

4   Etienne Brochot[1,2,*], Baptiste Demey[1,2], Antoine Touzé[3], Sandrine Belouzard[4], Jean Dubuisson[4],
5   Jean-Luc Schmit[1,2], Gilles Duverlie[1,2], Catherine Francois[1,2], Sandrine Castelain[1,2], Francois Helle[2,*]
6
7   [1] Department of Virology, Amiens University Medical Center, Amiens, France.
8   [2] AGIR Research Unit, UR UPJV 4294, Jules Verne University of Picardie, Amiens, France.
9   [3] ISP1282 INRA University of Tours, Tours, France.
10 [4] Université Lille, CNRS, INSERM, CHU Lille, Institut Pasteur de Lille, U1019-UMR 8204-CIIL-Center
11 for Infection and Immunity of Lille, F-59000 Lille, France.
12
Corresponding authors :
Dr Etienne Brochot
Laboratoire de Virologie
Centre de Biologie Humaine - CHU Amiens
F-80054 Amiens cedex 1, France
Phone: +33-322-080-764
Fax: +33-322-087-009
E-mail: etienne.brochot@u-picardie.fr
21
Dr Francois Helle
UR-UPJV 4294, Agents Infectieux, Résistance et Chimiothérapie
Centre Universitaire de Recherche en Santé,
30 Avenue de la Croix Jourdain,
80000 Amiens, France
Phone: +33-322-825-351
E-mail: francois.helle@u-picardie.fr
29
30
31

Abstract

Using longitudinal plasma samples from thirty COVID-19 patients, we observed that virus-specific

antibodies are detectable in 100% of patients two weeks after symptom onset. We also show that

these patients produced variable levels of neutralizing antibodies which reached a plateau two weeks

after symptom onset and then declined in the majority of patients. Furthermore, we report that

neutralizing antibodies were undetectable in 56% (14/25) of asymptomatic carriers.

Keywords

SARS-CoV-2; COVID-19, coronavirus, spike, nucleocapsid, RBD, neutralizing antibodies

The Severe Acute Respiratory Syndrome Coronavirus 2 (SARS-CoV-2) has recently emerged

and caused a human pandemic of coronavirus disease 2019 (COVID-19). Among the four SARS-CoV-

2 structural proteins, the Spike (S) and the Nucleocapsid (N) proteins are the main immunogens[1]. The

S protein consists of two subunits, S1 which contains the receptor-binding domain (RBD) and S2. The

kinetics of antibody detection is essential for the selection of commercial serological assays and the

interpretation of the results. Some manufacturers have decided to target the S1 and/or the S2

whereas others chose the RBD or the N protein. Furthermore, the neutralizing antibody (NAb)

response against the SARS-CoV-2 remains poorly understood and it is still unknown whether cured

patients are protected against new infection[2].

In order to accurately assess the time of seroconversion, we used 151 samples from 30

patients hospitalized at the Amiens University Hospital for a COVID-19 (see Supplementary Table 1)

to monitor the kinetics of detection of anti-S1, anti-S2, anti-RBD and anti-N antibodies with in-house

ELISAs. We observed that antibodies targeting the N protein and the RBD were the earliest to be

detected (Fig. 1a). Thirteen days post-symptom onset, 100% of patients had detectable antibodies to

both proteins. A similar profile was observed for anti-S2 antibodies but with a mean time lag of two

days. Antibodies to the S1 subunit were the last to be detected and remained undetectable for two

patients. High levels of anti-N and anti-RBD antibodies were detected in the large majority of samples

obtained fourteen days post-symptom onset whereas very heterogeneous levels of anti-S1 antibodies

were found in the same samples (Fig. 1b). The positive correlations between each ELISA are shown

in Extended Data Fig. 1. Significant differences were observed between intensive care versus non-

intensive care patients for anti-S1, anti-S2 and anti-N antibody levels, from eight days post-symptom

onset (Extended Data Fig. 2a). A slight difference was observed for anti-N antibody levels according to

the sex, from fourteen days post-symptom onset (Extended Data Fig. 2b). In contrast, no significant

difference was observed according to age, from fourteen days post-symptom onset (Extended Data

Fig. 2c).

We also monitored the presence of NAbs in all plasma samples using retroviral particles

pseudotyped with the S glycoprotein of the SARS-CoV-2 (SARS-CoV-2pp)[3]. The results obtained for

each patient are presented in Extended Data Fig. 3. One sample of each patient was also used to

perform dose-response curves with particles pseudotyped with the G glycoprotein of the Vesicular

Stomatitis Virus and no inhibition was observed, demonstrating that the neutralization observed with

the COVID-19 patient plasmas was specific to the SARS-CoV-2 (Extended Data Fig. 4a).

Furthermore, plasmas from twelve patients that had previously been infected with other coronaviruses

(OC43 (n=5), 229E (n=4), NL63 (n=2) or HKU1 (n=1)) did not have any effect on SARS-CoV-2

pseudotype infectivity (Extended Data Fig. 4b and Supplementary Table 2). As expected, our results

demonstrate that the production of NAbs correlates with the production of antibodies targeting the S1,

S2 and RBD domains and we detected NAbs in all COVID-19 patients fifteen days post-symptom

onset (Fig. 1a). The NAb titers increase from one week post-symptom onset and reaches a plateau

one week after (Fig. 2a, 2b and Extended Data Fig. 3). However, the NAb titers reached were variable

between patients, 17% generated low levels of NAbs (40 ≤ titers < 160), 73% intermediate levels (160

≤ titers < 1280) and 10% high levels (1280 ≤ titers) (Fig. 2a). Positive correlations between NAb titers

and anti-S1, anti-S2, anti-N or anti-RBD antibody levels, as well as white blood cells and lymphocytes

counts are shown in Extended Data Fig. 5. Significantly higher NAb titers were observed in patients

with severe forms (p=0.04) and in women (p=0.03) from 14 days post-symptoms onset (Extended

Data Fig. 6). In contrast, no significant difference was observed according to the age. We also had the

opportunity to monitor the presence of NAbs in late samples of twelve patients (≥40 days post-

symptom onset) and we observed that the NAb titer dropped to low or undetectable level in most of

these samples (Fig. 2b and Extended data Fig. 3).

Finally, we monitored the presence of NAbs in plasma samples from 25 asymptomatic carriers.

It is important to note that we could not establish when these patients had been infected since they

were asymptomatic but it probably occurred more than one week before sampling since they were

confirmed seropositive using commercial serological assays and our in-house ELISAs (Supplementary

Table 3). As shown in Fig. 2c and Extended Data Fig. 7, NAbs were below the detection limit of our

assay in the majority of these plasma samples (56%, 14/25). Low NAb levels (40 ≤ titers < 160) were

found in 28% of these patients (7/25). Three patients had intermediate NAb levels (160 ≤ titers < 1280)

and only one showed a high NAb titer (≥1280).

Commercial serological assays that are complementary to direct viral detection of the SARS-

CoV-2 by RT-PCR have recently become available but they need to be finely evaluated. With our four

in-house ELISAs, we showed that the detection of the RBD and the N protein may be more suitable

since it was highly or slightly more sensitive than the detection of S1 or S2, respectively. We only

tested IgG detection since recent data showed that anti-SARS-CoV-2 IgG levels increase at the same

time or earlier than IgM levels[4]. We also report that COVID-19 patients generate variable levels of

NAbs that are likely to drop a few weeks after infection and that most of asymptomatic carriers do not

generate NAbs. Thus, our results raise questions concerning the role played by NAbs in COVID-19

cure and protection against secondary infection. They also imply that anti-SARS-CoV-2 NAbs should

be titrated to optimize convalescent plasma therapy. Finally, they suggest that induction of NAbs is not

the only strategy to adopt for the development of a vaccine.

Methods

Study population and samples. Thirty patients diagnosed SARS-CoV-2 positive by RT-PCR on a nasopharyngeal swab sample, between 25 February and 23 March 2020, were enrolled in the study. The general information was extracted from electronic medical records and the clinical characteristics of the 30 patients are described in Supplementary Table 1. Samples from patients diagnosed positive for the human coronaviruses NL63, 229E, HKU1 and OC43 were also tested (Supplementary Table 2). Finally, we also used samples from 25 asymptomatic carriers (Supplementary Table 3) that were diagnosed positive using commercial serological tests (LIAISON® SARS-CoV-2 IgG from DiaSorin and/or ELISA SARS-CoV-2 (IgG) from EUROIMMUN). This study was approved by the institutional review board of Amiens University Hospital (number PI2020_843_0046, 21 April 2020).

In-house ELISAs. All plasmas were decomplemented at 56°C for 30 min. MaxiSorp Nunc-immuno 96-well plate were coated with a 1 µg/mL solution of SARS-CoV-2 S1, S2, RBD or N antigen (The Native Antigen Company, United Kingdom), overnight at 4°C. Wells were blocked with 1% fetal bovine serum for 1 hour at 37°C. Then 100 µL of diluted plasmas (1:100 for S1, S2, RBD and1:200 for N) were added and incubated for 1 h at 37°C. After washing 4 times, plates were incubated with peroxydase conjugated mouse anti-human IgG (Southern Biotech, 1/6000). After 4 washes, 100 µL of o-phenylenediamine peroxidase substrate was added at room temperature in the dark. After 15 minutes, the reaction was stopped with H_2SO_4 solution. The optical density was measured at 490 nm. All samples were run in triplicate. To establish the specificity of each assay, 40 pre-pandemic sera from 2019 were tested. Each cut-off values were defined as the means plus 3 standard deviations obtained with these samples.

Neutralization assay. SARS-CoV-2pp were produced as described previously[3] with a plasmid encoding a human codon-optimized sequence of the SARS-CoV-2 spike glycoprotein (accession

number: MN908947). Supernatants containing the pseudotyped particles were harvested at 48, 72

and 96 h after transfection, pooled and filtered through 0.45-μm pore-sized membranes. All plasmas

were decomplemented at 56°C for 30 min. Neutralization assays were performed by preincubating

SARS-CoV-2pp and diluted plasma for 1 h at room temperature before contact with Vero cells

(ATCC® CCL-81™) that were transiently transfected with the plasmids pcDNA3.1-hACE2 and

pcDNA3.1-TMPRSS2 48 h before inoculation. Luciferase activities were measured 72 h post-infection,

as indicated by the manufacturer (Promega). The NAbs titers were defined as the highest dilution of

plasma which resulted in a 90% decrease of the infectivity. Retroviral particles pseudotyped with the G

glycoprotein of the Vesicular Stomatitis Virus (VSVpp) were used to control the specificity of the

neutralization.

Statistical analysis. Quantitative variables were expressed as the median and compared using

Student's t-test. The Pearson correlation coefficient was used to measure the strength of a linear

association between two quantitative variables. Statistical analyses were performed using GraphPad

Prism 5. A two-sided P-value < 0.05 was considered statistically significant.

Acknowledgments

We thank the study participants who donated their blood samples for this project, and the healthcare

workers who are caring for COVID-19 patients. This work was supported by Flash Covid-19 funds

from the Amiens University Hospital and the French "Agence Nationale de la Recherche".

Author Contributions

EB, GD, CF, SC and FH designed the research and analyzed the data ; JLS collected clinical

samples ; BD extracted clinical information from electronic medical records ; AT, SB and JD provided

key reagents ; EB and FH performed the experiments ; EB and FH wrote the manuscript.

Competing Interests statement

The authors declare no competing interests.

References

1. Meyer, B., Drosten, C., Müller, M.A. Serological assays for emerging coronaviruses: challenges and pitfalls. *Virus Res.***194**, 175-83 (2014). https://doi.org/10.1016/j.virusres.2014.03.018

2. Ota, M. Will we see protection or reinfection in COVID-19? *Nat Rev Immunol.* (2020) https://doi.org/10.1038/s41577-020-0316-3

3. Millet, J.K., Whittaker, G.R. Murine Leukemia Virus (MLV)-based Coronavirus Spike-pseudotyped Particle Production and Infection. *Bio Protoc.* **6**(23), e2035 (2016). https://doi.org/10.21769/BioProtoc.2035

4. To, K.K., Tsang, O.T., Leung, W.S., Tam, A.R., Wu, T.C., Lung, D.C., Yip, C.C., Cai, J.P., Chan, J.M., Chik, T.S., Lau, D.P., Choi, C.Y., Chen, L.L., Chan, W.M., Chan, K.H., Ip, J.D., Ng, A.C., Poon, R.W., Luo, C.T., Cheng, V.C., Chan, J.F., Hung, I.F., Chen, Z., Chen, H., Yuen, K.Y. Temporal profiles of viral load in posterior oropharyngeal saliva samples and serum antibody responses during infection by SARS-CoV-2: an observational cohort study. *Lancet Infect Dis.* **20**(5), 565-574 (2020). https://doi.org/10.1016/S1473-3099(20)30196-1

Supplementary Table 1. Characteristics of the 30 hospitalized patients included in the study

Patient characteristics		Mild Disease (n=12)	Severe disease (n=18)
Female		6 (50%)	6 (33%)
Male		6 (50%)	12 (66%)
Median Age (Years)		77 (59-87)	63 (38-78)
Chronic comorbidities	Hypertension	7	9
	Chronic heart disease	3	3
	Chronic lung disease	3	0
	Chronic kidney disease	2	0
	Diabetes	3	4
	Hyperlipidemia	2	1

Supplementary Table 2. Patients previously infected with endemic coronaviruses used as control in the study

Patient	Coronavirus strain	Days post-diagnostic	Detection of IgG in our in house ELISA assays				SARS-CoV-2 NAb titer
			S1	S2	RBD	NP	
C1	OC43	174	-	-	+/-	-	<40
C2	229E	39	-	-	-	-	<40
C3	NL63	561	-	-	-	-	<40
C4	OC43	170	-	-	-	-	<40
C5	OC43	176	-	-	-	-	<40
C6	NL63	16	-	-	-	-	<40
C7	229E	14	-	-	+	+	<40
C8	HKU1	24	-	-	-	-	<40
C9	OC43	301	+/-	-	-	+/-	<40
C10	229E	56	-	-	-	-	<40
C11	229E	761	-	-	-	+/-	<40
C12	OC43	86	-	-	-	-	<40

+, positive ; -, negative ; +/-, equivocal

Supplementary Table 3. SARS-CoV-2 asymptomatic carriers included in the study

Patient	Detection of IgG in in house ELISAs			Diasorin assay	Euro-immun assay	SARS-CoV-2 NAb titer
	S1	S2	NP			
AC1	+	+	+	+	+/-	40
AC2	+	+/-	+	-	+	<40
AC3	+	+	+	+	+	40
AC4	+/-	+	+	+	-	<40
AC5	-	+	+	+	-	<40
AC6	+/-	+	+	+	+/-	<40
AC7	+	+	+	+	+	160
AC8	+	+	+	+	+	640
AC9	+	+	+	+	+	<40
AC10	+	+	+	+	+	80
AC11	+	+	+	-	+	<40
AC12	-	+	+	+	-	<40
AC13	+	+/-	+	-	+	<40
AC14	+	+	+	+	+	480
AC15	+	+	+	+/-	+	<40
AC16	+	+	+	+	+	120
AC17	+	+	+	+	+	80
AC18	+	+	+	+	+	5120
AC19	+	+	+	+	+	<40
AC20	+	+	+	+	+	<40
AC21	+	+	+	+	+	60
AC22	+	+	+	+	+	60
AC23	+	+	+	+	+	<40
AC24	+	+	+	+	+	<40
AC25	+	+	+	+	+	<40

+, positive ; -, negative ; +/-, equivocal

Figure Legends

Fig. 1 | Antibody response in COVID-19 patients. a, Kinetics of anti-S1, anti-S2, anti-RBD, anti-N

and NAb detection in 30 COVID-19 patients after symptom onset. **b**, Evolution of the anti-S1, anti-S2,

anti-RBD and anti-N antibody levels during the first month post-symptom onset.

Fig. 2 | NAb response in COVID-19 patients and SARS-CoV-2 asymptomatic carriers.

a, Evolution of the NAb titer in 30 COVID-19 patients during the first month post-symptom onset.

b, Evolution of the NAb titer in 12 COVID-19 patients after more than 40 days post-symptom onset.

The dashed line indicates the cut-off of the assay. **c**, Determination of the NAb titer in plasma samples

from 25 asymptomatic carriers.

Extended Data Fig. 1 | Correlations between anti-S1, anti-S2, anti-RBD and anti-N levels in in-

house ELISAs. a, anti-S1 versus anti-S2. **b**, anti-S1 versus anti-RBD. **c**, anti-S1 versus anti-N. **d**,

anti-S2 versus anti-RBD. **e**, anti-S2 versus anti-N. **f**, anti-RBD versus anti-N. Dashed lines indicate

assay cut-offs for positivity. OD, optical density.

Extended Data Fig. 2 | Temporal profiles of anti-S1, anti-S2, anti-RBD and anti-N antibody

levels. Patients samples were divided into three periods groups (day 0-7, day 8-14 and day >14). **a**,

The temporal profiles are presented according to the severity of the disease (SD, severe disease

requiring intensive care ; MD, mild disease). **b**, The temporal profiles are presented according to the

sex (M, male ; F, female). **c**, The temporal profiles are presented according to the age (< or > 60 years

old). Dashed lines indicate assays cut-offs for positivity and lines indicate the median for each assay.

OD, optical density. NS, not significant ; *, $p<0.05$; **, $p<0.01$.

Extended Data Fig. 3 | NAb response to SARS-CoV-2 in COVID-19 patients. SARS-CoV-2pp were

preincubated with serially diluted plasma obtained from 30 COVID-19 patients (#1 to #30) at different

days post-symptom onset (d2 to d58). Dose response curves represent the means of normalized

infectivity (%) from two independent experiments performed in duplicate. Error bars have been omitted

for clarity.

Extended Data Fig. 4 | Specificity of the neutralization assay. a VSVpp were preincubated with

serially diluted plasma obtained from 30 COVID-19 patients (#1 to #30). Dose response curves

represent the means of normalized infectivity (%) from two independent experiments performed in

duplicate. **b** SARS-CoV-2pp were preincubated with serially diluted plasma obtained from 12 patients

infected with 229E, NL63, HKU1 or OC43 coronaviruses (C1 to C12). Dose response curves

represent the means of normalized infectivity (%) from two independent experiments performed in

duplicate. Error bars have been omitted for clarity.

Extended Data Fig. 5 | Correlations between NAb titers and anti-S1, anti-S2, anti-RBD and anti-N antibody levels as well as white blood cells counts or lymphocyte counts. a, NAb titer versus anti-S1. **b**, NAb titer versus anti-S2. **c**, NAb titer versus anti-RBD. **d**, NAb titer versus anti-N. **e**, NAb titer versus white blood cells counts. **f**, NAb titer versus lymphocyte counts. Dashed lines indicate assay cut-offs for positivity. OD, optical density.

Extended Data Fig. 6 | Temporal profiles of NAb titers. Patients samples were divided into three periods groups (day 0-7, day 8-14 and day >14). **a**, The temporal profiles are presented according to the severity of the disease (SD, severe disease requiring intensive care ; MD, mild disease). **b**, The temporal profiles are presented according to the sex (M, male ; F, female). **c**, The temporal profiles are presented according to the age (< or > 60 years old). Dashed lines indicate assays cut-offs for positivity and lines indicate the median for each assay. NS, not significant ; *, p<0.05 ; **, p<0.01.

Extended Data Fig. 7 | SARS-CoV-2 NAbs in asymptomatic carrier samples. SARS-CoV-2pp were preincubated with serially diluted plasma obtained from 25 asymptomatic carriers that were confirmed with commercial serological assays (AC1 to AC25). Dose response curves represent the means of normalized infectivity (%) from two independent experiments performed in duplicate. Error bars have been omitted for clarity.

a

b

271   **Fig. 1**

a

b

c

Fig. 2

Extended Data Fig. 1

282   **Extended Data Fig. 2a**

286 **Extended Data Fig. 2b**

290 **Extended Data Fig. 2c**

Plasma dilution factor (Log$_{10}$)

294   **Extended Data Fig. 3**

297   **Extended Data Fig. 4**

301   **Extended Data Fig. 5**

a

b

c

305 **Extended Data Fig. 6**

308 **Extended Data Fig. 7**