Environmental indicator for effective control of COVID-19 spreading

Xinbo Lian[1], Jianping Huang[1,2,*], Li Zhang[1],

Chuwei Liu[1], Xiaoyue Liu[1], Lina Wang[3]

[1] Collaborative Innovation Center for Western Ecological Safety, College of Atmospheric Sciences, Lanzhou University, Lanzhou, 730000, China.

[2] CAS Center for Excellence in Tibetan Plateau Earth Sciences，Beijing 100101,China.

[3] Gansu Province Environmental Monitoring Center, Lanzhou, 730000, China.

Corresponding Author ：

Dr. Jianping Huang（hjp@lzu.edu.cn）

Abstract

Recently, a novel coronavirus (COVID-19) has caused viral pneumonia worldwide, spreading to more than 200 countries, posing a major threat to international health. To prevent the spread of COVID-19, in this study, we report that the city lockdown measure was an effective way to reduce the number of new cases, and the nitrogen dioxide (NO_2) concentration can be adopted as an environmental lockdown indicator. In China, after strict city lockdown, the average NO_2 concentration decreased 55.7% (95% confidence interval (CI): 51.5-59.6%) and the total number of newly confirmed cases decreased significantly. Our results also indicate that the global airborne NO_2 concentration steeply decreased over the vast majority of COVID-19-hit areas based on satellite measurements. We found that the total number of newly confirmed cases reached an inflection point about two weeks after the lockdown. The total number of newly confirmed cases can be reduced by about 50% within 30 days of the lockdown. The stricter lockdown will help newly confirmed cases to decline earlier and more rapidly. Italy, Germany and France are good examples. Our results suggest that NO_2 satellite measurement can help decision makers effectively monitor control regulations to reduce the spread of COVID-19.

1. Introduction

Large-scale COVID-19 viral pneumonia through human-to-human transmission poses a severe and acute public health emergency (Huang et al., 2020a; Li et al., 2020). As the epidemic worsened, most countries imposed city lockdown and quarantine measures to reduce transmission to control the epidemic. The Chinese government has gradually implemented a city-wide quarantine of Wuhan and several surrounding cities as of 23 January, flights and trains to and from Wuhan have been suspended, and public transport has been halted (Cyranoski et al., 2020; Wu et al., 2020). The entire northern Italy was quarantined since 9 March 2020, and three days later the government extended it to the whole country (Paterlini, 2020).The Spanish government declared a 15-day national emergency, starting on 15 March (Legido-Quigley et al., 2020). In the United States, on 19 March, California became the first state to order a lockdown (Baccini et al., 2020). In Germany, since 18 March, 16 states have closed, public gatherings of more than two people have been banned and most shops except supermarkets and pharmacies have closed (Dehning et al., 2020). On 23 March, the British government announced a new nationwide restriction allowing residents to only venture outside when absolutely necessary, e.g., to work, buy necessities (Iacobucci, 2020).

The worldwide lockdown, which was imposed to stop the spread of the novel coronavirus, not only caused an economic downturn but also appeared to result in cleaner air in urban areas usually heavily affected by pollution (Lian et al., 2020; Schiermeier, 2020). The most important measure of the lockdown policy was the reduction of traffic and control personnel flow, and traffic pollution is an important factor influencing air quality and public health. Vehicle exhaust and evaporation emissions are the main emission sources of ozone and secondary particle

precursors near the ground in cities and regions (Zhou et al., 2019; Yuan et al., 2013), and the

spatial variation of nitrogen dioxide (NO_2), fine particulate matter ($PM_{2.5}$) and black carbon (BC)

may also be significant affected by traffic flow density (Clougherty et al., 2013). A study in Los

Angeles showed that nitrogen oxides (NO_x) were identified as a source of pollution for light

vehicles, with NO_2, NO_x, carbon dioxide (CO_2), BC, and fine particle number (PN_{fine}) identified

as diesel exhaust sources (Tessum et al., 2018; Fan et al., 2018). In South Korea, source analysis

studies have shown that there is a high correlation between estimated traffic volume and NO_2

concentration (Kim et al., 2015). NO_2 levels can be used as a proxy for exposure to

traffic-related composite air pollution and to assess the impact of scenarios designed to reduce

traffic-related emissions (Brnnum-Hansen et al., 2018; Johansson et al., 2017).

In this report, we study the parameters of environmental indicators for city lockdown. Using

the automatic ground detection data and satellite data to analyze the trend of lockdown and the

total confirmed new cases in major cities in China, and using satellite data to further study the

impact of lockdown on virus transmission in countries mainly severely affected by the epidemic,

in order to help policymakers to formulate effective control measures to reduce the spread of

COVID-19.

2. Data and measurement

The ground observation daily data were provided by the China National Environmental

Monitoring Centre (http://www.cnemc.cn/). The data from January 24, 2020, to February 23,

2020, are selected as the representative data after the lockdown in Hubei, and the data from

December 24, 2019, to January 23, 2020, are selected as the representative data before the

lockdown (Fig. 1). The NO_2 ground observation data of China is from 1 January 2020 to 1

March 2020. The average concentration of major cities with severe epidemic diseases was

selected as the representative of NO_2 concentration of China, including Wuhan, Nanchang,

Guangzhou, Hangzhou, Changsha, Beijing, Shanghai, Hefei and Zhengzhou (Fig. 3). All

monitoring instruments of the air quality automatic monitoring system operate automatically 24

h a day. The monitoring items are $PM_{2.5}$, PM_{10}, SO_2, NO_2, and CO. The automatic monitoring of

$PM_{2.5}$ and PM_{10} adopts the micro-oscillating balance method and the β-absorption method,

respectively (ambient air quality standards, GB 3095-2012). SO_2 was determined by the

ultraviolet fluorescence method, NO_2 by the chemiluminescence method, CO by the

nondispersion infrared absorption method and gas filter correlation infrared absorption method.

This paper adopted the level 3 daily global gridded ($0.25°×0.25°$) nitrogen dioxide product

(OMNO2d) provided by the Ozone Monitoring Instrument (OMI) onboard the Aura satellite as

the daily NO_2 data, which can be obtained from GES DISC

(https://disc.gsfc.nasa.gov/datasets/OMNO2d_003). The Aura satellite was launched by NASA

on July 15, 2004, with its overall objective of monitoring the chemistry and dynamics of the

atmosphere from the ground to the mesosphere. The OMI is a nadir-viewing charge-coupled

device (CCD) spectrometer onboard the Aura satellite, whose observation band is

near-UV/visible. We selected the Column Amount NO_2 Trop product to calculate the changes in

the tropospheric NO_2 concentration impacted by the control measures in East Asia, Western

Europe and North America (Fig. 2):

65
$$VA = \frac{N_2 - N_1}{N_1} \times 100\% \qquad (1)$$

where VA is the relative variation ratio, N1 is the average NO_2 concentration in the

troposphere one month before the lockdown, and N_2 is the average NO_2 concentration in the

troposphere one month after the lockdown.

The data from 1 January 2020 to 3 March 2020, were selected to analyze the variation in

NO_2 over time in China (Fig. 3). The NO_2 satellite data of Italy, Germany, France, the United

States, Iran and Switzerland is from the time of the first case in each country to 20 April 2020

(Fig. 4). Due to the satellite orbit, default values occur among the daily data that were

determined via piecewise linear interpolation over time. Border data from the US Centers for

Disease Control and Prevention (CDC)

(https://www.cdc.gov/epiinfo/support/downloads/shapefiles.html) were selected to obtain the

borders of each country. To remove the influence of weather factors, a 7-day moving average

was calculated. To compare the relative changes among the different countries, the data for each

country were standardized.

The daily total number of new confirmed cases in each country and region was obtained

from the Center for Systems Science and Engineering (CSSE) of Johns Hopkins University

(https://github.com/CSSEGISandData/COVID-19). The daily total number of new confirmed

cases in China was retrieved from the Department Earth System Science of the Tsinghua

University shared case database (https://cloud.tsinghua.edu.cn/d/335fd08c06204bc49202/).

3. Results analysis

3.1. A spatial comparison of pollutants before and after lockdown in Hubei Province.

Fig. 1 shows the change in pollutant concentration in Hubei Province one month before and after the closure of major cities severely affected by the epidemic. Compared with before the lockdown, NO_2, sulphur dioxide (SO_2), particulate matter (PM_{10}), carbon monoxide (CO) and $PM_{2.5}$ concentrations all decreased to a certain extent, while NO_2 experienced the most notable decrease. Since biomass and coal combustion are major SO_2 and CO sources, they exhibit the lowest rate of improvement (Khalil et al., 1988; Huang et al., 2012). Both the $PM_{2.5}$ and PM_{10} concentrations decreased to a certain extent (31.2% and 34.3%, respectively) as a result of the reduction in fugitive dust, particulate matter and important precursors produced by motor vehicles and factories (Huang et al., 2019). The monthly average $PM_{2.5}/PM_{10}$ ratio was 0.81 (95% confidence interval (CI): 0.76-0.86), so $PM_{2.5}$ was the main particle pollutant after lockdown. Exhaust emissions contributed only moderately to local levels of the $PM_{2.5}$ total mass, which were mostly derived from other sources, such as biomass combustion and the remote transmission of secondary particles (Chen et al., 2012). Therefore, the impact of strict traffic control during the lockdown on $PM_{2.5}$ is not notable, and the spatial difference is large, so $PM_{2.5}$ is not suitable as a city lockdown indicator.

Although the NO_2 emissions per vehicle slightly decreased after the upgrading of the quality standards of petroleum products, the notable growth of vehicle ownership increased the proportion of NO_2 traffic source emissions, in addition, after the implementation of emission standards for coal-fired power plants, multiple technical improvements greatly controlled the NO_2 emissions from coal-fired sources, which all enhanced the correlation between NO_2 and city

lockdown effect (Liu et al., 2020). The effect of city closure on NO_2 was significantly greater

than that on the other pollutants, with an average concentration reduction of approximately 60.3%

(95% CI: 56.8-64.0%), which can be applied as an environmental indicator of the lockdown

effect.

3.2. Changes of airborne NO_2 plummets over COVID-19-hit area after lockdown.

In East Asia (Fig. 2a), satellite images show that compared to before the blockade, the total

emissions of NO_2 in eastern China have significantly decreased by approximately 56.6%. In

South Korea, the monthly NO_2 emissions have also been reduced by approximately 18.0 %. The

local government has implemented the most expansive testing programme and has isolated

people infected with the virus without locking down entire cities, and the sharp decrease in NO_2

may be linked to the reduction in local emissions and pollutant transport from surrounding areas

(Han, 2019; Bauwens et al., 2020). Japan has not imposed widespread lockdown policies, and a

4.8% increase in NO_2 may be linked to emissions from power generation and industrial

processes (Han, 2019). In western Europe (Fig. 2b), the monthly NO_2 concentrations have

decreased sharply in Italy by 47.5%, particularly in the north (82.4%), where the outbreak is the

most severe. This could be due to the reduction in road traffic and the decrease in economic

activities in the industrial heartland as a result of the widespread lockdown policy (Schiermeier,

2020). Other countries such as Germany, Denmark, and Poland also experienced notable

reductions. This is consistent with the results of the European Environment Agency (EEA) (EEA,

2020). However, in certain areas, such as northern and southern Spain, the NO_2 concentration

has risen, possibly because of lax closure measures and increased emissions from coal-fired

power plants (Curier et al., 2014). In the United States (Fig. 2c), one month after the lockdown,

the overall decline in NO_2 is relatively small. The worst affected states, such as New York,

Washington and California, still contain areas with increased NO_2 concentrations, and the NO_2

concentration is increasing significantly in the vast midwestern regions that have not yet been

locked down.

3.3. The temporal evolution of the NO_2 concentration and the total number of newly confirmed

cases in China.

After the strict city lockdown, the NO_2 concentration in the main virus-affected cities in

China decreased significantly (Fig. 3). Consistent with the satellite data, the ground monitoring

results showed that compared to the conditions before the closure, the monthly average NO_2

concentration after the lockdown decreased approximately 55.7% (95% CI: 51.5-59.6%). Since

the lockdown, the total number of confirmed new cases reaches an inflection point after

approximately two weeks (the incubation period of the virus is 14 days), and compared to the

period of 1-15 days after the closure, the total number of confirmed new cases in the 16-30 days

after the closure has decreased 73.6% (95% CI: 64.9-81.1%). The most significant improvement

was recorded in Hangzhou, where the NO_2 concentration decreased approximately 68.1%, and

the total number of confirmed new cases declined the most. Likewise, Zhengzhou, Changsha,

Guangzhou, and Nanchang are good examples. Wuhan, the worst virus-affected area in China,

also exhibited a downward trend. The total number of confirmed new cases reached 13,436 on

February 12 due to the inclusion of clinically diagnosed cases (Wei et al., 2020), resulting in a

new delayed peak in the figure.

The national emergency response has delayed the spread of the epidemic and greatly limited

its range. The suspension of intra-city public transport, the closure of entertainment venues and

the banning of public gatherings have been linked to a reduction in the incidence of cases.

Studies have shown that before emergency response initiation, the basic case reproduction

number (R_0) is 3.15, and after intervention measures were implemented in 95% of all places, the

average R_0 value has dropped to 0.04, the total number of actual cases has decreased 96% (Tian

et al., 2020).

3.4. The temporal evolution of the NO_2 concentration and the total number of newly confirmed

cases in the COVID-19-hit areas globally.

After the countries severely affected by COVID-19 implemented strict lockdown measures,

satellite data showed a significant decline in NO_2 emissions, and the total confirmed new cases

decreased after two weeks in most areas (Fig. 4). As a result, the strict lockdown of

COVID-19-hit areas other than those in China is also effective and easy to implement to prevent

the spread of the virus. The lockdown measures might have already prevented tens of thousands

of deaths in Europe (Flaxman et al., 2020). In Italy, where the epidemic is widespread, after the

lockdown the NO_2 emissions significantly decreased by an average of 36.6%, and the total

confirmed new cases reached an inflection point 12 days later, thus verifying that the spread of

the virus was effectively controlled. Studies have shown that 38,000 deaths have been averted in

this country due to the implemented intervention measures (Flaxman et al., 2020). The

occurrence time of the inflection point is mainly related to the magnitude of NO_2 decline. In

France, NO_2 declined less (27.1%), and the time for the total number of confirmed new cases to

reach an inflection point was delayed. In Germany, the NO_2 emissions decreased the most, by

54.7%, and the total number of confirmed new cases reached an inflection point within 8 days

after lockdown, which occurred earlier than in other countries. For Iran and Switzerland, due to

the relatively low number of confirmed cases, with the decline of NO_2 after the strict control,

confirmed new cases also reached the inflection point earlier. In the the worst-affected states,

United States, NO_2 emissions decreased by an average of 43.1% in New York, Washington and

California, and the total confirmed new cases dropped significantly, showing signs of easing.

3.5. The results after epidemic control in the COVID-19-hit areas globally.

Following the implementation of strict epidemic control measures, the newly confirmed

cases dropped by more than 50% within 30 days and NO_2 declined significantly in most

countries (Figure 5a). Especially in most countries in Europe and the Western Pacific region, the

number of newly confirmed cases dropped more markedly, with an average drop of more than

60%. The newly confirmed cases in Ireland, Austria, Japan fell by more than 80%. The reduction

in the number of newly confirmed cases is closely related to the intensity of interventions, and

NO_2 can quantify the effects and intensity of interventions. Among the top 50 countries with the

cumulative number of confirmed cases, the stricter the control measures, the higher the rate of

decrease in newly confirmed cases (Figure 5b). If the NO_2 reduction rate is only about 25%, the

control effect of government intervention measures on the epidemic situation is not significant

(Figure 5b).

In Germany, when large public events were canceled, the spreading rate (λ) dropped from

0.43 to 0.25. When schools, childcare facilities, and many stores were closed, λ dropped to 0.15.

Only when a contact ban was imposed by government authorities, λ dropped to 0.09, which is

lower than the recovery speed, and the exponential growth of new cases turned to decline

(Dehning et al., 2020). In Wuhan, China, due to the strict lockdown, the estimated effective

reproduction number Rt fell immediately after reaching a peak of 3.82 on January 24 and fell

below 1.0 on February 6, 2020 (NO_2 mutation time) (Pan et al., 2020). In summary, stringent

interventions help to reduce newly confirmed cases earlier and more rapidly. NO_2 can be used as

an environmental indicator for quantitative research to assess the efficacy and timing of

non-pharmacological interventions to guide current and future responses to global pandemics.

4. Discussion

Urbanization and rapid transportation system development accelerate the spread of

COVID-19, and only strict containment measures can effectively prevent a second and third

outbreak (Huang et al., 2020b; Huang et al, 2020c). The NO_2 concentration can be considered an

inexpensive indicator of virus transmission control. As a result of strict control measures and the

rapid implementation of first-level emergency measures, the NO_2 emissions and total number

confirmed new cases significantly decreased in China, especially in the strictly controlled cities.

In many European countries, a strict lockdown is also effective and easy to do to prevent the

spread of the virus. But there are also areas such as southern and northern Spain and parts of the

United States where the NO_2 level has increased.

Studies have shown that the likelihood of fewer cases in the gradual multi-stage policy is

zero, and that such a policy decision implies that the government is willing to risk an increase in

COVID19 cases and deaths in exchange for decreased economic and isolation impacts, which

may not be desirable from an objective point of view (Karnon, 2020). Although the immediate

adoption of a lockdown policy may lead to many people being adversely affected financially, in

the short term, the number of new confirmed cases will decline approximately 15 days after

policy implementation, and an earlier decline can occur with stricter lockdown measures.

International guidance supports a range of mandatory social isolation measures, extensive case

detection, and isolation and contact tracing. Compliance with quarantine directives is absolutely

critical to saving lives, protecting the most vulnerable in society, and ensuring that the national

security system can cope and care for the sick (Iacobucci, 2020). In such cases, an immediate

lockdown policy may be preferred, and NO_2, as an environmental indicator of virus control, can

help managers implement effective control measured to curb the spread of COVID-19. In

addition, NO_2 evolution can be used to predict the COVID-19 newly confirmed cases, using the

local NO_2 change rate to reflect the effects of the control measures of governments in various

countries. For details, please refer to the Global Prediction System of the COVID-19 Pandemic

(http://covid-19.lzu.edu.cn).

233   5. Conclusion

By combining historical COVID-19 data, NO_2 satellite data and government control

measures, we analyzed the quantification and control effects of epidemic intervention measures.

The inflection point and reduction rate of newly confirmed cases are closely related to the time

and intensity of epidemic control measures. The decline rate of NO_2 is influenced by the

emission characteristics of each country, on the other hand, it is largely related to the intensity

and implementation effect of control measures. Therefore, NO_2 can be used as an environmental

indicator to quantify the effectiveness and intensity of interventions and can be incorporated into

assessments of the implementation of existing outbreak control measures in countries and into

prediction of future scenarios and case numbers. In addition, NO_2 would help provide

policymakers with information on strengthening, easing and selecting appropriate measures to

contain COVID-19 and other future global pandemics.

Reference

Baccini L, Brodeur A. Explaining governors' response to the COVID-19 pandemic in the United States. 2020.

Bauwens M, Compernolle S, Stavrakou T, Müller JF, Van Gent J, Eskes H, Levelt PF, Vander R,Veefkind JP, Vlietinck J, Yu H, Zehner C. Impact of coronavirus outbreak on NO_2 pollution assessed using TROPOMI and OMI observations. Geophys Res Lett 2020; 47.

Brnnum-Hansen H, Bender AM, Andersen ZJ, Sørensen J, Bønløkke JH, Boshuizen H, Becker T, Diderichsen F, Loft S. Assessment of impact of traffic-related air pollution on morbidity and mortality in Copenhagen Municipality and the health gain of reduced exposure. Environ Int 2018; 121: 973-980.

Chen Y, Liu Q, Geng F, Zhang H, Cai C, Xu T, Ma X, Li H. Vertical distribution of optical and micro-physical properties of ambient aerosols during dry haze periods in Shanghai. Atmos Environ 2012; 50: 50-59.

Clougherty JE, Kheirbek I, Eisl HM, Ross Z, Pezeshki G, Gorczynski JE, Johnson S, Markowitz S, Kass D, Matte T. Intra-urban spatial variability in wintertime street-level concentrations of multiple combustion-related air pollutants: The New York City Community Air Survey (NYCCAS). J Expo Sci Env Epid 2013; 23: 232-240.

Curier RL, Kranenburg R, Segers AJS, Timmermans RMA, Schaap M. Synergistic use of OMI NO_2 tropospheric columns and LOTOS–EUROS to evaluate the NOx emission trends across Europe. Remote Sens of Environ 2014; 149: 58-69.

Dehning J, Zierenberg J, Spitzner FP, Wibral M, Neto JP, Wilczek M, Priesemann V. Inferring change points in the spread of COVID-19 reveals the effectiveness of interventions. Science 2020.

Fan X, Dawson J, Chen M, Qiu C, Khalizov A. Thermal stability of particle-phase monoethanolamine salts. Environ Sci Tech 2018; 52: 2409-2417.

EEA. Air pollution goes down as Europe takes hard measures to combat coronavirus. 2020.

Flaxman S, Mishra S, Gandy A. Estimating the number of infections and the impact of nonpharmaceutical interventions on COVID-19 in 11 European countries. Imperial College COVID-19 Response Team 2020.

Han KM. Temporal Analysis of OMI-Observed Tropospheric NO_2 Columns over East Asia
during 2006-2015. Atomos 2019; 10: 658.

Huang C, Wang Y, Li X, Ren L, Zhao J, Hu Y, Zhang L, Fan G, Xu J, Gu X, Cheng Z, Yu T,
Xia J, Wei Y, Wu W, Xie X, Yin W, Li H, Liu M, Xiao Y, Gao H, Guo L, Xie J, Wang G,
Jiang R, Gao Z, Jin Q, Wang J, Cao B. Clinical features of patients infected with 2019
novel coronavirus in Wuhan, China. Lancet (London, England) 2020a.

Huang F, Zhou J, Chen N, Li Y, Wu S. Chemical characteristics and source apportionment of
$PM_{2.5}$ in Wuhan, China. J Atmos Chem 2019; 76: 245-262.

Huang J, Liu X, Zhang Li, Yang K, Chen Y, Huang Z, Liu C, Lian X, Wang D. The amplified
second outbreaks of global COVID-19 pandemic. medRxiv 2020b.

Huang Q, Cheng S, Perozzi RE. Use of a MM5–CAMx–PSAT Modeling System to Study SO_2
Source Apportionment in the Beijing Metropolitan Region. Environ Model Assess 2012;
17: 527-538.

Huang Z, Huang J, Gu Q, Du P, Liang H, Dong Q. Optimal temperature zone for the dispersal of
COVID-19. Sci Total Environ 2020c: 139487.

Iacobucci G. Covid-19: UK lockdown is "crucial" to saving lives, say doctors and scientists.
Bmj-Brit Med J 2020; 368.

Johansson C, Lovenheim B, Schantz P, Wahlgren L, Almstrom P, Markstedt A, Stromgren M,
Forsberg B, Sommar J. Impacts on air pollution and health by changing commuting from
car to bicycle. Sci Total Environ 2017; 584: 55-63.

Karnon J. A Simple Decision Analysis of a Mandatory Lockdown Response to the COVID-19
Pandemic. Appl health econ hea 2020; 18: 329-331.

Khalil MAK, Rasmussen, R. A. Carbon Monoxide in an Urban Environment: Application of a
Receptor Model for Source Apportionment. Japca 1988; 38: 901-906.

Kim Y, Guldmann JM. Land-use regression panel models of NO_2 concentrations in Seoul, Korea.
Atmos Environ 2015; 107: 364-373.

Legido-Quigley H, Mateos-Garcia JT, Campos VR, Gea-Sanchez M, Muntaner C, McKee M.
The resilience of the Spanish health system against the COVID-19 pandemic. Lancet
Public heal 2020; 5: e251-e252.

Li Q, Guan X, Wu P, Wang X, Zhou L, Tong Y, Ren R, Kathy SML, Eric HYL, Jessica YW,
Xing X, Xiang N. Early Transmission Dynamics in Wuhan, China, of Novel Coronavirus
–Infected Pneumonia. New Engl J Med 2020.

Lian X, Huang J, Huang R, Liu C, Wang L, Zhang T. Impact of city lockdown on the air quality
of COVID-19-hit of Wuhan city. Sci Total Environ 2020: 140556.

Liu D, Deng Q, Ren Z, Zhou Z, Song Z, Huang J, Hu R. Variation trends and principal
component analysis of nitrogen oxide emissions from motor vehicles in Wuhan City from
2012 to 2017. Sci Total Environ 2020; 704: 134987.

Pan A, Liu L, Wang C, Guo H, Hao X, Wang Q, Huang J, He N, Yu H, Lin X. Association of
public health interventions with the epidemiology of the COVID-19 outbreak in Wuhan,
China. JAMA 2020; 323: 1915-1923.

Paterlini M. Lockdown in Italy: personal stories of doing science during the COVID-19
quarantine. Nature 2020.

Schiermeier Q. Why pollution is plummeting in some cities - but not others. Nature 2020; 580:
313.

Cyranoski D, Silver A. Wuhan scientists: What it's like to be on lockdown. Nature 2020.

Tessum MW, Larson T, Gould TR, Simpson CD, Yost MG, Vedal S. Mobile and Fixed-Site
Measurements To Identify Spatial Distributions of Traffic-Related Pollution Sources in
Los Angeles. Environ Sci Tech 2018; 52: 2844-2853.

Tian H, Liu Y, Li Y, Wu C, Chen B, Kraemer MUG, Li B, Cai J, Xu B, Yang Q. An
investigation of transmission control measures during the first 50 days of the COVID-19
epidemic in China. Science 2020; 368: 638-642.

Wei Y, Wei L, Jiang Y, Shen S, Zhao Y, Hao Y, Du Z, Tang J, Zhang Z, Jiang Q.
Implementation of Clinical Diagnostic Criteria and Universal Symptom Survey
Contributed to Lower Magnitude and Faster Resolution of the COVID-19 Epidemic in
Wuhan. Engineering 2020.

Wu JT, Leung K, Leung GM. Nowcasting and forecasting the potential domestic and
international spread of the 2019-nCoV outbreak originating in Wuhan, China: a
modelling study. Lancet (London, England) 2020.

Yuan B, Hu W, Shao M, Wang M, Chen W, Lu S, Zeng L, Hu M. VOC emissions, evolutions
and contributions to SOA formation at a receptor site in eastern China. Atmos Chem
Phys 2013; 13: 8815-8832.

Zhou Z, Tan Q, Liu H, Deng Y, Wu K, Liu C, Zhou X. Emission characteristics and
high-resolution spatial and temporal distribution of pollutants from motor vehicles in
Chengdu, China. Atmos Pollut Res 2019; 10: 749-75.

Corresponding Author

Dr. Jianping Huang（hjp@lzu.edu.cn）

Acknowledgements

This work was jointly supported by the National Science Foundation of China (41521004) and the Gansu Provincial Special Fund Project for Guiding Scientific and Technological Innovation and Development (Grant No. 2019ZX-06). The authors acknowledge the China National Environmental Monitoring Centre for providing the datasets. http://www.cnemc.cn/.

Additional information

Competing financial interests: The authors declare no competing financial interests.

Graphics software

All maps and plots were produced using licensed.

Fig. 1 The improvement rate of the major pollutants and the distribution of the accumulated epidemic numbers in each city of Hubei Province after the lockdown. One month after the blockade (23 February 2020), the cumulative number of confirmed cases ranged from 0-46607 (Wuhan), as indicated by the colour scale. The monthly mean improvement rate of NO_2, SO_2, PM_{10}, CO and $PM_{2.5}$ after the lockdown in the worst virus-affected areas is shown in the histogram, with SO_2, PM_{10}, CO and $PM_{2.5}$ indicated with blue columns, and NO_2 indicated with red columns. Red columns (NO_2) are significantly higher than the others, so it can characterize the locking measures.

Fig. 2 The relative variation in the monthly average tropospheric NO$_2$ concentration before and after the lockdown. a, Relative variation in East Asia. **b**, Relative variation in Western Europe. **c**, Relative variation in North America. The colour table represents the relative percentage of the monthly average change in NO$_2$ after the locksown compared to the before. The red areas indicate that the NO$_2$ concentration decreased, and the blue areas indicate that the NO$_2$ concentration increased. The lockdown in East Asia, Western Europe and North America began on January 23, March 10, and March 16, respectively. Source: Analysis of data from the NASA Ozone Monitoring Instrument (OMI).

Fig. 3 Temporal variation in the NO$_2$ concentration and number of new cases in China. The standardized 7-day smooth data of NO$_2$ is represented by blue lines, with satellite observations in dark blue and ground observations in light blue. The dotted line indicates the average value of NO$_2$ before and after the blockade. The 7-day smooth data of total confirmed new cases is indicated by red lines. ** indicates significant difference at the 0.01 level (bilateral), and * indicates significant difference at the 0.05 level (bilateral). Starting from the lockdown, the NO$_2$ concentration dropped significantly, and the number of new cases also showed a downward trend within two weeks.

Fig. 4 Temporal variation in the NO_2 concentration and number of new cases in the COVID-19-hit areas. Italy, Germany, France, the United States, Iran and Switzerland were selected as the representatives of COVID-19-hit areas. New York, Washington and California were selected as the representatives of the United States. From the time of the first case in each country to 20 April 2020, the standardized 7-day smooth satellite data of NO_2 is represented by the blue line. The dotted line indicates the average value of NO_2 before and after the blockade. The 7-day smooth data of total confirmed new cases is indicated by red line. ** indicates significant difference at the 0.01 level (bilateral), and * indicates significant difference at the 0.05 level (bilateral). The NO_2 in major countries showed a downward trend. The total confirmed new cases decreased significantly about 15 days after the lockdown.

Fig 5. The results of strict epidemic control measures. a, The reduction rate of new confirmed cases within 30 days at the NO_2 mutation point. Red represents Europe, dark blue represents Western Pacific, light blue represents Americas, and green represents Eastern Mediterranean. **b**, Changes in the rate of decline in NO_2 and the reduction rate of newly confirmed cases after the implementation of strict epidemic control measures.