1 **Clinical characteristics and early outcomes in patients with COVID-19 treated with**
2 **tocilizumab at a United States academic center**
3

4 Casey A. Rimland,[1a] Camille E. Morgan,[1,2a] Griffin J. Bell,[2] Min K. Kim,[3] Tanner
5 Hedrick,[4] Ashley Marx,[4] Brian Bramson,[3] Heidi Swygard,[3] Sonia Napravnik,[2,3] John L.
6 Schmitz,[5] Shannon S. Carson,[6] William A. Fischer,[6] Joseph J. Eron,[3] Cynthia L. Gay,[3]
7 Jonathan B. Parr[3*]
8 ([a]Contributed equally, *Corresponding)
9
[1] Medical Scientist Training Program, University of North Carolina at Chapel Hill (UNC),
Chapel Hill, NC, USA
[2] Department of Epidemiology, UNC Gillings School of Global Public Health, Chapel
Hill, NC, USA
[3] Division of Infectious Diseases, UNC School of Medicine, Chapel Hill, NC, USA
[4] Department of Pharmacy, UNC Medical Center, Chapel Hill, NC, USA
[5] Department of Pathology and Laboratory Medicine, UNC School of Medicine, Chapel
Hill, NC, USA
[6] Division of Pulmonary Diseases and Critical Care Medicine, UNC School of Medicine,
Chapel Hill, NC, USA
20
21

22 **Keywords:** SARS-CoV-2, tocilizumab, cytokine release syndrome, COVID-19,
23   interleukin 6 (IL-6)
24
25 **Running title:** Tocilizumab outcomes in severe COVID-19
26
27 **Abstract word count:** 48
28
29 **Manuscript word count:** 1368
30

* Corresponding author: Jonathan B. Parr, MD, MPH; 130 Mason Farm Rd, Chapel
Hill, NC 27599; jonathan_parr@med.unc.edu

Author contributions:
CAR, CEM extracted data, analyzed data, wrote first draft. GJB analyzed data. MKK extracted
data. JBP supervised the study. CAR, CEM, GJB, MKK, TH, AM, BB, SN, JLS, SSC, WAF, JJE,
CLG, JBP reviewed and edited the final draft.

Abstract

We describe early outcomes in 11 COVID-19 patients treated with the IL-6 receptor inhibitor tocilizumab. While C-reactive protein decreased, neither clinical improvement nor reduced temperature or oxygen requirements was observed in most patients. Our findings contrast with prior reports and raise questions about tocilizumab use in severe COVID-19.

INTRODUCTION

A hyperinflammatory state is a posited driver of severe COVID-19 [1]. Elevations of IL-6 have been described in acute respiratory distress syndrome and cytokine release syndrome (CRS) [2,3]. Recently, IL-6-driven inflammation in severe COVID-19 was suggested, as increased levels were associated with mortality [4]. IL-6 blockade via tocilizumab, a humanized monoclonal antibody that targets the IL-6 receptor, has been proposed as a treatment for COVID-19 [5].

Peer-reviewed reports of tocilizumab use for COVID-19 are largely limited to retrospective case series from China [6,7] and Italy [8,9], as well as case reports [10–13]. Xu *et al*. reported improved inflammatory markers, oxygen requirements, and fever in 21 patients after tocilizumab treatment [6]. Luo *et al.*'s description of 15 patients included more critically ill patients and more measured results, with 20% mortality [7]. A prospective, open-label study from Italy reported clinical improvement and increased survival after tocilizumab [8], while a second study in Italy found no significant clinical improvement or mortality benefit following tocilizumab [9]. Given the paucity of data on tocilizumab use in COVID-19 in the United States, we evaluated 11 patients treated with tocilizumab for COVID-19. We detail their clinical characteristics before and after treatment and reconsider tocilizumab use in severe COVID-19.

METHODS

We performed a retrospective cohort study of patients with COVID-19 treated with tocilizumab at UNC Medical Center between March 21st and April 25th, 2020. Clinical

data was extracted from the medical record. COVID-19 was diagnosed using

nasopharyngeal swabs and reverse-transcriptase PCR (RT-PCR). IL-6 testing was

conducted at Mayo Clinic Laboratories (Rochester, MN) using an

electrochemiluminescence assay. Baseline IL-6 values for two patients were from a multi-

cytokine panel (lower-limit of detection [LOD] of 5.0 pg/mL); all others were from a

standalone IL-6 assay (LOD 1.8 pg/ml). Disease severity was defined according to the

World Health Organization (WHO) COVID-19 clinical status scale with minor modification

[14]. Descriptive statistics were calculated using JMP 15 and Wilcoxon signed rank test

performed in GraphPad Prism 8. Data was visualized using the *tidyverse* package in R

3.6.0. The study was approved by the UNC Institutional Review Board.

RESULTS

Patient population

Eleven patients with confirmed SARS-CoV-2 infection treated with tocilizumab for COVID-

19 were identified and included. Demographic, clinical, and laboratory data are displayed

in **Table 1.** Six (55%) patients had an unknown exposure history. Median time from

symptom onset to admission was 8 days (IQR=5-11). Most patients were male (82%,

n=9); non-white (73%, n=8); non-substance users (82%) with a median age of 59 years

old (IQR=48-65). Patients had a median of 2 (IQR=1-6) comorbidities, including

hypertension (73%, n=8), type 2 diabetes mellitus (36%, n=4), and lung disease (27%,

n=3). Almost all were overweight (36%, n=4) or obese (55%, n=6). Five (46%) took an

angiotensin converting enzyme inhibitor (ACEI) or angiotensin receptor blocker (ARB) at

baseline. Seven (64%) were transferred from outside hospitals, all within one day of initial

presentation. Five (46%) were diagnosed with COVID-19 by SARS-CoV-2 RT-PCR as

outpatients prior to presentation.

Baseline laboratory and imaging characteristics

On admission, most patients had significantly elevated inflammatory markers including

elevated CRP (100%, n=11/11), fibrinogen (100%, n=9/9), ferritin (80%, n=8/10), LDH

(88.9%, n=8/9), and D-dimer (70%, n=7/10). Most patients had normal total white blood

cell counts (WBC) (81.82%, n=9/11); two patients had elevated total WBC (18.2%).

However, almost all patients presented with low absolute lymphocyte counts (90.9%,

n=10/11). Several patients presented with laboratory signs of additional organ damage.

Four patients had abnormal troponin I levels (36.4%), five patients had elevated pro-BNP

(45.5%), three patients had elevated creatinine (27.3%), and two patients had both

elevated AST and ALT (18.2%). Almost all patients had abnormal chest imaging on

admission (91%, n=10).

Hospital course

Most patients had severe or critical COVID-19 by modified WHO criteria (55%, n=6) at

the time of admission (**Table 1**). All patients required supplemental oxygen on admission,

including five (46%) via low-flow nasal cannula (LFNC, ≤ 10L/min), two (18%) via high-

flow nasal cannula (HFNC), and four (36%) via mechanical ventilation. During the

observation period, eight (73%) were critically ill requiring mechanical ventilation and

vasopressors, one (9%) was critically ill requiring mechanical ventilation alone, one (9%)

with do-not-intubate status received HFNC only, and one (9%) had mild disease requiring

LFNC. Ten of the 11 (91%) received at least one dose of an anti-microbial or

immunomodulating treatment during their hospitalization.

Outcomes after tocilizumab

Tocilizumab was administered a median of one day (IQR 1-4) from admission and nine

days (IQR 7-14) after symptom onset. Ten patients received a single tocilizumab dose,

and one received two doses. Median dosage was 7.9 mg/kg. Two of six patients with

available baseline IL-6 levels had low levels, one below the assay's LOD (≤5 pg/mL) and

another of 10.4 pg/mL. Four of five with available IL-6 levels after tocilizumab had IL-6

levels above the upper limit of quantification. IL-6 increased after tocilizumab in all four

patients with matched baseline and post-tocilizumab testing.

CRP and fibrinogen decreased significantly after tocilizumab (**Figure 1A,** median CRP

211.6 vs. 19.7 mg/L [*p*=0.001] and fibrinogen 767 vs. 420 mg/dL [*p*=0.002] pre- and 5

days post-tocilizumab, respectively). Other laboratory markers did not show clear trends

(**Figure 1B**). Temperature did not show obvious improvement (**Figure 1C**), and only

three (27%) showed improvement in oxygen requirements (**Figure 1D**). At end of follow-

up, patients were observed for a median of 17 days (range=7-34) after initial tocilizumab

dose. Three died (27%); five (45%) remained in the intensive care unit (ICU) on

mechanical ventilation, receiving vasopressors or hemodialysis; one (9%) was

transferred from the ICU to floor status and weaned to room air; and two (18%) were

discharged home without oxygen. Seven patients (64%) had minimally elevated liver

function tests after tocilizumab administration. Two were diagnosed with ileus and two

with bacterial pneumonia after tocilizumab administration; no serious adverse events

were observed.

DISCUSSION

While tocilizumab was recently described as a promising therapy for severe COVID-19,

our findings raise questions about its use in critically ill patients [7]. We report data from

patients with COVID-19 treated with tocilizumab at a United States academic center,

almost all of whom required ICU care. CRP and fibrinogen improved after tocilizumab

administration as previously reported [6–8], but improvement in disease severity or

mortality were not observed. These findings, in addition to low baseline IL-6 levels

observed in some patients, raise questions about tocilizumab use in patients presenting

with severe or critical COVID-19, especially without baseline IL-6 data.

Important differences exist between our cohort and those from China and Italy [6–9].

Three of four studies had few patients requiring mechanical ventilation (n=2/21, n=4/32,

and n=5/63 respectively) [6,8,9]; the fourth study did not report specific oxygen

requirements but did include more patients reported as critically ill (n=7/15) [7]. Our cohort

included higher rates of hypertension (73% versus 40% in Luo *et al*., 37% in Campochiaro

et al., and 38% in Sciascia *et al.*); a majority were overweight or obese (91%). We also

observed lower baseline IL-6 levels compared to those described in these cohorts,

although IL-6 assays were conducted in different laboratories. While Sciascia *et al.*

attribute observed survival benefit of tocilizumab treatment to early administration in their

Italian cohort [8], we did not observe the same clinical improvement, despite tocilizumab

administration within a median of one day of hospitalization. Our findings also

demonstrate the challenge of defining CRS in COVID-19, especially in the absence of

standardized IL-6 assays and limited data to support IL-6 thresholds for tocilizumab use.

Our findings contrast with most previous reports and do not provide clear evidence of

clinical improvement after tocilizumab treatment in patients with severe COVID-19.

Several explanations can be considered. First, our patients were more severely ill than

previous cohorts. It is possible tocilizumab was administered too late to offer benefit.

Second, our patient cohort was more racially and ethnically diverse and had more

comorbidities, notably obesity and hypertension, than those in China and Italy. Third,

improvements observed in prior, less severely ill cohorts, might have occurred as part of

the normal disease process, with or without tocilizumab.

In conclusion, tocilizumab reduces inflammatory markers (CRP, fibrinogen), but it is

unclear whether its activity leads to equally dramatic effects on clinical outcomes.

Randomized controlled trials are essential to determine when, or if, tocilizumab should be

used in severe COVID-19.

ACKNOWLEDGMENTS

The authors would like to thank the nurses; laboratory staff; and allied health providers

involved in the care of these patients. We also thank Carlton Moore and Hannah Little for

assistance with initial exploration of data.

Funding

The authors report no designated funding for the project. CAR and CEM receive training

support from the University of North Carolina at Chapel Hill Medical Scientist Training

Program. CAR additionally receives training support from the National Institutes of Health

MD/PhD Partnership Program.

Conflicts of interest

JBP reports research support outside the submitted work from Gilead Sciences and the

World Health Organization, and non-financial support from Abbott Laboratories. CLG

has received research support outside the submitted work from Viiv Healthcare and

Gilead Sciences. SSC reports research support outside of the submitted work from

Biomarck Pharmaceuticals.

References

1.   Ye Q, Wang B, Mao J. The pathogenesis and treatment of the `Cytokine Storm' in
COVID-19. J Infect **2020**; 80:607–613.

2.   Aisiku IP, Yamal J-M, Doshi P, et al. Plasma cytokines IL-6, IL-8, and IL-10 are
associated with the development of acute respiratory distress syndrome in patients
with severe traumatic brain injury. Crit Care Lond Engl **2016**; 20:288.

3.   Giavridis T, van der Stegen SJC, Eyquem J, Hamieh M, Piersigilli A, Sadelain M.
CAR T cell-induced cytokine release syndrome is mediated by macrophages and
abated by IL-1 blockade. Nat Med **2018**; 24:731–738.

4.   Zhou F, Yu T, Du R, et al. Clinical course and risk factors for mortality of adult
inpatients with COVID-19 in Wuhan, China: a retrospective cohort study. The
Lancet **2020**; 395:1054–1062.

5.   Liu B, Li M, Zhou Z, Guan X, Xiang Y. Can we use interleukin-6 (IL-6) blockade for
coronavirus disease 2019 (COVID-19)-induced cytokine release syndrome (CRS)?
J Autoimmun **2020**; :102452.

6.   Xu X, Han M, Li T, et al. Effective treatment of severe COVID-19 patients with
tocilizumab. Proc Natl Acad Sci **2020**; :202005615.

7.   Luo P, Liu Y, Qiu L, Liu X, Liu D, Li J. Tocilizumab treatment in COVID-19: A single
center experience. J Med Virol **2020**; :1–5.

8.   Sciascia S, Aprà F, Baffa A, et al. Pilot prospective open, single-arm multicentre
study on off-label use of tocilizumab in patients with severe COVID-19. Clin Exp
Rheumatol **2020**; 38:529–532.

9.   Campochiaro C, Della-Torre E, Cavalli G, et al. Efficacy and safety of tocilizumab in
severe COVID-19 patients: a single-centre retrospective cohort study. Eur J Intern
Med **2020**; 76:43–49.

10.   Ferrey AJ, Choi G, Hanna RM, et al. A Case of Novel Coronavirus Disease 19 in a
Chronic Hemodialysis Patient Presenting with Gastroenteritis and Developing
Severe Pulmonary Disease. Am J Nephrol **2020**; 51:337–342.

11.   Michot J-M, Albiges L, Chaput N, et al. Tocilizumab, an anti-IL6 receptor antibody,
to treat Covid-19-related respiratory failure: a case report. Ann Oncol **2020**;
Available at: https://linkinghub.elsevier.com/retrieve/pii/S0923753420363870.
Accessed 5 May 2020.

12.   Zhang X, Song K, Tong F, et al. First case of COVID-19 in a patient with multiple
myeloma successfully treated with tocilizumab. Blood Adv **2020**; 4:1307–1310.

13. Mihai C, Dobrota R, Schröder M, et al. COVID-19 in a patient with systemic
sclerosis treated with tocilizumab for SSc-ILD. Ann Rheum Dis **2020**; 79:668–669.

14. World Health Organization. WHO R&D Blueprint novel Coronavirus: COVID-19
Therapeutic Trial Synopsis. Geneva, Switzerland: 2020.

Figure 1. Laboratory and clinical outcomes of patients treated with tocilizumab. (**A-**

B) Laboratory trends following tocilizumab. Medians (blue lines), interquartile ranges

(IQR, blue shading), and reference ranges (dashed red lines). (**C-D**) Oxygen

requirements and temperature trends, displayed using locally estimated scatterplot

smoothing (orange lines). Patient #6 was treated with two doses of tocilizumab; time 0

refers to the time of initial administration.

Table 1. Patients with COVID19 treated with tocilizumab demographics and clinical characteristics (n=11).

Demographics and Clinical Characteristics	Median (IQR) / Percentage (n)
Age / Male sex	59 (48-65) / 82% (9)
Race	
Hispanic/Latino	46% (5)
Caucasian	27% (3)
African American	27% (3)
Tobacco / ethanol / substance use	9% (1) / 9% (1) / 9% (1)
BMI	31 (29-36)
Normal (18.5-25)	9% (1)
Overweight (25-30)	36% (4)
Obese (>30)	55% (6)
Comorbidities	
Hypertension	73% (8)
Type 2 diabetes mellitus	36% (4)
Lung disease	27% (3)
Cardiovascular disease	18% (2)
Liver disease	9% (1)
Renal disease	9% (1)
HIV	9% (1)
Active cancer	9% (1)
Solid organ transplant	9% (1)
ACE/ARB treatment	46% (5)
COVID-19 Clinical and Treatment Characteristics	
Time from symptom onset to admission	8 (5-11)
Abnormal CXR/CT on admission	91% (10)
Prone positioning while intubated	78% (7/9)
Dosage of tocilizumab (mg/kg)	7.9 (5.8-8.1)[*]

Other treatments

Azithromycin	64% (7)
Hydroxychloroquine	36% (4)
Ritonavir/Lopinavir	27% (3)
Steroids	27% (3)

Admission labs#	Median (IQR)	Percentage abnormal (n)
CRP (mg/L)	197.3 (146.8-324.9)	100% (11/11 elevated)
Fibrinogen (mg/dL)	662.0 (566-928.5)	100% (9/9 elevated)
Ferritin (ng/mL)	589.5 (389.8-1705)	80% (8/10 elevated)
LDH (U/L)	1203 (943-2122)	89% (8/9 elevated)
D-Dimer (ng/mL)	343.5 (199.5-1,203)	70% (7/10 elevated)
WBC (x10^9/L)	8.5 (5.1-9.9)	18% (2/11 elevated)
Absolute lymphocyte count (x10^9/L)	0.8 (0.6-1)	91% (10/11 reduced)
Platelet count (x10^9/L)	230 (162-382)	18% (2/11 reduced)
Troponin I (ng/mL)	<0.034 (<0.034-0.172)	36% (4/11 elevated)
Creatinine (mg/dL)	0.92 (0.7-2.53)	27% (3/11 elevated)
Pro-BNP (pg/mL)	69.7 (34.3-6,870)	46% (5/11 elevated)
AST (U/L)	51 (44-72)	46% (5/11 elevated)
ALT (U/L)	35 (30-60)	27% (3/11 elevated)

Clinical and Laboratory Outcomes

Disease severity	Admission	Highest Severity	Outcome
8 (Deceased)	---	---	27% (3)
7 (Critical, Ventilator + Organ Support)	36% (4)	73% (8)	46% (5)
6 (Critical, Ventilator)	0% (0)	9% (1)	0% (0)
5 (Severe, HFNC/NIPPV)	18% (2)***	9% (1)***	0% (0)
4 (Mild, LFNC)	36% (4)	9% (1)	0% (0)
3 (Mild, room air)	9% (1)	0% (0)	9% (1)
1-2 (Ambulatory – Discharged)	---	---	18% (2)

Other invasive therapies

Hemodialysis/CRRT	27% (3)
ECMO	9% (1)

Length of stay (days)[**]	18 (15-25)		
Length (days) on mechanical ventilation[**]	17 (12-29)		
Laboratory measures[#]	*Pre-Tocilizumab*	*Post-Tocilizumab*	*p-value*
CRP (mg/L)	211.6 (176.9-313.8)	19.7 (13.6-29)	0.001
Fibrinogen (mg/dL)	767 (625-862)	420 (386.3-473.5)	0.002
IL-6 (pg/mL)	30.65 (9.1–88.8)	>400 (249.4 - >400)	0.13[##]

[*] n=12 doses; one patient received tocilizumab twice.

[**] As of date of death, date of discharge, or end date of observation period.

[***] One patient was do not intubate status on admission, receiving only HFNC/NIPPV. Patient subsequently died.

[#] Not all patients had all laboratory values available, data representative of patients with data available.

[##] Only four patients had matched pre- and post-Tocilizumab IL-6 levels.

Abbreviations: OSH, outside hospital; BMI, body mass index; CXR, chest x-ray; CT, computed tomography; HIV,

human immunodeficiency virus; ACEI/ARB, angiotensin-converting enzyme inhibitor/angiotensin II receptor blocker;

CRP, c-reactive protein; LDH, lactate dehydrogenase; WBC, white blood cell count; BNP, b-type natriuretic peptide;

AST, aspartate aminotransferase; ALT, alanine aminotransferase; HFNC, high-flow nasal cannula; NIPPV, non-

invasive positive pressure ventilation; LFNC, low-flow nasal cannula; CRRT, continuous renal replacement therapy;

ECMO, extracorporeal membrane oxygenation; IL-6, interleukin-6.