Estimation of the true infection rate and infection fatality rate of COVID-19 in each country

Masahiro Sonoo[1], Takamichi Kanbayashi[1], Takayoshi Shimohata[2], Masahito Kobayashi[3], and Hideyuki Hayashi[4].

[1] Department of Neurology, Teikyo University School of Medicine, Kaga 2-11-1, Itabashi-ku, Tokyo, Japan.

[2] Department of Neurology, Gifu University Graduate School of Medicine, 1-1 Yanagido, Gifu-shi, Gifu, Japan.

3 Faculty of Echonomics, Yokohama National University, Tokiwadai 79-3, Hodogaya-ku, Yokohama-shi, Kanagawa, Japan

[4] Department of Chemistry, Faculty of Medicine, 2-7 Daigakumachi, Osaka Medical College Takatsuki-shi, Osaka, Japan.

Key Words: COVID-19; Infection Rate; Infection fatality rate; PCR Examination

Corresponding Author: Prof. Masahiro Sonoo, Department of Neurology, Teikyo University School of Medicine. Kaga 2-11-1, Itabashi-ku, Tokyo 1738605, Japan

e-mail: sonoom@med.teikyo-u.ac.jp

Abstract

The true infection rate of COVID-19 and the infection fatality rate in the whole population have been estimated for each country based on the open data. The estimate well coincided with local surveys. The fact that several to hundred times of identified cases are already infected might require reconsideration of the strategy to cope with COVID-19. Attenuation of the viral virulence may be considered to be a goal, especially in countries with already low fatality rate, mainly Asian countries.

Coronavirus disease 2019 (COVID-19) pandemic is now a worldwide peril. Recently, a few local surveys revealed surprisingly high infection rates in general population.[1–4] However, the True Infection Rate (TIR) in the whole population of each country is unknown. We devised a simple method to infer TIR, as well as the Infection Fatality Rate (IFR) based on the open data.

We picked up the data at a website 4 times from April 12th to May 24th, 2020.[5] Countries or regions having more than 1000 cases on April 12th were included. Counties fulfilling this condition have increased in later surveys, although we did not add such countries because we aimed to investigate the trend for the same population. Positive rate of PCR tests (Infection Rate; IR) and the rate of PCR tests among the population (Examination Rate; ER) were calculated. The cardinal assumption is that IR and ER are negatively correlated because PCR examinations will be restricted to cases with strong suspicion while ER is low, and vice versa. Then TIR of a specific region can be estimated using the linear regression. The ratio of the infected persons in the population to the already identified cases was calculated and named as True/Identified Case Ratio (TICR). IFR was estimated using the number of total deaths at the same website.[5] For each parameter, 95% confidence interval (CI) was calculated. All statistical calculation was done using Microsoft Excel for Macintosh.

Included were 66 countries or regions. ER and IR on logarithmic scales showed a significant negative correlation at all adopted dates. The correlation coefficients were -0.350, -0.308, -0.332, -0.338 on April 12th, April 26th, May 16th, and May 24th, respectively. The slope value was -0.211, -0.203, -0.248, and -0.268, respectively. The estimated TIR, TICR and IFR on May 24th of representative countries are presented, together with the plot of IR vs. ER on May 24th (Figure 1). The data of Tokyo prefecture in Japan are also added to the table. The data for the whole countries or regions investigated are presented as a supplementary table. TIR took on

values between 1% to 10% for most countries. TICR was around 7 to 20 in major western countries, and around 100 in some countries, including Japan. IFR also took on wide range of values, generally higher in western countries than in eastern countries.

A number of factors obscure the correlation between IR and ER. These include the variation of actual situations of conducting PCR-tests among different countries. However, these varieties will be averaged over many countries included in this study. The observed correlation is highly significant, indicating that the inverse relation of IR and ER is sufficiently robust. If countries with high TIR actively expand the PCR tests due to a sense of crisis, the negative correlation would reduce and the slope of the regression line would become shallower as the time lapses. However, such a trend was not observed, and therefore we used the newest result of the regression analysis for the estimation.

Comparing the present results with local surveys is interesting. Keio University in Tokyo found 6% IR using PCR tests for in-patients unrelated to COVID-19.[1] Our estimate of TIR in Tokyo is 5.7%. Reported IRs in USA using antibody tests were 1.5% in Santa Clara[2], 4.6% in Los Angeles county,[3] and 13.9% in New York state,[4] and our estimate of TIR in the whole USA is 5.0%. These results support the validity of our estimate. We of course understand that the present calculation remains a rough estimate. However, the reliability of the antibody test may not be always guaranteed,[2,6] and therefore our methods must have a certain role.

The trajectory analysis[7] reveals that the pandemic is still expanding in some countries, or temporarily controlled in others, most probably due to social distancing.[8] Only limited countries, including China, succeeded in almost completely containing the pandemic (present estimation is not applicable to these countries). If several to hundred times of identified cases are already infected, possibly often asymptomatic, reconsideration of the strategy in the management of

COVID-19 may be necessary. Finding out all infected persons and completely containing the infection would be unrealistic in such a situation. Other than complete containment, effective treatment/vaccines, and herd immunity, attenuation of the viral virulence is another way to make the pandemic less damaging.[9] This is achieved in animals by natural selection but in humans the prevention of in-hospital infection, especially from severe cases, would be the key.[10] This might be a choice, especially in countries with already low estimated IFR, mainly Asian countries such as Singapore, India, Indonesia or Japan.

Acknowledgments: We would like to thank Dr. Masashi Idogawa, Sapporo Medical University School of Medicine, for providing information on the trajectory analysis.

References

1. Keio University Hospital. Update on 2019 Novel Coronavirus Situation at Keio University Hospital. http://www.hosp.keio.ac.jp/en/oshirase/detail/40174/ [Accessed 26 April 2020].

2. Bendavid E, Mulaney B, Sood N, Shah S, Ling E, Bromley-Dulfano R, et al. COVID-19 Antibody Seroprevalence in Santa Clara County, California. medRxiv preprint doi: https://doi.org/10.1101/2020.04.14.20062463 2020 [Accessed 26 April 2020].

3. CBS Los Angeles. Antibody study estimates 4.6% percent of Angelenos have coronavirus. https://losangeles.cbslocal.com/2020/05/19/antibody-study-estimates-4-6-percent-of-angelenos-have-coronavirus/ [Accessed 26 May 2020].

4. BGR. New York coronavirus antibody study shows just how far the virus has spread. https://bgr.com/2020/04/23/coronavirus-new-york-spread-study-results/ [Accessed 26 April 2020].

5. Worldometers.info. Reported Cases and Deaths by Country, Territory, or Conveyance. Dover, Delaware, U.S.A. https://www.worldometers.info/coronavirus/ [Accessed 12 April, 26 April, 16 May, and 24 May 2020].

6. Kontou PI, Braliou GG, Dimou NL, Nikolopoulos G, Bagos PG. Antibody tests in detecting SARS-CoV-2 infection: a meta-analysis. medRxiv preprint doi: https://doi.org/10.1101/2020.04.22.20074914

7. Sapporo Medical University School of Medicine. Trajectory analysis of new coronavirus COVID-19 cases and deaths by country. https://web.sapmed.ac.jp/canmol/coronavirus/trajectory_e.html) [Accessed 25 May 2020].

8. Sonoo M, Idogawa M, Kanbayashi T, Shimohata T, Hayashi H. Correlation between PCR Examination Rate among the Population and the Containment of Pandemic of COVID-19.

Public Health 2020, in press. URL: https://doi.org/10.1016/j.puhe.2020.05.027

9. Hunter P. The spread of the COVID-19 coronavirus: Health agencies worldwide prepare for the seemingly inevitability of the COVID-19 coronavirus becoming endemic. EMBO Rep 2020;21(4):e50334. doi: 10.15252/embr.202050334.

10. Armengaud J, Delaunay‑Moisan A, Thuret JY, van Anken E, Acosta‑Alvear D, Aragón T, et al. The Importance Of Naturally Attenuated Sars‑Cov ‑2 In The Fight Against Covid ‑19. Environmental Microbiology 2020, in press. URL: https://doi.org/10.1111/1462-2920.15039

Figure Legend

Figure 1. The correlation of IR vs. ER and the estimated parameters.

Left: Two-dimensional plot of IR vs. ER on May 24th, 2020. There was a significant negative correlation (r = -0.338, p = 0.006). A dotted line indicates the linear regression line.

Right: Known and estimated parameters of 15 representative countries and Tokyo prefecture in Japan. The parameters for countries are adopted and calculated from the data on May 24th.[5] Those for Japanese prefectures are acquired from Toyo Keizai Online. Coronavirus Disease (COVID-19) Situation Report in Japan. https://toyokeizai.net/sp/visual/tko/covid19/en.html [Accessed 26 May 2020].

Abbreviations: IR, infection ratio; ER, examination ratio; IFR, infection fatality rate; TIR, true infection ratio; CI confidence interval; TICR, true/identified case ratio.

Supplementary Table. Parameters of the COVID-19 Pandemic in 67 countries and regions. (Supplementary Table is submitted as a separate word file)

Infection rate (IR, %)

r = -0.338
(p = 0.006)

May 24th, 2020

Examination rate (ER, %)

Country	Cases	ER (%)	IR (%)	TIR (%) (95%CI)	TICR (95%CI)	IFR (%) (95%CI)
USA	1,666,828	4.34	11.6	5.01 (2.79 − 8.99)	9.9 (5.5 − 17.8)	0.595 (0.332 − 1.068)
Brazil	347,398	0.35	47.3	10.36 (3.61 − 29.75)	63.4 (22.1 − 181.9)	0.100 (0.035 − 0.287)
Russia	335,882	5.76	4.0	1.86 (1.09 − 3.17)	8.1 (4.8 − 13.8)	0.125 (0.073 − 0.212)
Spain	282,370	6.45	9.3	4.47 (2.69 − 7.44)	7.4 (4.4 − 12.3)	1.372 (0.825 − 2.283)
UK	257,154	4.94	7.7	3.43 (1.96 − 6.01)	9.1 (5.2 − 15.9)	1.576 (0.900 − 2.759)
Italy	229,327	5.61	6.8	3.13 (1.83 − 5.35)	8.2 (4.8 − 14.1)	1.732 (1.013 − 2.961)
France	182,469	2.12	13.2	4.70 (2.29 − 9.62)	16.8 (8.2 − 34.4)	0.925 (0.451 − 1.894)
Germany	179,986	4.29	5.0	2.15 (1.20 − 3.87)	10.0 (5.6 − 18.0)	0.464 (0.258 − 0.833)
India	131,423	0.21	4.6	0.88 (0.28 − 2.80)	92.8 (29.3 − 293.5)	0.032 (0.010 − 0.100)
Canada	83,621	3.80	5.8	2.43 (1.32 − 4.47)	11.0 (6.0 − 20.2)	0.693 (0.377 − 1.274)
Belgium	56,810	6.64	7.4	3.57 (2.16 − 5.92)	7.3 (4.4 − 12.1)	2.231 (1.346 − 3.697)
Sweden	33,188	2.08	15.8	5.60 (2.72 − 11.53)	17.0 (8.3 − 35.1)	0.706 (0.343 − 1.451)
Singapore	31,068	5.04	10.6	4.74 (2.72 − 8.27)	8.9 (5.1 − 15.6)	0.008 (0.005 − 0.014)
Indonesia	21,745	0.09	9.1	1.38 (0.37 − 5.11)	173.1 (46.7 − 641.9)	0.036 (0.010 − 0.133)
Japan	16,536	0.21	6.2	1.18 (0.38 − 3.73)	90.6 (28.8 − 285.1)	0.054 (0.017 − 0.170)
Tokyo	5,152	0.11	35.2	5.65 (1.59 − 20.14)	148.1 (41.6 − 527.7)	0.037 (0.010 − 0.131)