# Transmission onset distribution of COVID-19

June Young Chun[1†], Gyuseung Baek[2†], Yongdai Kim[2]*

[1]Department of Internal Medicine, National Cancer Center, Goyang, South Korea

[2]Department of Statistics, Seoul National University, Seoul, South Korea

[†]June Young Chun and Gyuseung Baek contributed equally to this manuscript.

*Correspondence to:

Yongdai Kim, Ph.D.

Department of Statistics, Seoul National University, 56-1 Mountain, Sillim-dong, Gwanak-gu,

Seoul, 08826, South Korea

E-mail: ydkim0903@gmail.com, Tel: 82-2-880-9255; Fax: 82-2-883-6144

**Abstract**

**Objectives**

The distribution of the transmission onset of COVID-19 relative to the symptom onset is a key parameter for infection control. It is often not easy to study the transmission onset time, as is difficult to know who infected whom exactly when.

**Methods**

We inferred transmission onset time from 72 infector-infectee pairs in South Korea, either with known or inferred contact dates by means of incubation period. Combining this data with known information of infector's symptom onset, we could generate the transmission onset distribution of COVID-19, using Bayesian methods. Serial interval distribution could be automatically estimated from our data.

**Results**

We estimated the median transmission onset to be 1.31 days (standard deviation, 2.64 days) after symptom onset with peak at 0.72 days before symptom onset. The pre-symptomatic transmission proportion was 37% (95% credible interval [CI], 16–52%). The median incubation period was estimated to be 2.87 days (95% CI, 2.33–3.50 days) and the median serial interval to be 3.56 days (95% CI, 2.72–4.44 days).

**Conclusions**

Considering the transmission onset distribution peaked with the symptom onset and the pre-symptomatic transmission proportion is substantial, the usual preventive measure might be too late to prevent SARS-CoV-2 transmission.

37     **Keywords**: COVID-19, SARS-CoV-2, infectious disease transmission, infectious disease

38     incubation period

**Introduction**

SARS-CoV-2, a novel coronavirus first reported in Wuhan City, China in December 2019, has been spreading globally and the World Health Organization declared it as a pandemic on March 11, 2020 (Zhu et al., 2020). A total of 3,096,626 cases of coronavirus disease 2019 (COVID-19), and 217,896 confirmed deaths were counted world widely by April 30, 2020 .

Considering its global threat, the transmission dynamics of COVID-19 should be made explicit. There have been studies reporting the estimated serial interval and incubation period, based on the early epidemics in China (Backer et al., 2020, Lauer et al., 2020, Li et al., 2020, Nishiura et al., 2020). The serial interval is the duration between the symptom onset of successive cases, and the incubation period is the time since infection to the symptom onset. Those are key epidemiological parameters, giving essential insight to infer the transmission potential or to determine the quarantine duration (White and Pagano, 2008).

There have been multiple reports of pre-symptomatic transmission of SARS-CoV-2, highlighting the difficulty of containment and mitigation of the disease (He et al., 2020, Liu et al., 2020, Wei et al., 2020). However, transmission onset distribution of COVID-19 has never been studied before. It is not easy to study the transmission onset time, as is difficult to know who infected whom exactly when. In this report, we tried to estimate the transmission onset distribution relative to the symptom onset by means of solid epidemiologic data of infector-infectee pairs in South Korea.

**Methods**

**Data**

We searched public reports of confirmed COVID-19 patients by government and each

municipal website of South Korea . As of March 31, 9,887 cases have been confirmed of

which 8,260 (83.5%) cases were linked to certain clusters such as religious groups, hospitals,

or long-term healthcare facilities . Five-hundred and sixty cases (5.7%) were imported from

abroad . We screened all available cases between January 23 to March 31, 2020 and selected

the pairs with clearly defined contact history. We excluded cases from main clusters unless

the causal relationship between an infector and infectee pair was evident.

For each pair of cases, we collected information on the dates of symptom onset of both the

infector and the infectee, exposure dates of the infectee by the infector, and the dates of

confirmation of both. Considering the South Korean quarantine policy, the confirmation date

could be regarded as the date on which isolation started. Demographic information of age and

sex were also gathered.

**Procedures**

We tried to estimate the time difference between an infector's symptom onset and its

transmission onset time. We defined symptom onset date ($S$), an infectee's infection date ($I$),

an infector's transmission onset date ($P$), and contact time of the pair as ($C_L, C_R$), where

subscripts L and R denote the left and right boundaries. We put $i$ as an infectee and $j_i$ as

the corresponding infector, and observed ($S_{j_i}, C_{L,i}, C_{R,i}, S_i$). $I_i$ should be within interval

$[C_{L,i}, C_{R,i}]$, and transmission onset of an infector $(P_j)$ is described as $P_j = \min_{i:j_i=j} I_i$. Using

the pair $\{(S_j, P_j) | j : infector\}$, we were able to estimate the distribution $f(W)$ of

$W = P - S$.

There are three possible scenarios for $I_i$ in our data.

(i)      There is an exact calendar date of an exposure. In this case, we could directly

90                achieve the timing of transmission as $I_i = C_{L,i} = C_{R,i}$.

91        (ii)     There is a specified duration of exposure $(C_{L,i} < C_{R,i})$.

92        (iii)    There is a continuous exposure such as household members $(C_{L,i} = -\infty, C_{R,i} =$

93                $S_i)$.

Recall that the aim is to estimate the distribution $f(W)$ of $W$ based on the data

$(S_i, C_{L,i}, C_{R,i}, S_{j_i})$ for infectees. We consider the following parametric model for $f(W)$:

$$W \sim f(\cdot \,|\theta) \Rightarrow W + \mu \sim \pi \, g(\cdot \,|\phi) + (1 - \pi)\mathcal{U}(-D, 0)(\cdot)$$

where $\theta = (\phi, \mu, \pi)$, $\pi \in (0,1)$, $\mu > 0$, $g(\cdot \,|\phi)$ is the gamma distribution with the

parameter $\phi$ and $\mathcal{U}(-D, 0)(\cdot)$ is the uniform distribution with the support $(-D, 0)$. In the

model, $\mu$ is the shift parameter for cases with transmission prior to the symptom onset. We

introduced the term $(1 - \pi)\mathcal{U}(-D, 0)(\cdot)$ to make the effect of negative outliers of $W$ be

minimal to the estimation of the parameter. That is, we considered $W$ to be an outlier if

$W < -\mu$. In the model, $D > 0$ is an arbitrary limit for an outlier of $W$. In the analysis, we

used $D = 14$, since the usual incubation period is within 14 days.

A difficulty in estimating $\theta$ is that the exact $W$ is not observed for cases in categories (ii)

and (iii). However, conceptually we could impute the timing of transmission $I$ by use of the

estimated incubation period distribution (Reich et al., 2009). With our notation, $Z = S - I$ is

the incubation period. From our data of contact dates and symptom onset dates, we could

estimate the distribution of $Z$ which we utilized to impute $I$. That is, in case (ii), we could

generate $Z$ from conditional probability distribution $Z \,|\, Z \in [S - C_R, S - C_L]$, and then we

let $I = S - Z$. Similarly, we could generate $Z$ from conditional probability distribution

$Z \,|\, Z \leq S$ for case (iii). To accommodate this idea, we used a Bayesian method and obtained

the posterior distribution of $\theta$ based on the Markov chain Monte Carlo (MCMC) algorithm

in which $W$ is imputed for cases in categories (ii) and (iii). A detailed description of the

MCMC algorithm is provided in Supplementary Material.

For sensitivity analysis, we utilized the incubation period distribution of a previous study and

replicated the same analysis as above (Lauer et al., 2020). Moreover, we hold the parameter

$\mu$ as 4, and compared the result using the same procedure, since the minimum of $W$ from

our data was -4 days. Serial interval distribution could be automatically estimated from our

data.

All analyses were conducted using the $R$ statistical software version 3.6.3. All code and data

are available in the supplementary material. We present case numbers with randomization

process, instead of nationally designated identification numbers, in order to maintain

confidentiality.

**Ethics Approval**

All the data used in this study were publicly available and were approved from institutional

review board assessment of National Cancer Center (NCC2020-0119).

**Results**

We found 89 infectees with defined source of infection (infector) and contact history. Four

infectors (4.5%) were asymptomatic until diagnosed, and thus were excluded from the

analysis. Sixteen infectees (18.0%) were asymptomatic when diagnosed, among which 13

cases had no specified contact date and were unable to guess the infection time. In short, 72

infector-infectee pairs were included in the study (Fig 1).

Twenty-two cases documented a single point date of contact, and 16 cases had documented

duration of contact dates. Otherwise, 34 cases (32 household members and 2 colleagues) had

continuous exposure history (Supplementary Material). The median age of the 72 infectees

was 40 years (interquartile range [IQR], 24 to 54 years). Of these, 34 (47%) were male and

38 (53%) were female. The infectees had 40 unique infectors. The average number of

transmissions per infector was 1.8, with a maximum of four cases (Fig 2).

We estimated the incubation period distribution of COVID-19 using a lognormal model.

Among 35 pairs with given contact dates, the median incubation period was 2.87 days (95%

credible interval [CI], 2.33–3.50) (Table 1). The estimated median serial interval from 69

infector-infectee pairs with given symptom onset dates was 3.56 days (95% CI, 2.72–4.44),

with the best-fit using a lognormal distribution (Table 2).

Based on the 72 infecfor-infectee pairs, the Bayes estimate of the transmission onset

distribution after a smoothing procedure was given as Fig 3A. The mean and median values

were 1.31 days (95% CI, 0.38–2.55) and 0.68 days (95% CI, -0.09–1.73) after infectors'

symptom onset, respectively, with the peak at 0.72 days before symptom onset. The pre-

symptomatic transmission proportion was 37% (95% CI, 16–52%).

For sensitivity analysis, applying the incubation period of previous study, the transmission

onset distribution was inferred as Fig 3B. The mean and median values were 0.59 days (95%

CI, -0.33–1.79) and 0.12 days (95% CI, -0.75–1.06) after the infector's symptom onset,

respectively, with the peak at 0.41 days before symptom onset. The pre-symptomatic

transmission proportion was 48% (95% CI, 29–67%). With a fixed parameter of $\mu$ as 4, the

mean and median were 0.59 (95% CI, -0.34–1.79) and -0.07 days (95% CI, -0.91–0.99), with

the pre-symptomatic transmission proportion of 51% (95% CI, 37–64%).

**Discussion**

The transmission onset distribution is of utmost interest to clinicians and public health workers, but it has not been widely studied to date due to lack of epidemiological information. Based on concrete data, we hereby present that the transmission of COVID-19 could start before the symptom onset, and the probability peaked as the symptom start and declined thereafter. Interestingly, the trend of this distribution looks similar with published SARS-CoV-2 viral load kinetics though it could not include the viral load data before the symptom onset (Kim et al., 2020, Zou et al., 2020).

In South Korea, the first COVID-19 case was identified on January 19, 2020 in a traveler from Wuhan City, who was quarantined immediately at the airport screening. Until February 18, 2020, there were only imported cases from abroad or cases from their close contacts who were under the surveillance. The situation was totally changed after an outbreak with an unknown source of infection occurred within a religious community, resulting in a total of 5,210 confirmed cases in that single cluster as of April 12, 2020, a number which is almost half (49.6%) of the total 10,512 cases in South Korea . This study spanned from late January to the end of March, 2020, and encompassed all the phases observed in South Korea, thus enhancing the reliability of the results. Furthermore, in South Korea close contacts of confirmed patients were screened extensively with SARS-CoV-2 nucleic acid testing, regardless of whether the contact had symptoms. Given this situation, it can be assumed that cases of infection are unlikely to have been missed.

189 Among the 89 infectors, 4 cases (4.5%) were asymptomatic when diagnosed, but out of 89

190 infectees, 16 cases (18.0%) were asymptomatic when diagnosed. The difference is attributed

191 to the active surveillance of close contacts of COVID-19 patients, including the screening

192 tests of those asymptomatic, which in turn, pre-symptomatic. Our study revealed that

193 approximately 40% of individuals with COVID-19 infected others when they could not

194 acknowledge any symptom. This finding is in line with other recently published studies

195 revealing pre-symptomatic transmission ratio of 46% (95% CI, 21-46%), 44% (95% CI, 25-

196 69%) in China, and 48% (95% CI, 32-67%) in Singapore (Ganyani et al., 2020, He et al.,

197 2020, Liu et al., 2020).

199 This study has several limitations. First, the dates of symptom onset and contact time were

200 identified from epidemiological investigation and it could be false due to recall bias. Second,

201 we only collected data from definite infector-infectee pairs with defined contact times. Hence,

202 our estimation of transmission onset distribution is the upper bound of the true value. Third,

203 we excluded clustered data to promote the accuracy of causal relationships. The transmission

204 behavior of SARS-CoV-2 might have different characteristics within unique clusters, but this

205 could not be measured in this study. Fourth, we might not catch the late-onset transmission

206 cases, considering the pre-emptive screening and isolation measures implemented in South

207 Korea.

209 This study provides an important message in terms of public health practice. It indicates that

210 the usual preventive measure of isolating people when they become symptomatic, might be

211 too late to prevent SARS-CoV-2 transmission.

**Declarations**

**Funding**

None

**Conflict of Interests**

The authors declare that there are no competing interests.

**Data availability**

All code and data are available in the supplementary materials. We present case numbers with

randomization process, instead of nationally designated identification numbers, in order to

maintain confidentiality.

**Author contributions**

J.Y.C. and Y.K. conceived of the study, and G.B. performed the analysis. J.Y.C. and G.B.

wrote the first draft of the manuscript. Y.K. reviewed and edited the manuscript. All authors

interpreted the findings, contributed to writing the manuscript, and approved the final version

for publication.

**Acknowledgement**

We acknowledge public health authorities in South Korea for their innumerable efforts to

perform epidemiologic investigation to stop the COVID-19 outbreak.

**References**

Coronavirus Disease-19, Republic of Korea; Available from: http://ncov.mohw.go.kr/.

[Accessed April 12, 2020].

Coronavirus disease 2019 (COVID-19) Situation Report – 52; Available from:

https://www.who.int/docs/default-source/coronaviruse/situation-reports/20200312-sitrep-52-

covid-19.pdf?sfvrsn=e2bfc9c0_4. [Accessed March 30, 2020].

Coronavirus           disease          (COVID-19)         Pandemic;        Available         from:

https://www.who.int/emergencies/diseases/novel-coronavirus-2019. [Accessed May 7, 2020].

COVID-19               Response              Status;            Available            from:

https://blog.naver.com/gyeonggi_gov/221856148207. [Accessed April 10, 2020].

COVID-19           Situation         Report          (Gyeonggi);          Available          from:

https://www.gg.go.kr/bbs/board.do?bsIdx=722&menuId=2903#page=21. [Accessed April 10,

2020].

COVID-19          Situation         Report          (Seoul);          Available          from:

http://www.seoul.go.kr/coronaV/coronaStatus.do. [Accessed April 10, 2020].

Backer JA, Klinkenberg D, Wallinga J. Incubation period of 2019 novel coronavirus (2019-

nCoV) infections among travellers from Wuhan, China, 20-28 January 2020. Euro Surveill

2020;25(5).

Ganyani T, Kremer C, Chen D, Torneri A, Faes C, Wallinga J, et al. Estimating the generation

interval for COVID-19 based on symptom onset data. 2020:2020.03.05.20031815.

He X, Lau EHY, Wu P, Deng X, Wang J, Hao X, et al. Temporal dynamics in viral shedding

and transmissibility of COVID-19. Nat Med 2020.

Kim ES, Chin BS, Kang CK, Kim NJ, Kang YM, Choi JP, et al. Clinical Course and

Outcomes of Patients with Severe Acute Respiratory Syndrome Coronavirus 2 Infection: a

Preliminary Report of the First 28 Patients from the Korean Cohort Study on COVID-19. J

Korean Med Sci 2020;35(13):e142.

Lauer SA, Grantz KH, Bi Q, Jones FK, Zheng Q, Meredith HR, et al. The Incubation Period

of Coronavirus Disease 2019 (COVID-19) From Publicly Reported Confirmed Cases:

Estimation and Application. Ann Intern Med 2020.

Li Q, Guan X, Wu P, Wang X, Zhou L, Tong Y, et al. Early Transmission Dynamics in Wuhan,

China, of Novel Coronavirus-Infected Pneumonia. N Engl J Med 2020;382(13):1199-207.

Liu Y, null n, Funk S, Flasche S. The contribution of pre-symptomatic infection to the

transmission dynamics of COVID-2019 [version 1; peer review: 1 approved]. 2020;5(58).

Nishiura H, Linton NM, Akhmetzhanov AR. Serial interval of novel coronavirus (COVID-19)

infections. Int J Infect Dis 2020;93:284-6.

Reich NG, Lessler J, Cummings DA, Brookmeyer R. Estimating incubation period

distributions with coarse data. Stat Med 2009;28(22):2769-84.

Wei WE, Li Z, Chiew CJ, Yong SE, Toh MP, Lee VJ. Presymptomatic Transmission of

SARS-CoV-2 - Singapore, January 23-March 16, 2020. MMWR Morb Mortal Wkly Rep

2020;69(14):411-5.

White LF, Pagano M. A likelihood-based method for real-time estimation of the serial interval

and reproductive number of an epidemic. Stat Med 2008;27(16):2999-3016.

Zhu N, Zhang D, Wang W, Li X, Yang B, Song J, et al. A Novel Coronavirus from Patients

with Pneumonia in China, 2019. N Engl J Med 2020;382(8):727-33.

Zou L, Ruan F, Huang M, Liang L, Huang H, Hong Z, et al. SARS-CoV-2 Viral Load in

Upper Respiratory Specimens of Infected Patients. N Engl J Med 2020;382(12):1177-9.

Table 1. Estimated incubation period distributions for COVID-19 based on 35 infector-

infectee pairs with defined contact dates.

|  | Lognormal | Gamma | Weibull |
|---|---|---|---|
| Median (95% CI) | 2.87 (2.33–3.50) | 2.99 (2.44–3.60) | 3.10 (2.54–3.71) |
| Parameter 1 | 1.055 (0.845–1.254) | 4.544 (2.590–10.494) | 2.310 (1.747–3.743) |
| Parameter 2 | 0.493 (0.301–0.643) | 0.709 (0.312–1.178) | 3.635 (3.030–4.208) |
| Log-likelihood | -46.6 | -45.4 | -45.1 |

283    For the lognormal distribution, parameter 1 and parameter 2 are the mean and SD of the
284    natural logarithm and for all other distributions, parameter 1 and parameter 2 are the shape
285    and scale parameters, respectively.

287    Table 2. Estimated serial interval distributions for COVID-19 based on 69 infector-infectee

288    pairs with symptom onset dates.

|  | Lognormal | Gamma | Weibull |
|---|---|---|---|
| Median (95% CI) | 3.56 (2.72–4.44) | 3.75 (2.87–4.72) | 3.93 (3.04–4.92) |
| Parameter 1 | 3.18 (2.22–4.24) | 3.45 (2.56–5.19) | 1.96 (1.64–2.44) |
| Parameter 2 | 0.75 (0.47–1.03) | 2.19 (1.44–2.91) | 8.47 (7.53–9.53) |
| Log-likelihood | -77.5 | -192.8 | -189.8 |

For the lognormal distribution, parameter 1 and parameter 2 are the mean and SD of the
natural logarithm and for all other distributions, parameter 1 and parameter 2 are the shape
and scale parameters, respectively.

Figure legends

Fig 1. Transmission plot of infector-infectee pairs. For cases of continuous exposure, the

interval presentation was limited to 10 days prior to the symptom onset whoever came first,

considering the upper 95[th] percentile of pre-estimated incubation period is 10.1.

Fig 2. Number of secondary infections per infector in the reported data set.

Fig 3. Estimated transmission onset distribution of COVID-19 (A) using the incubation

period distribution in the current study, and (B) using the previous incubation period

distribution. (Lauer et al., 2020)