

1 ***Prevalence, intensity and control strategies of soil-transmitted helminth***
2 ***infections among pre-school age children after 10 years of preventive***
3 ***chemotherapy in Gamo Gofa zone, Southern Ethiopia: A call for action***

4

5

6 Mekuria Asnakew Asfaw^{1*}, Tigist Gezmu¹, Teklu Wegayehu², Alemayehu Bekele¹, Zeleke

7 Hailemariam³, Nebiyu Masresha⁴, Teshome Gebre⁵

8

9

10

11 ¹Collaborative Research and Training Centre for NTDs, Arba Minch University, Ethiopia,

12 ²Department of Biology, College of Natural Sciences, Arba Minch University, Ethiopia,

13 ³School of Public health, College of Medicine and Health Sciences, Arba Minch University,
14 Ethiopia,

15 ⁴Ethiopian Public Health Institute, Addis Ababa, Ethiopia,

16 ⁵The Task Force for Global Health, International Trachoma Initiative, Addis Ababa, Ethiopia

17

18

19 ***Corresponding author**

20 E-mail : maksambaramr23@gmail.com (MA)

21

22 **Abstract**

23 **Background:** Soil-transmitted helminths (STH) (*Ascaris lumbricoides*, *Trichuris trichiura*
24 and hookworm) are among the most prevalent neglected tropical diseases (NTDs) in
25 Ethiopia. Pre-school age children (PSAC) pay high morbidity toll associated with STH
26 infections. Site specific operational evidence is lacking on prevalence, intensity and control
27 strategies of STH among PSAC in Ethiopia. This study is, therefore, aimed to fill that
28 missing knowledge gap.

29 **Methods:** We did a community-based cross-sectional study in five districts of Gamo Gofa
30 zone; from December 2018 to January 2019. Data were collected using pre-tested
31 questionnaire, and the Kato-Katz technique was used to diagnose parasites egg in stool. Then
32 data were edited, coded and entered into EpiData 4.4.2, and exported to SPSS software (IBM,
33 version 25) for analysis.

34 **Results:** A total of 2462 PSAC participated in this study. Overall, the prevalence of STH was
35 23.5% (578/2462). *Ascaris lumbricoides* was the most prevalent, 18.6% (457/2462),
36 followed by *Trichuris trichiura*, 9.2% (226/2462), and hookworm, 3.1% (76/2462). The
37 prevalence of STH in Chench, Dita, Deremalo, Bonke and Demba Gofa districts were
38 33.8% (210/622), 26.4% (109/413), 21.3% (88/414), 20.6% (128/621), and 11% (43/392),
39 respectively. Of the total, 7.4% (181/2462) PSAC were infected with two STH species.
40 *Ascaris lumbricoides* infections had moderate intensity (15%), and the rest had low infections
41 intensity. It is found that 68.7% of PSAC were treated with Albendazole. This study also
42 revealed households level data as, 39.4% used water from hand-dug well, 52.5% of need to
43 move ≥ 30 minutes to collect water, 77.5% did not treat water, and 48.9% not had hand
44 washing facility.

45 **Conclusion:** This study showed that a significant proportion of pre-school age children are
46 suffering from STH infections across districts in the study area despite preventive

47 chemotherapy distribution for more than 10 years. Further, gaps in control strategies of STH
48 were highlighted, which calls for action.

49 **Key words/phrases:** Soil-transmitted helminths; Pre-school age children; Kato-Katz
50 technique; Prevalence; Intensity, Control strategies of STH; Operational research

51 **Author summary**

52 Infections with STH pose serious public health challenge among children; it causes anemia,
53 vitamin A deficiency, stunting, malnutrition, impaired development, and intestinal
54 obstruction. About 4 billion people are at risk for STH infection worldwide, and with over 2
55 billion already infected. In Ethiopia, about 81 million people are at risk for STH infection.
56 STH can be controlled, possibly eliminated by combined interventions such as preventive
57 chemotherapy (PC), large scale administration of anthelmintic drugs to at-risk population,
58 and improved water, sanitation and hygiene (WASH). Ethiopia has been working to reduce
59 the prevalence of moderate and heavy infections with soil transmitted helminth in pre-school
60 and school aged children to below 1%. Since 2013 remarkable progress has been made on
61 STH control strategies in terms of mapping and distribution of PC. However, evidence is
62 lacking to track progress towards control and elimination goal, particularly among PSAC.
63 The findings of our study highlight that achieving STH control and elimination goal in Gamo
64 Gofa zone by 2020 and even beyond could be challenging unless the current STH control
65 strategies are improved.

66 **Introduction**

67 Soil-transmitted helminth (*Ascaris lumbricoides*, *Trichuris trichiura* and hookworm)
68 infections are among the most common neglected tropical diseases (NTDs) [1]. It is prevalent
69 mainly in tropical and subtropical areas where water supply, hygiene and sanitation
70 infrastructure are inadequate [2-4] . Moderate and heavy intensity of STH infections are
71 associated with chronic harmful effects on vitamin A and iron status, physical, intellectual,

72 and cognitive development in pre-school age children (PSAC), these morbidities not only
73 take a huge toll on the health of children, but has also been shown to affect economic
74 development of a nation [5, 6].

75 Globally, over 2 billion people are affected with STH where ascariasis accounts for almost
76 1.2 billion infections while trichiuriasis and hookworm (*Ancylostoma duodenale* and *Necator*
77 *americanus*) responsible for over 800 million and 740 million infections, respectively [1, 7,
78 8]. The global burden of STH infection is estimated at between 5 and 39 million disability-
79 adjusted life years (DALYs) and in 2010, 5.18 DALYs were estimated as associated with
80 STH infections [9, 10]. The greater burden of STH infection is found in the tropical countries
81 including tropical South America, China, East Asia, and Sub-Saharan Africa [5]. According
82 to the WHO estimate 42 countries in Africa are endemic for STH with 284 million cases
83 occurring in both school age and pre-school age children. These children require periodic
84 administration of preventive chemotherapy [1, 11].

85 STH are widely prevalent parasites in Ethiopia, and exceeds over 85% prevalence in some
86 districts. The number of people living in STH endemic areas is estimated at 81 million, of
87 which pre-school age children account 9.1 million [12]. Pre-school age children (PSAC) and
88 school-age children (SAC) are the most exposed groups and who pay the highest toll of
89 morbidity associated with STH infections. Significant proportions of pre-school age children
90 are affected by STH, which account 10% to 20% of the 3.5 billion people living in STH-
91 endemic areas. The greatest numbers of intestinal worms harbored in these children resulting
92 in diarrhea, loss of appetite, weight loss, growth retardation, malnutrition, anemia &
93 cognitive defects [6, 13-16].

94 In 2012, the World health organization (WHO) endorsed road map to combat NTDs by 2020,
95 and substantial progress has been made in terms of controlling STH morbidity [3]. The WHO
96 goal is to reduce the prevalence of moderate and heavy infections with soil transmitted

97 helminths in preschool and school aged children to below 1%, to be no longer considered
98 public health problems, by 2020 [17]. In line with this, Ethiopia has also set a similar goal to
99 achieve by the same year [12]. To achieve these goals, in areas where prevalence of any soil-
100 transmitted infection is 20% or higher, periodic mass administration of preventive
101 chemotherapy (deworming) using annual or biannual single-dose Albendazole or
102 Mebendazole is recommended by WHO for all pre-school and school age children (10).
103 As part of the global action towards Universal Health Coverage (UHC), ending NTDs is
104 prioritized by 2030 in the Sustainable Development Goal (SDG) under target 3.3, to ensure
105 healthy lives and well-being for all ages groups [3]. Therefore, SDGs could be attained only
106 if the NTD goals are met. Moreover, working on NTDs helps the vision of universal health
107 coverage, which means that all individuals and communities access the health services they
108 need without suffering from financial suffering [18].
109 Since 2008 preventive chemotherapy (PC) against STH in PSAC has been implemented
110 alongside Vitamin A distribution in the study area as well as at national level through
111 community based drug distribution platforms [12]. In 2013, the first national master plan for
112 NTDs was launched, and then the government of Ethiopia has been collaborating with the
113 WHO and other partners for mapping all endemic districts to address SAC through the
114 school based mass drug administration [19]. This implies that PC was started before ten years
115 to combat STH in PSAC at study area. Targeting on SAC earlier studies have shown the
116 impact of deworming on prevalence and intensity of STH infections. On the other hand, the
117 impact of deworming on STH infection status among PSAC is not well monitored and
118 evaluated. Hence, evidence is lacking at national level on prevalence, intensity and control
119 strategies of STH infections among PSAC. This condition is challenging for planning of
120 deworming programs on STH. Furthermore, timely and reliable operational research
121 evidence is needed to inform decision-making, and also data could be used for further actions

122 to improve control strategies of soil-transmitted helminths among pre-school-age children.
123 Therefore, the present study is aimed to determine prevalence, intensity and control strategies
124 of STH infections among preschool children in Gamo Gofa zone.

125 **Methods**

126 **Study area and period**

127 This study was conducted in the former Gamo Gofa zone; from December 2018 to January
128 2019. The zone is found in Southern Nations, Nationalities, and Peoples' Regional State
129 (SNNPR) and it had 15 districts and two city administrations (CAs). All districts and CAs are
130 endemic for STH, 15 had moderate prevalence and two had low prevalence According to the
131 2007 estimate of Central Statistical Agency of Ethiopia, a total of 2,043,668 people live in the
132 zone, of which 1,013,533 are males and 1,030,135 are females [20]. Arba Minch is the capital
133 city of the zone, which is located at 437 km away from Addis Ababa, capital city of Ethiopia.

134 **Study design and populations**

135 Community-based cross-sectional study was conducted to determine prevalence, intensity
136 and control strategies of STH infections among PSAC in the study area. The study
137 population was all PSAC (1-5 years) in the selected STH endemic *Kebeles* (localities). PSAC
138 who were unable to give stool samples at the time of collection were excluded from the
139 study. Additionally, PSAC were excluded in case when they were seriously ill or parents
140 were unable to provide their information.

141 **Determining sample size and sampling technique**

142 The sample size was determined using single population proportion formula,

$$143 \mathbf{n} = \frac{\left(Z_{\left(\frac{\alpha}{2}\right)} \right)^2 \mathbf{P}(1-\mathbf{P})}{\mathbf{d}^2},$$

144 assuming p, **25.7%** (proportion of PSAC infected with STH) [21]; **Z**, 1.96 at significant
145 level of alpha (α) of **0.05**, and desired degree of precision (d) of **3 %**, and design effect=**3**.

146 The computed sample size was **2434** and by adding **10%** non-response rate, the total
147 computed sample size was **2678**.

148 Multi-stage cluster sampling technique was employed in order to select study participants
149 (Fig 1). First, districts and CAs STH endemicity status was identified based on results of
150 previous STH mapping survey conducted at national level (where 2 had low and 15 had
151 moderate prevalence level [22]). We excluded 2 districts with low endemicity status since
152 they were not eligible for preventive chemotherapy. Two, 5 districts and 12 *kebeles* were
153 selected from 15 districts by simple random sampling technique (SRS). Three, list of eligible
154 households in each *kebeles* which had children between 1 and 5 years of age were identified
155 by health extension workers (HEWs). Finally, children from each *kebeles* were selected by
156 taking probability proportionate to sample size into account.

157 **Figure 1. Sampling profile**

158 **Study variables and data collection**

159 Variables included in this study were STH infection status (positive or negative), intensity of
160 infections, socio-demographic characteristics of parent and guardians, social determinants of
161 health, wealth status of household and child related variables and implementation of STH
162 control strategies. Data on these socio-demographic and other variables were collected
163 through face-to-face interviews using pre-tested questionnaire with heads of households or
164 mothers or guardians using pre-tested questionnaire. Stool specimens were examined using
165 the WHO recommended Kato-Katz kit under microscopy [23].

166 **Stool collection and processing**

167 Fresh stool specimens were collected using clean, leak proof and screw cup container, and
168 then collected stool samples were transported in cool boxes to nearby health facility for
169 examination. The stool samples were processed within two hours of receipt or saved in
170 Icebox where travel time exceeded two hours. Samples were examined in the local health

171 center by Kato-Katz technique to determine the prevalence and intensity of STH. Duplicate
172 slides were prepared per stool sample in order to ensure reliability.

173 **Quality assurance**

174 Data quality was monitored through standard operational procedure (SOPs), recruiting
175 competent data collectors, pre-testing tools, training data collectors and supervisors, daily
176 checking consistency and accuracy of collected data. The quality of data collection was
177 closely monitored by supervisors. Stools were examined by a qualified laboratory
178 technologist. For the purpose of bench aid, pictures of parasites egg were displayed on wall in
179 front of microscopy examination for the purpose of internal quality control.

180 **Statistical data analysis and measurements**

181 First, data were edited, coded and entered into EpiData 4.4.2, and then exported to SPSS
182 software (IBM, version 25) for analysis. A difference in prevalence of STH between or
183 among categories of variables was analyzed using chi-square test (χ^2). Summary statistics
184 were computed, and data were presented with tables and graphs using excel spreadsheet.

185 Children between age 1 and 5 years were defined as PSAC [23]. Wealth analysis was
186 performed as, initially, reliability test was performed using the economic variables involved
187 in measuring the wealth of the households. The variables which were used to compute the
188 alpha value were entered into the principal component analysis. At the end of the principal
189 component analysis, the wealth index was obtained as a continuous scale of relative wealth.
190 Then, quintiles of the wealth index were created.

191 We calculated the prevalence by dividing the number of STH positive PSAC by the total
192 number of participants. Intensity of STH infection is the number of helminths (worms)
193 infecting an individual; for each parasite species it was analyzed as light, moderate and
194 heavy infections based on eggs per gram of stool (EPG) and was classified according to the
195 WHO guidelines [24], as follow (Table 1).

196 **Table 1. Criteria for classifying intensity of STH infection for each species.**

STH infection	Intensity of infection (EPG)		
	Light	Moderate	Heavy
<i>A.lumbricoides</i>	(1–4999)	(5000–49999)	(≥50000)
<i>T.trichiura</i>	(1–999)	(1000–9999)	(≥10000)
<i>Hookworms</i>	(1–1999)	(2000–3999)	(≥4000)

197

198 According to WHO STH endemicity mapping classifications, there are three categories in
199 line with implementation of mass drug administration (MDA): i) high transmission (where
200 prevalence is > 50%), ii) moderate transmission (where prevalence is between 20%- 50%),
201 and iii) low transmission (where prevalence is < 20%) [23, 25].

202 **Ethical considerations**

203 Ethical approval was obtained from Institutional and Research Ethics review board of Arba
204 Minch University, College of Medicine and Health Sciences, Arba Minch, Ethiopia. Oral and
205 written consent was sought from district administrators and heads of the household before
206 survey was conducted. Children tested positive for one or more STH were treated with
207 Albendazole or Mebendazole by health workers.

208

209

210

211

212

213

214

215

216 **Results**

217 **Socio-demographic characteristics of PSAC and heads of household**

218 A total of 2462 (92%) PSAC participated in this study. Of the total, 246 (10%) were under 2
 219 years of age, and slightly more males participated than females (52% versus 48%). The
 220 average number of people living in a household was 5.5 (standard deviation \pm 1.8)
 221 individuals. More than half of HHs (57.5%) did not attend any formal education (Table 2).

222 **Table 2. Socio-demographic characteristics of PSAC and HHs, 2019 (N=2462).**

Variables	Category	Frequency	%
Child sex	Male	1281	52
	Female	1181	48
Child age (years)	Less than 2yrs	246	10
	2-5yrs	2216	90
Age of HH (years)	Less than 20	22	0.89
	20 -29	404	16.41
	30-39	1284	52.15
	40-49	647	26.28
	55-59	59	2.4
	60 and above	46	1.87
Sex of HH	Male	2174	88.3
	Female	288	11.7
Educational status	Can't read and write	1073	43.58
	Can read and write	343	13.93
	Elementary	508	20.63
	Secondary	279	11.33
	Diploma and above	259	10.52
Occupation	Farming	1606	65.23
	Employed	312	12.67
	Merchant	303	12.31
	Unemployed	78	3.17
	Others*	163	6.62
Family size	Less than 4	288	11.7
	4-6	1503	61.05
	7 and more	671	27.25
Residence	Urban	628	25.5
	Rural	1834	74.5
Wealth quintile	Wealthiest	488	19.82
	Wealthy	498	20.22
	Middle income	482	19.58
	Poor	498	20.23
	Poorest	496	20.15
Water source	Pipe	1128	45.82
	Well	293	11.9
	Public bono	971	39.44
	Other**	70	2.84

223 *=Daily laborer; **=river, well and spring

224 **Prevalence of STH infections**

225 Of the children surveyed, 23.5% (578/2462) had at least one type of STH infection.
226 Ascariasis was the most prevalent (18.6%), followed by trichiuriasis (9.2%) and hookworms
227 (3.1%). Mixed STH infections (*Ascaris lumbricoides* and *Trichuris trichiura*) were found in
228 7.4% of PSAC. The highest prevalence of any one of STH infection was observed in
229 Chench district (33.8 %), as contrasted to the lowest prevalence that was found in Demba
230 Gofa district (11%). In Deremalo district, considerable amount of hookworm infections
231 (10%) were revealed (**Fig 2**).

232 **Fig 2. Prevalence of any STH infection by district among PSAC, Gamo Gofa zone, Southern**
233 **Ethiopia, 2019 (N=2462).**

234 The prevalence of STH infections across the age-group (1 – 5 years) was slightly higher at ≤ 2
235 years while comparing across the age-group (1-5 years). On the other hand, prevalence of any
236 STH infections among females (24.3%) was a little higher than the male's (22.7%), and a
237 higher prevalence of STH infections (24.6%) in rural area was noticed than urban (20.1%).
238 Children who did not receive PC in the last one year had slightly higher prevalence of STH
239 infections compared to those who received PC (23.9% versus 18.7%), (Table 3).

240

241

242

243

244

245

246

247 **Table 3. Prevalence of STH infections in association with different factors among PSAC in**
 248 **Gamo Gofa zone, 2019 (N=2462).**

Variable	Category	STH Infection status		X2-test	P-value
		Negative, n (%)	Positive, n (%)		
Child sex	Male	990(77.3)	291(22.7)	0.354	0.366
	Female	894(75.7)	287(24.3)		
Child age group (years)	Less than 2yrs	187(76)	59(24)	0.039	0.843
	2-5yrs	1697(76.6)	519(23.4)		
Place of residence	Urban	502(79.9)	126(20.1)	5.467	0.019
	Rural	1382(75.4)	452(24.6)		
Child age (years)	1	187(76)	59(24)	0.914	0.923
	2	414(75.1)	137(24.9)		
	3	563(77)	168(23)		
	4	522(77.2)	154(22.8)		
	5	198(76.7)	60(23.3)		
Dewormed in the last one year	No	608(78.9)	163(21.1)	3.408	0.065
	Yes	1276(75.5)	415(24.5)		
Child soil eating habit	No	1518(76.9)	457(23.1)	0.371	0.542
	Yes	261(78.4)	72(21.6)		
Family size	<4	220(76.4)	68(23.6)	2.219	0.330
	4-6	1164(77.4)	339(22.6)		
	7 and above	500(74.5)	171(25.5)		
Mothers' (guardians') educational status	Can't read and write	774(76.9)	233(23.1)	5.432	0.246
	Can read and write	291(73.9)	103(26.1)		
	Elementary	486(76.1)	153(23.9)		
	Secondary	225(76.8)	68(23.2)		
	Diploma and above	108(83.7)	21(16.3)		
Mothers' (guardians') occupation	Farming	974(74.6)	331(25.4)	16.288	0.003
	Employed (Gov.)	166(81.8)	37(18.2)		
	Merchant	348(73.7)	124(26.3)		
	Unemployed	334(82.3)	72(17.7)		
	Others*	62(81.6)	14(18.4)		
Wealth quintile	Wealthiest	379(77.7)	109(22.3)	1.372	0.712
	Wealthy	379(76.1)	119(23.9)		
	Middle income	374(77.6)	108(22.4)		
	Poor	369(74.1)	129(25.9)		
	Poorest	383(77.2)	113(22.8)		
Water source	Pipe	891(79)	237(21)	8.289	0.04
	Well	223(76.1)	70(23.9)		
	Public bono	721(74.3)	250(25.7)		
	Other**	49(70)	21(30)		

249 * = Daily laborer and housewife

** = River, well and spring

251 **Intensity of STH infections**

252 In majority of STH infections (85%) low infection intensity was found as a result of
253 *hookworms* and *Trichuris trichiura* infection while 15% of Ascariasis had moderate infection
254 intensity (Table 4). All of the moderate infections were from Chencha and Bonke district.

255 **Table 4. Intensity of STH infection among PSAC in Gamo Gofa zone, southern Ethiopia, 2019**
256 **(N=2462).**

Type of STH infection	Mean (EPG)	Intensity of infection		
		Light	Moderate	Total infected PSAC
Ascariasis	2152	388(84.90)	69(15.1)	457
Hookworm	154	76(100)	0	76
Trichiuriasis	135	226(100)	0	226

257 **Control strategies of STH**

258 **Treatment coverage of Albendazole**

259 The treatment coverage of Albendazole (ALB) in the last one year before the survey against
260 STH among PSAC was 68.7% (1691/2462) (**Fig 3**).

261 **Figure 3. Treatment coverage of Albendazole against STH in Gamo-Gofa Zone, 2019 (N=2462).**

262 **Knowledge and practice (KP) of mothers or guardians on STH**

263 Of the total surveyed mothers or guardians, the mean KP score (≤ 5) on transmission of STH
264 was 92% (2262/2462); 42.7% (1052/2462) did not wash hand after defecation, and 77.7%
265 (1913/2462) got information on STH from health extension workers (HEWs) (Table 5).

266

267

268

269

270

271

272

273 **Table 5. Knowledge and practice on STH transmission and prevention among mothers, 2019**
 274 **(N=2462).**

Variables	Category	Frequency	%
Know about STH	Yes	2208	89.7
	No	254	10.3
Know at least one STH transmission (n=2208)	Yes	2153	97.5
	No	55	2.5
Know at least one STH prevention way (n=2208)	Yes	2135	96.7
	No	73	13.5
Mean KP score on transmission STH	≤5	2262	91.9
	>5	200	8.1
Mean KP score on prevention STH	≤5	2283	92.7
	>5	179	7.3
Get information about STH through	Health facility	271	11
	HEWs	1913	77.7
	Radio or TV	24	1
	None	254	10.3
Wash hand after latrine	Yes	1410	57.3
	No	1052	42.7
Wash hand before meal	Yes	2347	95.3
	No	2462	4.7
Wash hand after cleaning child	Yes	2229	90.5
	No	233	9.5
Wash hand before cooking	Yes	2069	84
	No	393	16
Wash fruit before eating	Yes	1841	74.8
	No	621	25.2
Wash hand after work	Yes	2253	91.5
	No	209	8.5

275

276

277

278

279

280

281

282

283 **Water, Sanitation and Hygiene (WASH)**

284 This study also revealed households data as, 39.4% used water from hand-dug well, 52.5% of
 285 need to move ≥ 30 minutes to collect water, 77.5% did not use treat water, and 48.9% did not
 286 own hand washing facility (Table 6).

287 **Table 6. WASH characteristics of households in Gamo Gofa zone, 2019.**

Variables	Category	Frequency	%
Source of water	Pipe	1128	45.82
	Well	293	11.9
	Public bono	971	39.44
	Other*	70	2.84
Distance form water source	<30min	1170	47.5
	≥ 30 min	1292	52.5
Adequate water	No	536	21.8
	Yes	1926	78.2
Ever treat water	No	1909	77.5
	Yes	553	22.5
Latrine	Yes	2397	97.4
	No	65	2.6
Type of latrine (n=2397)	Pit	2360	98.5
	Improved pit latrine	37	1.5
Feces observed inside the latrine compound	No	1766	73.67
	Yes	631	26.32
Hand washing facility (functional)	No	1225	51.1
	Yes	1172	48.9
Soap or ash available (n=1172)	Yes	298	25.4
	No	874	74.6
Reason not latrine available (n=65)	No space	6	9.2
	No money	10	15.4
	No skill	9	13.8
	Did not know importance	40	61.5
If no latrine where you defecate (n=65)	Open field	56	86.2
	Public	9	13.8

288 *=river, well and spring

289

290

291

292

293

294 Discussion

295 This study provides operational evidence on prevalence, intensity and control strategies of
296 soil-transmitted helminth infections (STH) to improve control strategies of soil-transmitted
297 helminths in Gamo Gofa zone. Our study showed that a significant proportion of pre-school
298 age children are suffering from STH infections in the study area despite preventive
299 chemotherapy distribution for more than 10 years. Considerable variation in prevalence, and
300 intensity of infections across districts were also noticed. In addition, gaps in the control
301 strategies of STH to control and eliminate STH were highlighted.

302 In this study, the overall prevalence of STH infections with at least one STH parasite was
303 23.5%, would be classified into the moderate transmission category (where prevalence is
304 between 20% and 50%), and qualifies the requirement of annual STH mass drug
305 administration [17]. Ascariasis was the most prevalent infection (18.6%), followed by
306 trichiuriasis (9.2%) and hookworm (3.1%). All infections of *hookworm* and *Trichuris*
307 *trichiura* had low intensity while Ascariasis had moderate intensity (15 %). Despite initiation
308 of preventive chemotherapy about 10 years ago, the burden of STH did not show significant
309 reduction probably due to weak implementation of control strategies such as social behavioral
310 change communication (SBCC), inadequate mass drug administration coverage and WASH
311 as highlighted in the findings of this study.

312 The overall prevalence of STH infections observed in this study is comparable with studies
313 conducted in another parts of Ethiopia (Butajira and Wonji) and West China [26-28], but
314 slightly lower than the prevalence reported in Dembiya, northwest Ethiopia [21]. In addition,
315 the prevalence in our study is significantly lower than the findings of other studies conducted
316 in another part of Ethiopia and in Nigeria, Cameroon, Ecuador, Uganda, Kenya and Honduras
317 [29-36]. These differences observed from our study could be due to variation in socio-

318 cultural, social determinants, behavioral characteristics and implementation of prevention and
319 control measures.

320 In this study, Ascariasis was identified as the commonest species of STH, and this findings
321 supports studies done in other part of Ethiopia, Nigeria and China [21, 26, 28, 29]. On the
322 contrast, some other studies conducted in Ethiopia, Ecuador and Honduras showed high
323 prevalence of Trichiuriasis [32, 35, 37], and a study conducted in Uganda showed high
324 prevalence of hookworm [33]. These differences might be related to variation in
325 environmental factors such as climate, rainfall, topography, surface temperature, altitude, and
326 soil type which have a great impact on the distribution of STH [38]. Moreover, in this study
327 we found significant amounts of mixed infections (7.4%) of PSAC were infected with two
328 STH species (*Ascaris lumbricoides* and *Trichuris trichiura*); this finding is in line with a
329 study conducted in other areas of Ethiopia and Nigeria [26, 27, 29].

330 In our study, slightly higher prevalence at age ≤ 2 years was observed while comparing across
331 the age-group (1-5 years); the possible explanation related to this difference is due to the fact
332 that current mass drug administration among PSAC often do not include age ≤ 2 years. On the
333 contrast, other studies revealed numerical increase in prevalence of STH as age increase [26,
334 29]. In addition, in this study, prevalence of any STH infection among females (24.3%) was a
335 little higher than the male's prevalence (22.7%), this result supports the findings of a study
336 conducted in another part of Ethiopia [26]. These differences might be due to variation in
337 socio-cultural pattern, access and uptake of preventive chemotherapy.

338 In addition, significant proportion (15%) of moderate intensity Ascariasis was observed in
339 our study, and this finding is higher than results of a study conducted in Butajira, Ethiopia
340 and Honduras [26, 35]. The possible explanation related to this difference could be
341 consistency and frequency of mass drug administration that may affect intensity of infections
342 [23].

343 Furthermore, findings of our study showed inadequate implementation of control strategies.
344 The treatment coverage of Albendazole (ALB) in this survey against STH among PSAC was
345 (68.7%); which is lower than the national coverage (71%) of Ethiopia and WHO's target
346 (minimum of 75%) [10, 39]. Most importantly, study participants in this study were pre-
347 school age children, who might not be reached out by the deworming program, especially
348 those under 2 years old. On the other hand, obviously, school age children could have better
349 chance to be reached by school-based deworming.

350 The importance of WASH intervention to control and eliminate STH was reported in
351 different studies though site-specific data are required in our case [40-42]. However, in this
352 study inadequate WASH infrastructures were observed at household level where 39.4% were
353 using water from well, 52.5% were walking ≥ 30 minutes to collect water, 77.5 % did not treat
354 water, and 48.9% of not had hand washing facility. Of the total surveyed mothers or
355 guardians, the mean KP score (≤ 5) on transmission of STH was 92% (2262/2462); 42.7%
356 (1052/2462) did not wash hand after defecation, and 77.7% (1913/2462) got information on
357 STH from health extension workers (HEWs). The possible reason for these findings could be
358 related to weak social behavioral change communication (SBCC) intervention.

359 The most outstanding strength of our study is that it is addressing an important national
360 operational research priority which is focusing on parasitological monitoring and control
361 strategies of STH among pre-school age children.

362 In this study, there are limitations that need to be taken into account. There might be
363 underestimation of prevalence of STH due to the fact that [1] we collected single stool
364 specimen, which could cause variation in egg excretion over different times (hours) within a
365 day and across different days; [2] samples were collected from remote villages and there
366 might be rapid desiccation of hookworm eggs in the stool samples, and [3] even though the
367 Kato-Katz technique is sensitive in detecting moderate and high infection intensity (MHI), it

368 has lower detection power and therefore lower positive predictive values in low-prevalence
369 (and low-intensity) settings [28, 43].

370 **Conclusion**

371 Data from our study showed that substantive proportion of pre-school age children in the
372 study area are suffering from significant burden of STH infections despite provision of
373 preventive chemotherapy distribution for more than 10 years. Considerable variation in
374 prevalence and intensity of infections across districts and *kebeles* was noticed. Moreover, the
375 study highlights the need for implementing control strategies of STH in a coordinated and
376 regular manner in order to achieve the national goal of STH control and elimination. Further,
377 operational research needs to be conducted in different settings to monitor and evaluate
378 progress towards elimination and control of STH among PSAC.

379 **Funding**

380 This study is made possible by the generous research grant support of collaborative research
381 and training center for NTDs, Arba Minch University.

382 **Competing interests**

383 Authors declared that they have no conflicts of interest.

384 **Acknowledgements**

385 Authors would like to thank study participants, data collectors, supervisors, zonal health
386 office heads, district health office heads, NTDs focal persons in all selected districts of the
387 study area and Ethiopian public health institute for their support in the implementation of the
388 study.

389 **Author's contributions**

390 MA prepared the original draft. MA, TG, TW, AB, ZH, and TG⁵ involved in conception,
391 protocol development and design of the study. MA, TW and TG involved in supervision of

392 the project. MA and NM performed data cleaning and analysis. All authors were involved in
393 the interpretation of the data. MA, TW and TG⁵ critically reviewed the initial manuscript. All
394 authors reviewed and approved the final manuscript.

395 **References**

- 396 1. Montresor A, Trouleau W, Mupfasoni D, Bangert M, Joseph SA, Mikhailov A, et al.
397 Preventive chemotherapy to control soil-transmitted helminthiasis averted more than
398 500 000 DALYs in 2015. *Transactions of the Royal Society of Tropical Medicine and*
399 *Hygiene*. 2017 Oct 1; 111(10):457-63.
- 400 2. Hotez PJ, Fenwick A, Savioli L, Molyneux DH. Rescuing the bottom billion through
401 control of neglected tropical diseases. *The Lancet*. 2009 May 2; 373(9674):1570-5.
- 402 3. World Health Organization. Ending the neglect to attain the sustainable development
403 goals: A road map for neglected tropical diseases 2021–2030. World Health
404 Organization; 2020.
- 405 4. De Silva NR, Brooker S, Hotez PJ, Montresor A, Engels D, Savioli L. Soil-
406 transmitted helminth infections: updating the global picture. *Trends in parasitology*.
407 2003 Dec 1;19(12):547-51.
- 408 5. Bethony J, Brooker S, Albonico M, Geiger SM, Loukas A, Diemert D, et al. Soil-
409 transmitted helminth infections: ascariasis, trichuriasis, and hookworm. *The lancet*.
410 2006 May 6; 367(9521):1521-32.
- 411 6. Albonico M, Allen H, Chitsulo L, Engels D, Gabrielli AF, Savioli L. Controlling soil-
412 transmitted helminthiasis in pre-school-age children through preventive
413 chemotherapy. *PLoS neglected tropical diseases*. 2008 Mar; 2(3).
- 414 7. Schistosomiasis WH. soil-transmitted helminth infections—preliminary estimates of
415 the number of children treated with albendazole or mebendazole. *Wkly Epidemiol*
416 *Rec*. 2006 Apr 21; 81(16):145-64.


- 417 8. Pullan RL, Brooker SJ. The global limits and population at risk of soil-transmitted
418 helminth infections in 2010. *Parasites & vectors*. 2012 Dec 1; 5(1):81.
- 419 9. Brooker S. Estimating the global distribution and disease burden of intestinal
420 nematode infections: adding up the numbers—a review. *International journal for*
421 *parasitology*. 2010 Aug 15; 40(10):1137-44.
- 422 10. World Health Organization. Guideline: preventive chemotherapy to control soil-
423 transmitted helminth infections in at-risk population groups. World Health
424 Organization; 2017.
- 425 11. World Health Organization. Intestinal worms: Soil-transmitted helminthes World
426 Health Organization; 2011.
- 427 12. Federal Democratic Republic of Ethiopia Ministry of Health. Second Edition of
428 Ethiopia National Master Plan For Neglected Tropical Diseases. Addis Ababa,
429 Ethiopia. 2016.
- 430 13. Tchuente LA. Control of schistosomiasis and soil-transmitted helminthiasis in sub-
431 Saharan Africa: challenges and prospects. *Current topics in tropical medicine*. 2012
432 Mar 16:359-76.
- 433 14. Stephenson LS, Latham MC, Ottesen EA. Malnutrition and parasitic helminth
434 infections. *Parasitology*. 2000 Oct; 121(S1):S23-38.
- 435 15. Soeripto N. Reinfection and infection rates of soil-transmitted-helminths in Kemiri
436 Sewu, Yogyakarta, Indonesia. *Southeast Asian J Trop Med Public Health*. 1991 Jun;
437 22(2):216-21.
- 438 16. Schistosomiasis WH. soil-transmitted helminth infections—preliminary estimates of
439 the number of children treated with albendazole or mebendazole. *Wkly Epidemiol*
440 *Rec*. 2006 Apr 21; 81(16):145-64.


- 441 17. World Health Organization. Soil-transmitted helminthiasis: eliminating as public
442 health problem soil-transmitted helminthiasis in children: progress report 2001-2010
443 and strategic plan 2011-2020.
- 444 18. World Health Organization. Universal health coverage. WHO fact sheet. Geneva.
445 World Health Organization; 2019.
- 446 19. Federal Democratic Republic of Ethiopia Ministry of Health. Ethiopian National
447 Master Plan for Neglected Tropical Diseases (2013-2015). Addis Ababa, Ethiopia;
448 2013.
- 449 20. Central statistical agency (2013) Population Projection for Ethiopia 2007-2037. Addis
450 Ababa, Ethiopia.
- 451 21. Gizaw Z, Adane T, Azanaw J, Addisu A, Haile D. Childhood intestinal parasitic
452 infection and sanitation predictors in rural Dembiya, northwest Ethiopia.
453 Environmental health and preventive medicine. 2018 Dec; 23(1):26.
- 454 22. Leta GT, Mekete K, Wuletaw Y, Gebretsadik A, Sime H, Mekasha S, et al. National
455 mapping of soil-transmitted helminth and schistosome infections in Ethiopia.
- 456 23. World Health Organization. Preventive chemotherapy in human helminthiasis.
457 Coordinated use of anthelmintic drugs in control interventions: a manual for health
458 professionals and programme managers. World Health Organization; 2006.
- 459 24. World Health Organization. Prevention and control of schistosomiasis and soil-
460 transmitted helminthiasis: report of a WHO expert committee.
- 461 25. Helminthiasis WS. Eliminating soil-transmitted helminthiasis as a public health
462 problem in children: progress report 2001–2010 and strategic plan 2011–2020.
463 France: World Health Organization. 2012:19-29.

- 464 26. Shumbej T, Belay T, Mekonnen Z, Tefera T, Zemene E. Soil-transmitted helminths
465 and associated factors among pre-school children in Butajira Town, South-Central
466 Ethiopia: a community-based cross-sectional study. *PloS one*. 2015; 10(8).
- 467 27. Degarege A, Erko B. Prevalence of intestinal parasitic infections among children
468 under five years of age with emphasis on *Schistosoma mansoni* in Wonji Shoa Sugar
469 Estate, Ethiopia. *PloS one*. 2014; 9(10).
- 470 28. Wang X, Zhang L, Luo R, Wang G, Chen Y, Medina A, et al. Soil-transmitted
471 helminth infections and correlated risk factors in preschool and school-aged children
472 in rural southwest China. *PLoS One*. 2012; 7(9).
- 473 29. Sowemimo OA, Asaolu SO. Current status of soil-transmitted helminthiasis among
474 pre-school and school-aged children from Ile-Ife, Osun State, Nigeria. *Journal of*
475 *helminthology*. 2011 Sep; 85(3):234-8.
- 476 30. Nkengazong L, Njiokou F, Wanji S, Teukeng F, Enyong P, Asonganyi T. Prevalence
477 of soil transmitted helminths and impact of Albendazole on parasitic indices in Kotto
478 Barombi and Marumba II villages (South-West Cameroon). *African Journal of*
479 *Environmental Science and Technology*. 2010; 4(3).
- 480 31. Dana D, Mekonnen Z, Emanu D, Ayana M, Getachew M, Workneh N, et al.
481 Prevalence and intensity of soil-transmitted helminth infections among pre-school age
482 children in 12 kindergartens in Jimma Town, southwest Ethiopia. *Transactions of the*
483 *Royal Society of Tropical Medicine and Hygiene*. 2015 Mar 1; 109(3):225-7.
- 484 32. Moncayo AL, Lovato R, Cooper PJ. Soil-transmitted helminth infections and
485 nutritional status in Ecuador: findings from a national survey and implications for
486 control strategies. *BMJ open*. 2018 Apr 1; 8(4):e021319.
- 487 33. Ojja S, Kisaka S, Ediau M, Tuhebwe D, Kisakye AN, Halage AA, et al. Prevalence,
488 intensity and factors associated with soil-transmitted helminths infections among

- 489 preschool-age children in Hoima district, rural western Uganda. *BMC infectious*
490 *diseases*. 2018 Dec 1; 18(1):408.
- 491 34. Davis SM, Worrell CM, Wiegand RE, Odero KO, Suchdev PS, Ruth LJ, et al. Soil-
492 transmitted helminths in pre-school-aged and school-aged children in an urban slum:
493 a cross-sectional study of prevalence, distribution, and associated exposures. *The*
494 *American journal of tropical medicine and hygiene*. 2014 Nov 5; 91(5):1002-10.
- 495 35. Torres RE, Garcia DN, Sandoval GA, Santana AH, Singh P, Bucheli ST, et al.
496 Prevalence and intensity of soil-transmitted helminthiasis, prevalence of malaria and
497 nutritional status of school going children in Honduras. *PLoS neglected tropical*
498 *diseases*. 2014 Oct; 8(10).
- 499 36. Tekalign E, Bajiro M, Ayana M, Tiruneh A, Belay T. Prevalence and Intensity of
500 Soil-Transmitted Helminth Infection among Rural Community of Southwest Ethiopia:
501 A Community-Based Study. *BioMed Research International*. 2019; 2019.
- 502 37. Nyantekyi LA, Legesse M, Belay M, Tadesse K, Manaye K, Macias C, et al.
503 Intestinal parasitic infections among under-five children and maternal awareness
504 about the infections in Shesha Kekele, Wondo Genet, Southern Ethiopia. *Ethiopian*
505 *Journal of Health Development*. 2010; 24(3).
- 506 38. Hotez PJ, Brindley PJ, Bethony JM, King CH, Pearce EJ, Jacobson J. Helminth
507 infections: the great neglected tropical diseases. *The Journal of clinical investigation*.
508 2008 Apr 1; 118(4):1311-21.
- 509 39. Asfaw M, Zerdo Z, Churko C, Seife F, Yihune M, Chisha Y, et al. Preventive
510 chemotherapy coverage against Soil-transmitted helminthiasis among school age
511 children in vertical versus integrated treatment approaches: Implications from
512 coverage validation survey in Ethiopia. *medRxiv*. 2020 Jan 1.

- 513 40. Freeman MC, Ogden S, Jacobson J, Abbott D, Addiss DG, Amnie AG, et al.
514 Integration of water, sanitation, and hygiene for the prevention and control of
515 neglected tropical diseases: a rationale for inter-sectoral collaboration. PLoS
516 neglected tropical diseases. 2013 Sep; 7(9).
- 517 41. Ziegelbauer K, Speich B, Mäusezahl D, Bos R, Keiser J, Utzinger J. Effect of
518 sanitation on soil-transmitted helminth infection: systematic review and meta-
519 analysis. PLoS Med. 2012 Jan 24; 9(1):e1001162.
- 520 42. Becker SL, Liwanag HJ, Snyder JS, Akogun O, Belizario Jr V, Freeman MC,
521 Gyorkos TW, Imtiaz R, Keiser J, Krolewiecki A, Levecke B. Toward the 2020 goal of
522 soil-transmitted helminthiasis control and elimination. PLoS neglected tropical
523 diseases. 2018 Aug; 12(8).
- 524 43. Freeman MC, Akogun O, Belizario Jr V, Brooker SJ, Gyorkos TW, Imtiaz R, et al.
525 Challenges and opportunities for control and elimination of soil-transmitted helminth
526 infection beyond 2020. PLoS neglected tropical diseases. 2019 Apr; 13(4).





Swallow ALB (%)

