The Lancet Infectious Diseases

The Presence of COVID-19 in Urine: A Systematic Review and Meta-analysis of the Literature
--Manuscript Draft--

Manuscript Number:	
Article Type:	Article (Meta-analysis)
Keywords:	COVID-19; SARS-Cov-2; urine; review
Corresponding Author:	Maryam Vaezjalali, Associate Professor Shaheed Beheshti University of Medical Sciences Tehran, IRAN, ISLAMIC REPUBLIC OF
First Author:	Amir H Kashi, M.D.
Order of Authors:	Amir H Kashi, M.D.
	Morteza Fallah-karkan
	Erfan Amini, Associate Professor
	Maryam Vaezjalali, Associate Professor
Manuscript Region of Origin:	IRAN, ISLAMIC REPUBLIC OF
Abstract:	Purpose: To investigate the literature on the presence of COVID-19 virus in urine of infected patients and evaluate the attributes and clinical significance of COVID-19 in urine including probability of infection transmission through urine. Data sources: A systematic review of literature from December 2019 to 6 th May 2020 was conducted on Pubmed, google scholar, ovid, scopus and ISI web of science. Study eligibility criteria: Studies which investigated urinary viral shedding of COVID-19 in infected patients were included. Study appraisal and synthesis methods: Two reviewers selected relative studies and performed quality assessment of individual studies. Meta-analysis was performed the pooled case reports and case series. Fixed-effects model was used for analysis as no significant heterogeneity was observed between studies. Results: Thirty three studies were finally included in the systematic review including 12 case reports, 20 case series, and one cohort. Urinary samples from 430 patients were investigated. Ten studies reported the presence of COVID-19 in urinary samples from 16 patients. The rate of COVID-19 presence in urinary samples was 3.7%. Urinary viral load was low in most reports. The presence of virus in urine was not related to the disease course of the illness. Urinary COVID-19 was mostly detected from patients with moderate to severe disease (13 pts) but was also isolated from two children (one neonate and one 7 year-old girl) and one adult with mild disease. The pathogenicity of virus isolated from urine has been demonstrated in cell culture media in one study. Conclusions: This review highlights the low frequency of COVID-19 presence in urine of infected individuals and the potential of isolated virus for cytopathic effects. Therefore the probability of infection transmission through urine can be suggested. Caution must be exerted when dealing with urine of patients infected with COVID-19 including medical interventions like endoscopy and urethral catheterization.

The Presence of COVID-19 in Urine: A Systematic Review and Meta-analysis of the Literature

Amir H Kashi[1], Morteza Fallah-karkan[2], Erfan Amini[3], Maryam Vaezjalali[4]

1 Urology and Nephrology Research Center (UNRC), Shahid Labbafinejad Hospital, Shahid Beheshti University of Medical Sciences, Tehran, Iran.

2 Department of Urology, Shohada Tajrish Hospital, Shahid Beheshti University of Medical Sciences (SBMU), Tehran, Iran.

3 Uro-Oncology Research Center, Tehran University of Medical Sciences, Tehran, Iran.

4 Department of Microbiology, School of Medicine, Shahid Beheshti University of Medical Sciences, Tehran, Iran.

Address for correspondence:

Maryam Vaezjalali, Department of Microbiology, School of Medicine, Shahid Beheshti University of Medical Sciences, Tehran, Iran. Maryam.vaezjalali@gmail.com

ABSTRACT

Purpose: To investigate the literature on the presence of COVID-19 virus in urine of infected patients and evaluate the attributes and clinical significance of COVID-19 in urine including probability of infection transmission through urine.

Data sources: A systematic review of literature from December 2019 to 6[th] May 2020 was conducted on Pubmed, google scholar, ovid, scopus and ISI web of science.

Study eligibility criteria: Studies which investigated urinary viral shedding of COVID-19 in infected patients were included.

Study appraisal and synthesis methods: Two reviewers selected relative studies and performed quality assessment of individual studies. Meta-analysis was performed the pooled case reports and case series. Fixed-effects model was used for analysis as no significant heterogeneity was observed between studies.

Results: Thirty three studies were finally included in the systematic review including 12 case reports, 20 case series, and one cohort. Urinary samples from 430 patients were investigated. Ten studies reported the presence of COVID-19 in urinary samples from 16 patients. The rate of COVID-19 presence in urinary samples was 3.7%. Urinary viral load was low in most reports. The presence of virus in urine was not related to the disease course of the illness. Urinary COVID-19 was mostly detected from patients with moderate to severe disease (13 pts) but was also isolated from two children (one neonate and one 7 year-old girl) and one adult with mild disease. The pathogenicity of virus isolated from urine has been demonstrated in cell culture media in one study.

Conclusions: This review highlights the low frequency of COVID-19 presence in urine of infected individuals and the potential of isolated virus for cytopathic effects. Therefore the probability of infection transmission through urine can be suggested. Caution must be exerted when dealing with urine of patients infected with COVID-19 including medical interventions like endoscopy and urethral catheterization.

Keywords: COVID-19; SARS-Cov-2; urine; review

INTRODUCTION

Novel coronavirus disease (COVID-19) first reported form Wuhan is a new disease caused by severe acute respiratory syndrome- coronavirus-2 (SARS-CoV-2), manifesting as an acute respiratory illness, however the involvement of multiple organs including kidney has been reported[1]. The pathophysiological mechanisms for SARS-CoV-2 infection and organ invasion are not yet fully discovered, which leads to difficulties in understanding routes of transmission, clinical diagnosis and treatment[2].

Genomic sequence analysis indicated that SARS-CoV-2 has 75%-80% genomic similarity to coronavirus causative agent for severe acute respiratory syndrome (SARS) namely SARS-CoV[3]. In previous reports of SARS and the Middle East respiratory syndrome coronavirus (MERS-CoV) infections, acute kidney injury was observed in 5% to 15% cases and was associated with a high (60%–90%) mortality rate[4].

The angiotensin-converting enzyme 2 (ACE2), known to be a cell receptor for human SARS-CoV, is also reported to play the same role for cellular entry of SARS-CoV-2 [5]. In addition to respiratory organs, upregulation of ACE2 expression was also identified in urogenital system including kidney proximal tubule cells, bladder urothelial cells [6] and genital organs including testis[7,8].

The widely accepted routes of human to human transmission for COVID-19 are through respiratory droplets and direct contact, however viral shedding in urine has been reported and infection transmission through infected urine remains a possibility. The idea of virus transmission thorough urine originated form the homogeneity of viral SARS-CoV-2 genome with SARS virus and the abundant previous evidence on the presence of SARS virus in urine[9,10]. Original protocols for sample collections from COVID-19 patients included urine sample collection[11]. Nevertheless,

the mechanism of virus shedding urine is unclear. Two suggested mechanisms for SARS-CoV-2 shedding in urine have been proposed: Firstly, Sepsis and cytokine storm results in renal dysfunction and subsequent leakage of SARS-CoV-2 from circulation into urine; Secondly, virus may directly invade the urinary system via binding to ACE2 receptors and shed into the urine [12].

Although the virus shedding into the urine is hypothetically probable, most studies showed that virus is absent in the urine of infected patients[6,13-15]. However, contradicting results exist in the literature as virus shedding was found in some studies [1,16,17].

Therefore, we performed a systematic review on the published literature to provide a summary of evidence on detection of SARS-CoV-2 in urinary samples.

METHODS

Search strategy and data sources

We conducted a comprehensive systematic literature review of online databases, including Web of Science, PubMed, Scopus, Ovid, and google scholar from 1st December 2019 till 6th May 2020. The search was performed by two independent investigators. The search terms used was: "(covid-19 OR ncovid-19 OR sars-cov-2 OR covid OR ncovid) AND urine".

Database searching was started on March 29th 2020 and was regularly updated during extraction and analysis of retrieved studies to find newly published articles. The latest electronic search on cited databases was performed on May 6th, 2020.

References of retrieved articles were manually searched to find eligible studies. The search and selection criteria were restricted to English language.

Study selection

The title and abstract of retrieved studies were screened through two different researchers independently. After removing duplicates and irrelevant studies, the full texts of articles were examined for presence of original data on the presence of COVID-19 in urine (Figure 1). Any disagreement was resolved by a third person. Personal viewpoints, opinion articles, correspondence, and letters not presenting original data were excluded as well as studies which did not report their result of urinary testing for COVID-19. Locations of studies was noted to identify duplicate case reports/series from the same area. When there was reports from the same area or suspicion of reports from same population of patients, authors were contacted to confirm independence in population of patients. All the search results were evaluated in accordance with the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) statement. The results of the search revealed 33 studies suitable for meta-analysis.

Data Extraction

The main outcome in this study is the evaluation of virus shedding into the urine of patients infected with COVID-19. Data were extracted from the eligible manuscripts into pre-defined data-fields including study location, sample size, mean or median age, gender of patients, illness category, total number of patients and/or urine samples tested, urine assessment technique, total number of positive urine samples, and sampling time.

Quality assessment of included studies

The included studies were evaluated in terms of quality according to the quality assessment tool for case series reported by the National Heart, Lung, and Blood Institute form the National

Institutes of Health[18]. This tools evaluates the quality based on a 9 item questionnaire. The questions focus on study population description, case definition, methods of including cases, comparability of included cases, description of interventions or assessments, follow-up and statistical methods used. The total score ranges from 0-9. Scores ≤ 3 were considered poor. Score of 4-6 was considered fair and score of 7-9 was considered good in terms of quality (Table 1).

Statistical methods

Extracted data from 33 studies were used for statistical analysis. Case reports and case series with a sample size of 6 or less were cumulated into one case series with sample size of 37 including 12 case reports and 7 case series. Thirteen case series and one cohort study with sample size of 9 or higher were also included in meta-analysis.

The effect size of individual studies was calculated by weighting each one of them by its inverse variance, and a confidence interval (CI) was thus obtained[19]. Each study was weighted inversely proportional to its variance. To calculate the variance of each study, a binomial distribution was used. To investigate heterogeneity, the Q statistics and I2 index with α significance level of less than 10% were used. In this study, the random-effects model was considered, when there is heterogeneity among the studies (I2> 50%). The authors used the Egger's test to check publication bias. In this study, Metaprop command in STATA used to stabilize the variances[20]. STATA software (version 16) was used to analyze the data.

The relationship between disease severity in each study and frequency of viral shedding in urine was investigated by weighting each study according to its sample size and performing spearman correlation.

RESULTS

A total of 1112 articles were retrieved using the search strategy mainly through google scholar search engine. After studying the title and abstract of studies, and removing duplicate studies, the number was reduced to 146 studies. Full text of these 146 studies were studied and non-original studies including communications without original data, personal reviews and letters were excluded resulting in 33 articles. (Figure 1) These 33 articles include one formally unpublished article retrieved by google scholar search engine from a database of unpublished studies (medRxiv).

There were 12 case reports, 20 case series, and one cohort study from 12 different countries. The characteristics of the included studies are shown in Table 1. Out of the total reported population of 1870 patients in these 33 studies, urinary testing has been performed on 430 patients during admission and up to day 52 after illness onset. Positivity of urinary specimens were reported in 10 studies ranging from 1 to 4 positive urinary samples in each study summing to a total of 16 patients (3.7% frequency of viral shedding in patients' urine). Positive urinary samples were reported form China (8 studies, 14 patients) and Korea (2 studies, 2 patients). The time of urine sampling in positive patients were reported as admission day+day 10, day 7, day 9, days 6 through 17, days 9 through 12, day 30, and day 52 after illness onset. The patient urine sample that was considered

positive in day 52 in this review was not positive on urine but was positive on urinary sediment. Excluding studies with a sample size of 1-2 patients, the percent positivity of COVID-19 in urine samples of different included studies varies from zero in most reports to as high as 11 percent in a 2 reports of 9 patients from Guangdong, and Guangzhou[2,21] on adults and children.

The meta-analysis forest plot which takes into account the weight of each study according to the inverse of its variance revealed a pooled estimate of 1.23% (CI 95%: 0.12 − 3.06%) for viral shedding in urine of patients (Figure 2). Meta-regression analysis revealed that the positivity rate of urinary viral shedding was not related to sample size of studies (graph not shown). The Begg's funnel plot failed to reveal publication bias (Figure 3).

One of the studies which confirmed the presence of viral RNA of the COVID-19 in urine of a patient stated that the positivity of the urine did not meet the reference for positivity in rRT-PCR (real time reverse transcriptase PCR), however in the diagram of patient diagnostic tests the level of detected E and RdRp genes at days 9 and 12 of illness were higher than cut off value in rRT-PCR. This patient urine sample was considered positive in the current review. The detection of COVID-19 RNA was observed from the end of the first week after illness till patient improvement[22].

Peng et al. reported the presence of COVID-19 in urine of one of the 9 studied patients on day 7 after symptom onset. The patient urine sample turned negative on day 10 after symptom onset. In this study, virus quantity in urine sample was lower than rectal and oropharyngeal samples. The patient with positive urinary PCR for COVID-19 did not complain of any urinary symptoms[2].

Wang et al. investigated urinary samples of patients with chronic kidney disease (CKD) versus patients with normal renal function. Urinary PCR was positive in one of 5 patients with CKD versus 3 out of 48 patients with normal renal function. In this study, the clinical course of COVID-19 disease and characteristics of patients with COVID-19 in urine were not different compared with patients without COVID-19 in their urine[16].

Ling and colleagues reported 66 patients with COVID-19 from Shanghai, China. Urine samples of 4 patients (6.9%) was positive for COVID-19. In 3 patients, urinary samples were positive even after clearance of virus in oropharyngeal samples[17].

Han et al. reported the presence of COVID-19 virus in urine of a neonate born from an infected mother. The virus was discovered in samples form oropharynx, saliva, urine and feces. However, the urine viral load was relatively low. Nevertheless, urine viral load was above the diagnostic cut off for days 6 through 17 after illness onset (11 days). The urine viral load was still positive after clearance of virus from nasopharyngeal and plasma samples[23].

In another study on children, the urine sample of 1 out of nine infected children was positive for COVID-19 by real time RT-PCR. All children in this study suffered from asymptomatic or mild disease. The urine was positive in a 7 year-old girl who presented only with fever (38.7 ° C) without cough or respiratory symptoms [21].

In a landmark study, Sun and colleagues investigated the urine samples of a 72 year-old male with severe COVID-19 infection. Urinary sampling was performed on days 12, 30 and 42 after symptom onset. The viral load was above diagnostic threshold **only** on day 30. The authors inoculated Vero E6 cells with urine of patient on day 12 (with viral load **below** diagnostic

threshold). Interestingly cytopathic effects were observed after 3 days. Electron microscopy revealed the presence of virus in inoculated cells by demonstration of spherical-shaped particles with distinct surface projections, resembling spikes. The authors furthermore used serum sample from this patient (who had high IgM and IgG against SARS-CoV-2) and a healthy candidate and demonstrated staining of inoculated cells in immunofluorescent assay only with patient serum and not with control serum[24].

Yang et al. reported *urine sediment* positivity in a 44 year-old man who had initially recovered with negative throat swab test but then revealed throat, and salivary positive COVID-19 rRT-PCR results. The urine was negative for COVID-19 RT-PCR on day 52 after illness onset however, urinary sediment was positive for COVID-19 RT_PCR in the same day [25].

The severity of disease has been reported in some case reports and case series. The severity of disease was correlated with urine positivity (spearman correlation coefficient: 0.26, P<0.001). In most reports (7 reports including 13 patients) the presence of COVID-19 in urine was detected in patients with moderate to severe disease. However, in three reports the virus has been isolated from urine of patients with mild disease (one neonate, one 7 year-old girl and a 44 year-old man) [21,23,25].

DISCUSSION

This is the first report to provide a comprehensive overview of the available evidence from December 2019 to 30[th] April 2020 for detection of SARS-CoV-2 in urine samples. Thirty three studies are included with a total of 430 patients in whom results of urinary testing for COVID-19 is reported across 12 countries. Initially during the COVID-19 outbreak, evaluation and investigation of urinary samples were considered part of routine sampling as stated by the World

Health Organization interim guideline for laboratory testing in COVID-19[11]. Later publications pointed to the rarity of viral presence in urine or totally rejected the presence of COVID-19 in urine[26]. Then, quite recently several publications revealed the presence of COVID-19 in urine. Herein, we reported the detection of COVID-19 in urinary samples in 8 studies.

COVID-19 is a rapidly emerging pandemic which threatens the security and biosafety at the world level. According to the latest World Health Organization situation report number 107 released on 6[th] May, more than 3.5 million people have been infected worldwide leaving more than 240000 deaths [27].

Preparedness against such a vast pandemic requires efforts in diagnosis, epidemiology, prevention and treatment. Clinical, laboratory, imaging, and prognostic studies as well as disease outcome analysis builds up the core knowledge necessary for any new disease. Several questions in the context of COVID-19 have been addressed including spectrum of disease severity (asymptomatic to fatal)[28] and the bodily organs or secretions in which the virus in present.

ACE2 is the cellular receptor for SARS-CoV, and high structural similarity has been demonstrated between cellular binding sites for SARS-CoV and 2019-nCoV[29]. Virus usual initial binding site is through ACE2 receptor at ciliated bronchial epithelial cells, and then spread of infection happened to other organs. COVID-19 infection induces a cytokine storm in the body incorporating a cascade of immune responses. ACE2 expression is tissue-specific, and is mainly expressed in the cardiovascular, respiratory, renal and gastrointestinal systems. A 1% positivity of ACE2 receptor in cells has been suggested for organ involvement in COVID-19 by Zhou et al. Based on this assumption, heart, lung, esophagus, ileum, bladder, and kidneys has been suggested as potential

organs for COVID-19 invasion[26]. Also in another study, the expression of ACE2 in testis has been demonstrated[7]. Therefore, the possibility of viral shedding in urine and semen can be postulated[7].

Urinary samples were not routinely collected in earliest infected patients in the initial outbreak of COVID-19 in many countries due to delay in proper diagnosis of COVID-19 in patients referring with respiratory symptoms[30].

The presence of COVID-19 in urine samples was not related to clinical course of disease in 53 patients reported from Wuhan[16]. Nevertheless, in most studies patients with positive urine for COVID-19 were in severe clinical disease or needed oxygen supplement. On the other hand, no association has been reported between urinary symptoms and presence of virus in urine [2] while some investigators have reported a positive association between urine analysis and clinical severity of illness [28]. The presence of proteinuria and microscopic hematuria has been associated with greater clinical severity of COVID-19 disease[31]. Combination of previous findings can point toward kidney involvement and urinary alteration in COVID-19 as a result of cytokine storm [12] in comparison with direct invasion of kidney parenchyma by the virus.

Han et al. reported a relatively long duration of urinary viral shedding in a neonate. Interestingly, the virus was also detected in saliva from this patient and urinary shedding was observed for 11 days during infection in comparison with 1-2 days of viral shedding in adult patients suggesting a higher potential for infection transmission in case of neonatal infection. Also, Lu et al.[21] reported one infected urine sample in 9 children with asymptomatic to mild COVID-19 raising further concerns over infection transmission through urine of asymptomatic children.

In most reports that have reported the presence of virus in urinary samples, the quantity of virus in urinary samples have been low in comparison with rectal or pharyngeal specimens[2] or has been marginally above the diagnostic threshold of the PCR assay[22]. However, Sun et al. isolated COVID-19 from urine of a patient with viral load below the diagnostic threshold of rRT-PCR and showed cytopathic effect of isolated virus in cell culture. This observation challenges the results of several studies that have reported negative urinary rRT-PCR based on readings below or marginally above test cut off[22] and suggests the possibility that the real frequency of COVID-19 in urine could be higher than reports based on diagnostic cut off of conventional real time RT-PCR assays.

The timeline for positivity of urinary samples in not consistent between reports. While Xu and colleagues reported urine positivity on day 7 after onset of symptoms and clearance of virus in urinary sample on the 10[th] day[32], Ling et al. reported the persistence of COVID-19 in urinary samples of 3 patients after clearance of virus in their nasopharyngeal samples[17].

Collectively, COVID-19 was reported in 3.7 percent of urinary samples from 430 patients. The highest frequency of infected urinary samples (in studies with more than 3 patients) belongs to the report of Peng et al. and Lu et al. from Guangdong, and Guangzhou who reported one infected urinary sample within urinary samples of 9 patients (11%). This rate is greatly lower in comparison with urinary infection rates of up to 42% which were previously reported for SARS-CoV[33]. One of the possible reasons for the low detection rate of COVID-19 can be short duration of virus presence in urine. Kim et al. investigated urinary samples from two Korean patients. They evaluated urinary samples from day 3 through day 14 of the illness. The PCR for RdRp was marginally positive only on day 12 and for gene E was again marginally positive only on day 9.

Another cause can be low quantity of virus in urinary samples which makes its detection in real time PCR assays difficult. However, as indicated above Sun et al. suggested the pathogenicity of low urinary viral load in cell cultures.

This review investigated the presence of COVID-19 in urine of infected patients or evaluated its pathogenic effects in cell cultures. The clinical significance low urinary viral load of patients and the potential of urine to be a source of infection transmission remains to be further investigated.

The significance of the findings of the current systematic review relies on the reported low overall positive rate of COVID-19 infection in patients' urinary samples which has been confirmed in several studies. We think that medical intervention like urethral catheter insertion and urinary endoscopic operations should be performed with caution considering the possibility of urine infection and possibility of COVID-19 transmission through this route.

CONCLUSIONS

We performed a systematic review of literature on the presence of COVID-19 in urine of infected patients. The results of this review revealed an average positivity rate of 3.7% for COVID-19 in patients' urine samples. The quantity of virus in most reports were lower than rectal or oropharyngeal samples. Nevertheless, we emphasize the low rate of urinary infection in COVID-19 and propose regular cautions in dealing with urinary samples in patients with COVID-19 infection and when performing medical interventions including urethral catheter passage and endoscopy in patients with COVID-19.

DECLARATION OF INTERESTS

He authors report on conflict of interest.

FUNDING INFORMATION

No funding was available for the current meta-analysis.

AUTHORS' CONTRIBUTIONS

AHK concepted of the study, helped in data gathering, helped in data analysis, and drafted the article.

MF helped in data gathering and in revising the manuscript.

EA helped in data gathering and in manuscript revision.

MV helped in data gathering, drafting the manuscript, and revising manuscript.

All authors approved the final version of the manuscript.

REFERENCES

1.   Guan W-j, Ni Z-y, Hu Y, et al. Clinical Characteristics of Coronavirus Disease 2019 in China. *New England Journal of Medicine* 2020.
2.   Peng L, Liu J, Xu W, et al. 2019 Novel Coronavirus can be detected in urine, blood, anal swabs and oropharyngeal swabs samples. *medRxiv* 2020.
3.   Lu R, Zhao X, Li J, et al. Genomic characterisation and epidemiology of 2019 novel coronavirus: implications for virus origins and receptor binding. *Lancet (London, England)* 2020; **395**(10224): 565-74.
4.   Petrosillo N, Viceconte G, Ergonul O, Ippolito G, Petersen E. COVID-19, SARS and MERS: are they closely related? *Clinical microbiology and infection : the official publication of the European Society of Clinical Microbiology and Infectious Diseases* 2020.
5.   Hoffmann M, Kleine-Weber H, Schroeder S, et al. SARS-CoV-2 Cell Entry Depends on ACE2 and TMPRSS2 and Is Blocked by a Clinically Proven Protease Inhibitor. *Cell* 2020; **181**(2): 271-80.e8.

6.   Zou X, Chen K, Zou J, Han P, Hao J, Han Z. Single-cell RNA-seq data analysis on the receptor ACE2 expression reveals the potential risk of different human organs vulnerable to 2019-nCoV infection. *Frontiers of medicine* 2020.

7.   Song C, Wang Y, Li W, et al. Absence of 2019 Novel Coronavirus in Semen and Testes of COVID-19 Patients. *Biology of reproduction* 2020.

8.   Kashi AH. COVID-19 and Semen: An Unanswered Area of Research. *Urol J* 2020.

9.   Kashi AH. COVID-19, Urologists and Hospitals. *Urol J* 2020.

10.   Jin Y, Yang H, Ji W, et al. Virology, Epidemiology, Pathogenesis, and Control of COVID-19. *Viruses* 2020; **12**(4).

11.   Organization WH. Laboratory testing for coronavirus disease (COVID-19) in suspected human cases. World Health Organization; 19 March 2020.

12.   Naicker S, Yang C-W, Hwang S-J, Liu B-C, Chen J-H, Jha V. The novel coronavirus 2019 epidemic and kidneys. *Kidney International* 2020.

13.   Wölfel R, Corman VM, Guggemos W, et al. Virological assessment of hospitalized patients with COVID-2019. *Nature* 2020: 1-10.

14.   Wang W, Xu Y, Gao R, et al. Detection of SARS-CoV-2 in Different Types of Clinical Specimens. *Jama* 2020.

15.   Pan Y, Zhang D, Yang P, Poon LL, Wang Q. Viral load of SARS-CoV-2 in clinical samples. *The Lancet Infectious Diseases* 2020.

16.   Wang L, Li X, Chen H, et al. Coronavirus Disease 19 Infection Does Not Result in Acute Kidney Injury: An Analysis of 116 Hospitalized Patients from Wuhan, China. *American Journal of Nephrology* 2020.

17.   Ling Y, Xu S-B, Lin Y-X, et al. Persistence and clearance of viral RNA in 2019 novel coronavirus disease rehabilitation patients. *Chinese Medical Journal* 2020.

18.   National Heart LaBI. Quality Assessment Tool fo rCase Series Studies. accessed April 2020. https://www.nhlbi.nih.gov/health-topics/study-quality-assessment-tools2020).

19.   Sanchez-Meca J, Marin-Martinez F. Confidence intervals for the overall effect size in random-effects meta-analysis. *Psychological methods* 2008; **13**(1): 31-48.

20.   Nyaga VN, Arbyn M, Aerts M. Metaprop: a Stata command to perform meta-analysis of binomial data. *Archives of public health = Archives belges de sante publique* 2014; **72**(1): 39.

21.   Lu Y, Wen H, Rong D, Zhou Z, Liu H. Clinical characteristics and radiological features of children infected with the 2019 novel coronavirus. *Clinical Radiology* 2020.

22.   Kim JY, Ko J-H, Kim Y, et al. Viral load kinetics of SARS-CoV-2 infection in first two patients in Korea. *Journal of Korean Medical Science* 2019; **35**(7).

23.   Han MS, Seong MW, Heo EY, et al. Sequential analysis of viral load in a neonate and her mother infected with SARS-CoV-2. *Clinical infectious diseases : an official publication of the Infectious Diseases Society of America* 2020.

24.   Sun J, Zhu A, Li H, et al. Isolation of Infectious SARS-CoV-2 from Urine of a COVID-19 Patient. *Emerging microbes & infections* 2020: 1-8.

25.   Yang JR, Deng DT, Wu N, Yang B, Li HJ, Pan XB. Persistent viral RNA positivity during recovery period of a patient with SARS-CoV-2 infection. *Journal of Medical Virology* 2020.

26.   Puliatti S, Eissa A. COVID-19 and Urology: A Comprehensive Review of the Literature. 2020.

27.   Organization WH. Coronavirus disease 2019 (COVID-19) Situation Report – 107. World Health Organization; 6th May 2020.

28.   Liu R, Ma Q, Han H, et al. The value of urine biochemical parameters in the prediction of the severity of coronavirus disease 2019. *Clinical Chemistry and Laboratory Medicine (CCLM)* 2020; **1**(ahead-of-print).

29.   Li F, Li W, Farzan M, Harrison SC. Structure of SARS coronavirus spike receptor-binding domain complexed with receptor. *Science (New York, NY)* 2005; **309**(5742): 1864-8.
30.   Zheng S, Fan J, Yu F, et al. Viral load dynamics and disease severity in patients infected with SARS-CoV-2 in Zhejiang province, China, January-March 2020: retrospective cohort study. 2020; **369**: m1443.
31.   Zhang N, Zhang R, Yao H, et al. Severity Detection For the Coronavirus Disease 2019 (COVID-19) Patients Using a Machine Learning Model Based on the Blood and Urine Tests. 2020.
32.   Xu Y, Li X, Zhu B, et al. Characteristics of pediatric SARS-CoV-2 infection and potential evidence for persistent fecal viral shedding. *Nature Medicine* 2020: 1-4.
33.   Peiris JS, Chu CM, Cheng VC, et al. Clinical progression and viral load in a community outbreak of coronavirus-associated SARS pneumonia: a prospective study. *Lancet (London, England)* 2003; **361**(9371): 1767-72.
34.   Chan JF-W, Yuan S, Kok K-H, et al. A familial cluster of pneumonia associated with the 2019 novel coronavirus indicating person-to-person transmission: a study of a family cluster. *The Lancet* 2020; **395**(10223): 514-23.
35.   Thevarajan I, Nguyen TH, Koutsakos M, et al. Breadth of concomitant immune responses prior to patient recovery: a case report of non-severe COVID-19. *Nature Medicine* 2020: 1-3.
36.   Park JY, Han MS, Park KU, Kim JY, Choi EH. First Pediatric Case of Coronavirus Disease 2019 in Korea. *Journal of Korean Medical Science* 2020; **35**(11).
37.   Ngoc NM, That BTT, Hong NTT, et al. Duration of viral detection in throat and rectum of a patient with COVID-19. *medRxiv* 2020.
38.   Hill KJ, Russell CD, Clifford S, et al. The index case of SARS-CoV-2 in Scotland: a case report. *Journal of Infection* 2020.
39.   Caly L, Druce J, Roberts J, et al. Isolation and rapid sharing of the 2019 novel coronavirus (SAR-CoV-2) from the first diagnosis of COVID-19 in Australia.
40.   Cai J, Xu J, Lin D, et al. A Case Series of children with 2019 novel coronavirus infection: clinical and epidemiological features. *Clinical Infectious Diseases* 2020.
41.   Young BE, Ong SWX, Kalimuddin S, et al. Epidemiologic features and clinical course of patients infected with SARS-CoV-2 in Singapore. *Jama* 2020.
42.   To KK-W, Tsang OT-Y, Leung W-S, et al. Temporal profiles of viral load in posterior oropharyngeal saliva samples and serum antibody responses during infection by SARS-CoV-2: an observational cohort study. *The Lancet Infectious Diseases* 2020.
43.   Schwartz DA. An Analysis of 38 Pregnant Women with COVID-19, Their Newborn Infants, and Maternal-Fetal Transmission of SARS-CoV-2: Maternal Coronavirus Infections and Pregnancy Outcomes. *Archives of Pathology & Laboratory Medicine* 2020.
44.   Kam K-q, Yung CF, Cui L, et al. A well infant with coronavirus disease 2019 (COVID-19) with high viral load. *Clinical Infectious Diseases* 2020.
45.   Cui Y, Tian M, Huang D, et al. A 55-Day-Old Female Infant infected with COVID 19: presenting with pneumonia, liver injury, and heart damage. *The Journal of Infectious Diseases* 2020.
46.   Cheng S-C, Chang Y-C, Chiang Y-LF, et al. First case of Coronavirus Disease 2019 (COVID-19) pneumonia in Taiwan. *Journal of the Formosan Medical Association* 2020.
47.   Chan JF-W, Yip CC-Y, To KK-W, et al. Improved molecular diagnosis of COVID-19 by the novel, highly sensitive and specific COVID-19-RdRp/Hel real-time reverse transcription-polymerase chain reaction assay validated in vitro and with clinical specimens. *Journal of Clinical Microbiology* 2020.
48.   Chen Y, Chen L, Deng Q, et al. The Presence of SARS-CoV-2 RNA in Feces of COVID-19 Patients. *Journal of Medical Virology* 2020.
49.   Paoli D, Pallotti F, Colangelo S, et al. Study of SARS-CoV-2 in semen and urine samples of a volunteer with positive naso-pharyngeal swab. *Journal of endocrinological investigation* 2020.

50.   Clinical and virologic characteristics of the first 12 patients with coronavirus disease 2019 (COVID-19) in the United States. *Nat Med* 2020.

51.   Lescure FX, Bouadma L, Nguyen D, et al. Clinical and virological data of the first cases of COVID-19 in Europe: a case series. *The Lancet Infectious diseases* 2020.

52.   Yu F, Yan L, Wang N, et al. Quantitative Detection and Viral Load Analysis of SARS-CoV-2 in Infected Patients. *Clinical infectious diseases : an official publication of the Infectious Diseases Society of America* 2020.

53.   Lo IL, Lio CF, Cheong HH, et al. Evaluation of SARS-CoV-2 RNA shedding in clinical specimens and clinical characteristics of 10 patients with COVID-19 in Macau. *International journal of biological sciences* 2020; **16**(10): 1698-707.

Table 1. Characteristics of included studies.

No.	Location	Journal	Study type	Quality Assessment	Technique	Total study population	Mean age (range) or [Median, IQR]; years	Sex ratio or M/F	Frequency of severe illness	Total number of urine samples (patients) tested	Age of patient/patients with urine test (year)	Total number of positive urine samples (patients) †	Urine Sampling timing*
14	Hubei and Shandong provinces and Beijing, China	JAMA	Case series	Good	rRT-PCR	205	44 (5-67)	68% male	19%	72	NR	(0)	NR
15	NR	The Lancet	Case series	Good	N-gene-specific quantitative RT-PCR	82	NR	NR	NR	2	NR	(0)	3-15 days
34	Shenzhen, China	The Lancet	Case series	Good	In-house real-time RT-PCR assay	6	NR (36-66)	NR	NR	6	NR	(0)	NR
35	Melbourne, Australia	Nature Medicine	Case report	Good	real-time RT-PCR	1	47	female	0%	1	47	0 (0)	NR
36	Seoul, Korea	J Kor Med Sci	Case report	Good	Real-time RT-PCR	1	10	female	0%	1	10	0 (0)	Day 3 and day 8
37	Ho Chi Minh City, Vietnam	The Lancet	Case report	Good	real time RT-PCR	1	73	male	0%	1	73	0 (0)	Days 4 to 24
38	Scotland	J Inf	Case report	Good	real time RT-PCR	1	51	male	0%	1	51	0 (0)	NR
1	31 provinces in China	N Eng J Med	Case series	Good	real time RT-PCR	1099	[47, NR]	41.90% female	173/1099	4	NR	1 (1)	Day 9 *
39	Melbourne, Australia	Med J Aust	Case report	Good	Real time RT-PCR	1	57	Male	NR	6 (1)	57	0 (0)	Days 1-8
40	Anhui and Shandong, China	Clin Inf Dis	Case series	Good	Duplex one step real time RT-PCR	10	6 (0-11)	4/6	NR	(6)	(3-11)	0	Day 3

30	Hangzhou, China	BMJ	Case series	Good	qRT-PCR	96	55 [IQR: 44-64]	58/38	74/96	180 (96)	NR	1	Admission, day 10 *
41	Singapore	JAMA	Case series	Good	Real time RT-PCR	18	47 (31-73)	9/9	6/18	10	NR	0	0-14
16	Wuhan, China	Am J Nephrol	Case series	Good	Real time RT-PCR	53	54 (20-95)	58% male	46/116	(53)	NR	(4)	NR
42	Hong Kong	Lancet Inf Dis	Cohort	Good	RT-qPCR	23	62 (37-75)	57% male	50%	(18)	NR	0	NR
43	US	Arch Path Lab Med	Case series	Fair	rt-PCR	3	34, 34, 30	Female	NR	(2)	34, 30	(0)	NR
2	Guangdong, China	medRxiv	Case series	Fair	qRT-PCR	9	38.9 (27-62)	4/5	NR	(9)	31	(1)	Day 7*
17	Shanghai, China	Chin Med J	Case series	Good	Dual fluorescence PCR	66	44 [IQR: 34-62]	38/28	NR	(58)	NR	(4)	NR
22	Korea	J Kor Med Sci	Case series	Good	Real time RT-PCR	2	35, 55	1/1	0%	(2)	35	(1)	Day 9-12*
44	Singapore	Clin Inf Dis	Case report	Good	rRT-PCR	1	0.5	male	NR	2 (1)	0.5	0	Day 2, 9
45	Guizhou, China	J Inf Dis	Case report	Fair	Real time RT-PCR	1	NR	Female	0%	(1)	0.1	0	NR
46	Taiwan	J Formofosan Med Ass	Case report	Good	rRT-PCR	1	55	Female	NR	(1)	55	0	Day 25
47	Hong Kong	J Clin Microbiol	Case series	Fair	rRT-PCR	23	NR	NR	NR	33 (15)	NR	0	NR
48	Zhongnan Hosp., Wuhan, China	J Med Virol	Case series	Fair	RT-PCR	42	51 (42-62)	15/27	11/42	(10)	NR	0	NR
13	Munich, Germany	Nature	Case Series	Fair	rRT-PCR	9	NR		0%	27 (9)	NR	0	Days 2-4
24	Guangzhou, China	Emerg Microbes Infec	Caes Report	Good	rRT-PCR	1	72	Male	100%	3(1)	72	(1)	Day 30*

49	Italy	J Endcorinol Invest	Case Report	Good	rRT-PCR	1	31	Male	0%	1 (1)	31	0	Day 8
50	US	Nat Med	Case Series	Good	rRT-PCR	12	[53, 21-68]	8/4	1/12	(10)	NR	0	Days 4-32
23	Seoul, Korea	Clin Infect Dis	Case Report	Good	rRT-PCR	1	0.1	Female	0%	7 (1)	0.1	6 (1)	Days 6-17 *
51	France	Lancet Infec Dis	Case Series	Good	rRT-PCR	5	47 (30-80)	3/2	3/5	(4)	47	0	Days 2-13
52	Beijing, China	Clin Infect Dis	Case Series	Fair	droplet digital PCR and RT-PCR	76	[40, 32-63]	38/38	22%	14	NR	0	NR
53	Macau, China	Int J Biol Sci	Case Series	Good	qRT-PCR	10	[54, 27-64]	3/7	40%	49 (10)	NR	0	Days 2-18
25 ‡	Wuhan, China	J Med Virol	Case Report	Fair	RT-qPCR	1	44	Male	0%	(1)	44	(1)	Day 52‡
21	Guangzhou, China	Clin Radiol	Case Series	Fair	Real time RT-PCR	9	7.8 (0.2-15)	5/4	0%	(9)	7	(1)	NR

NR: not reported; y/o: year old; IQR: interquartile range; rRT-PCR: real time RT-PCR; qRT-PCr: quantitative RT-PCR

*In case urine sample is reported positive for COVID-19, the sampling time of patients with positive samples or the sampling time of positive samples from a patient is reported.

† Number in parenthesis reveals the number of patients with a positive urinary result for SARS-Cov-2, and number outside parenthesis indicates the number of urinary samples positive for SARS-CoV-2.

‡ The virus was positive in urinary sediment on day 52 in a patient who had recovered from COVID-19 in addition to viral positivity in throat and saliva.

FIGURE LEGENDS:

Figure 1. Flow diagram of included studies.

Figure 2. Forest plot of the frequency of urinary viral shedding in each study and the pooled estimate.

Figure 3. Funnel plot for evaluation of publication bias.

Figure 1. Flow diagram of included studies.

Identification

Screening

Eligibility

Included

Records identified through
database searching
(n =1089)

Additional records identified
through other sources
(n =23   )

Records screened
(n =1112)

Records excluded
(n = 938)

174 records,
28 duplicates

Full-text articles assessed
for eligibility
(n = 146)

Full-text articles excluded,
with reasons
(n = 113)
Not including original data
on urinary shedding of
COVID-19 (n=110)

Duplicate patient source
(n=3)

Studies included in
qualitative synthesis
(n = 33)

Studies included in
quantitative synthesis
(meta-analysis)
(n = 33)

Figure 2. Forest plot of the frequency of urinary viral shedding in each study and the pooled estimate.

Figure 3. Funnel plot for evaluation of publication bias.